University of Groningen

Can the circadian system of a diurnal and a nocturnal rodent entrain to ultraviolet light?

Hut, R.A.; Scheper, A.; Daan, S.

Published in:

Journal of comparative physiology a-Sensory neural and behavioral physiology

DOI:

[10.1007/s003590000124](https://doi.org/10.1007/s003590000124)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2000

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Hut, R. A., Scheper, A., & Daan, S. (2000). Can the circadian system of a diurnal and a nocturnal rodent entrain to ultraviolet light? *Journal of comparative physiology a-Sensory neural and behavioral physiology*, 186(7-8), 707-715. <https://doi.org/10.1007/s003590000124>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Supplemental Material

Fig. 1

The left panel shows the yellow colour of the eye lens of the European ground squirrel (diameter 5.0 mm) indicating the strong absorption at lower wavelengths, including UV light and extending into the blue band of the human visible spectrum. The right panel shows the colourless appearance of the hamster lens (diameter 4 mm) which does transmit UV light. The transmittance spectra for both types of lenses are plotted in Fig. 5 of the manuscript.

Fig. 2

Typical upright position of the European ground squirrel. The visual system of the European ground squirrel is well adapted for use during daytime and plays an important role in predator recognition (see discussion section of the manuscript).