University of Groningen

Highly enantioselective Cu-catalysed allylic substitutions with Grignard reagents

Lopez, F; van Zijl, AW; Minnaard, AJ; Feringa, BL

Published in:
Chemical Communications

DOI:
[10.1039/b513887f](https://doi.org/10.1039/b513887f)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2006

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Lopez, F., van Zijl, AW., Minnaard, AJ., & Feringa, BL. (2006). Highly enantioselective Cu-catalysed allylic substitutions with Grignard reagents. *Chemical Communications*, (4), 409-411.
<https://doi.org/10.1039/b513887f>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

ELECTRONIC SUPPLEMENTARY INFORMATION

Highly Enantioselective Cu-catalysed Allylic Substitutions with Grignard Reagents

Fernando López, Anthoni W. van Zijl, Adriaan J. Minnaard, Ben L. Feringa*

*Department of Organic Chemistry and Molecular Inorganic Chemistry, Stratingh Institute,
University of Groningen, Nijenborgh 4, 9747 AG, Groningen, The Netherlands*

General Procedures: ^1H NMR spectra were recorded at 300 or 400 MHz with CDCl_3 as solvent. ^{13}C NMR spectra were obtained at 75.4 or 100.59 MHz in CDCl_3 , (Varian VXR300 or AMX400 spectrometers). Carbon types were determined from APT ^{13}C NMR experiments. Chemical shifts were determined relative to the residual solvent peaks (CHCl_3 , $\delta = 7.26$ ppm for hydrogen atoms, $\delta = 77.0$ for carbon atoms). The following abbreviations are used to indicate signal multiplicity: s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; br, broad. Progress and conversion of the reaction was determined by GC-MS (GC, HP6890; MS HP5973) with an HP1 or HP5 columns (Agilent Technologies, Palo Alto, CA). Enantio- and regioselectivities were determined by capillary GC analysis (HP 6890, ChiralDEX G-TA column (30 m x 0.25 mm), CP-Chiralsil-Dex-CB (25 m x 0.25 mm)) using flame ionization detector (in comparison with racemic products). Optical rotations were measured in CHCl_3 on a *Schmidt + Haensdch* polarimeter (Polartronic MH8) with a 10 cm cell (*c* given in g/100 mL). Absolute configuration of the products was determined by comparison with compounds previously published. Thin-layer chromatography (TLC) was performed on commercial Kieselgel 60F₂₅₄ silica gel plates, and components were visualized with KMnO_4 reagent. Flash chromatography was performed on silica gel. Drying of solutions was performed with Na_2SO_4 . Concentrations were conducted with a rotary evaporator.

Ligands **1a-1b** were generously donated by Solvias.¹ Ligand Taniaphos (**2**) was prepared according to literature procedures.² CuCl , CuI , and CuBr SMe_2 were purchased from Aldrich or Acros, and used without further purification. CuTC refers to copper thiophene-2-carboxylate.³ Hoveyda-Grubbs 2nd generation catalyst,⁴ (*E*)-cinnamyl bromide (**3a**) and (*E*)-cinnamyl chloride (**3b**) were purchased from Aldrich. The substrates **3b**,⁵ **3d-3f**,⁶ **3g**,⁷ and **3h**⁸ were prepared according to literature procedures. Grignard reagents were purchased from Aldrich (EtMgBr , MeMgBr) or prepared from the corresponding alkyl bromides and magnesium turnings in Et_2O following standard procedures. Grignard reagents were titrated using $^t\text{BuOH}$ and catalytic amounts of 1,10-phenanthroline. $^t\text{BuOMe}$ was purchased as anhydrous grade, stored over 4 Å MS and used without further purification. Et_2O was distilled from Na/benzophenone. CH_2Cl_2 was distilled from CaH_2 . All reactions were conducted under argon atmosphere using standard Schlenk techniques.

Racemic products **4** and regioisomers **5** were obtained by reaction of the bromides **3** with the corresponding Grignard reagent (5.0 equiv) at $-25\text{ }^\circ\text{C}$ in CH_2Cl_2 in the presence of CuCN (100 mol %). In some cases, the racemic products were also obtained by using *racemic-2* ligand, following the general procedure described in the next page.

Spectroscopic and analytical data of products **4b**, **4e-h** were obtained from their mixtures with **5**. The products **4a-4l**, and **6** have been previously described (see appropriate references in the following pages).

¹ A. Togni, C. Breutel, A. Schnyder, F. Spindler, H. Landert, A. Tijani, *J. Am. Chem. Soc.*, 1994, **116**, 4062.

² T. Ireland, G. Grossheimann, C. Wieser-Jeunesse, P. Knochel, *Angew. Chem. Int. Ed.*, 1999, **38**, 3212.

³ G. D. Allred, L. S. Liebeskind, *J. Am. Chem. Soc.*, 1996, **118**, 2748.

⁴ S. B. Garber, J. S. Kingsbury, B. L. Gray, A. H. Hoveyda, *J. Am. Chem. Soc.*, 2000, **122**, 8168.

⁵ C. A. Luchaco-Cullis, H. Muzitani, K. E. Murphy, A. H. Hoveyda, *Angew. Chem. Int. Ed.* 2001, **40**, 1456.

⁶ A. W. van Zijl, L. A. Arnold, A. J. Minnaard, B. L. Feringa, *Adv. Synth. Catal.*, 2004, **346**, 413 and references therein.

⁷ M. R. Binns, R. K. Haynes, D. E. Lambert, P. A. Schober, S. G. Turner, *Aust. J. Chem.*, 1987, **40**, 281.

⁸ G. Kottirsch, G. Koch, R. Feifel, U. Neumann *J. Med. Chem.*, 2002, **45**, 2289.

General Procedure for the Enantioselective Cu-catalysed Allylic Alkylation with Grignard Reagents

In a Schlenk tube equipped with septum and stirring bar, CuBr SMe₂ (15.0 μmol, 3.08 mg) and ligand **2** (18 μmol, 12.4 mg) were dissolved in CH₂Cl₂ (3.0 mL) and stirred under argon at room temperature for 10 min. The mixture was cooled to – 75 °C and the corresponding Grignard reagent (solution in Et₂O, 1.73 mmol) was added dropwise. Allylic bromide **3** (1.50 mmol) was then added dropwise as a solution in CH₂Cl₂ at that temperature over 15 min *via* a syringe pump. Once the addition was complete the resulting mixture was further stirred at – 75 °C for 4-12h. The reaction was quenched by addition of MeOH (0.5 mL) and the mixture was allowed to reach rt. Then, aqueous NH₄Cl solution (1M, 2 mL) was added to the mixture. The organic phase was separated, and the resulting aqueous layer is extracted with Et₂O (0.5 mL, 3x). The combined organic phases were dried and concentrated to a yellow oil which was flash chromatographed (2 : 98 Et₂O/pentane) to yield the corresponding allylic substrates as a mixture of S_N2' (**4**) and S_N2 (**5**) regioisomers.

Note: GC analysis was carried out on a sample obtained after aqueous extraction with Et₂O, which has been passed through a short plug of silica gel to remove transition metal residues.

(+)-1-((S)-but-3-en-2-yl)benzene (4a):⁹ Reaction time: 4h. Purification by column chromatography (2 : 98 Et₂O/pentane) afforded a 97 : 3 mixture of **4a** and **5a** as a colorless oil. [91% yield, **4a**: 98% ee, [α]_D = +5.4 (*c* 1.2, CHCl₃), lit.^{9a} (81% ee) [α]_D = + 4.8 (neat), lit.^{9b} for (*R*)-**4a** (60% ee) [α]_D = –2.2 (*c* 0.7, CHCl₃)]. **4a**: ¹H-NMR δ 7.28-7.14 (m, 5H), 6.02-5.93 (m, 1H), 5.04-4.98 (m, 2H), 3.47-3.40 (m, 1H), 1.33 (d, *J* = 7.0 Hz, 3H); ¹³C-NMR δ 145.5 (C), 143.2 (CH), 128.3 (CH), 127.2 (CH), 126.0 (CH), 113.0 (CH₂), 43.1 (CH), 20.7 (CH₃). Enantioselectivity determined by chiral GC analysis, CP-Chiralsil-Dex-CB (25 m x 0.25 mm), initial temp. 75 °C, retention times (min): 15.6 (minor) and 15.8 (major); Retention time **5a**: 23.7 min.

(+)-1-((S)-pent-1-en-3-yl)benzene (4b):¹⁰ Reaction time: 4h. Purification by column chromatography (2 : 98 Et₂O/pentane) afforded a 81 : 19 mixture of **4b** and **5b** as a colorless oil.

⁹ (a) T. Hayashi, M. Konishi, M. Fukushima, K. Kanehira, T. Hioki, M. Kumada, *J. Org. Chem.*, 1983, **48**, 2195. (b) M. Kawatsura, Y. Uozumi, M. Ogasawara, T. Hayashi, *Tetrahedron*, 2000, **56**, 2247.

¹⁰ (a) A. W. van Zijl, L. A. Arnold, A. J. Minnaard, B. L. Feringa, *Adv. Synth. Catal.*, 2004, **346**, 413. (b) K. Tissot-Croset, D. Polet, A. Alexakis, *Angew. Chem., Int. Ed.*, 2004, **43**, 2426. (c) A. Alexakis, K. Tissot-Croset, *Org. Lett.*, 2002, **4**, 4147. (d) M. A. Kacprzyński, A. H. Hoveyda, *J. Am. Chem. Soc.*, 2004, **126**, 10676.

[92% yield, **4b**: 95% ee, $[\alpha]_D = +47$ (c 1.0, CHCl_3), lit.^{10b} (96% ee) $[\alpha]_D = +55$ (c 1.1, CHCl_3); lit.^{10d} for (**R**)-**4b** (95% ee) $[\alpha]_D = -51$ (c 0.5, CHCl_3)]. **4b**: $^1\text{H-NMR}$ δ 7.32-7.12 (m, 5H), 5.91 (m, 1H), 5.01-4.97 (m, 2H), 3.11 (q, $J = 7.4$ Hz, 1H), 1.74-1.64 (m, 2H), 0.83 (t, $J = 7.31$ Hz, 3H); $^{13}\text{C-NMR}$ δ 144.4, 142.2, 128.4, 127.6, 126.1, 114.0, 51.7, 28.3, 12.1.. LRMS (EI) m/z 146 (M^+ , 32), 128 (10), 117 (100), 91 (62). Enantioselectivity determined by chiral GC analysis, CP-Chiralsil-Dex-CB (25 m x 0.25 mm), initial temp. 75 °C for 30 min, then 5 °C/min to 100 °C (final temp), retention times (min): 26.0 (minor) and 26.2 (major); Retention time **5b**: 38.0 min.

(R)-3-methylhept-1-ene (4c):¹¹ Reaction carried out using 5.0 mol% $\text{CuBr}\cdot\text{SMe}_2$ and 6.0 mol% **2**. Reaction time: 12 h. Purification by column chromatography (2 : 98 Et_2O /pentane) afforded a 100 : 0 mixture of **4c** and **5c** as a colorless oil. [99% conversion,¹² 92% ee].¹³ **4c**: $^1\text{H-NMR}$ δ 5.67-5.58 (m, 1H), 4.83 (dd, $J = 10.4$ and 7.3 Hz, 2H), 2.05-2.01 (m, 1H), 1.28-1.19 (m, 6H), 0.91 (d, $J = 6.7$ Hz, 3H), 0.82 (t, $J = 7.0$ Hz, 3H); $^{13}\text{C-NMR}$ δ 144.9 (CH), 112.1 (CH_2), 37.7 (CH), 36.3 (CH_2), 29.4 (CH_2), 22.8 (CH_2), 20.1 (CH_3), 14.0 (CH_3). Enantioselectivity determined by chiral GC analysis, CP-Chiralsil-Dex-CB (25 m x 0.25 mm), initial temp. 55°C for 20 min, retention times (min): 5.4 (minor) and 5.5 (major).

(R)-3-ethylhept-1-ene (4d):¹⁴ Reaction carried out using 5.0 mol% $\text{CuBr}\cdot\text{SMe}_2$ and 6.0 mol% **2**. Reaction time: 12 h. Purification by column chromatography (2 : 98 Et_2O /pentane) afforded a 100 : 0 mixture of **4d** and **5d** as a colorless oil. [99% conversion,¹² 93% ee].¹⁵ **4d**: $^1\text{H-NMR}$ δ 5.51-5.42 (m, 1H), 4.92-4.86 (m, 2H), 1.80-1.77 (m, 1H), 1.37-1.15 (m, 8H), 0.85-0.78 (m, 6H); $^{13}\text{C-NMR}$ δ 143.4 (CH), 113.9 (CH_2), 45.8 (CH), 34.4 (CH_2), 29.4 (CH_2), 27.7 (CH_2), 22.8 (CH_2), 14.1 (CH_3), 11.6 (CH_3); LRMS (EI) m/z 126 (M^+ , 3), 97 (18), 84 (72), 69 (81), 55 (100). Enantioselectivity determined by chiral GC analysis, CP-Chiralsil-Dex-CB (25 m x 0.25 mm), initial temp. 55°C for 20 min, retention times (min): 9.8 (minor) and 9.9 (major).

(-)-1-((S)-pent-1-en-3-yl)naphthalene (4e):^{10a,16} Reaction time: 4h. Purification by column chromatography (2 : 98 Et_2O /pentane) afforded a 87 : 13 mixture of **4e** and **5e** as a colorless oil. [86% yield, 90% ee, $[\alpha]_D = -26$ (c 1.0, CHCl_3)]. **4e**: $^1\text{H-NMR}$ δ 8.12 (d, $J = 8.1$ Hz, 1H), 7.84 (d, $J = 7.9$ Hz, 1H), 7.70 (d, $J = 7.9$ Hz, 1H), 7.51-7.36 (m, 4H), 6.11-6.02 (m, 1H), 5.11-5.07 (m, 2H), 4.01 (q, $J = 7.1$ Hz, 1H), 1.95-1.88 (m, 2H), 0.95 (t, $J = 7.4$ Hz, 3H); $^{13}\text{C-NMR}$ δ 142.8 (CH), 141.5 (C), 135.3 (C), 132.9 (C), 130.0 (CH), 127.7 (CH), 126.7 (CH), 126.6 (CH), 126.4 (CH), 125.0 (CH), 124.5 (CH), 115.7 (CH_2), 47.1 (CH), 29.1 (CH_2), 13.5 (CH_3); LRMS (EI) m/z 196 (M^+ , 26),

¹¹ T. L. Underiner, S. D. Paisley, J. Schmitter, L. Lesheski, H. L. Goering, *J. Org. Chem.*, 1989, **54**, 2369.

¹² Conversion based on GC. The high volatility of the products **4c** and **4d** did not allow to completely remove the solvents after the chromatography, impeding the calculation of an accurate isolated yield.

¹³ With 1.0 mol% $\text{CuBr}\cdot\text{SMe}_2$ and 1.12 mol% **2** the product **4c** was obtained with 99% conversion, a regioselectivity of 100: 0, but a slightly lower enantioselectivity (84% vs. 92%ee).

¹⁴ A. Pelter, K. Smith, S. Elgendy, M. Rowlands, *Tetrahedron Lett.*, 1989, **30**, 5647.

¹⁵ With 1.0 % $\text{CuBr}\cdot\text{SMe}_2$ and 1.1 % **2**, **4d** was obtained with 99% conversion, 97 : 3 regioselectivity and 88% ee.

¹⁶ A. O. Larsen, W. Leu, C. N. Oberhuber, J. E. Campbell, A. H. Hoveyda, *J. Am. Chem. Soc.*, 2004, **126**, 11130.

152 (42), 139 (11), 115 (14). Enantioselectivity determined by chiral HPLC analysis, Chiralcel OD-H (99.75% heptane/*i*PrOH), 40°C, retention times (min): 20.4 (major) and 23.2 (minor).

1-Chloro-4-((*S*)-pent-1-en-3-yl)benzene (4f):^{10a} Reaction time: 4h. Purification by column chromatography (2 : 98 Et₂O/pentane) afforded a 82 : 18 mixture of **4f** and **5f** as a colorless oil. [80% yield, 96% ee] **4f**: ¹H-NMR δ 7.25 (d, *J* = 8.4 Hz, 2H), 7.09 (d, *J* = 8.4 Hz, 2H), 5.88 (m, 1H), 5.00 (m, 2H), 3.10 (dt, *J* = 7.7 and 7.3 Hz, 1H), 1.69 (m, 2H), 0.83 (t, *J* = 7.3 Hz, 3H). Enantioselectivity determined by chiral GC analysis, CP-Chiralsil-Dex-CB (25 m x 0.25 mm), initial temp. 50°C, then 10°C/min to 120 °C (final temp) for 30 min, retention times (min): 19.7 (minor) and 20.1 (major).

(+)-1-((*S*)-hept-1-en-3-yl)benzene (4g):^{10a,17} Reaction time: 4h. Purification by column chromatography (2 : 98 Et₂O/pentane) afforded a 87 : 13 mixture of **4g** and **5g** as a colorless oil. [92% yield, 94% ee, [α]_D = +47 (*c* 0.5, CHCl₃), lit.^{10a} (88% ee) [α]_D = + 44 (*c* 0.1, CHCl₃)]. **4g**: ¹H-NMR δ 7.21 (m, 5H), 5.95 (m, 1H), 5.02 (m, 2H), 3.32 (q, *J* = 7.5 Hz, 1H), 1.70 (q, *J* = 7.4 Hz, 2H), 1.39-1.02 (m, 4H), 0.87 (t, *J* = 7.0 Hz, 3H); ¹³C-NMR δ 144.7, 142.5, 128.4, 127.6, 126.0, 113.8, 49.9, 35.1, 29.7, 22.6, 14.0. Enantioselectivity determined by chiral GC analysis, CP-Chiralsil-Dex-CB (25 m x 0.25 mm), initial temp. 80 °C for 60 min, then 10 °C/min to 140 °C (final temp), retention times (min): 53.7 (minor) and 54.2 (major); retention time **5g**: 70.4 min.

(+)-1-((*S*)-hepta-1,6-dien-3-yl)benzene (4h):^{10b,c} Reaction time: 4h. Purification by column chromatography (2 : 98 Et₂O/pentane) afforded a 91 : 9 mixture of **4h** and **5h** as a colorless oil. [93% yield, 95% ee, [α]_D = +36 (*c* 1.5, CHCl₃), lit.^{10b} (92% ee) [α]_D = + 33 (*c* 1.0, CHCl₃)]. **4h**: ¹H-NMR δ 7.30-7.16 (m, 5H), 5.95 (m, 1H), 5.79 (m, 1H), 5.03-4.92 (m, 4H), 3.26 (q, *J* = 7.5 Hz, 1H), 2.10-1.94 (m, 2H), 1.81-1.75 (m, 2H); ¹³C-NMR δ 144.1 (C), 142.1 (CH), 138.4 (CH), 128.4 (CH), 127.6 (CH), 126.1 (CH), 114.6 (CH₂), 114.1 (CH₂), 49.1 (CH), 34.4 (CH₂), 31.5 (CH₂); LRMS (EI) *m/z* 172 (M⁺, 13), 159 (6), 130 (33), 117 (100), 115 (41), 91 (48); HRMS Calcd. for C₁₃H₁₆ 172.12520, found 172.12554. Enantioselectivity determined by chiral GC analysis, CP-Chiralsil-Dex-CB (25 m x 0.25 mm), initial temp. 90°C for 45 min, then 5 °C/min to 140 °C (final temp), retention times (min): 30.0 (minor) and 30.3 (major); retention time **5h**: 56.0 min.

(-)-1-((*S*)-but-3-en-2-yl)naphthalene (4i):^{9b,16,18} Reaction time: 4h. Purification by column chromatography (2 : 98 Et₂O/pentane) afforded a 100 : 0 mixture of **4i** and **5i** as a colorless oil. [87% yield, 96% ee, [α]_D = - 29.8 (*c* 1.1, CHCl₃); lit.¹⁶ [α]_D = - 29 (*c* 1.0, CHCl₃); lit.^{18a} [α]_D = - 37 (neat); lit.^{9b} for (*R*)-**4i** (90% ee) +16.3 (*c* 0.4, CHCl₃)]. **4i**: ¹H-NMR δ 8.15 (d, *J* = 8.3 Hz, 1H), 7.87 (d, *J* = 7.8 Hz, 1H), 7.74 (d, *J* = 8.0 Hz, 1H), 7.54-7.41 (m, 4H), 6.23-6.15 (m, 1H), 5.17-5.13

¹⁷ (a) A. Yanagisawa, N. Nomura, H. Yamamoto, *Tetrahedron* 1994, **50**, 6017. (b) J. J. Van Veldhuizen, J. E. Campbell, R. E. Giudici, A. H. Hoveyda, *J. Am. Chem. Soc.*, 2005, **127**, 6877.

¹⁸ (a) R. Menicagli, O. Piccolo, L. Lardicci, M. L. Wis, *Tetrahedron*, 1979, **35**, 1301. (b) Y. Obora, Y. Tsuji, *J. Org. Chem.* 1995, **60**, 4647.

(m, 2H), 4.36-4.29 (m, 1H), 1.54 (d, $J = 7.0$ Hz, 3H); ^{13}C -NMR δ 142.9 (CH), 141.4 (C), 134.0 (C), 131.4 (C), 128.9 (CH), 126.8 (CH), 125.7 (CH), 125.6 (CH), 125.3 (CH), 123.6 (CH), 123.5 (CH), 113.6 (CH₂), 37.8 (CH), 20.2 (CH₃); LRMS (EI) m/z 182 (M⁺, 50), 167 (100), 165 (31), 152 (24), 84 (27), 51 (11); HRMS Calcd. for C₁₄H₁₄ 182.10955, found 182.11070. Enantioselectivity determined by chiral GC analysis, CP-Chiralsil-Dex-CB (25 m x 0.25 mm), initial temp. 120°C for 60 min, retention times (min): 35.8 (minor) and 36.3 (major); retention time **5i**: 56.3 min.

(+)-1-((S)-but-3-en-2-yl)-4-chlorobenzene (4j):¹⁹ Reaction time: 4h. Purification by column chromatography (2 : 98 Et₂O/pentane) afforded a 99 : 1 mixture of **4j** and **5j** as a colorless oil. [95% yield, 97% ee, $[\alpha]_{\text{D}} = +12$ (c 1.6, CHCl₃)]. **4j**: ^1H -NMR δ 7.29 (d, $J = 8.5$ Hz, 2H), 7.07 (d, $J = 8.5$ Hz, 2H), 6.00-5.90 (m, 1H), 5.06-4.97 (m, 2H), 3.44-3.36 (m, 1H), 1.32 (d, $J = 7.0$ Hz, 3H); LRMS (EI) m/z 166 (M⁺, 47), 165 (9), 151 (46), 139 (10), 131 (68), 91 (100); HRMS Calcd. for C₁₀H₁₁Cl 166.05492, found 166.05570. Enantioselectivity determined by chiral GC analysis, CP-Chiralsil-Dex-CB (25 m x 0.25 mm), initial temp. 95°C for 45 min, then 2°C/min to 140 °C (final temp), retention times (min): 25.8 (minor) and 26.1 (major); retention time **5j**: 40.8 min.

(+)-Methyl 4-((S)-but-3-en-2-yl)benzoate (4k):²⁰ Reaction time: 4h. Purification by column chromatography (2 : 98 Et₂O/pentane) afforded a 98 : 2 mixture of **4k** and **5k** as a colorless oil. [94% yield, 97% ee, $[\alpha]_{\text{D}} = +12$ (c 0.9, CHCl₃)]. **4k**: ^1H -NMR δ 7.91 (d, $J = 8.30$ Hz, 2H), 7.22 (d, $J = 8.30$ Hz, 2H), 5.97-5.90 (m, 1H), 5.03-4.98 (m, 2H), 3.84 (s, 3H), 3.49-3.44 (m, 1H), 1.32 (d, $J = 7.0$ Hz, 3H); ^{13}C -NMR δ 167.0 (C), 150.9 (C), 142.2 (CH), 129.7 (CH), 128.0 (C), 127.2 (CH), 113.8 (CH₂), 51.9 (CH₃), 43.1 (CH), 20.5 (CH₃); LRMS (EI) m/z 190 (M⁺, 44), 159 (33), 131 (100), 115 (25), 91 (22), 59 (8); HRMS Calcd. for C₁₂H₁₄O₂ 190.09937, found 190.09969. Enantioselectivity determined by chiral GC analysis, CP-Chiralsil-Dex-CB (25 m x 0.25 mm), initial temp. 105 °C for 70 min, then 1°C/min to 140 °C (final temp), retention times (min): 71.7 (minor) and 72.1 (major); retention time **5k**: 93.8 min.

(-)-(((S)-2-methylbut-3-enyloxy)methyl)benzene (4l):²¹ Reaction time: 12 h. Purification by column chromatography (2 : 98 Et₂O/pentane) afforded a 100 : 0 mixture of **4l** and **5l** as a colorless oil. [93% yield, 92% ee, $[\alpha]_{\text{D}} = -6$ (c 1.1, CHCl₃); lit.^{21a} $[\alpha]_{\text{D}} = -3$ (c 1.0, CHCl₃)]; **4l**: ^1H -NMR δ 7.32 (d, $J = 4.5$ Hz, 4H), 7.29-7.21 (m, 1H), 5.83-5.74 (m, 1H), 5.07-4.99 (m, 2H), 4.50 (s, 2H), 3.36 (dd, $J = 9.1$ and 6.5 Hz, 1H), 3.28 (dd, $J = 9.1$ and 6.8 Hz, 1H), 2.52-2.44 (m, 1H), 1.03 (d, $J = 6.8$ Hz, 3H); ^{13}C -NMR δ 141.3 (CH), 138.6 (C), 128.3 (CH), 127.5 (CH), 127.4 (CH), 114.0 (CH₂), 75.0 (CH₂), 72.9 (CH₂), 37.8 (CH), 16.6 (CH₃); LRMS (EI) m/z 176 (M⁺, 12), 175 (5), 92 (11), 91 (100), 65 (7); HRMS Calcd. for C₁₂H₁₆O 176.12011, found 176.12262. Enantioselectivity

¹⁹ (a) G. Franciò, F. Faraone, W. Leitner, *J. Am. Chem. Soc.*, 2002, **124**, 736. (b) K. Tissot-Croset, A. Alexakis, *Tetrahedron Lett.*, 2004, **45**, 7375.

²⁰ P. Gomes, C. Gosmini, J. Périchon, *J. Org. Chem.*, 2003, **68**, 1142.

²¹ (a) H. Lebel, V. Paquet *J. Am. Chem. Soc.*, 2004, **126**, 320. (b) J. C. Grandguillot, F. Pouessac, *Tetrahedron*, 1991, **47**, 5133.

determined by chiral HPLC analysis, Chiralcel OD-H (99.75% heptane/*i*PrOH), 40°C, retention times (min): 10.9 (minor) and 11.2 (major).

(+)-1-(((*S*)-2-ethylbut-3-enyloxy)methyl)benzene (4m): Reaction time: 12 h. Purification by column chromatography (2 : 98 Et₂O/pentane) afforded a 98 : 2 mixture of **4m** and **5m** as a colorless oil. [97% yield, 94% ee, [α]_D = + 19 (*c* 1.1, CHCl₃)]. **4m**: ¹H-NMR δ 7.30-7.21 (m, 5H), 5.66-5.57 (m, 1H), 5.06-5.01 (m, 2H), 4.47 (s, 2H), 3.35 (d, *J* = 6.5 Hz, 2H), 2.22 (m, 1H), 1.56-1.47 (m, 1H), 1.27-1.20 (m, 1H), 0.83 (t, *J* = 7.5 Hz, 3H); ¹³C-NMR δ 140.0 (CH), 138.6 (C), 128.3 (CH), 127.5 (CH), 127.4 (CH), 115.6 (CH₂), 73.6 (CH₂), 73.0 (CH₂), 45.7 (CH), 24.0 (CH₂), 11.4 (CH₃); LRMS (EI) *m/z* 190 (M⁺, 11), 189 (11), 123 (21), 105 (100), 91 (79), 77 (30); HRMS Calcd. for C₁₃H₁₈O 190.13576, found 190.13505. Enantioselectivity determined by chiral HPLC analysis, Chiralcel OD-H (99.75% heptane/*i*PrOH), 40°C, retention times (min): 9.9 (minor) and 10.9 (major).

(-)-(*S,E*)-methyl 4-phenylpent-2-enoate (6):²² Following the general procedure, in a Schlenk tube equipped with septum and stirring bar, CuBr SM₂ (15.0 μ mol, 3.08 mg) and ligand **2** (18.0 μ mol, 12.4 mg) were dissolved in CH₂Cl₂ (3.0 mL) and stirred under argon at room temperature for 10 min. The mixture was cooled to – 75 °C and MeMgBr (3.0 M solution in Et₂O, 1.73 mmol, 0.575 ml) was added dropwise. Cinnamyl bromide **3a** (296 mg, 1.50 mmol) was then added dropwise over 15 min *via* a syringe pump at that temperature. Once the addition was complete the resulting mixture was further stirred at – 75 °C for 4h. The reaction was quenched by addition of MeOH (0.5 mL) and the mixture was allowed to reach rt. Then, aqueous NH₄Cl solution (1M, 2 mL) was added to the mixture. The organic phase was separated, and the resulting aqueous layer is extracted with Et₂O (0.5 mL, 3x). The combined organic phases were dried and concentrated to a yellow oil which was dissolved in CH₂Cl₂ (3 mL) in a dried Schlenk tube. Methyl acrylate (645 mg, 7.5 mmol) and Hoveyda-Grubbs 2nd generation catalyst (18 mg, 0.03 mmol) were sequentially added producing a light green solution which was stirred for 36 h at rt. The mixture was then concentrated *in vacuo* to a dark brown oil. Purification of this residue by silica gel chromatography (2 : 98 to 5/ 95 Et₂O/pentane) affords **6** as a colorless oil [66% yield, [α]_D = – 20 (*c* 0.3, CHCl₃)]; ¹H-NMR δ 7.29-7.25 (m, 2H), 7.21-7.14 (m, 3H), 7.07 (dd, *J* = 15.7 and 6.7 Hz, 1H), 5.77 (dd, *J* = 15.7 and 1.5 Hz, 1H), 3.67 (s, 3H), 3.61-3.54 (m, 1H), 1.38 (d, *J* = 7.1 Hz, 3H); ¹³C-NMR δ 167.1, 152.9, 143.2, 128.6, 127.3, 126.7, 119.6, 51.4, 42.0, 20.1; LRMS (EI) *m/z* 190 (M⁺, 40), 159 (18), 131 (100), 91 (22), 51 (13); HRMS Calcd. for C₁₂H₁₄O₂ 190.09937, found 190.09953.

Enantioselective Catalytic Conjugate Addition of EtMgBr to (*S*)-**6**.²³

²² (a) J. M. Concellón, E. Bardales, *Org. Lett.*, 2002, **4**, 189. (b) D. B. Barma, A. Kundu, A. Bandyopadhyay, A. Kundu, B. Sangras, A. Briot, C. Mioskowski, J. R. Falck, *Tetrahedron Lett.* 2004, **45**, 5917.

²³ F. López, S. R. Harutyunyan, A. Meetsma, A. J. Minnaard, B. L. Feringa, *Angew. Chem., Int. Ed.*, 2005, **44**, 2752.

(+)-(3*S*,4*S*)-methyl 3-ethyl-4-phenylpentanoate (7): In a Schlenk tube CuBr SMe_2 (8.0 μmol , 1.62 mg) and ligand (*R,S*)-**1b** (9.4 μmol , 5.60 mg) were dissolved in CH_2Cl_2 (1.5 mL) and stirred under argon at room temperature for 10 min. The mixture was cooled to -75°C and EtMgBr (3.0 M in Et_2O , 0.78 mmol) was added dropwise. After stirring for 5 min at that temperature a solution of **6** (30 mg, 0.16 mmol) in CH_2Cl_2 (0.25 mL) was added dropwise over 10 min. After stirring at -75°C for 22 h, MeOH (0.25 mL) and NH_4Cl (1M, 2 mL) were sequentially added, and the mixture was warmed to rt. After extraction with Et_2O (0.5 mL, 3x), the combined organic phases were dried and concentrated to a yellow oil which was flash chromatographed (2 : 99 Et_2O /pentane) to yield **7** as a colourless oil [81% yield, 98% de, $[\alpha]_{\text{D}} = +25$ (*c* 0.2, CHCl_3)]; $^1\text{H-NMR}$ δ 7.26-7.12 (m, 5H), 3.57 (s, 3H), 2.82-2.73 (m, 1H), 2.30-2.15 (m, 2H), 2.08-1.97 (m, 1H), 1.38-1.27 (m, 1H), 1.17 (d, $J = 7.1$ Hz, 3H), 1.14-1.06 (m, 1H), 0.81 (t, $J = 7.4$ Hz, 3H); $^{13}\text{C-NMR}$ δ 174.1 (C), 145.6 (C), 128.2 (CH), 127.7 (CH), 126.0 (CH), 51.4 (CH_3), 42.8 (CH), 41.4 (CH), 35.3 (CH_2), 24.4 (CH_2), 17.1 (CH_3), 11.1 (CH_3); LRMS (EI) m/z 220 (M^+ , 20), 189 (11), 146 (43), 105 (100) 57 (21); HRMS Calcd. for $\text{C}_{14}\text{H}_{20}\text{O}_2$ 220.14632, found 220.14572. Diastereoselectivity determined by Chiraldex G-TA column (30 m x 0.25 mm), 100°C , retention times (min): 59.6 (minor: *3R,4S* and *3S,4R*) and 62.4 (major, *3S,4S*). Alternatively, the diastereoselectivity could also be determined by $^1\text{H-NMR}$, by integration of the 3.5 ppm signals corresponding to the methyl ester group.

(+)-(3*R*,4*S*)-methyl 3-ethyl-4-phenylpentanoate (8): Same procedure as above but using (*S,R*)-**1b** instead of (*R,S*)-**1b**. [84% yield, 92% de, $[\alpha]_{\text{D}} = +6$ (*c* 0.6, CHCl_3)]; $^1\text{H-NMR}$ δ 7.24-7.20 (m, 2H), 7.14-7.11 (m, 3H), 3.53 (s, 3H), 2.71-2.67 (m, 1H), 2.16-2.01 (m, 3H), 1.44-1.29 (m, 2H), 1.19 (d, $J = 7.2$ Hz, 3H), 0.83 (t, $J = 7.4$ Hz, 3H); $^{13}\text{C-NMR}$ δ 174.0 (C), 145.6 (C), 128.2 (CH), 127.8 (CH), 126.1 (CH), 51.3 (CH_3), 42.5 (CH), 41.8 (CH), 36.3 (CH_2), 23.1 (CH_2), 18.2 (CH_3), 10.4 (CH_3); LRMS (EI) m/z 220 (M^+ , 18), 189 (12), 146 (58), 105 (100); HRMS Calcd. for $\text{C}_{14}\text{H}_{20}\text{O}_2$ 220.14632, found 220.14621. Diastereoselectivity determined by Chiraldex G-TA column (30 m x 0.25 mm), 100°C , retention times (min): 59.6 (major: *3R,4S*), 62.4 (minor: *3S,4S*) and 63.3 (minor: *3R,4R*). Alternatively, the de could also be determined by $^1\text{H-NMR}$, by integration of the 3.57 and 3.53 ppm signals corresponding to the methyl ester groups of **7** and **8**.

Note: The conjugate addition of EtMgBr to (*S*)-**6**, following the procedure described above, but using *racemic*-**1b** instead of (*R,S*)-**1b** led to a 84 : 16 mixture of **7** and **8**, as deduced by GC [Chiraldex G-TA column (30 m x 0.25 mm), 100°C , retention times (min): 59.6 (minor: *3R,4S* and *3S,4R*) and 62.4 (major, *3S,4S*)] and $^1\text{H-NMR}$ [by integration of the 3.57 and 3.53 ppm signals corresponding to the methyl ester groups of **7** and **8** respectively], indicating a strong preference for the formation of the 1,2-*anti* product **7**.

Influence of the copper source and Grignard halide in the enantioselective allylic alkylation of cinnamyl halides catalyzed by Taniaphos 2.