University of Groningen

Excitation energies, for a benchmark set of molecules obtained within time-dependent current-density functional theory using the Vignale-Kohn functional

van Faassen, M.; de Boeij, P. L.

Published in:
Journal of Chemical Physics

DOI:
[10.1063/1.1697372](https://doi.org/10.1063/1.1697372)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2004

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

van Faassen, M., & de Boeij, P. L. (2004). Excitation energies, for a benchmark set of molecules obtained within time-dependent current-density functional theory using the Vignale-Kohn functional. *Journal of Chemical Physics*, 120(18), 8353 - 8363. <https://doi.org/10.1063/1.1697372>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Publisher's Note: "Excitation energies for a benchmark set of molecules obtained within time-dependent current-density functional theory using the Vignale-Kohn functional" [J. Chem. Phys. 120, 8353 (2004)]

M. van Faassen and P. L. de Boeij

*Theoretical Chemistry, Materials Science Centre, Rijksuniversiteit Groningen, Nijenborgh 4,
9747 AG Groningen, The Netherlands*

[DOI: 10.1063/1.1762871]

This article was originally published with an error in a value within Table V on p. 8362. The second entry in the final column (LDA/VK) should have been 5.63 instead of 6.63. AIP apologizes for this error. All online versions of the article have been corrected.