University of Groningen

Limited concordance between teachers, parents and healthcare professionals on the presence of chronic diseases in ID-adolescents

Oeseburg, B.; Jansen, D. E. M. C.; Reijneveld, S. A.; Dijkstra, G. J.; Groothoff, J. W.

Published in:
Research in Developmental Disabilities

DOI:
[10.1016/j.ridd.2010.04.015](https://doi.org/10.1016/j.ridd.2010.04.015)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2010

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Oeseburg, B., Jansen, D. E. M. C., Reijneveld, S. A., Dijkstra, G. J., & Groothoff, J. W. (2010). Limited concordance between teachers, parents and healthcare professionals on the presence of chronic diseases in ID-adolescents. *Research in Developmental Disabilities*, 31(6), 1645-1651.
<https://doi.org/10.1016/j.ridd.2010.04.015>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Limited concordance between teachers, parents and healthcare professionals on the presence of chronic diseases in ID-adolescents

B. Oeseburg^{a,b,*}, D.E.M.C. Jansen^{a,c}, S.A. Reijneveld^a, G.J. Dijkstra^a, J.W. Groothoff^a

^a Department of Health Sciences and Graduate School for Health Research (SHARE), University Medical Center Groningen, University of Groningen, The Netherlands

^b Wenckebach Institute, School of Nursing and Health, University Medical Center Groningen, Groningen, The Netherlands

^c Department of Sociology, Faculty of Behavioral and Social Sciences, University of Groningen, The Netherlands

ARTICLE INFO

Article history:

Received 16 April 2010

Accepted 19 April 2010

Keywords:

Mental retardation

Chronic diseases

Adolescents

Inter-rater agreement

ABSTRACT

Evidence on teachers' knowledge about somatic and mental chronic diseases among ID-adolescent compared to the knowledge parents and healthcare professionals have, is limited. The aim of this study is: (1) to assess the knowledge of teachers on the presence of chronic diseases in ID-adolescents; (2) to compare teachers with parents and healthcare professionals and parents with healthcare professionals regarding the knowledge on the presence of chronic diseases in ID-adolescents. We obtained data on 1044 ID-adolescents attending secondary schools, fully covering one region of the Netherlands. Teachers, parents and general practitioners (GPs) of the adolescents completed a questionnaire about the occurrence of chronic diseases in their child during the previous 12 months. The questionnaire was derived from the Dutch National Permanent Survey on Living Conditions questionnaire periodically administered in a representative population sample ($n \approx 10,000$). Concordance between teachers, parents and healthcare professionals on the presence of chronic diseases in ID-adolescents was relatively low. In about half of all 66 dyads the concordance was for the most part fair and just in 10 dyads good to very good; nine of these latter cases concerned somatic chronic diseases. In addition, teachers reported mostly lower prevalence rates of chronic diseases in ID-adolescents compared to the parents, in particular on mental chronic diseases. Although prevalence rates of chronic diseases among ID-adolescents are very high, knowledge on this among teachers is limited. While information on chronic diseases in ID-adolescents is available among different informants, the disagreement between them reflects different points of view between the informants and probably indicates a lack of communication. The communication among teachers, parents and GPs should be improved to combine the knowledge and information on the presence of chronic diseases in ID-adolescents. This may provide opportunities to improve the support of these adolescents in their school career and in their transition from school to work.

© 2010 Elsevier Ltd. All rights reserved.

1. Introduction

Chronic diseases are more prevalent among adolescents with intellectual disability (ID-adolescents) than among adolescents without ID (Emerson & Hatton, 2007; Emerson, 2003; Magnusson & Saemundsen, 2001; Oeseburg, Jansen,

* Corresponding author at: University Medical Center Groningen (UMCG), Department of Health Sciences and Graduate School for Health Research (SHARE), P.O. Box 30.001, 9700 RB Groningen, The Netherlands. Tel.: +31 050 363 8914; fax: +31 050 363 62.

E-mail address: b.oeseburg@med.umcg.nl (B. Oeseburg).

Dijkstra, Groothoff, & Reijneveld, 2010; Petterson, Bourke, Leonard, Jacoby, & Bower, 2007; Voigt, Barbaresi, Colligan, Weaver, & Katusic, 2006). Several studies have indicated that chronic diseases in *ID-adolescents*, in particular mental chronic diseases, largely increase the likelihood of emotional and behavioral problems (Bradley, Summers, Wood, & Bryson, 2004; Brereton, Tonge, & Einfeld, 2006; Buelow et al., 2003; Cormack, Brown, & Hastings, 2000; Hill & Furniss, 2006; Lewis et al., 2000; Oeseburg, Jansen, Groothoff, Dijkstra, & Reijneveld, 2010). Both, emotional and behavioral problems and disease specific problems, such as pain, fatigue or deficits in attention or hyperactivity, have a profound effect on participation of ID-adolescents in educational programs, occupational opportunities, and also on the potential to live in the community (Einfeld et al., 2006; Emerson & Hatton, 2007; Kanne, Abbacchi, & Constantino, 2009; Oeseburg, Jansen, Dijkstra, et al., 2010; Oeseburg, Jansen, Groothoff, et al., 2010; Reijneveld et al., 2003; Turk, Graham, & Verhulst, 2007).

Adequate school support highly contributes to the educational achievements of adolescents and their subsequent transition to employment (Clay, Cortina, Harper, Cocco, & Drotar, 2004; Nabors, Little, Akin-Little, & Iobst, 2008; Patel, Flisher, Hetrick, & McGorry, 2007; Patton & Viner, 2007; Sawyer, Drew, Yeo, & Britto, 2007; Taggart & McMullan, 2007; Turk et al., 2007). Teachers play a pivotal role in the challenging process to prepare adolescent successfully for the workforce. In case of ID-adolescents, teachers are more challenged since most ID-adolescents have chronic diseases which affect educational success (Clay et al., 2004; Mukherjee, Lightfoot, & Sloper, 2000; Nabors et al., 2008; Taggart & McMullan, 2007). Therefore, information on chronic diseases among ID-adolescents is highly needed by teachers, since it may help teachers to manage the effects of chronic diseases on the adolescents' emotional and behavioral functioning and their disabilities.

The limited evidence as available shows that teachers' knowledge about chronic diseases of their ID-adolescent students is much less than needed. This may be due to the fact that parents and healthcare professional do not share this information with teachers (Bishop & Boag, 2006; Clay et al., 2004; Nabors et al., 2008; Taggart & McMullan, 2007). Diagnostic overshadowing among teachers may be the result, i.e. teachers attribute the adolescents' problem behavior and disabilities to their ID rather than to other, potentially manageable causes related to their chronic disease(s) (Jopp & Keys, 2001). This misattribution may lead to the maintenance or aggravation of problem behavior because effective interventions to prevent symptoms of the disease that elicit problem behavior will not be used.

The aim of this study is therefore: (1) to assess the knowledge of teachers on the presence of chronic diseases in ID-adolescents; (2) to compare teachers with parents and healthcare professionals and parents with healthcare professionals regarding this knowledge on the presence of chronic diseases in ID-adolescents.

2. Methods

2.1. Participants

2.1.1. Sample

We collected data in 2006–2007 on adolescents with a borderline, mild, moderate or severe ID aged 12–18 years living in two provinces in the north of the Netherlands, Groningen and Drenthe (total population about 1.1 million people). Nearly all adolescents of the target population attended secondary schools (schools for practical training) or special secondary schools (regional expertise centers). ID-adolescents attending schools for practical training can be classified as mainly educable and have IQs between 60 and 80. ID-adolescents attending regional expertise centers can be classified as mainly trainable and have IQs between 30 and 60 (Dekker, Koot, van der Ende, & Verhulst, 2002). ID-adolescents not attending secondary schools, most of them with profound ID, were not included. The target population had been officially classified as having ID by an independent committee established by The Dutch Ministry of Education, Culture and Science. The classification of ID is based on information from validated intelligence tests (Dutch Eurydice Unit, 2007).

In the current school-based cross-sectional study, 88% of the schools for practical training and regional expertise centers in both provinces participated. Non-participating schools did not differ from participating schools regarding urbanization of the catchment area and number of students. All parents of the 2156 adolescents aged 12–18 years received a questionnaire and a reminder when they did not respond. One thousand forty four parents (48%) returned the questionnaire. Adolescents in the response and non-response group did not differ regarding age (t -test = 1.751, ns), but the response group had a higher proportion of girls ($\chi^2 = 5.9$; $p < 0.05$) and a higher proportion of adolescents with borderline or mild ID ($\chi^2 = 9.8$; $p < 0.05$). However, the effect sizes for both variables were trivial; Cohen's w were 0.06 and 0.07, respectively (Cohen, 1988).

2.1.2. Subsample with data from parents, teachers and general practitioners

Additionally, parents were asked for informed consent to derive information on chronic diseases of their child from their teacher and healthcare professionals, in this case general practitioners (GPs). GPs are in the Netherlands the best setting to provide valid information on common and lifelong health problems (Westert et al., 2005). Eight hundred ninety one (85%) parents gave informed consent. Teachers and GPs returned the questionnaires of 767 (86%) and 724 (81%) adolescents, respectively. Data on the presence of chronic diseases in ID-adolescents from all three informants were available for 539 adolescents. These 539 adolescents did not differ regarding gender ($\chi^2 = 0.4$; $p = 0.54$) and IQ ($\chi^2 = 3.4$; $p = 0.06$), but they were older (t -test = 2.2, $p < 0.05$) compared to the non-response group within this sub sample. However, the effect size was trivial; Cohen's d was 0.12 (Cohen, 1988).

The study protocol was approved by the Medical Ethics Committee of the University Medical Center Groningen, the Netherlands.

Table 1
Demographic characteristics of the sample.

Age (N = 539)	Years
Mean	15.7
SD	1.7
Range	12–18
Gender (N = 538)	N (%)
Boys	319 (59.3)
Girls	219 (40.7)
Level of ID ^a (N = 539)	N (%)
IQ 60–80	416 (77.2)
IQ 30–59	123 (22.8)

^a As measured by school type.

2.2. Measures

2.2.1. Chronic diseases in ID-adolescents

Chronic diseases in ID-adolescents were measured by the National Permanent Survey on Living Conditions questionnaire (POLS); module health and labor, part chronic diseases in children (Statistics Netherlands, 2003). POLS was developed by Statistics Netherlands and is yearly used in a representative sample ($n \approx 10,000$) of the Dutch population (Otten & Winkels, 1998). POLS part chronic diseases in children covers the most prevalent chronic diseases such as: ear, eye, skin diseases, diseases of the nervous, musculoskeletal, blood and circulatory, respiratory, digestive, and endocrine, nutritional and metabolic systems and attention deficit hyperactivity disorder (ADHD). Questions were added about the presence of pervasive developmental disorders (PDD). Parents, and if informed consent was obtained from the parents, also teachers and GPs were asked to report the presence or absence of each specific chronic disease in the last 12 months for their child, pupil, or patient, respectively. They could also report the presence of chronic diseases that were not listed in the questionnaire.

2.2.2. Background characteristics

The questionnaire comprised questions on age, gender and school type of the adolescent. School type was used as proxy for severity of ID.

2.3. Analysis

First, we computed prevalence rates of the separate chronic diseases, based on teachers', parents' and GPs' report in 539 ID-adolescents. Second, we compared the prevalence rates teachers reported on the separate chronic diseases in ID-adolescents with that of the parents and GPs, respectively. In addition, the prevalence rates parents and GPs reported on the separate chronic diseases in ID-adolescents were compared. Differences were tested using chi-square tests and for all differences effect sizes according to Cohen, Cohen's h , were calculated (Cohen, 1988). Cohen (Cohen, 1988) defines an effect size of <0.20 as trivial; of ≥ 0.20 to <0.50 as small; of ≥ 0.50 to <0.80 as moderate; and of ≥ 0.80 as large effect. Finally, the concordance between the knowledge of the three informants on the presence of chronic diseases in ID-adolescents was measured using Cohen's kappa for three dyads: teacher–parent, teacher–GP and parent–GP. We used the guidelines for interpretation of kappa proposed by Landis and Koch (Landis & Koch, 1977). Landis and Koch (Landis & Koch, 1977) defined a kappa of <0.21 as poor agreement; of ≥ 0.21 to <0.41 as fair agreement; of ≥ 0.41 to <0.61 as moderate agreement; of ≥ 0.61 to <0.81 as good agreement; and of ≥ 0.81 to 1.00 as very good agreement.

3. Results

Table 1 shows the background characteristics of the adolescents. The gender ratio, 59.3% boys and 40.7% girls, was similar to the ratio boys and girls with ID in the Netherlands (Schrojenstein Lantman-de Valk, Heurn-Nijsten, & Wullink, 2002).

3.1. Differences between the informants

Table 2 shows that teachers reported statistically significant lower prevalence rates for eight chronic diseases compared to the adolescents' parent. The effect sizes were trivial for four chronic diseases (asthma, chronic bronchitis and COPD, chronic eczema, diseases of neck, shoulder and upper extremities, ADHD), and small for the four other ones (migraine or chronic headache, psoriasis, autistic disorder, dyslexia). Moreover, teachers reported statistically significant higher prevalence rates for two chronic diseases (migraine or chronic headache, dyslexia) compared to the GPs. Both effect sizes were small. Teachers reported statistically significant lower prevalence rates for two chronic diseases compared to the GPs. The effect size was small for asthma, chronic bronchitis, COPD and trivial for congenital malformations nervous system.

Table 2Prevalence rates teachers, parents and GPs reported on the separate chronic diseases in ID-adolescents ($N = 539$) and the differences between the three informants.

	Informants						Differences					
	Teachers		Parents		GPs		Teachers–parents		Teachers–GPs		Parents–GPs	
	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>	<i>p</i> -Value	<i>h</i> ^a	<i>p</i> -Value	<i>h</i>	<i>p</i> -Value	<i>h</i>
Somatic chronic diseases												
Asthma, chronic bronchitis, COPD	4.3	23	9.3	50	8.9	48	0.001^b	0.20	0.002	0.19	0.832	0.01
Cerebral palsy	0.6	3	0.7	4	0.6	3	0.704	0.02	1.000	0.00	0.704	0.02
Chronic eczema	1.7	9	4.8	26	2.4	13	0.003	0.18	0.389	0.05	0.033	0.13
Congenital malformations circulatory system	0.9	5	1.9	10	2.2	12	0.194	0.08	0.087	0.11	0.667	0.03
Congenital malformations nervous system	2.0	11	2.4	13	5.6	30	0.676	0.03	0.002	0.19	0.008	0.16
Congenital malformations eye	1.1	6	2.0	11	1.5	8	0.217	0.08	0.583	0.03	0.487	0.03
Congenital malformations ear	1.9	10	1.3	7	1.5	8	0.463	0.04	0.635	0.03	0.795	0.02
Chromosome abnormalities	2.8	15	3.5	19	3.2	17	0.486	0.15	0.720	0.02	0.735	0.02
Diabetes	1.1	6	0.4	2	0.2	1	0.156	0.09	0.058	0.13	0.563	0.04
Epilepsy	3.7	20	4.3	23	5.4	29	0.641	0.03	0.189	0.08	0.394	0.05
Gastrointestinal and liver diseases	1.7	9	1.3	7	0.7	4	0.614	0.03	0.163	0.04	0.363	0.06
Heart- and blood diseases	2.2	12	2.6	14	1.5	8	0.691	0.02	0.367	0.05	0.196	0.08
Migraine or chronic headache	7.1	38	14.7	79	1.3	7	0.000	0.25	0.000	0.31	0.000	0.56
Muscular diseases	0.6	3	1.1	6	0.2	1	0.315	0.06	0.316	0.06	0.058	0.13
Musculoskeletal diseases												
Diseases of the back	2.2	12	2.8	15	2.2	12	0.559	0.04	1.000	0.00	0.559	0.04
Inflammatory polyarthropathies	0.6	3	0.7	4	0.4	2	0.705	0.02	0.654	0.03	0.413	0.05
Diseases of neck, shoulder and upper extremities	2.2	12	5.9	32	1.3	7	0.002	0.19	0.247	0.07	0.000	0.26
Psoriasis ^c	0.0	0	1.1	6	0.2	1	0.014	0.21	0.317	0.09	0.058	0.13
Mental chronic diseases												
Attention deficit/hyperactivity disorder (ADHD)	16.7	90	22.8	123	16.5	89	0.012	0.15	0.935	0.01	0.009	0.16
Autistic disorder	3.5	19	10.8	58	2.2	12	0.000	0.28	0.202	0.08	0.000	0.37
Dyslexia	5.9	32	14.1	76	1.5	8	0.000	0.28	0.000	0.25	0.000	0.53
Pervasive developmental disorder not otherwise specified (PDD-NOS)	11.5	62	13.9	75	10.2	55	0.235	0.07	0.493	0.04	0.061	0.11

^a *h* = Cohen's *h*.^b Bold differences are statistically significant.

Table 3

Concordance between the knowledge of the three informants on the presence of chronic diseases in ID-adolescents measured by Cohen's kappa.

	Teachers–parents Cohen's kappa	Teachers–GPs Cohen's kappa	Parents–GPs Cohen's kappa
Somatic chronic diseases			
Asthma, chronic bronchitis, COPD	0.35	0.24	0.39
Cerebral palsy	0.57	0.33	0.28
Chronic eczema	0.27	0.26	0.29
Congenital malformations circulatory system	0.53	0.11	0.35
Congenital malformations nervous system	0.74^a	0.42	0.45
Congenital malformations eye	0.22	0.01	0.09
Congenital malformations ear	0.58	0.55	0.53
Chromosome abnormalities	0.82	0.87	0.83
Diabetes	0.25	0.28	0.67
Epilepsy	0.59	0.55	0.72
Gastrointestinal and liver diseases	0.24	0.30	0.17
Heart- and blood diseases	0.53	0.39	0.44
Migraine or chronic headache	0.24	0.11	0.10
Muscular diseases	0.44	0.50	0.28
Musculoskeletal diseases			
Diseases of the back	0.67	0.49	0.51
Inflammatory polyarthropathies	0.57	0.80	0.67
Diseases of neck, shoulder and upper extremities	0.33	0.10	0.09
Psoriasis	– ^b	–	–
Mental chronic diseases			
Attention deficit/hyperactivity disorder (ADHD)	0.59	0.59	0.57
Autistic disorder	0.30	0.44	0.20
Dyslexia	0.39	0.03	0.00
Pervasive developmental disorder not otherwise specified (PDD-NOS)	0.67	0.51	0.53

^a Bold kappas are good to very good.^b Value of kappa was not meaningful to report because teachers did not report any case of psoriasis in ID-adolescents and GPs just one case of psoriasis in ID-adolescents.

We also analyzed the dyad parents–GPs. GPs reported statistically significant higher prevalence rates for congenital malformations of the nervous system compared to the parents. However, the effect size was trivial. Parents reported statistically significant higher prevalence rates for six chronic diseases compared to the GPs. The effect sizes were trivial for chronic eczema and ADHD, small for diseases of neck, shoulder and upper extremities and autistic disorder, and moderate for migraine or chronic headache and dyslexia.

3.2. Concordance between the informants

Table 3 shows that in about half of all 66 dyads concordance was poor to fair. In 10 of all dyads the kappas ranged from good to very good. This occurred four times in the dyad teachers–parents, four times in the dyad parents–GPs and two times in the dyad teachers–GPs. In nine cases, it concerned somatic chronic diseases and in one case a mental chronic disease.

4. Discussion

This study shows that the concordance between teachers, parents and healthcare professionals on the presence of chronic diseases in ID-adolescents was relatively low. In about half of all 66 dyads the concordance was at most fair and just in 10 dyads good to very good.

4.1. Fit with other studies

To our knowledge, no previous studies have examined the knowledge of teachers on the presence of a wide range of chronic diseases in ID-adolescents, and compared this with the knowledge parents and GPs have. However, we can state that our findings are mostly in line with studies reporting on a lack of information among teachers on issues associated with chronic diseases. These studies showed that teachers have limited knowledge regarding chronic disease-related information and limited confidence on their ability to work with children and adolescents with chronic diseases (Bishop & Boag, 2006; Brook & Galili, 2001; Clay et al., 2004; Mukherjee et al., 2000; Nabors et al., 2008; Taggart & McMullan, 2007). Moreover, our results are in line with studies on inter-informant agreement. These studies also reported limited concordance between different informants. Disconcordance may be due to variations in situation of informants and due the nature of the problems: parents versus teachers, internalizing versus externalizing problems, visible versus not visible problems (Achenbach, McConaughy, & Howell, 1987; Meester-Delver et al., 2008; Reijneveld, de Meer, Wiefferink, & Crone, 2008).

Our study shows that teachers reported mostly lower prevalence rates of chronic diseases in ID-adolescents than their parents, in particular on mental chronic diseases. In addition, this study shows that teachers and GPs and parents and GPs also reported different prevalence rates of chronic diseases in ID-adolescent. Thus, when information on chronic diseases in ID-adolescents is available among different informants and the disagreement between them reflects different points of view between the informants and probably indicates a lack of communication between them.

4.2. Strengths and limitations

Important strengths of this study are that it examined the prevalence rates of a wide range of chronic diseases in ID-adolescents in a community-based sample representative for about 90% of the adolescents with ID. Another strength is the use of multi-informant information and the comparison of the similarly obtained data and the high response rates of teachers and GPs in our sub sample. A limitation is the relatively low response rate of the total sample (48%). This could have lead to selection bias, but non-response analyses revealed no major differences in response by age, gender and educational level.

4.3. Implications for practice

Teachers need to be fully aware of the presence of chronic diseases in ID-adolescents and the impact on their functioning in order to meet their needs and to support them successfully in the transition from school to work. They need professional advice to handle the problems of adolescents with ID and chronic diseases in their classroom (Bishop & Boag, 2006; Brook & Galili, 2001; Clay et al., 2004; Mukherjee et al., 2000; Nabors et al., 2008). Information from parents or healthcare professionals on the presence of chronic diseases in ID-adolescents could improve teachers' awareness of chronic diseases in ID-adolescents and the impact on their functioning, and could improve their confidence to support them adequately. However, arrangements must be found to improve the communication among teachers, parents and GPs to combine the knowledge and information on the presence of chronic diseases in ID-adolescents (Clay et al., 2004; Mukherjee et al., 2000; Nabors et al., 2008; Perrin, Lewkowicz, & Young, 2000; Wolraich et al., 2004). Parents are in this communication process crucial because they are in the position to inform both teacher and GP about the health condition of their child. Moreover, teachers and GPs need their informed consent to transfer information on the health condition of the adolescent.

4.4. Implications for research

Our study is the first to examine the knowledge of teachers, parents and GP's on the presence of a wide range of chronic diseases in ID-adolescents. Therefore, our findings need confirmation including an assessment of the validity of the knowledge of each informant. Moreover, future research should examine whether arrangements to improve the communication among teachers, parents and GPs will be effective (Clay et al., 2004). Our results show very high prevalence rates of chronic diseases among ID-adolescents, and thus a high need for informed and skilled teachers to support these ID-adolescents in their school career and in their transition from school to work.

References

- Achenbach, T. M., McConaughy, S. H., & Howell, C. T. (1987). Child/adolescent behavioral and emotional problems: Implications of cross-informant correlations for situational specificity. *Psychological Bulletin*, 101, 213–232.
- Bishop, M., & Boag, E. M. (2006). Teachers' knowledge about epilepsy and attitudes toward students with epilepsy: Results of a national survey. *Epilepsy and Behavior*, 8, 397–405.
- Bradley, E., Summers, J., Wood, H., & Bryson, S. (2004). Comparing Rates of Psychiatric and Behavior Disorders in Adolescents and Young Adults with Severe Intellectual Disability with and without Autism. *Journal of Autism and Developmental Disorders*, 34, 151–161.
- Brereton, A. V., Tonge, B. J., & Einfeld, S. L. (2006). Psychopathology in children and adolescents with autism compared to young people with intellectual disability. *Journal of Autism and Developmental Disorders*, 36, 863–870.
- Brook, U., & Galili, A. (2001). Knowledge and attitudes of high school teachers towards pupils suffering from chronic diseases. *Patient Education and Counseling*, 43, 37–42.
- Buelow, J. M., Austin, J. K., Perkins, S. M., Shen, J., Dunn, D. W., & Fastenau, P. S. (2003). Behavior and mental health problems in children with epilepsy and low IQ. *Developmental Medicine and Child Neurology*, 45, 683–692.
- Clay, D. L., Cortina, S., Harper, D. C., Cocco, M., & Drotar, D. (2004). Schoolteachers' experiences with childhood chronic illness. *Children's Health Care*, 33, 227–239.
- Cohen, J. (1988). *Statistical power analysis for the behavioural sciences* (2nd ed.). New York: Academic Press.
- Cormack, K. F., Brown, A. C., & Hastings, R. P. (2000). Behavioural and emotional difficulties in students attending schools for children and adolescents with severe intellectual disability. *Journal of Intellectual Disability Research*, 44(Pt 2), 124–129.
- Dekker, M. C., Koot, H. M., van der Ende, E. J., & Verhulst, F. C. (2002). Emotional and behavioral problems in children and adolescents with and without intellectual disability. *Journal of Child Psychology and Psychiatry*, 43, 1087–1098.
- Dutch Eurydice Unit. (2007). *The Education System in the Netherlands 2007*. The Hague: Ministry of Education, Culture and Science.
- Einfeld, S. L., Piccinini, A. M., Mackinnon, A., Hofer, S. M., Taffe, J., Gray, K. M., et al. (2006). Psychopathology in young people with intellectual disability. *JAMA*, 296, 1981–1989.
- Emerson, E. (2003). Prevalence of psychiatric disorders in children and adolescents with and without intellectual disability. *Journal of Intellectual Disability Research*, 47, 51–58.
- Emerson, E., & Hutton, C. (2007). Mental health of children and adolescents with intellectual disabilities in Britain. *British Journal of Psychiatry*, 191, 493–499.
- Hill, J., & Furniss, F. (2006). Patterns of emotional and behavioural disturbance associated with autistic traits in young people with severe intellectual disabilities and challenging behaviours. *Research in Developmental Disabilities*, 27, 517–528.
- Jopp, D. A., & Keys, C. B. (2001). Diagnostic overshadowing reviewed and reconsidered. *American Journal of Mental Retardation*, 106, 416–433.

- Kanne, S. M., Abbacchi, A. M., & Constantino, J. N. (2009). Multi-informant ratings of psychiatric symptom severity in children with autism spectrum disorders: The importance of environmental context. *Journal of Autism and Developmental Disorders*, 39, 856–864.
- Landis, J. R., & Koch, G. G. (1977). The measurement of observer agreement for categorical data. *Biometrics*, 33, 159–174.
- Lewis, J. N., Tonge, B. J., Mowat, D. R., Einfeld, S. L., Siddons, H. M., & Rees, V. W. (2000). Epilepsy and associated psychopathology in young people with intellectual disability. *Journal of the Paediatrics and Child Health*, 36, 172–175.
- Magnusson, P., & Saemundsen, E. (2001). Prevalence of autism in Iceland. *Journal of Autism and Developmental Disorders*, 31, 153–163.
- Meester-Delver, A., Beelen, A., Folmer, K., Medema, D., Hadders-Algra, M., & Nollet, F. (2008). How well do care providers know the children with developmental disabilities they care for? *Acta Paediatrica*, 97, 608–612.
- Mukherjee, S., Lightfoot, J., & Sloper, P. (2000). The inclusion of pupils with a chronic health condition in mainstream schools: What does it mean for teachers? *Educational Research*, 42, 59–72.
- Nabors, L. A., Little, S. G., Akin-Little, A., & Iobst, E. A. (2008). Teacher knowledge of and confidence in meeting the needs of children with chronic medical conditions: Pediatric psychology's contribution to education. *Psychology in the Schools*, 45, 217–226.
- Oeseburg, B., Jansen, D. E., Dijkstra, G. J., Groothoff, J. W., & Reijneveld, S. A. (2010). Prevalence of chronic diseases in adolescents with intellectual disability. *Research in Developmental Disabilities*, 31, 698–704.
- Oeseburg, B., Jansen, D. E., Groothoff, J. W., Dijkstra, G. J., & Reijneveld, S. A. (2010). Emotional and behavioural problems in adolescents with intellectual disability with and without chronic diseases. *Journal of Intellectual Disability Research*, 54, 81–89.
- Otten, F., & Winkels, J. (1998). Explanation of the permanent research on living conditions (in Dutch). *Maandbericht Gezondheidsstatistiek*, 11–15.
- Patel, V., Flisher, A. J., Hetrick, S., & McGorry, P. (2007). Mental health of young people: A global public-health challenge. *Lancet*, 369, 1302–1313.
- Patton, G. C., & Viner, R. (2007). Pubertal transitions in health. *Lancet*, 369, 1130–1139.
- Perrin, E. C., Lewkowicz, C., & Young, M. H. (2000). Shared vision: Concordance among fathers, mothers, and pediatricians about unmet needs of children with chronic health conditions. *Pediatrics*, 105, 277–285.
- Peterson, B., Bourke, J., Leonard, H., Jacoby, P., & Bower, C. (2007). Co-occurrence of birth defects and intellectual disability. *Paediatric and Perinatal Epidemiology*, 21, 65–75.
- Reijneveld, S. A., de Meer, G., Wiefferink, C. H., & Crone, M. R. (2008). Parents' concerns about children are highly prevalent but often not confirmed by child doctors and nurses. *BMC Public Health*, 8, 124.
- Reijneveld, S. A., Vogels, A. G., Brugman, E., van Ede, J., Verhulst, F. C., & Verloove-Vanhorick, S. P. (2003). Early detection of psychosocial problems in adolescents: How useful is the Dutch short indicative questionnaire (KIVPA)? *The European Journal of Public Health*, 13, 152–159.
- Sawyer, S. M., Drew, S., Yeo, M. S., & Britto, M. T. (2007). Adolescents with a chronic condition: Challenges living, challenges treating. *Lancet*, 369, 1481–1489.
- Schrotenstein Lantman-de Valk, H. M. J. v., Heurn-Nijsten, E. W. A. v., & Wullink, M. (2002). *The prevalence of intellectual disability in the Netherlands (in Dutch)*. Maastricht: Universiteit Maastricht, capaciteitsgroep huisartsgeneeskunde.
- Statistics Netherlands. (2003). *Permanent Survey on Living Conditions (POLS); Health 2004 (in Dutch)*. Heerlen: Statistics Netherlands.
- Taggart, L., & McMullan, P. (2007). An exploratory study of teachers' knowledge about the symptoms of depression in young people with and without intellectual disabilities. *Journal of Intellectual Disability Research*, 11, 183–195.
- Turk, J., Graham, P., & Verhulst, F. (2007). *Child and Adolescent Psychiatry a Developmental Approach* (4th ed.). Oxford: Oxford University Press.
- Voigt, R. G., Barbaresi, W. J., Colligan, R. C., Weaver, A. L., & Katusic, S. K. (2006). Developmental dissociation, deviance, and delay: Occurrence of attention-deficit-hyperactivity disorder in individuals with and without borderline-to-mild intellectual disability. *Developmental Medicine and Child Neurology*, 48, 831–835.
- Westert, G. P., Schellevis, F. G., De Bakker, D. H., Groenewegen, P. P., Bensing, J. M., & van der Zee, J. (2005). Monitoring health inequalities through general practice: The Second Dutch National Survey of General Practice. *The European Journal of Public Health*, 15, 59–65.
- Wolraich, M. L., Lambert, E. W., Bickman, L., Simmons, T., Doffing, M. A., & Worley, K. A. (2004). Assessing the impact of parent and teacher agreement on diagnosing attention-deficit hyperactivity disorder. *Journal of Developmental & Behavioral Pediatrics*, 25, 41–47.