University of Groningen

Prevalence of chronic diseases in adolescents with intellectual disability

Oeseburg, B.; Jansen, D. E. M. C.; Dijkstra, G. J.; Groothoff, J. W.; Reijneveld, S. A.

Published in:
Research in Developmental Disabilities

DOI:
[10.1016/j.ridd.2010.01.011](https://doi.org/10.1016/j.ridd.2010.01.011)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2010

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Oeseburg, B., Jansen, D. E. M. C., Dijkstra, G. J., Groothoff, J. W., & Reijneveld, S. A. (2010). Prevalence of chronic diseases in adolescents with intellectual disability. *Research in Developmental Disabilities*, 31(3), 698-704. <https://doi.org/10.1016/j.ridd.2010.01.011>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Contents lists available at ScienceDirect

Research in Developmental Disabilities

Prevalence of chronic diseases in adolescents with intellectual disability

B. Oeseburg^{a,b,*}, D.E.M.C. Jansen^{a,c}, G.J. Dijkstra^a, J.W. Groothoff^a, S.A. Reijneveld^a

^a Department of Health Sciences and Graduate School for Health Research (SHARE), University Medical Center Groningen, University of Groningen, The Netherlands

^b Wenckebach Institute, School of Nursing and Health, University Medical Center Groningen, Groningen, The Netherlands

^c Department of Sociology, Faculty of Behavioral and Social Sciences, University of Groningen, The Netherlands

ARTICLE INFO

Article history:

Received 1 January 2010

Accepted 26 January 2010

Keywords:

Mental retardation

Chronic diseases

Adolescents

Epidemiology

ABSTRACT

Valid community-based data on the prevalence of chronic diseases in adolescents (12–18 years) with intellectual disability (ID-adolescents) are scarce. The aim of this study was to assess the prevalence rates and the nature of chronic diseases in a population of ID-adolescents and to compare them with the rates among adolescents in the general population. Therefore, we obtained data on 1083 ID-adolescents attending secondary schools, day care centers or living in residential centers fully covering one region of the Netherlands. Parents of the adolescents completed a questionnaire about the occurrence of chronic diseases in their child during the previous 12 months and about background characteristics. The questionnaire was derived from the Dutch National Permanent Survey on Living Conditions questionnaire periodically administered in a representative population sample ($n \approx 10,000$). Prevalence rates of chronic diseases in ID-adolescents were compared with those in adolescents in the Dutch general population. Among ID-adolescents, high prevalence rates of a wide range of chronic diseases were found. The five most prevalent were: ADHD (21.1%), PDD-NOS (14.0%), dyslexia (13.9%), migraine or chronic headache (12.7%), and autistic disorder (10.9%). These prevalence rates were all higher ($p < 0.05$) than among adolescents in the general population. Of all ID-adolescents, 62.9% was reported to have at least one chronic disease. The burden of chronic diseases among ID-adolescents is very high, showing a high need for adequate care. These high prevalence rates should alert policymakers and clinicians regarding the widespread of chronic diseases among ID-adolescents.

© 2010 Elsevier Ltd. All rights reserved.

1. Introduction

Valid community-based data on the prevalence rates of the full range of chronic diseases in adolescents (12–18 years) with intellectual disability (ID-adolescents) are scarce (Sawyer, Drew, Yeo, & Britto, 2007; Van Schrojenstein Lantman-de Valk & Walsh, 2008). A number of studies have reported on the prevalence of specific chronic diseases in young people with ID (Airaksinen et al., 2000; Bradley & Bolton, 2006; Bradley & Isaacs, 2006; Bryson, Bradley, Thompson, & Wainwright, 2008; Cans et al., 1999; Christianson et al., 2002; de Bildt, Sytema, Kraijer, & Minderaa, 2005; Dekker & Koot, 2003; Emerson & Hatton, 2007; Emerson, 2003; Fernell, 1998; Gothelf et al., 2008; Hou, Wang, & Chuang, 1998; Jelliffe-Pawlowski, Shaw, Nelson, & Harris, 2003; Koskentausta, Iivanainen, & Almqvist, 2002; Lewis et al., 2000; Magnusson & Saemundsen, 2001; Merrick & Morag, 2000; Molteno, Molteno, Finchilescu, & Dawes, 2001; Morgan, Baxter, & Kerr, 2003; Nielsen, Skov, & Jensen,

* Corresponding author at: University Medical Center Groningen (UMCG), Department of Health Sciences and Graduate School for Health Research (SHARE), P.O. Box 30.001, 9700 RB Groningen, The Netherlands. Tel.: +31 0 50 363 8914; fax: +31 0 50 363 6251.

E-mail address: b.oeseburg@med.umcg.nl (B. Oeseburg).

2007a; Nielsen, Skov, & Jensen, 2007b; Nordin & Gillberg, 1996; Petterson, Bourke, Leonard, Jacoby, & Bower, 2007; Steffenburg, Gillberg, & Steffenburg, 1996; Stromme & Diseth, 2000; Stromme & Hagberg, 2000; Van Schrojenstein Lantman-de Valk, van den Akker, Maaskant, & Haveman, 1997; Voigt, Barbaresi, Colligan, Weaver, & Katusic, 2006; Yeargin-Allsopp, Murphy, Cordero, Decoufle, & Hollowell, 1997; Zhang & Ji, 2005). However, only few studies have reported such prevalence rates of a wide range of chronic diseases associated with ID (Fernell, 1998; Stromme & Diseth, 2000; Stromme & Hagberg, 2000; Van Schrojenstein Lantman-de Valk et al., 1997). Moreover, only some studies compared their results with the prevalence rates of chronic diseases in the general population (Emerson & Hatton, 2007; Emerson, 2003; Magnusson & Saemundsen, 2001; Petterson et al., 2007; Voigt et al., 2006), and none of these studies reported prevalence rates of chronic diseases among ID-adolescents.

Adolescence is a specific stage of life between child- and adulthood with specific health needs. It is a time of life marked by physical, emotional, behavioural and social changes, but also by relatively high risks for the onset of (chronic) health problems (Patton & Viner, 2007; Sawyer et al., 2007; Turk, Graham, & Verhulst, 2007). Literature about young people with ID suggests that adolescents have a greater risk on chronic diseases compared to adolescents without ID (Cooper, Melville, & Morrison, 2004; Emerson & Hatton, 2007; Emerson, 2003; Jansen, Krol, Groothoff, & Post, 2004; Kolaitis, 2008; Magnusson & Saemundsen, 2001; Petterson et al., 2007; Van Schrojenstein Lantman-de Valk & Walsh, 2008; Voigt et al., 2006), but inclusive data on this are lacking.

Age-specific community-based data are thus needed to support policymakers and professionals in the adequate provision and planning of care to ID-adolescents. Policymakers need these data for the planning and financing of adequate care arrangements (e.g. health, education, work) to enhance the well being and societal participation of ID-adolescents and their families. Professionals need these data to know who are at risk for chronic diseases and to prevent chronic diseases, to support the early detection and adequate treatment of chronic diseases and their consequences among ID-adolescents and their families (American Association on Mental Retardation, 2002; Goddard, Davidson, Daly, & Mackey, 2008; McDermott, Durkin, Schupf, & Stein, 2007; Newacheck, Rising, & Kim, 2006).

The aim of this study is: (1) to assess the prevalence and the nature of chronic diseases in a population of ID-adolescents; (2) to compare the prevalence rates of chronic diseases in ID-adolescents with that among adolescents in the general population.

2. Methods

2.1. Participants

2.1.1. Adolescents with ID

We collected data in 2006–2007 from adolescents with a borderline, mild, moderate, severe or profound ID aged 12–18 years living in two provinces in the north of the Netherlands, Groningen and Drenthe (total population of about 1.1 million people). Adolescents of the target population attended secondary schools, special secondary schools, day care centers or lived in residential centers. ID-adolescents attending secondary schools can be classified as mainly educable and have IQs between 60 and 84. ID-adolescents attending special secondary schools can be classified as mainly trainable and have IQs between 30 and 59. ID-adolescents not attending school, most of them with IQs < 30, attend day care centers or live in residential centers (Dekker, Koot, van der, & Verhulst, 2002). The target population had been officially classified as having ID by two independent committees. The Dutch Ministry of Education, Culture and Science established a committee for adolescents attending secondary or special secondary schools and the Dutch Ministry of Health, Welfare and Sport established a committee for adolescents attending day care centers or those living in residential centers. The classification of ID is based on the information from validated intelligence tests obtained by both committees (Care Assessment Centre, 2009; Dutch Eurydice Unit, 2007).

In the current community-based cross-sectional research project, 70% of the schools and centers in both the provinces participated, with a total of 2271 adolescents. Non-participating schools and centers did not differ from participating schools and centers with regard to urbanization of the catchment area and number of adolescents. All parents of the 2271 adolescents aged 12–18 years received a questionnaire and a reminder when they did not respond; 1083 parents returned the questionnaire (48%). Adolescents in the response and non-response group did not differ with regard to age (t -test = 1.86, $p > 0.05$), but in the response group girls ($\chi^2 = 4.35$; $p < 0.05$) and adolescents with borderline or mild ID ($\chi^2 = 8.96$; $p < 0.05$) were somewhat overrepresented. However, the effect sizes for both the variables were negligible; Cohen's w were 0.05 and 0.07, respectively (Cohen, 1988).

The study protocol was approved by the Medical Ethics Committee of the University Medical Center Groningen, the Netherlands.

2.1.2. Adolescents in the general population

Statistics Netherlands conducts yearly the National Permanent Survey on Living Conditions (POLS) questionnaire in a representative sample ($n \approx 10,000$) of the Dutch population (Otten & Winkels, 1998). We used the data, adjusted for non-response, on adolescents aged 12–18 years in 2007 (Statistics Netherlands, 2009) (response: 64%). Prevalence data on chronic diseases in adolescents in the general population that were not available via Statistics Netherlands were derived from the National Institute for Public Health and the Environment (NIPHE). The NIPHE provides national and international

data to support policymakers and professionals in various fields of work such as prevalence rates of chronic diseases (National Institute for Public Health and the Environment, 2009).

2.2. Measures

2.2.1. Chronic diseases in ID-adolescents

Chronic diseases in ID-adolescents were measured by the National Permanent Survey on Living Conditions (POLS) questionnaire; module health and labor, part chronic diseases in children (Statistics Netherlands, 2003). POLS part chronic diseases in children covers the most prevalent chronic diseases such as ear, eye, skin diseases, diseases of the nervous, musculoskeletal, blood and circulatory, respiratory, digestive, and endocrine, nutritional and metabolic systems and attention deficit hyperactivity disorder (ADHD). Questions were added about the presence of pervasive developmental disorders (PDD). Parents were asked to fill in the presence or absence of each specific chronic disease in the last 12 months for their child. Parents were also offered the possibility to mention chronic diseases that were not listed in the questionnaire.

2.2.2. Chronic diseases in adolescents in the general population

The prevalence rates of chronic diseases in adolescents in the general population were measured in the same way as in our study, using the POLS questionnaire. Prevalence rates of chronic diseases in adolescents in the general population that were not available in POLS were obtained from the NIPHE. The NIPHE data are based on national and international research (National Institute for Public Health and the Environment, 2009).

2.2.3. Background characteristics of ID-adolescents

The questionnaire in the ID-adolescents sample comprised questions on age, gender and type of school or institution the adolescent is attending. We used type of school or institution as a proxy for severity of ID.

2.3. Analysis

First, we computed prevalence rates and the 95% confidence intervals (95%CI) of having one or more chronic diseases, overall and by type of chronic diseases. The association between the number of chronic diseases and ID severity and gender were assessed via univariate analysis of variance parameter estimates. Second, we performed multinomial logistic regression analyses. Odds ratios (OR) and their 95% confidence intervals (95%CI) were computed to assess the association between the occurrence of chronic diseases by type, and ID severity and gender. Third, we computed prevalence rates of separate chronic diseases in ID-adolescents, and compared these rates with rates among adolescents in the general population (12–18 years) (Statistics Netherlands, 2009). We tested differences using chi-square tests. If national data were not available from Statistics Netherlands, we used data from the NIPHE instead (National Institute for Public Health and the Environment, 2009). Differences were tested via single proportion tests. For all the differences in proportions, effect sizes according to Cohen were calculated (Cohen, 1988). Cohen (1988) defines an effect sizes of <0.20 as trivial effects; of ≥ 0.20 to <0.50 as small effects; of ≥ 0.50 to <0.80 as moderate effects; and of ≥ 0.80 as large effects.

3. Results

Table 1 shows the background characteristics of the adolescents. The gender ratio, 58.3% boys and 41.7% girls, was similar to the ratio of boys and girls with ID in the Netherlands (Van Schrojenstein Lantman-de Valk et al., 2002).

Table 2 shows that 63% of the ID-adolescents had at least one chronic disease and 34% had two or more chronic diseases (mean 1.30; SD 1.47). Thirty-nine per cent of the ID-adolescents had one or more somatic chronic diseases and 41% had one or more mental chronic diseases. Out of these, 20% had a combination of somatic and mental chronic diseases.

Table 1
Demographic characteristics of the sample.

Age in years (N = 1066)	Years
Mean (SD)	15.4 (1.6)
Range	12–18
Gender (N = 1074)	n (%)
Boys	626 (58.3)
Girls	448 (41.7)
Severity of ID ^a (N = 1077)	n (%)
IQ 60–80 (mild ID)	785 (72.9)
IQ 30–59 (moderate ID)	253 (23.5)
IQ < 30 (severe ID)	39 (3.6)

^a Derived from the type of institution at which they were allocated.

Table 2

Number and type of chronic disease(s) and 95% confidence intervals in ID-adolescents.

	ID-adolescents (N = 1083)		
	n	%	95%CI
Number of chronic diseases			
0	402	37.1	(34.29; 40.04)
1	313	28.9	(26.28; 31.67)
2	178	16.4	(14.35; 18.77)
3	97	9.0	(7.40; 10.81)
4	56	5.2	(4.00; 6.65)
5–11	37	3.4	(2.49; 4.68)
Type of chronic diseases			
None	402	37.1	(34.29; 40.04)
Somatic chronic disease(s)	235	21.7	(19.35; 24.25)
Mental chronic disease(s)	245	22.6	(20.23; 25.21)
Somatic & mental chronic diseases in combination	201	18.6	(16.36; 20.99)

Gender and severity of ID were associated with these prevalence rates. Gender was not associated with the number of chronic diseases, but boys were more likely than girls to be diagnosed with mental chronic diseases (OR 2.23; 95%CI 1.57–3.16) and with a combination of somatic and mental chronic diseases (OR 1.47; 95%CI 1.03–2.10). Severity of ID was positively associated with the number of chronic diseases ($F=48.12$, $p<0.000$). With regard to the type of the chronic diseases, adolescents with severe and moderate ID were more likely than adolescents with mild ID to be diagnosed with somatic chronic diseases (OR (95%CI) 12.35 (2.70–55.56), and 3.31 (2.04–5.00), respectively), mental chronic diseases (9.00 (1.91–41.67), and 1.88 (1.22–2.86), respectively) and a combination of somatic and mental chronic diseases (25.00 (5.62–111.11), and 3.85 (2.56–5.88), respectively). In addition, adolescents with severe ID were more likely than adolescents with moderate ID to be diagnosed with a combination of somatic and mental chronic diseases (6.44 (1.43–29.14)).

Table 3 shows that the prevalence rates of chronic diseases among ID-adolescents were statistically significant higher for 8 of the 17 chronic diseases compared to the prevalence rates of these chronic diseases in adolescents in the general population. The effect sizes were negligible for two (asthma, chronic bronchitis and COPD, and psoriasis) of the four somatic chronic diseases that differed with statistical significance, and small for the three other ones (epilepsy, heart diseases, and blood diseases). With regard to mental chronic diseases, the effect sizes for the statistically significant differences were small for dyslexia and medium for ADHD, PDD-NOS and autistic disorder.

4. Discussion

This study shows high prevalence rates of a wide range of chronic diseases in ID-adolescents. For 8 of the 17 chronic diseases that we assessed the prevalence rates were statistically significant higher among ID-adolescents than among adolescents in the general population. Differences were particularly large with regard to ADHD, autistic disorder, and PDD-NOS, and smaller for some somatic diseases. Moreover, differences in prevalence rates were larger if ID was more severe, with regard to any mental chronic diseases and with regard to a combination of any somatic and any mental chronic diseases.

4.1. Fit with other studies

Our findings regarding adolescents cannot be compared with previous ones on adolescents because our study was the first that examined the prevalence rates of a wide range of chronic diseases in ID-adolescents, and compared them with the prevalence rates among adolescents in the general population. Previous studies on youth with ID also found higher prevalence rates of a wide range of mental chronic diseases (Emerson & Hatton, 2007; Emerson, 2003; Magnusson & Saemundsen, 2001; Voigt et al., 2006) and a wide range of congenital malformations (Pettersson et al., 2007) though. However, these studies did not separately report on adolescents, thus failing to recognize adolescence as a specific developmental stage. It is likely that this explain the differences in the prevalence rates of mental chronic diseases and congenital malformations these studies found compared to our study.

Three factors have been proposed to explain the high rates of chronic diseases in the ID population in general. First factor is biological/genetic, i.e. genetic and chromosomal disorders that cause both ID and a wide range of chronic somatic and mental diseases (Dykens, 2000; Emerson & Hatton, 2007; Koskentausta et al., 2002). Second, the association of ID with socio-economic disadvantage may lead to adverse health outcomes and to higher rates of chronic diseases (Cooper et al., 2004; Emerson & Hatton, 2007; Emerson & Hatton, 2008; Walsh, 2008). Third, mental chronic diseases may be associated with ID because they share diagnostic characteristics (Dykens, 2000; Emerson & Hatton, 2007).

Table 3

Prevalence rates of chronic diseases in adolescents with ID and in the general population.

	ID-adolescents		General population		Differences	
	%	n/N	%	n/N	p-Value	Cohen's h^a
Somatic chronic diseases						
Asthma, chronic bronchitis, COPD ^b	9.9	107/1083	6.2	30/481	0.019	0.13
Chronic eczema ^b	4.3	47/1083	2.6	14/536	n.s. ^c	0.09
Diabetes ^b	0.5	5/1083	0.4	2/498	n.s.	0.01
Gastrointestinal and liver diseases ^b	1.8	19/1083	1.0	4/396	n.s.	0.06
Migraine or chronic headache ^b	12.7	138/1083	11.5	52/452	n.s.	0.04
Musculoskeletal diseases^b						
Diseases of the back ^b	2.6	28/1083	2.7	14/517	n.s.	0.01
Inflammatory polyarthropathies ^b	0.8	9/1083	0.4	2/442	n.s.	0.05
Diseases of neck, shoulder and upper extremities ^b	5.3	57/1083	3.3	14/430	n.s.	0.10
Psoriasis ^b	1.3	14/1083	0.1	1/519	0.030	0.14
Heart and blood diseases ^b	2.4	26/1083	0.1	1/524	0.001	0.22
Congenital malformations circulatory system ^d	2.1	23/1083	0.62 ^e	–	n.s.	0.13
Congenital malformations nervous system ^d	2.0	22/1083	0.21 ^e	–	n.s.	0.19
Epilepsy ^d	5.3	57/1083	0.27 ^f	–	0.000	0.36
95%CI						
Somatic chronic diseases no reference data available						
Congenital malformations eye	3.0		32/1083			(2.10; 4.14)
Congenital malformations ear	1.8		20/1083			(1.20; 2.84)
Chromosome abnormalities	3.5		38/1083			(2.57; 4.78)
Cerebral palsy	0.5		5/1083			(0.20; 1.07)
Muscular diseases	0.7		8/1083			(0.38; 1.45)
Other congenital malformations	6.1		66/1083			(4.81; 7.68)
Other somatic disorders	3.7		40/1083			(2.72; 4.99)
	%	n/N	%	n/N	p-Value	Cohen's h^a
Mental chronic diseases						
Dyslexia ^b	13.9	151/1083	3.8 ^g	87/2285	0.000	0.37
Attention deficit/hyperactivity disorder (ADHD) ^d	21.1	229/1083	1.3 ^h	–	0.000	0.73
Autistic disorder ^d	10.9	118/1083	0.1 ⁱ	–	0.000	0.61
Pervasive developmental disorder not otherwise specified (PDD-NOS) ^d	14.0	152/1083	0.2 ⁱ	–	0.000	0.68
95%CI						
Mental chronic diseases no reference data available						
Other psychiatric disorders	1.8		19/1083			(1.12; 2.72)

Bold results are statistically significant.

^a Cohen thresholds: negligible effect (≤ 0.20); small effect (≥ 0.20 and < 0.50); medium effect (≥ 0.50 and < 0.80); large effect (≥ 0.80).^b Reference data on adolescents aged 12–18 from Statistics Netherlands (POLS).^c Not significant.^d Reference data from the National Institute for Public Health and the Environment.^e Only data at birth available.^f Only data on the age-group 15–24 years available.^g Only data on the age-group 4–12 years available.^h Only data on the age-group 13–17 years available.ⁱ Prevalence data estimated on the basis of the review of Fombonne (2005) on autistic disorder and other pervasive developmental disorders.

4.2. Strengths and limitations

Important strengths of this study are that it examined the prevalence rates of a wide range of chronic diseases in ID-adolescents in a community-based sample representative for about 90% of the adolescents with ID, and compared most outcomes with similarly obtained data on adolescents without ID. A limitation may be that data collection procedures may have slightly differed between adolescents with ID and in the general population. POLS did not cover mental disorders other than ADHD, we used data of the NIPHE instead. However, the differences in prevalence rates were so large that they are unlikely to be fully explained by methodological differences. A second limitation is the relatively low response rate of our study (48%). This could lead to selection bias, but non-response analyses revealed no major differences with regard to age, gender and educational level.

4.3. Implications

We found disconcertingly high prevalence rates for some chronic diseases among ID-adolescents which should alert policymakers and clinicians to these diseases among ID-adolescents. This shows a need for effective care arrangements to handle this huge burden of morbidity, both with regard to its prevention and its treatment. As such, it provides a challenge to

both clinicians and policy (American Association on Mental Retardation, 2002; Goddard et al., 2008; McDermott et al., 2007; Newacheck et al., 2006).

Our study is the first to examine the prevalence rates of a wide range of chronic diseases in ID-adolescents and compare the results with data on adolescents without ID. Therefore, our findings need confirmation, including an assessment of the pathways leading to such high prevalence rates. Anyhow, our results show a very high burden of chronic diseases among ID-adolescents, and thus adequate care is highly needed (Cooper et al., 2004; Van Schrojenstein Lantman-de Valk & Walsh, 2008).

References

- Airaksinen, E. M., Matilainen, R., Mononen, T., Mustonen, K., Partanen, J., Jokela, V., et al. (2000). A population-based study on epilepsy in mentally retarded children. *Epilepsia*, 41, 1214–1220.
- American Association on Mental Retardation. (2002). *Mental Retardation: Definitions, classification and systems of supports* (10th ed.). Washington: American Association on Mental Retardation.
- Bradley, E., & Bolton, P. (2006). Episodic psychiatric disorders in teenagers with learning disabilities with and without autism. *British Journal of Psychiatry*, 189, 361–366.
- Bradley, E. A., & Isaacs, B. J. (2006). Inattention, hyperactivity, and impulsivity in teenagers with intellectual disabilities, with and without autism. *Canadian Journal of Psychiatry*, 51, 598–606.
- Bryson, S. E., Bradley, E. A., Thompson, A., & Wainwright, A. (2008). Prevalence of autism among adolescents with intellectual disabilities. *Canadian Journal of Psychiatry*, 53, 449–459.
- Cans, C., Wilhelm, L., Baille, M. F., du-Mazaubrun, C., Grandjean, H., & Rumeau-Rouquette, C. (1999). Aetiological findings and associated factors in children with severe mental retardation. *Developmental Medicine and Child Neurology*, 41, 233–239.
- Care Assessment Centre. (2009). *CAC indication guide version: 2.0*. Eindhoven: Roto Smeets Graf Services (in Dutch).
- Christianson, A. L., Zwane, M. E., Manga, P., Rosen, E., Venter, A., Downs, D., et al. (2002). Children with intellectual disability in rural South Africa: Prevalence and associated disability. *Journal of Intellectual Disability Research*, 46, 179–186.
- Cohen, J. (1988). *Statistical power analysis for the behavioural sciences* (2nd ed.). New York: Academic Press.
- Cooper, S. A., Melville, C., & Morrison, J. (2004). People with intellectual disabilities. *British Medical Journal*, 329, 414–415.
- de Bildt, A., Sytema, S., Kraijer, D., & Minderaa, R. (2005). Prevalence of pervasive developmental disorders in children and adolescents with mental retardation. *Journal of Child Psychology and Psychiatry*, 46, 275–286.
- Dekker, M. C., & Koot, H. M. (2003). DSM-IV disorders in children with borderline to moderate intellectual disability. I. Prevalence and impact. *Journal of the American Academy of Child and Adolescent Psychiatry*, 42, 915–922.
- Dekker, M. C., Koot, H. M., van der, E. J., & Verhulst, F. C. (2002). Emotional and behavioral problems in children and adolescents with and without intellectual disability. *Journal of Child Psychology and Psychiatry*, 43, 1087–1098.
- Dutch Eurydice Unit. (2007). *The Education System in the Netherlands 2007*. The Hague: Ministry of Education, Culture and Science.
- Dykens, E. M. (2000). Psychopathology in children with intellectual disability. *Journal of Child Psychology and Psychiatry*, 41, 407–417.
- Emerson, E. (2003). Prevalence of psychiatric disorders in children and adolescents with and without intellectual disability. *Journal of Intellectual Disability Research*, 47, 51–58.
- Emerson, E., & Hatton, C. (2007). Mental health of children and adolescents with intellectual disabilities in Britain. *British Journal of Psychiatry*, 191, 493–499.
- Emerson, E., & Hatton, C. (2008). Socioeconomic disadvantage, social participation and networks and the self-rated health of English men and women with mild and moderate intellectual disabilities: Cross sectional survey. *European Journal of Public Health*, 18, 31–37.
- Fernell, E. (1998). Aetiological factors and prevalence of severe mental retardation in children in a Swedish municipality: The possible role of consanguinity. *Developmental Medicine and Child Neurology*, 40, 608–611.
- Fombonne, E. (2005). Epidemiology of autistic disorder and other pervasive developmental disorders. *Journal of Clinical Psychiatry*, 10, 3–8.
- Goddard, L., Davidson, P. M., Daly, J., & Mackey, S. (2008). People with an intellectual disability in the discourse of chronic and complex conditions: An invisible group? *Australian Health Review*, 32, 405–414.
- Gothelf, D., Goral, O., Avni, S., Stawski, M., Hartmann, I., Basel-Vanagaite, L., et al. (2008). Psychiatric morbidity with focus on obsessive-compulsive disorder in an Israeli cohort of adolescents with mild to moderate mental retardation. *Journal of Neural Transmission*, 115, 929–936.
- Hou, J. W., Wang, T. R., & Chuang, S. M. (1998). An epidemiological and aetiological study of children with intellectual disability in Taiwan. *Journal of Intellectual Disability Research*, 42, 137–143.
- Jansen, D. E., Krol, B., Groothoff, J. W., & Post, D. (2004). People with intellectual disability and their health problems: A review of comparative studies. *Journal of Intellectual Disability Research*, 48, 93–102.
- Jelliffe-Pawlowski, L. L., Shaw, G. M., Nelson, V., & Harris, J. A. (2003). Risk of mental retardation among children born with birth defects. *Archives of Pediatrics and Adolescent Medicine*, 157, 545–550.
- Kolaitis, G. (2008). Young people with intellectual disabilities and mental health needs. *Current Opinion in Psychiatry*, 21, 469–473.
- Koskentausta, T., Iivanainen, M., & Almqvist, F. (2002). Psychiatric disorders in children with intellectual disability. *Nordic Journal of Psychiatry*, 56, 126–131.
- Lewis, J. N., Tonge, B. J., Mowat, D. R., Einfeld, S. L., Siddons, H. M., & Rees, V. W. (2000). Epilepsy and associated psychopathology in young people with intellectual disability. *Journal of the Paediatrics and Child Health*, 36, 172–175.
- Magnusson, P., & Saemundsen, E. (2001). Prevalence of autism in Iceland. *Journal of Autism and Developmental Disorders*, 31, 153–163.
- McDermott, S., Durkin, M. S., Schupf, N., & Stein, Z. A. (2007). Epidemiology and etiology of mental retardation. In J. W. Jacobsen, J. A. Mulick, & J. Rojahn (Eds.), *Handbook of intellectual and developmental disabilities* (pp. 3–40). New York: Springer.
- Merrick, J., & Morag, A. (2000). Human immunodeficiency virus (HIV) in institutions for the mentally retarded in Israel. *International Journal of Rehabilitation Research*, 23, 173–175.
- Molteno, G., Molteno, C. D., Finchilescu, G., & Dawes, A. R. (2001). Behavioural and emotional problems in children with intellectual disability attending special schools in Cape Town, South Africa. *Journal of Intellectual Disability Research*, 45, 515–520.
- Morgan, C., Baxter, H., & Kerr, M. (2003). Prevalence of Epilepsy and Associated Health Service Utilization and Mortality Among Patients With Intellectual Disability. *American Journal of Mental Retardation*, 108, 293–300.
- National Institute for Public Health the Environment. (2009). *National compass for public health and health care; health and diseases* http://www.rivm.nl/vtv/object_class/kom_ziekaandoen.html. Retrieved September 2009 (in Dutch).
- Newacheck, P. W., Rising, J. P., & Kim, S. E. (2006). Children at risk for special health care needs. *Pediatrics*, 118, 334–342.
- Nielsen, L. S., Skov, L., & Jensen, H. (2007). Visual dysfunctions and ocular disorders in children with developmental delay. I. Prevalence, diagnoses and aetiology of visual impairment. *Acta Ophthalmologica Scandinavica*, 85, 149–156.
- Nielsen, L. S., Skov, L., & Jensen, H. (2007). Visual dysfunctions and ocular disorders in children with developmental delay. II. Aspects of refractive errors, strabismus and contrast sensitivity. *Acta Ophthalmologica Scandinavica*, 85, 419–426.
- Nordin, V., & Gillberg, C. (1996). Autism spectrum disorders in children with physical or mental disability or both. I. Clinical and epidemiological aspects. *Developmental Medicine and Child Neurology*, 38, 297–313.
- Otten, F., & Winkels, J. (1998). Explanation of the permanent research on living conditions. *Maandbericht Gezondheidsstatistiek*, 11–15 (in Dutch).
- Patton, G. C., & Viner, R. (2007). Pubertal transitions in health. *Lancet*, 369, 1130–1139.

- Petterson, B., Bourke, J., Leonard, H., Jacoby, P., & Bower, C. (2007). Co-occurrence of birth defects and intellectual disability. *Paediatric and Perinatal Epidemiology*, 21, 65–75.
- Sawyer, S. M., Drew, S., Yeo, M. S., & Britto, M. T. (2007). Adolescents with a chronic condition: Challenges living, challenges treating. *Lancet*, 369, 1481–1489.
- Statistics Netherlands. (2003). *Permanent survey on living conditions (POLS); Health 2004*. Heerlen: Statistics Netherlands (in Dutch).
- Statistics Netherlands. (September 16, 2009). *Statline: Self-reported medical consumption, health and life style* <http://statline.cbs.nl/StatWeb/publication/?DM=SLNL&PA=03799&D1=94-147&D2=0-17&D3=0&D4=a&VW=T>, Retrieved September 2009 (in Dutch).
- Steffenburg, S., Gillberg, C., & Steffenburg, U. (1996). Psychiatric disorders in children and adolescents with mental retardation and active epilepsy. *Archives of Neurology*, 53, 904–912.
- Stromme, P., & Diseth, T. H. (2000). Prevalence of psychiatric diagnoses in children with mental retardation: Data from a population-based study. *Developmental Medicine and Child Neurology*, 42, 266–270.
- Stromme, P., & Hagberg, G. (2000). Aetiology in severe and mild mental retardation: A population-based study of Norwegian children. *Developmental Medicine and Child Neurology*, 42, 76–86.
- Turk, J., Graham, P., & Verhulst, F. (2007). *Child and Adolescent Psychiatry a Developmental Approach* (4th ed.). Oxford: Oxford University Press.
- Van Schrojenstein Lantman-de Valk, H. M. J., van den Akker, M., Maaskant, M. A., & Haveman, M. J. (1997). Prevalence and incidence of health problems in people with intellectual disability. *Journal of Intellectual Disability Research*, 41, 42–51.
- Van Schrojenstein Lantman-de Valk, H. M. J., van Heurn-Nijsten, E. W. A., & Wullink, M. (2002). *The prevalence of intellectual disability in the Netherlands*. Maastricht: Universiteit Maastricht, capaciteitsgroep huisartsgeneeskunde (in Dutch).
- Van Schrojenstein Lantman-de Valk, H. M. J., & Walsh, P. N. (2008). Managing health problems in people with intellectual disabilities. *British Medical Journal*, 337, 1408–1412.
- Voigt, R. G., Barbaresi, W. J., Colligan, R. C., Weaver, A. L., & Katusic, S. K. (2006). Developmental dissociation, deviance, and delay: Occurrence of attention-deficit-hyperactivity disorder in individuals with and without borderline-to-mild intellectual disability. *Developmental Medicine and Child Neurology*, 48, 831–835.
- Walsh, P. N. (2008). Health indicators and intellectual disability. *Current Opinion in Psychiatry*, 21, 474–478.
- Yeargin-Allsopp, M., Murphy, C. C., Cordero, J. F., Decoufle, P., & Hollowell, J. G. (1997). Reported biomedical causes and associated medical conditions for mental retardation among 10-year-old children, metropolitan Atlanta, 1985 to 1987. *Developmental Medicine and Child Neurology*, 39, 142–149.
- Zhang, X., & Ji, C. Y. (2005). Autism and mental retardation of young children in China. *Biomedical and Environmental Sciences*, 18, 334–340.