University of Groningen

Tuning energy transfer in switchable donor-acceptor systems

Hurenkamp, Johannes H.; de Jong, Jaap J. D.; Browne, Wesley R.; van Esch, Jan H.; Feringa, Ben L.

Published in:
Organic and Biomolecular Chemistry

DOI:
[10.1039/b719095f](https://doi.org/10.1039/b719095f)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2008

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Hurenkamp, J. H., de Jong, J. J. D., Browne, W. R., van Esch, J. H., & Feringa, B. L. (2008). Tuning energy transfer in switchable donor-acceptor systems. *Organic and Biomolecular Chemistry*, 6(7), 1268-1277.
<https://doi.org/10.1039/b719095f>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Electronic Supporting information for:

Controlling Energy Transfer in Switchable Donor-Acceptor Systems

Johannes H. Hurenkamp, Jaap J.D. de Jong, Wesley R. Browne, Jan H. van Esch and Ben L. Feringa

Photochromic switching of CSC 5 and PSC 10 over several cycles

Left: Absorption spectrum of CSC 5 before and after photochemical ring closure ($\lambda_{exc} > 312$ nm) and opening ($\lambda_{exc} > 400$ nm). Irradiation was carried out at 220 K in CH_2Cl_2 . Right: Absorption at $\lambda = 493$ nm plotted against number of times switched.

Left: Absorption spectra of PSC 10 before and after switching from PSC open to PSC PSS, irradiated with $\lambda = 312$ nm at 220K and $\lambda > 400$ nm light at RT in CH_2Cl_2 respectively. Right: Absorption at 493 nm plotted over three switching cycles of PSC using $\lambda = 312$ nm light at 220K to close and $\lambda > 400$ nm at RT to ring-open the dithienylcyclopentene unit. Measurements were performed in CH_2Cl_2 and spectra were recorded at RT.

Effect of irradiation of PSC **10** at $\lambda = 254$ nm

Left: The change in the absorption spectrum of PSC **10** upon irradiation over 20 min with $\lambda = 254$ nm light in CH_2Cl_2 at RT. Right: The change in emission spectra ($\lambda_{\text{ex}} = 322$ nm) of PSC **10** by irradiation over 18 min with $\lambda = 254$ nm light in CH_2Cl_2 at RT.