University of Groningen

The influence of solar ultraviolet radiation on the photochemical production of H₂O₂ in the equatorial Atlantic Ocean

Gerringa, LJA; Rijkenberg, MJA; Timmermans, KR; Buma, AGJ

Published in:
Journal of Sea Research

DOI:
[10.1016/j.seares.2003.03.002](https://doi.org/10.1016/j.seares.2003.03.002)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2004

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Gerringa, LJA., Rijkenberg, MJA., Timmermans, KR., & Buma, AGJ. (2004). The influence of solar ultraviolet radiation on the photochemical production of H₂O₂ in the equatorial Atlantic Ocean. *Journal of Sea Research*, 51(1), 3-10. <https://doi.org/10.1016/j.seares.2003.03.002>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

The influence of solar ultraviolet radiation on the photochemical production of H_2O_2 in the equatorial Atlantic Ocean

L.J.A. Gerringa^{a,*}, M.J.A. Rijkenberg^{a,b}, K.R. Timmermans^a, A.G.J. Buma^b

^aRoyal Netherlands Institute for Sea Research, PO Box 59, 1790 AB Den Burg, The Netherlands

^bCentre for Ecological and Evolutionary Sciences, University of Groningen, PO Box 14, 9750 AA Haren, The Netherlands

Received 28 May 2002; accepted 28 March 2003

Abstract

Hydrogen peroxide (H_2O_2) was measured in marine surface waters of the eastern Atlantic Ocean between 25°N and 25°S . H_2O_2 concentrations decreased from 80 nM in the north to 20 nM in the south, in agreement with earlier observations. A diel cycle of H_2O_2 production as a function of sunlight in surface waters was followed twice whilst the ship steamed southward. Around 23°N a distinct diel cycle could be measured which correlated well with irradiance conditions.

The wavelength dependency of H_2O_2 formation was studied near the equator. For 16 hours, water samples were incubated with wavelength bands of the solar spectrum, i.e. visible (VIS: 400–700 nm), VIS and ultraviolet A radiation (UVAR: 320–400 nm) and VIS, UVAR and ultraviolet B radiation (UVBR: 280–320 nm). A significant relationship was found between wavelength band and the production of H_2O_2 . In addition, a clear positive relationship between intensity and production was found. UVAR was 6.5 times more efficient than VIS in producing 1 nM of H_2O_2 , whereas UVBR was 228 times more efficient than VIS. When these data were weighted with respect to the energy of the solar spectrum at zenith hour, 28% of the H_2O_2 was formed by VIS, 23% was formed by UVAR and 48% was formed by UVBR. Considering the strong attenuation of UVBR in marine waters as compared with UVAR and VIS radiation, the role of UVAR deeper in the water column is recognised. Furthermore results of this research emphasise the importance of VIS radiation in the formation of H_2O_2 .

© 2003 Elsevier B.V. All rights reserved.

Keywords: Hydrogen peroxide; Organic peroxides; Atlantic Ocean; UV radiation; Photochemical production

1. Introduction

Hydrogen peroxide (H_2O_2) is an intermediate in the reduction of oxygen to water. It is a potential toxicant and it can affect the distribution and redox-chemistry of biologically active metals such as iron, copper and manganese. For example H_2O_2 is the dominant oxida-

tion pathway of Fe(II) (Moffett and Zika, 1987; Miller and Kester, 1994).

H_2O_2 can be formed by photochemical reactions and (to a lesser extent) by biota (Palenik et al., 1987; Palenik and Morel, 1988; Hanson et al., 2001). It can be introduced to the marine environment by dry and wet deposition (Miller and Kester, 1994; Hanson et al., 2001; Kieber et al., 2001). The focus of the present study was on the photochemical origin of H_2O_2 . Samples of natural surface and ground waters showed a rapid increase in H_2O_2 concentration after exposure

* Corresponding author.

E-mail address: loes@nioz.nl (L.J.A. Gerringa).

to sunlight (Cooper and Zika, 1983). H_2O_2 is photochemically generated from organic constituents present in water. Humic materials are believed to be the primary agent producing the H_2O_2 (Cooper and Zika, 1983). Distinct diel variation of the H_2O_2 concentration was observed with highest concentrations in the late afternoon. The observations are consistent with patterns of H_2O_2 formation due to photo-oxidation of dissolved organic matter (Zika et al., 1985; Palenik and Morel, 1988; Moore et al., 1993). The H_2O_2 concentrations were not only related to light intensity, but also to wavelength (Plane et al., 1987). Plane et al. (1987) demonstrated that production occurs primarily between 300 and 400 nm. This wavelength dependence of H_2O_2 formation is related to the light absorbance of organic substances in the water (Plane et al., 1987; Cooper et al., 1988; Sikorski and Zika, 1993). The absorbance of organic substances is highest in the UVBR region of the solar spectrum (280–320 nm), decreasing in the UVA region (320–400 nm) and lower at higher wavelengths (>400 nm). The weighting function, relating wavelength band and H_2O_2 formation was assessed by Yocis et al. (2000).

Published data of H_2O_2 concentrations in the Atlantic Ocean are scarce (Weller and Schrems, 1993; Miller and Kester, 1994; Obernosterer, 2000; Obernosterer et al., 2001). Recently, diel cycles in H_2O_2 concentrations were reported by Yuan and Shiller (2001) in the western central Atlantic Ocean. They observed a decrease of H_2O_2 concentrations with latitude from north to south.

In the present study we investigated factors affecting H_2O_2 formation in marine waters, such as solar irradiance and wavelength dependency. We measured surface H_2O_2 concentrations in the eastern Atlantic, from 25°N to 25°S, encountering various water masses and irradiance conditions.

Furthermore, during the cruise two diel cycles were measured. Finally, water taken near the equator (2°N) was incubated on deck under different wavelength conditions during which H_2O_2 formation was followed.

2. Material and methods

Measurements were executed during a cruise from Bremerhaven (Germany) to Cape Town (South Africa) aboard RV 'Polarstern' in October 2000

(Fig. 1). Data on salinity, temperature, and the position of the ship were obtained from the 'Polarstern' PODEV data acquisition system. Samples were taken under way by a FISH towed just under the sea surface, alongside the ship. The samples were filtered on-line through a 0.2 μm filter cartridge. Global radiation was measured by an artificially ventilated pyranometer (model CM11, Kipp and Zonen) mounted on the mast of the 'Polarstern'. It was fixed horizontally with respect to the ship. Rolling and pitching will have had no significant influence on averaged values, but on individual measurements. The values given are 1-min averages (Figs. 2 and 3). Light measurements used for the deck incubations were measured every hour using a high accuracy UV-Visible spectroradiometer (model OL 752, Optronic Laboratories). The OL 752 system consisted of an Optics Head (OL 752-O-PMT) and an OL 752-C

Fig. 1. Cruise track of cruise ANT18/1 of RV 'Polarstern' from Bremerhaven (Germany) to Cape Town (South Africa). The positions of H_2O_2 measurements are indicated by crosses, the positions of the experiments by numbers.

Fig. 2. (A) Global solar radiation in W m^{-2} ; (B) H_2O_2 (■) and organic peroxide (○) concentrations (all in nM) in sea surface waters; (C) salinity in psu; (D) temperature in $^{\circ}\text{C}$ against latitude. In (A) the three days on which experiments were executed (9, 15 and 20 October 2000) are indicated by arrows. Where no data are shown, no measurements were made.

Controller. One measurement of the solar spectrum, 280–750 nm, took ~ 28 min.

VIS is defined as light between 400 and 700 nm, UVAR between 320 and 400 nm, and UVBR between 280 and 320 nm. Diel cycles of H_2O_2 formation were

followed on 9 October near $23^{\circ}\text{N } 18^{\circ}\text{W}$ (1 in Fig. 1) and on 20 October near $23^{\circ}\text{S } 8.5^{\circ}\text{E}$ (3 in Fig. 1). Unfortunately, on 20 October the sky was cloudy, resulting in a global radiation at noon that was half the global radiation at noon on 9 October.

A sample of 50 L of filtered seawater was taken on 14 October, near $2^{\circ}\text{N}, 12^{\circ}\text{W}$ (2 in Fig. 1). The salinity of the water was 35.6 ppt. The water was stored in the dark at ambient seawater temperature (24°C) during the afternoon. After sunset, the water was sub-sampled in 2-L home-made polymethylmetacrylate (PMMA) bottles, which have a high transmission in the UV region (Steeneken et al., 1995). To assure minimal shading the bottles were, immediately after filling, put in UV transparent PMMA incubators on the highest deck of the ship. The bottles were kept at a constant ambient temperature of 24°C using running seawater from the ship's pumping system. Two PMMA bottles packed in light-tight black plastic functioned as dark controls. Two PMMA bottles received the full solar spectrum (UVBR, UVAR and VIS), two PMMA bottles were screened by glass, which served as a UVBR cut-off filter, receiving only UVAR and VIS, two PMMA bottles were screened by UV-opaque PMMA, only permeable for VIS and thus serving as a UVBR and UVAR cut-off filter. Total irradiance was varied by aluminium neutral density gauze, painted black, causing a 50% reduction in the full solar spectrum. 80% light reduction was obtained by wrapping two bottles in black plastic with tiny (1 mm) holes. Light reduction of these materials was measured before use. Sampling during the deck incubation started at 5:00 h UTC in the morning of 15 October and continued until 23:00 h UTC. Each bottle was sampled at least every two hours, and all light conditions were sampled at least once every hour. The dark bottles were only sampled three times during the day.

H_2O_2 was measured as fluorescence (Waters fluorometer, type 470) after enzyme-catalysed dimerisation of (*p*-hydroxyphenyl)acetic acid (POHPAA) (Miller and Kester, 1988, 1994). The POHPAA (Merck, purified by recrystallization) stock solution (25 mM in milliQ) was kept at 4°C . The peroxidase stock solution (horseradish, 10 000 U/val, Merck; 3.6 mg in 100 ml 0.25 M TRIS, pH 8.8) was stored in 2 ml portions at -20°C . A fluorometric reagent of POHPAA and peroxidase was prepared daily in 0.25 M TRIS (pH 8.8). Concentrations and enzyme activities

Fig. 3. H₂O₂ concentrations (nM) and global solar radiation (W m⁻²) of two diel cycles between 8 and 10 October 2000 at 23°N, and around 20 and 21 October 2000 at 23°S.

in the samples were $5.1 \cdot 10^{-6}$ M POHPAA and 0.153 units/ml (U/ml) peroxidase.

The fluorometric reagent was immediately added to the sample to fix the hydrogen peroxide. The samples (in triplicate) were stored at 4 °C and analysis was completed within 12 h. The fluorescence response is a combination of the signal of the hydrogen peroxide and the organic peroxides. To distinguish between these forms, 4 min before the addition of the fluorometric reagent, catalase (65 U/ml) was added to the samples to remove the hydrogen peroxide. The resulting fluorescence signal represented the organic peroxides, whereas subtraction from the total signal resulted in the hydrogen peroxide concentrations. Corrections were made for the natural fluorescence of seawater (no reactants added to the seawater) and the fluorescence of catalase (only catalase added to the seawater).

The precision of the method was 3% (st dev after 15 analyses of the same sample). Its detection limit was smaller than 2 nM (three times the standard deviation of open ocean water blank). The H₂O₂ standard was checked by measuring the reaction with KI at 353 nm (Cotton and Dunford, 1973).

3. Results and discussion

3.1. Surface samples from the transect

The H₂O₂ concentrations in the surface samples decreased from 70–80 nM at 25°N to 20–30 nM at 25°S in samples taken at noon (Fig. 2B). Obernosterer (2000) found surface values between 125 and 50 nM in the subtropical northern Atlantic Ocean in 1996. In 1992, Weller and Schrems (1993) measured concen-

trations from 150 nM at 30°N to 100 nM at 30°S over approximately the same cruise track as shown in Fig. 1. A decrease from north to south was also observed by Yuan and Shiller (2001) in the central and southern Atlantic Ocean. They suggest that this latitude dependence is due to differences in precipitation. Rain is a well-known source of H_2O_2 in oceans (Weller and Schrems, 1993; Miller and Kester, 1994; Yuan and Shiller 2000; Hanson et al., 2001).

Organic peroxides were measured irregularly; they varied between 1 and 6 nM (Fig. 2B). To our knowledge no other data on organic peroxides in the Atlantic Ocean have been published, making comparison impossible.

In the Northern Hemisphere the H_2O_2 concentrations did show a clear positive relationship with the global radiation (Fig. 3). A decrease in H_2O_2 concentrations, seen the night before, was followed by a sharp increase of more than 30 nM H_2O_2 with increasing radiation. Obernosterer (2000) found diel variations of 42 nM in the northern subtropical Atlantic Ocean, whereas Yuan and Shiller (2001) found a variation of 25 nM in the central and southern Atlantic Ocean. A slight decrease in concentration in our data occurred after sunset on 9 October (Fig. 3). Johnson et al. (1989) observed a similar slow decrease in peroxide concentrations with time and with depth in the western Mediterranean Sea, where they used H_2O_2 as a tracer for vertical advection. Johnson et al. (1989) calculated a decay rate of 3.8 nM h^{-1} during darkness. The decrease in the present data in Fig. 3 for the Northern Hemisphere, measured from the slope through the middle of the data points, would give a rate of 1 nM h^{-1} . Johnson et al. (1989) did not observe a decline in concentration until after sunset as is also the case in our data. This is especially the case for seawater compared to a relatively rapid decrease observed in fresh water (Cooper and Lean, 1989).

The second diel cycle observation in the Southern Hemisphere gave a different result (Fig. 3). A potential diel variation of 5–10 nM in H_2O_2 concentration disappeared in the noise of the measurements during the day and the following night. Since the global radiation was lower due to clouds, a lower production of H_2O_2 was expected. In order to check whether the difference in H_2O_2 production could be explained by the difference in the solar intensity, a

linear relationship between H_2O_2 production and the intensity of the global radiation of 9 and 20 October was calculated. The slope of the linear relation in the data of 9 October was 2.7 times higher than that of 20 October. Thus the lower magnitude in the diel cycle must have had an additional cause, most likely a lower concentration in organic matter in the Southern Hemisphere sampling station (Scully et al., 1996). Both the northern and the southern experiments were performed in subtropical gyres, which presumably have relatively nutrient-poor surface waters (Antoine et al., 1996; Field et al., 1998). The northern experiment was conducted in warmer and more saline waters ($S=36.5$, $T=23.1^\circ\text{C}$ versus $S=35.4$, $T=16.2^\circ\text{C}$, Fig. 2). Both locations are relatively close to deserts, known to release dust into the adjacent ocean. Both positions are close to a zone of coastal upwelling (Krauss, 1996; Shannon and Nelson, 1996). However, the northern position is closer to the coast and is thus more likely to receive organic-rich water from the coastal upwelling. Since the formation of H_2O_2 results principally from the excitation of humic substances (Cooper et al., 1988; Scully et al., 1996) differences in H_2O_2 can be expected from differences in water characteristics such as content of organic material. Moore et al. (1993) related peroxide production with organic matter fluorescence in the eastern Caribbean. Fluorescence was measured continuously aboard the 'Polarstern', but values were very low and had the same order of magnitude for both regions. No rain fell during the cruise, giving no additional explanation for the relatively high diel cycle of H_2O_2 at the northern position or for the decrease in H_2O_2 concentrations from north to south in this part of the Atlantic Ocean.

3.2. Deck incubations

The results of the deck incubations showed a clear relationship between H_2O_2 concentration and the wavelength and cumulative dose of sunlight (Figs. 4 and 5). More H_2O_2 was produced with decreasing wavelength (Fig. 5A). Even when only VIS was allowed through the bottles, H_2O_2 concentrations increased during the day compared to the dark control bottles. The net H_2O_2 production rate at noon was 7 nM h^{-1} in the bottles with the

Fig. 4. (A) UVAR and UVBR (W m^{-2}) of the solar spectrum of 15 October 2000 against time (UTC); (B) the cumulative energies for UVAR and UVBR against time (UTC).

full solar spectrum, 4 nM h^{-1} in the bottles without UVBR and 2 nM h^{-1} in the bottles in which only VIS was allowed. In the dark control bottles, $0.4 \text{ nM h}^{-1} \text{ H}_2\text{O}_2$ formed. Taking into account that 88% of the energy of the solar spectrum consisted of VIS, 11% consisted of UVAR, and less than 1% of UVBR, the relationship between wavelength and H_2O_2 was much stronger than the production rates show.

A net production rate under the full solar spectrum of 7 nM h^{-1} compares well with other studies. Yuan and Shiller (2001) found 8.3 nM h^{-1} at local noon in the central Atlantic Ocean. Obernosterer (2000) found a net production of 5.5 nM h^{-1} in the northern subtropical region of the Atlantic Ocean. Yocis et al. (2000) measured 4.5 nM h^{-1} in Antarctic waters and attributed this relatively low production to the high latitude, lower temperatures and lower UV irradiances in polar waters.

In order to calculate the wavelength dependency of the net H_2O_2 production by UVAR, the concentrations of H_2O_2 formed under the full solar spectrum were subtracted from those formed under UVAR and VIS. By subtracting concentrations formed under UVAR and VIS from those under VIS we obtained the production caused by UVAR.

The light intensities of the UVBR, UVAR and the VIS portion of the solar spectrum were fitted into a model in order to explain the increases in H_2O_2 . It was assumed that the relation between H_2O_2 production and light intensity is linear (Cooper et al., 1994). It was assumed that the increase in peroxide concentration C at time t_{x+1} was due to the energy of the incoming

Fig. 5. Peroxide concentrations (nM) in the incubated bottles against time (UTC). Two bottles per light condition were used, distinguished by filled and open symbols. (A) Peroxide production related to wavelength of the incoming light; (B) peroxide production related to light intensity. In both figures the data of the dark control bottles are presented. 100% light intensity in (B) is similar to the data of UVBR, UVAR and VIS in (A).

light E between t_x and t_{x+1} multiplied by a factor L , according to:

$$C_{x+1} = C_x + ((E_x + E_{x+1})/2) * (t_x - t_{x+1}) * L \quad (1)$$

L ($\text{nM min}^{-1} \text{ W}^{-1} \text{ m}^2$) was found by minimising the root mean square (RSQ) of the differences between measured and calculated concentrations. To account for the oxidation of peroxides, a loss function was included in the model, assuming $dC/dt = LE - KC$. However, this loss factor (K) did not improve the model and was left out.

A 50% light reduction indeed gives half of the peroxide production, a reduction to 20% of the original light causes the production to decrease to 30% compared to full light conditions (Fig. 5B, Table 1). This discrepancy between 20 and 30% is difficult to explain. The light reduction to 20% by the plastic was valid for the entire solar spectrum, independent of wavelength.

The importance of UVBR in the production of H_2O_2 is evident, confirming the results of Cooper et al. (1994). This radiation is 227 times more efficient for the production of H_2O_2 than VIS and 35 times more efficient than UVAR (Table 1). The result of Yocis et al. (2000) showed a comparable ratio in the UVBR-UVAR region, demonstrating a factor 100 in the relationship between wavelength and peroxide production for the wavelengths at 290 and at 410 nm. However, some care must be taken in directly comparing the results from Yocis et al. (2000) and our results. We did not incorporate

wavelength-related differences in absorption into our results.

The energy distribution of the solar spectrum at noon on 14 October near the equator consisted of 88% VIS, 11.4% UVAR and 0.65% UVBR (280–700 nm: 391 W m^{-2} , 400–700: 346 W m^{-2} , 320–400 nm: 43 W m^{-2} , 280–320 nm: 2.8 W m^{-2}). Considering the contribution of the energies in the solar spectrum and using the L factors from Table 1, we obtained a H_2O_2 production rate at noon of which 28% originated from the VIS region, 24% from UVAR and 48% from UVBR.

UVAR penetrates to greater depths than UVBR, so the role of UVAR may become increasingly important at greater water depths (Scully et al., 1996; Obernosterer, 2000; Yocis et al., 2000). The decrease with depth found paralleled decrease in UV penetration. However, the data from our study showed an important contribution of VIS to H_2O_2 formation as well. This was also concluded by Sikorski and Zika (1993). They constructed a model in which the optical properties of light in water play an important role. They found that the attenuation of light was very sensitive to the solar angle of incident light and that this effect has a large wavelength dependency. They concluded that due to these optical properties of the solar spectrum, the visible part of the irradiance penetrates much deeper into the water column compared to the UV part of the spectrum than was thought before. They therefore concluded that part of the solar spectrum above 400 nm caused a significant production of hydrogen peroxides, especially at larger depths and at smaller zenith angles.

Table 1

The factor L indicating the relation between light intensity (W m^{-2}) and peroxide increase (nM min^{-1})

Experimental conditions	Factor $L * 10^{-3}$ ($\text{nM min}^{-1} \text{ W}^{-1} \text{ m}^2$)	RSQ(C_{x+1}, C_x)
20% of full spectrum	0.112	0.76
50% of full spectrum	0.175	1.46
100% of full spectrum	0.347	1.77
VIS	0.121	1.2
UVAR	0.785	1.29
UVBR	27.5	2

L was obtained by fitting the data in equation (1). The obtained root mean square of the differences between measured and calculated concentrations (RSQ C_{x+1}, C_x) are given.

Acknowledgements

We are very grateful to captain Keil and his crew of RV ‘Polarstern’. We thank the cruise leader Saad el Naggat and the Alfred-Wegener-Institut für Polar- und Meeresforschung (AWI) for facilities and hospitality. Hendrik van Aken (NIOZ) was so kind as to enlighten us on the circulation of the sub-equatorial regions of the Atlantic Ocean and Ingrid Obernosterer (NIOZ) introduced us to H_2O_2 analysis in seawater. We wish to thank Rick Tax and Willem Willemsen from the Dutch National Institute of Public Health and the Environment (RIVM), Wim

Boot from the Institute for Marine and Atmospheric Research Utrecht (IMAU), Foeke Kuik from the Royal Netherlands Meteorological Institute (KNMI), and Saad el Naggar from AWI for the equipment and advice on the collection of the UVR data. This research is funded by NWO/NAAP grant number 85120004.

References

- Antoine, D., André, J.-M., Morel, A., 1996. Oceanic primary production. 2. Estimation at global scale from satellite (coastal zone color scanner) chlorophyll. *Global Biogeochem. Cycles* 10, 57–69.
- Cooper, W.J., Lean, D.R.S., 1989. Hydrogen peroxide concentration in a Northern Lake: photochemical formation and diel variability. *Environ. Sci. Technol.* 23, 1425–1428.
- Cooper, W.J., Zika, R.G., 1983. Photochemical formations of hydrogen peroxide in surface and ground waters exposed to sunlight. *Science* 220, 711–712.
- Cooper, W.J., Zika, R.G., Petasne, R.G., Plane, J.C., 1988. Photochemical formation of H_2O_2 in natural waters exposed to sunlight. *Environ. Sci. Technol.* 22, 1156–1160.
- Cooper, W.J., Shao, C., Lean, D.R.S., Gordon, A.S., Scully Jr., F.E., 1994. Factors affecting the distribution of H_2O_2 in surface waters. In: Baker, L.A. (Ed.), *Environmental Chemistry in Lakes and Reservoirs*. Adv. Chem. Ser., vol. 237, pp. 393–422.
- Cotton, M.L., Dunford, H.B., 1973. Studies on horseradish peroxidase. XI. On the nature of compounds I and II as determined from the kinetics of oxidation of ferrocyanide. *Can. J. Chem.* 51, 582–587.
- Field, C.B., Behrenfeld, M.J., Randerson, J.T., Falkowski, P., 1998. Primary production of the biosphere: integrating terrestrial and oceanic components. *Science* 281, 237–240.
- Hanson, A.K., Tindale, N.W., Abdel-Moati, M.A.R., 2001. An Equatorial Pacific rain event: influence on the distribution of iron and hydrogen peroxide in surface waters. *Mar. Chem.* 75, 69–88.
- Johnson, K.S., Williamson, S.W., Wiesenburg, D.A., Lohrenz, S.E., Arnone, R.A., 1989. Hydrogen peroxide in the Western Mediterranean Sea: a tracer for vertical advection. *Deep-Sea Res.* 36, 241–254.
- Kieber, R.J., Cooper, W.J., Willey, J.D., Avery Jr., G.B., 2001. Hydrogen peroxide at the Bermuda Atlantic time series station. Part I: temporal variability of atmospheric hydrogen peroxide and its influence on seawater concentrations. *J. Atm. Chem.* 39, 1–13.
- Krauss, W., 1996. Comments on the development of our knowledge of the general circulation of the North Atlantic Ocean. In: Krauss, W. (Ed.), *The Warm Water Sphere of the North Atlantic Ocean*. Borntraeger, Berlin, pp. 1–32.
- Miller, W.L., Kester, D., 1988. Hydrogen peroxide measurement in seawater by (*p*-hydroxyphenyl) acetic acid demerization. *Anal. Chem.* 60, 2711–2715.
- Miller, W.L., Kester, D., 1994. Peroxide variations in the Sargasso Sea. *Mar. Chem.* 48, 17–29.
- Moffett, J.W., Zika, R.G., 1987. Reaction kinetics of hydrogen peroxide with copper and iron in seawater. *Environ. Sci. Technol.* 21, 804–810.
- Moore, C.A., Farmer, C.T., Zika, R.G., 1993. Influence of the Orinoco river on the hydrogen peroxide distribution and production in the Eastern Caribbean. *J. Geophys. Res.* 98, 2289–2298.
- Obernosterer, I.B., 2000. Photochemical transformations of dissolved organic matter and its subsequent utilization by marine bacterioplankton. Ph.D. Thesis, University of Groningen, The Netherlands. 133 pp.
- Obernosterer, I.B., Ruardij, P., Herndl, G.J., 2001. Spatial and diurnal dynamics of dissolved organic matter (DOM) fluorescence and H_2O_2 and the photochemical oxygen demand of surface water DOM across the subtropical Atlantic Ocean. *Limnol. Oceanogr.* 46, 632–643.
- Palenik, B., Morel, F.M.M., 1988. Dark production of H_2O_2 in the Sargasso Sea. *Limnol. Oceanogr.* 33, 1606–1611.
- Palenik, B., Zafiriou, O.C., Morel, F.M.M., 1987. Hydrogen peroxide production by a marine phytoplankter. *Limnol. Oceanogr.* 32, 1365–1369.
- Plane, J.M.C., Zika, R.G., Zepp, R.G., Burns, L.A., 1987. Photochemical modeling applied to natural waters. In: Zika, R.G., Cooper, W.J. (Eds.), *Photochemistry of Environmental Aquatic Systems*. American Chemical Society Group Washington D.C, pp. 250–267.
- Scully, N.M., McQueen, D.J., Lean, D.R.S., Cooper, W.J., 1996. Hydrogen peroxide formation: the interaction of UV radiation and dissolved organic carbon in lake waters along a 43–75 degrees N gradient. *Limnol. Oceanogr.* 41, 540–548.
- Shannon, L.V., Nelson, G., 1996. The Benguela: large scale features and processes and system variability. In: Wefer, G., Berger, W.H., Siedler, G., Webb, D.J. (Eds.), *The South Atlantic: Present and Past Circulation*. Springer Verlag, Berlin, pp. 163–210.
- Sikorski, R.J., Zika, R.G., 1993. Modeling mixed-layer photochemistry of H_2O_2 —optical and chemical modeling of production. *J. Geophys. Res.* 98, 2315–2328.
- Steeneken, S.F., Buma, A.G.J., Gieskes, W.W.C., 1995. Changes in transmission characteristics of polymethylmethacrylate and cellulose (III) acetate during exposure to ultraviolet light. *Photochem. Photobiol.* 61, 276–280.
- Weller, R., Schrems, O., 1993. H_2O_2 in the marine troposphere and seawater of the Atlantic Ocean (48°N–63°W). *Geophys. Res. Lett.* 20, 125–128.
- Yocis, B.H., Kieber, D.J., Mopper, K., 2000. Photochemical production of hydrogen peroxide in Antarctic waters. *Deep. Sea Res.* I 47, 1077–1099.
- Yuan, J., Shiller, A.M., 2000. The variation of hydrogen peroxide in rainwater over the South and Central Atlantic Ocean. *Atm. Environm.* 34, 3973–3980.
- Yuan, J., Shiller, A.M., 2001. The distribution of hydrogen peroxide in the southern and central Atlantic Ocean. *Deep-Sea Res.* II 48, 2947–2970.
- Zika, R.G., Saltzman, E.S., Cooper, W.J., 1985. Hydrogen peroxide concentrations in the Peru upwelling area. *Mar. Chem.* 17, 265–275.