University of Groningen

Mutual influence of backbone proline substitution and lipophilic tail character on the biological activity of simplified analogues of caspofungin

Mulder, Monique P. C.; Fodran, Peter; Kemmink, Johan; Breukink, Eefjan J.; Kruijtzer, John A. W.; Minnaard, Adriaan J.; Liskamp, Rob M. J.

Published in:
Organic & Biomolecular Chemistry

DOI:
[10.1039/c2ob25951f](https://doi.org/10.1039/c2ob25951f)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2012

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Mulder, M. P. C., Fodran, P., Kemmink, J., Breukink, E. J., Kruijtzer, J. A. W., Minnaard, A. J., & Liskamp, R. M. J. (2012). Mutual influence of backbone proline substitution and lipophilic tail character on the biological activity of simplified analogues of caspofungin. *Organic & Biomolecular Chemistry*, 10(37), 7491-7502. <https://doi.org/10.1039/c2ob25951f>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Supplementary data

Mutual Influence of Backbone Proline Substitution and Lipophilic Tail Character on the Biological Activity of Simplified Analogues of Caspofungin

Monique P.C. Mulder^[a], Peter Fodran^[b], Johan Kemmink^[a], Eefjan Breukink^[c], John A.W. Kruijtz^[a],
Adriaan J. Minnaard^[b] and Rob M.J. Liskamp^[a]

^a Medicinal Chemistry and Chemical Biology, Utrecht Institute for Pharmaceutical Sciences, Utrecht University, PO Box 80.082, 3508 TB Utrecht, The Netherlands

^b Department of Bio-organic Chemistry,, Stratingh Institute for Chemistry, University of Groningen, Nijenborgh 7, 9747 AG, Groningen, The Netherlands

^c Biochemistry of Membranes, Bijvoet Center for Biomolecular Research and Institute of Biomembranes, Utrecht University, Padualaan 8, 3584 CH Utrecht, The Netherlands

Additional CD spectra:

Figure S 1. 22-membered ring size analogues **22** and **23** with a terphenyl and acetyl tail, respectively.

Figure S 2. CD spectra of analogues (**22** and **23**). A) Measured in MeCN/H₂O (1/1, v/v); B) Measured in TFE/H₂O (1/1, v/v). All peptides were measured at 0.1 mM concentration.

A distinct different pattern in CD of analogue **22** with a terphenyl tail was observed, at wavelengths below 210 nm, compared to analogue **23** (Figure S 2). This corroborates the observation regarding the influence of the terphenyl side chain on CD spectra. And may be explained by a large contribution of the aromatic residues present in the terphenyl side chain.

Modelling:

Modeling of the structures **3** and **6** was accomplished using the YASARA Structure 10.5.2.1 software package. Structures were energy minimized using the simulated annealing protocol employing the AMBER99¹ force field. A 2500 ps MD in water was run of each structure. After 250 ps equilibration, the structure with the lowest energy trajectory (between 250-2500 ps) was saved as a yob file. Molecules were superimposed, with the reported crystal structure of ECBN, by minimizing the rmsd between the backbone atoms in the ring.

Figure S 3. Superimposition of ECBN (carbons colored green) with analogues **3** (right) and **6** (left) (carbons colored orange). The backbone atoms have been used as fixed coordinates for superimposition.

¹ Ponder, J.W.; Case, D.A. *Adv. Protein Chem.*, **2003**, 66, 27-85.

NMR

Cyclo[-(DMT-Myr)-Orn-Thr-Hyp-hTyr-Orn-3-Hyp] (4)

Cyclo[-(Me-Myr)-Orn-Thr-Hyp-hTyr-Orn-3-Hyp] (5)

(E)-S-ethyl pent-2-enethioate (10)

(E)-S-ethyl pent-2-enethioate (10)

(*S*)-*S*-ethyl 3-methylpentanethioate (11)

(*S*)-*S*-ethyl 3-methylpentanethioate (11)

(3*S*,5*S*)-*S*-ethyl 3,5-dimethylheptanethioate (13)

(3*S*,5*S*)-*S*-ethyl 3,5-dimethylheptanethioate (13)

(10*S*,12*S*)-10,12-dimethyltetradec-7-enoic acid (14)

(10*S*,12*S*)-10,12-dimethyltetradec-7-enoic acid (14)

(10*R*,12*S*)-10,12-dimethyltetradecanoic acid (8)

(10*R*,12*S*)-10,12-dimethyltetradecanoic acid (8)

(*S*)-12-methyltetradec-9-enoic acid (15)

(*S*)-12-methyltetradec-9-enoic acid (15)

Cyclo[-(Ter)-Orn-Thr-Hyp-hTyr-Orn-3-Hyp] (19)

Cyclo[-(DMT-Myr)-Orn-Thr-Hyp-pTyr-Orn-Hyp] (20)