University of Groningen

Catalytic Enantioselective Carbon-Carbon Bond Formation by Addition of Dialkylzinc Reagents to Cyclic 1,3-Diene Monoepoxides.

Badalassi, F.; Crotti, P.; Macchia, F.; Pineschi, M.; Arnold, L.A.; Feringa, B.L.

Published in:
Tetrahedron Letters

DOI:
[10.1016/S0040-4039\(98\)01704-3](https://doi.org/10.1016/S0040-4039(98)01704-3)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1998

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Badalassi, F., Crotti, P., Macchia, F., Pineschi, M., Arnold, L. A., & Feringa, B. L. (1998). Catalytic Enantioselective Carbon-Carbon Bond Formation by Addition of Dialkylzinc Reagents to Cyclic 1,3-Diene Monoepoxides. *Tetrahedron Letters*, 39(42), 7795 - 7798. [https://doi.org/10.1016/S0040-4039\(98\)01704-3](https://doi.org/10.1016/S0040-4039(98)01704-3)

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Pergamon

Tetrahedron Letters 39 (1998) 7795–7798

TETRAHEDRON
LETTERS

Catalytic Enantioselective Carbon-Carbon Bond Formation by Addition of Dialkylzinc Reagents to Cyclic 1,3-Diene Monoepoxides.

Fabrizio Badalassi,^a Paolo Crotti,^a Franco Macchia,^a Mauro Pineschi,^{*a}
Alexander Arnold^b and Ben L. Feringa^{*b}

^a Dipartimento di Chimica Bioorganica, Università di Pisa, Via Bonanno 33, 56126 Pisa, Italy

^b Department of Organic and Molecular Inorganic Chemistry, University of Groningen, Nijenborgh 4, NL, AG Groningen, The Netherlands.

Fax: +39-50-43321; E-mail: pineschi@farm.unipi.it. E-mail: feringa@chem.rug.nl

Received 10 July 1998; accepted 11 August 1998

Abstract. Chiral copper complexes of 2,2'-binaphthyl-based phosphorus amidites are shown to be highly effective catalysts for the conjugate addition of dialkylzinc reagents to vinyloxiranes. The corresponding allylic alcohol reaction products (S_N2' -pathway) were obtained with moderate to high regioselectivity. Both direct- (S_N2 -pathway) and conjugate-opening addition (S_N2' -pathway) seem to proceed with complete *anti* stereoselectivity. The enantioselectivity of these addition reactions according to a kinetic resolution protocol turned out to be high (>90% ee) with 1,3-cyclohexadiene and 1,3-cycloheptadiene monoepoxides.

© 1998 Elsevier Science Ltd. All rights reserved.

The catalytic enantioselective formation of carbon-carbon bonds is one of the more challenging areas in organic synthesis.¹ The enantioselective nucleophilic ring opening of oxiranes by the use of a chiral catalyst has become an attractive method for the preparation of optically active compounds.² Despite the fact that the catalyzed enantioselective ring opening of epoxides with several nucleophiles has been investigated,³ the catalyzed enantioselective addition of organometallics to oxiranes is an almost unexplored area,⁴ although this reaction may offer a powerful tool for the stereocontrolled synthesis of complex organic compounds. The S_N2' displacements of vinyloxiranes with cuprates afford synthetically useful allylic alcohols.⁵ In this context, the reactions of 1,3-cycloalkadiene monoepoxides have received considerable attention.⁶ The ratio of S_N2'/S_N2 products in the nucleophilic addition to cyclic vinyloxiranes turned out to be very sensitive to the reagent type and highly regio- and stereoselective additions of mixed cyanocuprates have been developed.⁶ Only recently Lipshutz et al. have reported an unprecedented copper-catalyzed cross-coupling of functionalized zinc reagents with vinyloxiranes.⁷ However a catalytic asymmetric version of this synthetically useful reaction has not been described to our knowledge.

In the present work we wish to report our preliminary results regarding the catalytic enantioselective addition of simple commercially available organozinc reagents to cycloaliphatic vinyloxiranes. The remarkable ligand-accelerated catalysis⁸ and the high enantioselectivities previously obtained in the addition of organozinc reagents to enones,⁹ prompted us to verify the possibility of using chiral copper complexes of 2,2'-binaphthyl-based phosphorus amidites, such as **1** and **2**, in the conjugate addition of organozinc reagents to vinyloxiranes (see Figure).

The reaction of Me_2Zn (1.5 eq.) with **4** (-30°C, 3h) catalyzed by $Cu(OTf)_2$ (3 mol%) afforded a complex reaction mixture containing a small amount (8 % combined yield) of a 2:1 mixture of allylic alcohols **10** (*anti*- S_N2' pathway) and **11** (*anti*- S_N2 pathway), respectively (see Scheme).

Figure

The preventive addition (see typical procedure)¹⁰ of a catalytic amount (6 mol%) of phosphorus amidite ligand (+)-1 to the reaction mixture gave a dramatic increase in the conjugate addition pathway. In this case we obtained a 9:1 mixture of unsaturated alcohols **10** and **11**, with a substantial increase in the regioisomer **10** (55% isolated yield, data not shown in the Table). To our surprise, we found that allylic alcohol **10** showed a 60% ee despite the fact that the reaction was not performed under kinetic resolution conditions (Me_2Zn excess). The use of 0.50 eq. of Me_2Zn (-70°C , 1h), together with the chiral ligand **1** (6 mol%) and $\text{Cu}(\text{OTf})_2$ (3 mol%) gave practically the same enantioselectivity (entry 2, Table). This clearly indicated the preferential reactivity of one enantiomer of the racemic epoxide **4** with a high stereoselectivity factor.¹¹

Scheme

The same reaction carried out with 0.50 eq. of Et_2Zn gave the corresponding allylic alcohol **12** in a low yield (18%) but with a substantial increase in both enantioselectivity and regioselectivity (entry 3, Table). The addition of Me_2Zn to 1,3-cycloheptadiene monoepoxide **5** showed similar levels of enantioselectivity and regioselectivity compared with vinyloxirane **4** (entry 4, Table). The cyclopentadiene monoepoxide **3** was found to be less reactive under these reaction conditions (-70°C , 1h) affording the corresponding conjugate adduct **6** with a low yield and regioselectivity and a moderate enantioselectivity (entry 1, Table).

Having established the ability of phosphorus amidite **1** to strongly accelerate the rate of addition of dialkyl zinc reagents to vinyloxiranes **3-5**, we tested the chiral ligand (+)-2, derived from (*S*)-2,2'-binaphthol and bis-(*R*)-1-phenylethylamine (entries 5-9, Table).^{9b}

While the chiral ligand **2** showed a similar level of enantioselectivity if compared to **1** in the addition of organozinc reagents to cyclopentadiene monoepoxide **3** (entries 5 and 6, Table), high enantioselectivities (>90%ee) were obtained in the cases of the six- and seven-membered analogs **4** and **5**. Once again we found an increased regioselectivity for 1,4-addition when Et_2Zn was used (entries 6 and 8, Table). The highest

Table. Enantioselective Conjugate Addition of Dialkylzinc Reagents to Vinyloxiranes 3-5 Catalyzed by Cu(OTf)₂/ (+)-1 and (+)-2.^a

Entry	substrate	ligand	R ₂ Zn	yield (%) ^b	ee (%) ^c	S _N 2'/S _N 2 ^d
1	3	1	R=Me	10 ^e	46	4
2	4	1	R=Me	30	62 (-)	13
3	4	1	R=Et	18	86 (-)	>20
4	5	1	R=Me	32	60 (-)	10
5	3	2	R=Me	12 ^e	50	3
6	3	2	R=Et	8 ^e	54	12
7	4	2	R=Me	33	92(-)	13
8	4	2	R=Et	32	91(-)	59
9	5	2	R=Me	38	96(-)	16

^a All reactions were run as described in ref.10 (0.50 eq. R₂Zn). ^b Isolated yield of allylic alcohols after chromatographic purification (SiO₂), unless otherwise stated (maximum yield 50%). ^c All ee values (compounds 6, 8, 10, 12, 14) were determined by chiral GC (Chromopak fused silica 50 m X 0.25 mm, coated with CP-Cyclodextrin-B-236-M-19). The sign of rotation of the predominant enantiomer is indicated in parentheses. Absolute configuration not determined. ^d Composition determined by GC capillary analysis of the crude reaction mixture. ^e GC yield using 2-cyclohexen-1-ol as the internal quantitative standard.

enantioselectivity was observed for the addition of Me₂Zn to 5 (96% ee, see typical procedure).¹⁰ In this case, the regioselectivity (S_N2'/S_N2 ratio) is even better than that obtained for the addition of MeCuCNLi to vinyloxirane 5 (entry 9, Table).¹²

In conclusion, the present work represents the first report demonstrating a successful combination of an organometallic reagent and an external chiral ligand in a kinetic resolution protocol for the nucleophilic displacement of racemic vinyloxiranes. Chiral copper complexes of 2,2'-binaphthyl-based phosphorus amidites were found to exhibit a strong ligand accelerated catalysis effect in the conjugate addition of dialkylzinc reagents to vinyloxiranes. The high selection of enantiotopic reaction sites shown by 1,3-cyclohexadiene- and 1,3-cycloheptadiene-monoepoxides 4 and 5 with organozinc reagents can allow a versatile entry for optically active 4-alkyl-2-cycloalken-1-ols of six- and seven-membered rings. Additional studies of the scope and mechanism of this process are in progress.

References and Notes

- (1) *Catalytic Asymmetric Synthesis*; Ojima, I., Ed.; VCH Publishers: New York, 1993.
- (2) Paterson, I.; Berrisford, D.J. *Angew. Chem. Int. Ed. Engl.* **1992**, *31*, 1179.
- (3) For leading references see: (a) Martinez, L.E.; Leighton, J.L.; Carsten, D.H.; Jacobsen, E.N. *J. Am. Chem. Soc.* **1995**, *117*, 5897. (b) Tokunaga, M.; Larrow, J.F.; Kakiuchi, F.; Jacobsen, E.N. *Science* **1997**, *277*, 936. (c) Cole, B.M.; Shimizu, K.D.; Kruger, C.A.; Harrity, J.P.A.; Snapper, M.; Hoveyda, A.H. *Angew. Chem. Int. Ed. Engl.* **1996**, *35*, 1668. (d) Iida, T.; Yamamoto, N.; Sasai, H.; Shibasaki, M. *J. Am. Chem. Soc.* **1997**, *119*, 4783. (e) Hodgson, D.M.; Gibbs, A.R.; Lee, G.L. *Tetrahedron* **1996**, *52*, 14361.

- (4) (a) Davies, S.G.; Wollowitz, S. *Tetrahedron Lett.* **1980**, *21*, 4175. (b) Mizuno, M.; Kanai, M.; Iida, A.; Tomioka, K. *Tetrahedron: Asymmetry* **1996**, *7*, 2483. (c) Crotti, P.; Di Bussolo, V.; Favero, L.; Macchia, F.; Pineschi, M. *Gazz. Chim. Ital.* **1997**, *127*, 273.
- (5) Marshall, J.A. *Chem. Rev.* **1989**, *89*, 1503.
- (6) Marino, J.P.; Fernandez de la Pradilla, R.; Laborde, E. *J. Org. Chem.* **1987**, *52*, 4898 and references therein.
- (7) Lipshutz, B.H.; Woo, K.; Gross, T.; Buzard, D.J.; Tirado, R. *Synlett*, **1997**, 47, 7.
- (8) Berrisford, D.J.; Bolm, C.; Sharpless, K.B. *Angew. Chem. Int. Ed. Engl.* **1995**, *34*, 1059.
- (9) (a) de Vries, A.H.M.; Meetsma, A.; Feringa, B.L. *Angew. Chem. Int. Ed. Engl.* **1996**, *35*, 2374. (b) Feringa, B.L.; Pineschi, M.; Arnold, L.A.; Imbos, R.; de Vries, A.H.M. *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 2620. For a brief review, see: Krause, N. *Angew. Chem. Int. Ed. Engl.* **1998**, *37*, 283.
- (10) Typical procedure as follows (entry 9, Table): A solution of Cu(OTf)₂ (0.022 g, 0.06 mmol) and 2 (0.070 g, 0.12 mmol) in anhydrous toluene (6.0 ml) was stirred at rt for 40 min. The colorless solution was cooled to -70°C, and subsequently, a solution of 5 (0.220 g, 2.0 mmol) in toluene (1.0 ml) and 0.50 ml of a 2.0 M Me₂Zn in toluene, were added. The temperature was allowed to warm slowly to -30°C (1h) and the mixture was quenched with saturated aqueous NH₄Cl (3.0 ml). Extraction with Et₂O and evaporation of the dried (MgSO₄) organic phase gave a crude product which was subjected to flash chromatography (SiO₂, 8:2 hexanes:EtOAc) to afford (-)-trans-4-methyl-2-cyclohexen-1-ol¹² (**14**) (0.094 g, 38%) as a pale yellow liquid: [α]_D = -106.5 (c=0.15, CH₂Cl₂).
- (11) Throughout this work the ee values of the unreacted vinyloxiranes were not determined. For a discussion of kinetic resolution see Kagan, H.B.; Fiaud, J.C. *Top. Stereochem.* **1988**, *18*, 249.
- (12) Marino, J.P.; Floyd, D.M. *Tetrahedron Lett.* **1979**, *20*, 675.

Acknowledgments. This work was supported by the Consiglio Nazionale delle Ricerche (CNR) and the Ministero della Universita' e della Ricerca Scientifica e Tecnologica (MURST), Roma. We acknowledge generous financial support from the European Community (EU contract no. ERBFMBICT961635) in the early stages of this project, and from Merck 1998 Academic Development Program (ADP) Chemistry Award to P.C..