University of Groningen

Stereoselective synthesis of syn and anti 1,2-hydroxyalkyl moieties by Cu-catalyzed asymmetric allylic alkylation

Fañanás-Mastral, Martín; Horst, Bjorn ter; Minnaard, Adriaan J.; Feringa, Bernard

Published in:
Chemical Communications

DOI:
[10.1039/c0cc05161f](https://doi.org/10.1039/c0cc05161f)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2011

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Fañanás-Mastral, M., Horst, B. T., Minnaard, A. J., & Feringa, B. (2011). Stereoselective synthesis of syn and anti 1,2-hydroxyalkyl moieties by Cu-catalyzed asymmetric allylic alkylation. *Chemical Communications*, 47(20), 5843-5845. <https://doi.org/10.1039/c0cc05161f>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

ELECTRONIC SUPPLEMENTARY INFORMATION

Stereoselective synthesis of *syn* and *anti* 1,2-hydroxyalkyl moieties by Cu-catalyzed asymmetric allylic alkylation

Martín Fañanás-Mastral, Bjorn ter Horst, Adriaan J. Minnaard and Ben L. Feringa*

*Stratingh Institute for Chemistry, University of Groningen, Nijenborgh 4, 9747 AG, Groningen,
The Netherlands.*

Table of Contents:

General Procedures	S3
General procedure for the stereoselective Cu-catalyzed synthesis of 1,2-hydroxyalkyl compounds 5	S3
Characterization of <i>anti</i> and <i>syn</i> compounds 5	S4
General procedure for the synthesis of compounds 7 and α,β-unsaturated δ-lactones 8	S8
Characterization of compounds 7 and 8	S9
NMR spectra	S12
Proposed mechanism for the copper-catalyzed AAA of allyl bromide 4 with Grignard reagents	S25
GC traces for the temperature-depending experiments for the AAA of allyl bromide 4 with MeMgBr catalyzed by Cu-TaniaPhos	S26

General Procedures:

Chromatography: Merck silica gel type 9385 230-400 mesh, TLC: Merck silica gel 60, 0.25 mm. Components were visualized by UV and cerium/molybdenum staining. Progress and conversion of the reaction were determined by GC-MS (GC, HP6890; MS HP5973) with an HP1 or HP5 column (Agilent Technologies, Palo Alto, CA). Mass spectra were recorded on a AEI-MS-902 mass spectrometer (EI+) or a LTQ Orbitrap XL (ESI+). ^1H - and ^{13}C -NMR were recorded on a Varian AMX400 (400 and 100.59 MHz, respectively) or a Varian VXR300 (300 and 75 MHz, respectively) using CDCl_3 as solvent. Chemical shift values are reported in ppm with the solvent resonance as the internal standard (CHCl_3 : δ 7.26 for ^1H , δ 77.0 for ^{13}C). Data are reported as follows: chemical shifts, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, br = broad, m = multiplet), coupling constants (Hz), and integration. Carbon assignments are based on APT ^{13}C -NMR experiments. Optical rotations were measured on a *Schmidt + Haensch* polarimeter (Polartronic MH8) with a 10 cm cell (c given in g/100 mL).

All reactions were carried out under a nitrogen atmosphere using oven dried glassware and using standard Schlenk techniques. CH_2Cl_2 was dried and distilled over calcium hydride. Substrate **4** was prepared according to literature procedures.¹ $\text{CuBr}\cdot\text{SMe}_2$, Grubbs 2nd generation and Hoveyda-Grubbs 2nd generation catalysts, ligands **L1-4** and commercially available reagents were purchased from Aldrich, and used without further purification. Grignard reagents were purchased from Aldrich (MeMgBr , EtMgBr , $n\text{-HexMgBr}$, $c\text{-C}_5\text{H}_9\text{MgBr}$) or prepared from the corresponding alkyl bromides and magnesium turnings in Et_2O following standard procedures ($\text{PhCH}_2\text{CH}_2\text{MgBr}$). Grignard reagents were titrated using $s\text{-BuOH}$ and catalytic amounts of 1,10-phenanthroline.

General procedure for the stereoselective Cu-catalyzed synthesis of 1,2-hydroxyalkyl compounds **5**

In a Schlenk tube equipped with septum and stirring bar, $\text{CuBr}\cdot\text{SMe}_2$ (10 μmol , 2.06 mg) and ligand **L1** (12 μmol , 8.24 mg) were dissolved in CH_2Cl_2 (2 mL) and stirred under nitrogen at room temperature for 15 min. The mixture was cooled to -80°C and the

¹ (a) J. A. Marshall, J. D. Trometer, D. G. Cleary, *Tetrahedron*, 1989, **45**, 391; (b) A. R. Ellwood, A. J. P. Mortimer, D. A. Tocher, M. J. Porter, *Synlett*, 2008, 2199.

corresponding Grignard reagent (solution in Et₂O, 0.75 mmol) was added dropwise. Allyl bromide **4** (0.5 mmol, 110 mg) was then added dropwise as a solution in CH₂Cl₂ (0.8 mL) at that temperature over 1 h *via* a syringe pump. Once the addition was complete the resulting mixture was further stirred at -80 °C for 4 h. The reaction was quenched by addition of MeOH (0.5 mL) and the mixture was allowed to reach rt. Then, saturated aqueous NH₄Cl solution (2 mL) was added to the mixture. The organic phase was separated, and the resulting aqueous layer was extracted with Et₂O (3 x 5 mL). The combined organic layers were dried with anhydrous Na₂SO₄, filtered and concentrated *in vacuo*. The crude product was purified by flash chromatography on silica gel using a mixture of pentane:Et₂O as eluent to yield the corresponding 1,2-hydroxyalkyl compound **5**.

Note: GC analysis was carried out on a sample obtained after aqueous extraction with Et₂O, which has been passed through a short plug of silica gel to remove transition metal residues.

(-)-(S)-4-((S)-but-3-en-2-yl)-2,2-dimethyl-1,3-dioxolane (*anti*-5a): Purification by column chromatography (SiO₂, pentane/Et₂O 30:1) afforded *anti*-**5a** (71 mg, 91%) as a colourless oil. $[\alpha]_D^{20} = -2.8$ ($c = 0.5$ in CHCl₃) [lit.² $[\alpha]_D^{20} = -2.5$ ($c = 1.21$ in CHCl₃)]. ¹H NMR (400 MHz, CDCl₃) δ 5.87 – 5.78 (m, 1H), 5.09 – 5.04 (m, 2H), 4.00 – 3.94 (m, 2H), 3.65 – 3.60 (m, 1H), 2.33 (sext., $J = 6.5$ Hz, 1H), 1.40 (s, 3H), 1.35 (s, 3H), 1.00 (dd, $J = 6.8, 1.0$, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 140.2, 115.2, 109.2, 79.5, 67.6, 41.0, 26.8, 25.7, 15.7. HRMS (APCI+, m/z): calculated for C₉H₁₇O₂ [M+H⁺]: 157.1223, found: 157.1219.

² R. W. Hoffmann, W. Helbig, *Chem. Ber.* 1981, **114**, 2802.

(+)-(S)-4-((R)-but-3-en-2-yl)-2,2-dimethyl-1,3-dioxolane (*syn*-5a): Purification by column chromatography (SiO₂, pentane/Et₂O 30:1) afforded *syn*-5a (66 mg, 85%) as a colourless oil. $[\alpha]_D^{20} = +7.6$ (c = 0.65 in CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ 5.74 – 5.65 (m, 1H), 5.08 (d, *J* = 17.4 Hz, 1H), 5.04 (d, *J* = 10.3 Hz, 1H), 3.97 – 3.87 (m, 2H), 3.64 (t, *J* = 7.3 Hz, 1H), 2.32 (sext., *J* = 6.8 Hz, 1H), 1.41 (s, 3H), 1.35 (s, 3H), 1.09 (d, *J* = 6.7, 4H). ¹³C NMR (101 MHz, CDCl₃) δ 139.7, 115.6, 109.2, 79.6, 67.9, 41.9, 26.9, 25.7, 16.6. HRMS (APCI+, *m/z*): calculated for C₉H₁₇O₂ [M+H⁺]: 157.1223, found: 157.1220.

(+)-(S)-2,2-dimethyl-4-((S)-pent-1-en-3-yl)-1,3-dioxolane (*anti*-5b): Purification by column chromatography (SiO₂, pentane/Et₂O 30:1) afforded *anti*-5b (76 mg, 89%) as a colourless oil. $[\alpha]_D^{20} = +6.0$ (c = 1.0 in CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ 5.67 – 5.58 (m, 1H), 5.13 (dd, *J* = 10.3, 0.6 Hz, 1H), 5.05 (dd, *J* = 17.2, 0.6 Hz, 1H), 4.06 – 3.96 (m, 2H), 3.63 (t, *J* = 7.4 Hz, 1H), 2.04 – 1.97 (m, 1H), 1.47 – 1.40 (m, 1H), 1.38 (s, 3H), 1.33 (s, 3H), 1.32 – 1.20 (m, 1H), 0.87 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 138.3, 117.3, 108.9, 78.5, 67.7, 49.3, 26.7, 25.7, 23.9, 11.9. HRMS (ESI+, *m/z*): calculated for C₁₀H₁₉O₂ [M+H⁺]: 171.1370, found: 171.1379.

(+)-(S)-2,2-dimethyl-4-((R)-pent-1-en-3-yl)-1,3-dioxolane (*syn*-5b): Purification by column chromatography (SiO₂, pentane/Et₂O 30:1) afforded *syn*-5b (68 mg, 80%) as a colourless oil. $[\alpha]_D^{20} = +3.0$ (c = 1.0 in CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ 5.54 – 5.42 (m, 1H), 5.12 – 5.00 (m, 2H), 3.95 – 3.90 (m, 2H), 3.61 (t, *J* = 10.2 Hz, 1H), 2.08 – 1.99 (m, 1H), 1.80 – 1.71 (m, 1H), 1.39 (s, 3H), 1.34 (s, 3H), 1.32 – 1.18 (m, 1H), 0.87 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 138.0, 117.5, 109.2, 78.5, 68.3, 50.3,

27.0, 25.9, 24.1, 11.5. HRMS (ESI⁺, m/z): calculated for C₁₀H₁₉O₂ [M+H⁺]: 171.1370, found: 171.1378.

(+)-(S)-2,2-dimethyl-4-((S)-5-phenylpent-1-en-3-yl)-1,3-dioxolane (*anti*-5c):

Purification by column chromatography (SiO₂, pentane/Et₂O 20:1) afforded *anti*-5c (103 mg, 84%) as a colourless oil. $[\alpha]_D^{20} = +25.4$ ($c = 1.0$ in CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ 7.31 – 7.27 (m, 2H), 7.21 – 7.17 (m, 3H), 5.78 – 5.69 (m, 1H), 5.25 (d, $J = 10.2$ Hz, 1H), 5.13 (d, $J = 17.2$ Hz, 1H), 4.08 (dd, $J = 12.5, 6.4$ Hz, 1H), 4.01 – 3.97 (m, 1H), 3.64 (t, $J = 7.6$ Hz, 1H), 2.77 – 2.70 (m, 1H), 2.57 – 2.49 (m, 1H), 2.22 – 2.15 (m, 1H), 1.79 – 1.64 (m, 2H), 1.41 (s, 3H), 1.36 (s, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 142.4, 138.1, 128.7, 128.6, 126.0, 118.0, 109.1, 78.7, 67.6, 47.0, 33.6, 32.9, 26.7, 25.7. HRMS (APCI⁺, m/z): calculated for C₁₆H₂₃O₂ [M+H⁺]: 247.1693, found: 247.1690.

(-)-(S)-2,2-dimethyl-4-((R)-5-phenylpent-1-en-3-yl)-1,3-dioxolane (*syn*-5c):

Purification by column chromatography (SiO₂, pentane/Et₂O 25:1) afforded *syn*-5c (89 mg, 72%) as a colourless oil. $[\alpha]_D^{20} = -1.4$ ($c = 1.0$ in CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ 7.30 – 7.26 (m, 2H), 7.23 – 7.10 (m, 3H), 5.63 – 5.54 (m, 1H), 5.18 – 5.12 (m, 2H), 3.98 – 3.92 (m, 2H), 3.67 – 3.62 (m, 1H), 2.77 – 2.70 (m, 1H), 2.57 – 2.49 (m, 1H), 2.23 – 2.15 (m, 1H), 2.13 – 2.05 (m, 1H), 1.62 – 1.53 (m, 1H), 1.38 (s, 3H), 1.35 (s, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 142.5, 138.1, 128.7, 128.5, 125.9, 118.0, 109.3, 78.6, 68.2, 48.2, 33.3, 32.8, 27.0, 25.9. HRMS (APCI⁺, m/z): calculated for C₁₆H₂₃O₂ [M+H⁺]: 247.1693, found: 247.1691.

(+)-(S)-2,2-dimethyl-4-((S)-non-1-en-3-yl)-1,3-dioxolane (*anti*-5d): Purification by column chromatography (SiO₂, pentane/Et₂O 40:1) afforded ***anti*-5d** (99 mg, 87%) as a colourless oil. $[\alpha]_D^{20} = +33.8$ ($c = 1.0$ in CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ 5.64 (dt, $J = 17.2, 10.2$ Hz, 1H), 5.12 (dd, $J = 10.2, 1.1$ Hz, 1H), 5.04 (dd, $J = 17.2, 1.1$ Hz, 1H), 4.05 – 3.97 (m, 2H), 3.63 (t, $J = 7.2$ Hz, 1H), 2.15 – 2.05 (m, 1H), 1.39 (s, 3H), 1.34 (s, 3H), 1.33 – 1.18 (m, 10H), 0.87 (t, $J = 6.5$ Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 138.6, 117.1, 109.0, 78.8, 67.7, 47.6, 32.0, 31.0, 29.5, 27.3, 26.7, 25.7, 22.8, 14.3. HRMS (APCI+, m/z): calculated for C₁₄H₂₇O₂ [M+H⁺]: 227.2006, found: 227.2005.

(+)-(S)-2,2-dimethyl-4-((R)-non-1-en-3-yl)-1,3-dioxolane (*syn*-5d): Purification by column chromatography (SiO₂, pentane/Et₂O 40:1) afforded ***syn*-5d** (88 mg, 79%) as a colourless oil. $[\alpha]_D^{20} = +2.0$ ($c = 1.0$ in CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ 5.54 – 5.45 (m, 1H), 5.09 – 5.02 (m, 2H), 3.94 – 3.88 (m, 2H), 3.65 – 3.60 (m, 1H), 2.16 – 2.07 (m, 1H), 1.72 – 1.64 (m, 1H), 1.40 (s, 3H), 1.35 (s, 3H), 1.33 – 1.18 (m, 9H), 0.87 (t, $J = 6.8$, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 138.4, 117.3, 109.2, 78.7, 68.2, 48.7, 32.0, 31.2, 29.5, 27.1, 25.9, 22.9, 14.3. HRMS (APCI+, m/z): calculated for C₁₄H₂₇O₂ [M+H⁺]: 227.2006, found: 227.2005.

(+)-(S)-4-((R)-1-cyclopentylallyl)-2,2-dimethyl-1,3-dioxolane (*anti*-5e): Purification by column chromatography (SiO₂, pentane/Et₂O 30:1) afforded *anti*-5e (84 mg, 80%) as a colourless oil. $[\alpha]_D^{20} = +13.8$ ($c = 1.0$ in CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ 5.79 – 5.70 (m, 1H), 5.12 (d, $J = 10.3$ Hz, 1H), 5.00 (d, $J = 17.2$ Hz, 1H), 4.23 – 4.17 (m, 1H), 3.97 (t, $J = 7.0$ Hz, 1H), 3.63 (t, $J = 7.7$ Hz, 1H), 1.93 – 1.79 (m, 3H), 1.66 – 1.48 (m, 4H), 1.37 (s, 3H), 1.34 (s, 3H), 1.23 – 1.07 (m, 2H). ¹³C NMR (101 MHz, CDCl₃) δ 137.2, 117.4, 108.8, 77.4, 67.9, 52.3, 41.5, 30.9, 30.8, 26.6, 25.7, 25.2, 25.1. HRMS (ESI+, m/z): calculated for C₁₃H₂₃O₂ [M+H⁺]: 211.1693, found: 211.1690.

General procedure for the synthesis of compounds 7 and α,β -unsaturated δ -lactones 8.

To a solution of **5** (1 mmol) in water (2 mL) at room temperature was added AcOH (5 mL). The solution was stirred at this temperature during 16 h. After some coevaporations with toluene (4 x 10 mL), the residue was dissolved in DMF (2 mL) and imidazole (75 mg, 1.1 mmol), DMAP (1 mg, 0.08 mmol) and *tert*-butyl(chloro)diphenylsilane (0.29 mL, 1.1 mmol) were added at 0 °C. The mixture was warmed to room temperature and was stirred during 16 h. Then it was poured into water (5 mL) and extracted with CH₂Cl₂ (3 x 5 mL). The combined organic layers were washed with water and brine and dried

over Na₂SO₄. The resulting product **S1** was used in the next step without further purification.

To a solution of **S1** (1 mmol) and DIPEA (0.34 mL, 2 mmol) in CH₂Cl₂ (5 mL), acryloyl chloride (0.13 mL, 1.5 mmol) was added at 0 °C. The mixture was stirred at this temperature for 2 h. Then it was quenched with saturated aqueous solution of NaHCO₃ (2 mL), extracted with CH₂Cl₂ (3 x 5 mL) and dried over Na₂SO₄. The product was purified by flash chromatography on silica gel using a mixture of Pentane:Et₂O as eluent to yield the corresponding compound **7**.

(-)-(2*S*,3*R*)-1-((*tert*-butyldiphenylsilyl)oxy)-3-ethylpent-4-en-2-yl acrylate (7a**):**

Purification by column chromatography (SiO₂, pentane/Et₂O 30:1) afforded **7a** (337 mg, 80%, over 3 steps) as a colourless oil. $[\alpha]_D^{20} = -25.8$ ($c = 1.0$ in CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ 7.67 – 7.64 (m, 4H), 7.45 – 7.34 (m, 6H), 6.43 (dd, $J = 17.3, 1.5$ Hz, 1H), 6.17 (dd, $J = 17.3, 10.4$ Hz, 1H), 5.85 (dd, $J = 10.4, 1.5$ Hz, 1H), 5.55 – 5.46 (m, 1H), 5.10 – 5.00 (m, 3H), 3.80 – 3.72 (m, 2H), 2.40 (ddd, $J = 18.3, 9.4, 3.5$ Hz, 1H), 1.59 – 1.49 (m, 1H), 1.28 – 1.17 (m, 1H), 1.03 (s, 9H), 0.84 (t, $J = 7.4$ Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 165.8, 137.6, 135.6, 135.5, 133.4, 133.4, 130.5, 129.6, 129.6, 128.8, 127.6, 117.5, 76.4, 63.6, 46.6, 26.7, 22.9, 19.2, 11.4. HRMS (ESI+, m/z): calculated for C₂₆H₃₄O₃SiNa [M+Na⁺]: 445.2169, found: 445.2178.

(+)-(2*S*,3*S*)-1-((*tert*-butyldiphenylsilyl)oxy)-3-vinylnonan-2-yl acrylate (7b**):**

Purification by column chromatography (SiO₂, pentane/Et₂O 40:1) afforded **7b** (85%, over 3 steps) as a colourless oil. $[\alpha]_D^{20} = +9.0$ ($c = 0.98$ in CHCl₃). ¹H NMR (400 MHz,

CDCl₃) δ 7.67 – 7.64 (m, 4H), 7.45 – 7.34 (m, 6H), 6.38 (dd, J = 17.3, 1.5 Hz, 1H), 6.11 (dd, J = 17.3, 10.4 Hz, 1H), 5.81 (dd, J = 10.4, 1.5 Hz, 1H), 5.58 (dt, J = 17.1, 9.8 Hz, 1H), 5.12 (q, J = 5.1 Hz, 1H), 5.06 – 4.98 (m, 2H), 3.75 (dd, J = 10.8, 6.3 Hz, 1H), 3.68 (dd, J = 10.8, 4.8 Hz, 1H), 2.47 – 2.40 (m, 1H), 1.45 – 1.38 (m, 1H), 1.33 – 1.18 (m, 9H), 1.02 (s, 9H), 0.87 (t, J = 6.7 Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 165.6, 137.8, 135.6, 135.53, 133.4, 133.4, 130.4, 129.6, 129.6, 128.8, 127.6, 127.6, 117.2, 76.0, 63.5, 44.9, 31.7, 30.7, 29.1, 27.0, 26.7, 22.6, 19.2, 14.1. HRMS (ESI+, m/z): calculated for C₃₀H₄₃O₃Si [M+H⁺]: 479.2976, found: 479.2969.

General procedure for the RCM

Grubbs 2nd generation catalyst (4.3 mg, 0.005 mmol) was added to a degassed solution of **7** (0.1 mmol) in CH₂Cl₂ (10 mL), and the mixture was refluxed during 14 h. The solvent was removed and the residue was purified by flash chromatography on silica gel using a 5:1 mixture of Pentane:Et₂O as eluent to yield the corresponding compound **8**.

(-)-(5R,6S)-6-(((tert-butyldiphenylsilyl)oxy)methyl)-5-ethyl-5,6-dihydro-2H-pyran-2-one (8a): Purification by column chromatography (SiO₂, pentane/Et₂O 5:1) afforded **8a** (39 mg, 97%) as a colourless oil. $[\alpha]_D^{20}$ = - 136.2 (c = 1.0 in CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ 7.68 – 7.63 (m, 4H), 7.47 – 7.37 (m, 6H), 7.04 (dd, J = 9.8, 6.0 Hz, 1H), 6.03 (dd, J = 9.8, 0.9 Hz, 1H), 4.56 (ddd, J = 8.0, 5.9, 3.8 Hz, 1H), 3.92 (dd, J = 10.6, 5.9 Hz, 1H), 3.81 (dd, J = 10.6, 8.0 Hz, 1H), 2.56 – 2.50 (m, 1H), 1.69 – 1.59 (m, 1H), 1.45 (ddq, J = 14.7, 9.8, 7.5 Hz, 1H), 1.07 (s, 9H), 0.94 (t, J = 7.5 Hz, 1H). ¹³C NMR (101 MHz, CDCl₃) δ 163.8, 150.3, 135.5, 132.9, 132.8, 129.9, 127.8, 120.9, 79.5, 62.1, 36.1, 26.8, 20.5, 19.2, 11.0. HRMS (ESI+, m/z): calculated for C₂₄H₃₀O₃SiNa [M+Na⁺]: 417.1856, found: 417.1863.

(+)-(5S,6S)-6-(((*tert*-butyldiphenylsilyl)oxy)methyl)-5-hexyl-5,6-dihydro-2H-pyran-2-one (8b): Purification by column chromatography (SiO₂, pentane/Et₂O 5:1) afforded **8b** (38 mg, 84%) as a colourless oil. $[\alpha]_D^{20} = + 84.2$ ($c = 0.92$ in CHCl₃). ¹H NMR (400 MHz, CDCl₃) δ 7.67 – 7.64 (m, 4H), 7.46 – 7.37 (m, 6H), 6.73 (dd, $J = 9.9, 4.0$ Hz, 1H), 5.93 (dd, $J = 9.9, 1.8$ Hz, 1H), 4.32 – 4.28 (m, 1H), 3.86 (dd, $J = 11.0, 4.9$ Hz, 1H), 3.83 (dd, $J = 11.0, 4.2$ Hz, 1H), 2.77– 2.70 (m, 1H), 1.58 – 1.50 (m, 1H), 1.46 – 1.26 (m, 9H), 1.07 (s, 9H), 0.90 (t, $J = 6.9$ Hz, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 163.3, 149.0, 135.6, 135.5, 132.9, 132.7, 129.9, 129.9, 127.8, 127.8, 119.9, 81.4, 63.4, 34.0, 31.6, 31.2, 29.2, 26.8, 26.3, 22.6, 19.2, 14.0. HRMS (ESI+, m/z): calculated for C₂₈H₃₈O₃SiNa [M+Na⁺]: 473.2482, found: 473.2459.

syn-5a

Proposed mechanism for the copper-catalyzed AAA of allyl bromide **4** with Grignard reagents

In analogy with the proposed mechanism by Goering and co-workers,³ the catalytic cycle depicted above can be proposed for the Cu-taniaphos catalyzed AAA of allyl bromide **4**. In this mechanism the precatalyst **A** and the Grignard reagent form the active catalyst **B**.⁴ Subsequently, the interaction with allyl bromide **4** forms the Cu^{I} π -complex **C**. Oxidative addition and allylic rearrangement from **C** gives the Cu^{III} σ -complex **D** which leads to the $\text{S}_{\text{N}}2'$ product via reductive elimination.

The formation of the competing $\text{S}_{\text{N}}2$ product can be explained through an isomerization of the Cu^{III} σ -complex **D** into the Cu^{III} π -complex **E**. It has been shown that this conversion of the Cu^{III} σ -complex into the Cu^{III} π -complex is faster at higher

³ C. C. Tseng, S. D. Paisley, H. L. Goering, *J. Org. Chem.* 1986, **51**, 2884.

⁴ S. Harutyunyan, F. López, W. R. Brown, A. Correa, D. Peña, R. Badorrey, A. Meetsma, A. J. Minnaard, B. L. Feringa, *J. Am. Chem. Soc.* 2006, **128**, 9103.

temperatures.⁵ Intermediate **E** would evolve to the S_N2 product via reductive elimination (initially Goering and co-workers^{ref1} proposed the formation of a **F** type Cu^{III} σ-complex before the reductive elimination can take place). Recent calculations by Nakamura and co-workers have shown that reductive elimination will proceed directly from the π-complex.⁶

GC traces for the temperature-depending experiments for the AAA of allyl bromide **4 with MeMgBr catalyzed by Cu-TaniaPhos (Table 1).**

Product ratio was determined by GC analysis of the reaction crude, (GC, HP6890: MS HP5973) with an HP5 column, initial temp. 50°C then 10 °C/min to 270 °C (hold for 3 min, final temp).

Retention times (min): 4.2 (*syn*-**5a**), 4.4 (*anti*-**5a**); and 5.3 (**6**).

⁵ E. R. Bartholomew, S. H. Bertz, S. Cope, M. Murphy, C. A. Ogle, *J. Am. Chem. Soc.* 2008, **130**, 11244.

⁶ M. Yamanaka, S. Kato, E. Nakamura, *J. A. Chem. Soc.* 2004, **126**, 6287.

- L = (R,R)-(+)-taniaphos, T = -75 °C (Table 1, entry 1):

- L = (R,R)-(+)-taniaphos, T = -80 °C (Table 1, entry 11):

- L = (S,S)-(-)-taniaphos, T = -75 °C (Table 1, entry 2):

- L = (S,S)-(-)-taniaphos, T = -80 °C (Table 1, entry 12):

- L = (S,S)-(-)-taniaphos, T = -50 °C: