University of Groningen

Monotonicity and phase diagram for multirange percolation on oriented trees

de Lima, Bernardo N. B.; Rolla, Leonardo T.; Valesin, Daniel

Published in:
Random structures & algorithms

DOI:
[10.1002/rsa.20805](https://doi.org/10.1002/rsa.20805)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2019

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

de Lima, B. N. B., Rolla, L. T., & Valesin, D. (2019). Monotonicity and phase diagram for multirange percolation on oriented trees. *Random structures & algorithms*, 55(1), 160-172.
<https://doi.org/10.1002/rsa.20805>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

RESEARCH ARTICLE

WILEY

Monotonicity and phase diagram for multirange percolation on oriented trees

Bernardo N. B. de Lima¹ | Leonardo T. Rolla^{2,3} | Daniel Valesin⁴

¹Department of Mathematics, UFMG-Federal University of Minas Gerais, Belo Horizonte, Brazil

²Argentina National Research Council at the University of Buenos Aires, Buenos Aires,

³NYU-ECNU Institute of Mathematical Sciences at NYU Shanghai, Shanghai, China

⁴Bernoulli Institute for Mathematics, Computer Science and Artificial Intelligence, University of Groningen, Groningen, The Netherlands

Correspondence

Bernardo N. B. de Lima, UFMG-Federal University of Minas Gerais, Belo Horizonte, Brazil.

Email: bnblima@mat.ufmg.br

Funding information

This research was supported by the CNPq; 309468/2014-0. FAPEMIG (Programa Pesquisador Mineiro). PIP; 11220130100521CO. PICT; 2015-3154; 2013-2137; 2012-2744. Conicet; 45955. MinCyT; BR-13/14.

Abstract

We consider Bernoulli bond percolation on oriented regular trees, where besides the usual short bonds, all bonds of a certain length are added. Independently, short bonds are open with probability p and long bonds are open with probability q . We study properties of the critical curve which delimits the set of pairs (p, q) for which there are almost surely no infinite paths. We also show that this curve decreases with respect to the length of the long bonds.

KEYWORDS

critical curve, long range percolation, monotonicity of connectivity

1 | INTRODUCTION

Consider the graph having $\mathbb{Z}^d$ as vertex set and all edges of the form $\{x, x \pm e_i\}$ and $\{x, x \pm k \cdot e_i\}$ for some $k \geq 2$. It was shown in [9] that the critical probability for Bernoulli bond percolation on this graph converges to that of $\mathbb{Z}^{2d}$ as $k \rightarrow \infty$. This result, later generalized in [15], is a particular instance of Schramm's conjecture [7] that the percolation threshold for transitive graphs is a local property. The convergence proved in [9] is conjectured to be monotone, that is, the percolation threshold for the above graph should be decreasing in the length k of long edges.¹

¹In support of this conjecture, simulations [5] confirm that increasing k decreases the critical parameter, and the proof of [9, Lemma 2] shows that replacing k by a multiple of k does not increase it.

FIGURE 1 Phase space and critical curve separating the percolative region P_k from the non-percolative region N_k . (a) The curve stays between the three dotted lines. In the gray region, infinite paths necessarily use both short and long edges. (b) Critical curves for different ranges k meet at one point

Monotonicity questions are often intriguing for being extremely simple to ask and hard to answer. A good example [18] is the following: for Bernoulli bond percolation on the usual graph $\mathbb{Z}^d$, prove that the probability of the origin being connected to $(n, 0, \dots, 0)$ monotone in n . This problem is still open, except when the parameter is close to 0 or 1 [10]. For oriented percolation on $\mathbb{Z}_+^2$, the probability of the origin being connected to $(m - n, m + n)$ is decreasing in $n \in \{0, \dots, m\}$ for fixed m ; this may be obvious but the proof is not straightforward [4]. In the same spirit, for unoriented percolation on $\mathbb{Z}_+^2$, if the parameter is smaller for horizontal edges than for vertical ones, the above probability should be larger than the probability of the origin being connected to $(m + n, m - n)$. This has only been proved under the assumption that the ratio between horizontal and vertical parameters is small enough [8].

For first-passage percolation, it was conjectured [12] that the expected minimum travel time from $(0, 0)$ to $(n, 0)$ along paths contained in the strip $\{(x, y) : 0 \leq x \leq n\}$ is nondecreasing in n . This question is still open, with a number of partial results [1, 3, 11, 13]. In the negative direction, for first-passage percolation on $\mathbb{Z}_+ \times \mathbb{Z}$, there is a counter-example [17] where the expected passage time from the origin to $(2, 0)$ is less than the expected passage time from the origin to $(1, 0)$. Another context where strict monotonicity is expected to happen is in the case of essential enhancements as introduced in [2], see also [6].

In this paper we consider percolation on $\mathbb{T}_{d,k}$, the graph given by the oriented rooted d -ary tree ($d \geq 2$) with a root at the top, bearing the usual “short” downward edges plus the addition of all downward edges of length k , called “long” edges. This is an oriented version of Trofimov’s grandfather graph for $k = 2$, or the great ^{k} -grandfather graph for larger k . We let short and long edges be open independently with probability p and q , respectively. The phase space $[0, 1]^2$ is decomposed in two regions: a set P_k of pairs (p, q) for which a.s. there are infinite open paths, and a set N_k of pairs for which a.s. there are none, see Figure 1a.

For $p > \frac{1}{d}$ there are a.s. infinite open paths of short edges, and for $q > \frac{1}{d^k}$ there are a.s. infinite open paths of long edges. For $dp + d^k q \leq 1$, a simple comparison with a branching process (to be given in section 3) shows that a.s. there are no infinite open paths. By monotonicity of the process with respect to p and q , there is a *critical curve* γ_k joining the points $(\frac{1}{d}, 0)$ and $(0, \frac{1}{d^k})$ which separates N_k and P_k , as depicted in Figure 1a. Define

$$q_c(p, k) = \inf\{q : (p, q) \in P_k\} \quad \text{and} \quad p_c(q, k) = \inf\{p : (p, q) \in P_k\}. \quad (1)$$

Let k be fixed. We show that q_c is continuous and strictly decreasing in p (equivalent formulations are that p_c is strictly decreasing and continuous in q , that both p_c and q_c are continuous, or that γ contains neither vertical nor horizontal segments). In particular, γ_k is described by $q = q_c(p, k)$ as well as by $p = p_c(q, k)$, and there is a non-trivial subregion of $\mathcal{P}_k$ at which infinite open paths necessarily use both long and short edges, see Figure 1a. A similar description is given in [14] for percolation with a defect plane.

We also show that γ_{k+1} stays strictly below γ_k for $p < d^{-1}$, and they meet only at the critical point $(d^{-1}, 0)$. This means that $q_c(p, k)$ is strictly decreasing in k for as long as it is positive, and analogously for $p_c(q, k)$, see Figure 1b.

In section 2, we present the model and the above statements more formally.

In section 3, we prove that $q_c(k, p)$ is continuous and decreasing in p . In the proof, we tile $\mathbb{T}_{d,k}$ by layers and consider a construction of the process where the state of tiles are sampled independently. We then couple configurations with different values of p and q so that some advantage in q compensates for small decreases in p and vice-versa. Each comparison is done by finding one particular tile that makes no useful connections without extra open edges and at the same time makes all possible connections with their help. We learned this idea from [16].

In section 4, we show that $q_c(p, k+1) < q_c(p, k)$ for $p < d^{-1}$. Together with the results of section 3, this inequality completes the previous description illustrated by Figure 1b. The proof involves a joint exploration of a percolation “cluster” in $\mathbb{T}_{d,k}$ and a percolation cluster in $\mathbb{T}_{d,k+1}$. The joint exploration is an algorithm in which parts of both clusters are revealed simultaneously using the same random variables. After each step of the algorithm is concluded, there is an injective function from the revealed portion of the cluster in $\mathbb{T}_{d,k}$ to the one in $\mathbb{T}_{d,k+1}$. When trying to ensure this, one might run into collisions, that is, situations where an edge that could potentially grow the cluster in $\mathbb{T}_{d,k}$ has as a counterpart an edge which does not grow the cluster in $\mathbb{T}_{d,k+1}$. The challenge is thus to design the algorithm so that collisions do not occur. We succeed in doing so by introducing a recursive procedure which alternately reveals clusters of short edges and then groups long edges, in a way that allows the comparison between the k and $k+1$ scenarios. This gives $q_c(p, k+1) \leq q_c(p, k)$. Strict inequality is obtained by extending the idea mentioned in the previous paragraph to a dynamic, hybrid construction. When revealing the state of a whole batch of long edges at once we can use the increase in k to compensate for a small decrease in q .

As a final remark, there seems to be no obvious way to adapt the argument just described to the graph with the vertices in $\mathbb{Z}^d$ and edges of the form $\{x, x \pm e_i\}$ and $\{x, x \pm k \cdot e_i\}$. A similar joint exploration would lead to collisions as illustrated in Figure 2. Proving the inequality $p_c(k+1) \leq p_c(k)$ mentioned in the previous footnote remains open, let alone strict inequality.

2 | DEFINITIONS AND RESULTS

Let $d \in \{2, 3, 4, \dots\}$ be fixed. We denote $[d] = \{1, \dots, d\}$, and we will make frequent use of the set

$$[d]_\star = \bigcup_{0 \leq n < \infty} [d]^n;$$

the set $[d]^0$ is understood to consist of a single point o . Points of $[d]_\star \setminus \{o\}$ are represented as sequences $u = (u_1, \dots, u_n)$. In case $u = (u_1, \dots, u_m)$ and $v = (v_1, \dots, v_n)$, we define the concatenation $u \cdot v = (u_1, \dots, u_m, v_1, \dots, v_n)$.

FIGURE 2 Situation where the natural coupling of explorations which maps short edges to short edges and long edges to long edges leads to a “collision” in the graphs given by adding edges of length 3 and 4 to $\mathbb{Z}$. Dashed lines represent closed edges and full lines represent open edges. The bold edge being open increases the cluster of the first graph by one vertex, but has no effect on the cluster of the second graph

Given $k \in \mathbb{N} = \{1, 2, 3, \dots\}$, we define the oriented graph $\mathbb{T}_{d,k}$ as the graph with vertex set $\mathbb{V}_{d,k} = [d]_\star$ and edge set $\mathbb{E}_{d,k} = \mathbb{E}_{d,k}^\delta \cup \mathbb{E}_{d,k}^\ell$, where

$$\mathbb{E}_{d,k}^\delta = \{\langle u, u \cdot a \rangle : u \in \mathbb{V}_{d,k}, a \in [d]\},$$

$$\mathbb{E}_{d,k}^\ell = \{\langle u, u \cdot r \rangle : u \in \mathbb{V}_{d,k}, r \in [d]^k\}.$$

These will be referred to as the sets of short and long edges of $\mathbb{T}_{d,k}$.

As the above notation suggests, we will normally use the letters a, b for elements of $[d]$, the letters r, s for elements of $[d]^k$ and the letters u, v, w, x for general vertices of $\mathbb{T}_{d,k}$.

Consider the process in which, independently, short edges are open with probability p and long edges are open with probability q . Let $\mathbb{P}_{p,q}$ denote the corresponding probability measure.

We define the event $u \rightsquigarrow v$ that there exist $u^0, u^1, \dots, u^{n-1}, u^n$ such that $u^0 = u$, $u^n = v$ and the edge $\langle u^j, u^{j+1} \rangle$ is open for all $j < n$. The event $u \rightsquigarrow \infty$ means that $u \rightsquigarrow v$ for infinitely many v . Let $\mathcal{P}_k = \{(p, q) : \mathbb{P}_{p,q}(o \rightsquigarrow \infty) > 0\}$, $\mathcal{N}_k = [0, 1]^2 \setminus \mathcal{P}_k$, and let $p_c(q, k)$ and $q_c(p, k)$ be given by (1).

We prove the following monotonicity property.

Theorem 2.1 *The inequality $q_c(p, k+1) < q_c(p, k)$ holds unless $q_c(p, k) = 0$.*

This says that γ_{k+1} stays under γ_k , and they can only intersect each other at the boundary $\{pq = 0\}$, except maybe where one of them contains a vertical segment. The next result rules out the latter possibility, thus completing the picture provided in Figure 1b.

Theorem 2.2 *For each fixed $k \in \mathbb{N}$, the function $p \mapsto q_c(p, k)$ is continuous on $[0, 1]$ and strictly decreasing on $[0, d^{-1}]$.*

We observe that, as a consequence of the above results, defining

$$p_c(k) = \inf\{p : (p, p) \in \mathcal{P}_k\},$$

we have $p_c(k+1) < p_c(k)$, as the diagonal $\{(p, p) : 0 \leq p \leq 1\}$ intersects the critical curves γ_k at distinct points for different values of k . However, for $k \geq 2$ this conclusion can be drawn from the simpler observation that the curves γ_k are delimited by the dotted lines in Figure 1b.

The next result says that there is no percolation along the critical curves γ_k .

Theorem 2.3 For (p, q) on the critical curve γ_k , $\mathbb{P}_{p,q}(o \rightsquigarrow \infty) = 0$.

Proof It is enough to prove that $\mathcal{P}_k$ is an open set in $[0, 1]^2$. Define

$$N_n = \# \{u \in [d]^{kn} : \text{there exists } v \in \cup_{i=0}^{k-1} [d]^i \text{ with } o \rightsquigarrow u \cdot v\}, \quad n \in \mathbb{N}.$$

We claim that $N_n \rightarrow \infty$ a.s. on the event $o \rightsquigarrow \infty$. Indeed, assuming $p < 1$ and $q < 1$, for each $j \in \mathbb{N}$ we have

$$\mathbb{P}_{p,q}(N_m = 0 \text{ for all } m > n \mid N_1, \dots, N_n) > \sigma_j$$

on the event that $N_n \leq j$, where σ_j is a positive constant depending on j and also on p, q, d, k , but not on n . This shows that $\mathbb{P}_{p,q}(N_n = j \text{ i.o.}) = 0$ thus a.s. either $N_n \rightarrow 0$ or $N_n \rightarrow \infty$. The case $p = 1$ or $q = 1$ being trivial, the claim is proved.

Suppose $\theta_{p,q} := \mathbb{P}_{p,q}(o \rightsquigarrow \infty) > 0$ and let $\zeta < \theta_{p,q}$. By the previous claim, there exists n^* such that $\mathbb{P}_{p,q}(N_{n^*} > \frac{2k^2}{\zeta}) > \zeta$. Now observe that this probability is continuous in (p, q) , thus for (p', q') close enough to (p, q) it is still larger than ζ . From this observation, using the definition of N_n and reverse union bound, there is $\ell \in \{kn^*, \dots, kn^* + k - 1\}$ such that, with probability larger than $\frac{\zeta}{k}$, there are at least $\frac{2k}{\zeta}$ sites $u \in [d]^\ell$ such that $o \rightsquigarrow u$.

Therefore, the process $(N_{\ell i})_{i \in \mathbb{N}}$ dominates a supercritical branching process with offspring assuming values on $\{0, \lceil \frac{2k}{\zeta} \rceil\}$ and mean larger than 2. This implies that $\mathbb{P}_{p',q'}(o \rightsquigarrow \infty) > 0$, proving that $(p', q') \in \mathcal{P}_k$. ■

3 | LONG AND SHORT EDGE COMPENSATION

The goal of this section is to prove Theorem 2.2. We will need the following elementary fact.

Lemma 3.1 Let P_α denote probability measures on a given finite space S , parametrized by $\alpha \in [0, 1]$, and such that $P_\alpha(x)$ is continuous in α for every $x \in S$. Let κ and y be such that $P_\kappa(y) > 0$. Then for any α, β close enough to κ , there exists a coupling (X, Y) such that $X \sim P_\alpha$, $Y \sim P_\beta$ and such that, almost surely, $X = Y$ unless $X = y$ or $Y = y$.

Proof Sample the pair (X, Y) as

$$(X, Y) = \begin{cases} (z, z) & \text{w.p. } P_\alpha(z) \wedge P_\beta(z), \\ (y, z) & \text{w.p. } [P_\beta(z) - P_\alpha(z)]^+, \\ (z, y) & \text{w.p. } [P_\alpha(z) - P_\beta(z)]^+, \end{cases}$$

for $z \neq y$, and

$$(X, Y) = (y, y) \text{ w.p. } 1 - \sum_{z \neq y} P_\alpha(z) \vee P_\beta(z).$$

The last term is positive for α and β are close to κ because it is positive when $\alpha = \beta = \kappa$. This sampling only include pairs for which $X = Y$ unless $X = y$ or $Y = y$. From the first equation we have $\mathbb{P}(X = z) = P_\alpha(z)$ for all $z \neq y$, which all together imply $\mathbb{P}(X = y) = P_\alpha(y)$, and similarly for Y . ■

We define the progeny of a vertex $u \in \mathbb{V}_{d,k}$ as the set

$$\text{prog}(u) = \{u \cdot v \in \mathbb{V}_{d,k} : v \in [d]_\star\},$$

that is, it is the subtree started at u . The progeny of an edge is defined as the progeny of its endpoint, that is, if $e = \langle u, v \rangle$, then $\text{prog}(e) = \text{prog}(v)$.

We now turn to the proof of Theorem 2.2. Recall that k is fixed, $q_c(0) = d^{-k}$ and $q_c(p) = 0$ for $p > d^{-1}$. Let $\mathcal{C}_{p,q,k}$ denote the percolation cluster of the root in $\mathbb{T}_{d,k}$ under the measure $\mathbb{P}_{p,q}$. (We use the word “cluster” to denote the set of sites which can be reached from the root, so unlike unoriented percolation it does not define an equivalence class.) We observe that, under this measure, the expected number of open edges having o as an extremity is equal to $dp + d^k q$. If such expectation is less than one, we can embed $\mathcal{C}_{p,q,k}$ in a subcritical branching process to conclude that $P_{p,q}(o \rightsquigarrow \infty) = 0$. Therefore, $q_c(p, k) \geq d^{-k} - d^{-k+1}p$. This implies that $q_c(p, k) > 0$ for $p < d^{-1}$. Since $q_c(p, k) \leq q_c(0, k) = d^{-k}$, we also conclude that $p \mapsto q_c(p, k)$ is continuous at $p = 0$.

The proof of Theorem 2.2 will thus be complete once we establish the following two facts:

$$\begin{aligned} &\text{for all } p_0, q, q' \in (0, 1) \text{ with } q < q', \text{ there exist } p, p' \text{ with } p' < p_0 < p \\ &\text{such that } \mathbb{P}_{p',q'}(o \rightsquigarrow \infty) \geq \mathbb{P}_{p,q}(o \rightsquigarrow \infty); \end{aligned} \quad (2)$$

$$\begin{aligned} &\text{for all } q_0, p, p' \in (0, 1) \text{ with } p < p', \text{ there exist } q, q' \text{ with } q' < q_0 < q \\ &\text{such that } \mathbb{P}_{p',q'}(o \rightsquigarrow \infty) \geq \mathbb{P}_{p,q}(o \rightsquigarrow \infty). \end{aligned} \quad (3)$$

Indeed, condition (2) rules out jump discontinuities in the curve $q = q_c(p, k)$ for $p > 0$, and condition (3) rules out horizontal segments in this curve for $p < d^{-1}$.

We start the proof of (2) by introducing some notation. We let $\tilde{\mathbb{E}}_{d,k} = \tilde{\mathbb{E}}_{d,k}^{\mathcal{J}} \cup \tilde{\mathbb{E}}_{d,k}^{\mathcal{L}}$, where

$$\tilde{\mathbb{E}}_{d,k}^{\mathcal{J}} = \left\{ e = \langle u, v \rangle \in \mathbb{E}_{d,k}^{\mathcal{J}} : u \in \bigcup_{n=0}^{2k-1} [d]^n \right\},$$

$$\tilde{\mathbb{E}}_{d,k}^{\mathcal{L}} = \left\{ e = \langle u, v \rangle \in \mathbb{E}_{d,k}^{\mathcal{L}} : u \in \bigcup_{n=0}^{2k-1} [d]^n \right\}.$$

Configurations in $\bar{\Omega} = \bar{\Omega}_{\mathcal{J}} \times \bar{\Omega}_{\mathcal{L}} = \{0, 1\}^{\tilde{\mathbb{E}}_{d,k}^{\mathcal{J}} \cup \tilde{\mathbb{E}}_{d,k}^{\mathcal{L}}}$ are written as $\bar{\omega} = (\bar{\omega}_{\mathcal{J}}, \bar{\omega}_{\mathcal{L}})$.

Given $A \subseteq \bigcup_{n=0}^{k-1} [d]^n$ and $\bar{\omega} = (\bar{\omega}_{\mathcal{J}}, \bar{\omega}_{\mathcal{L}})$, we define

$$J_{\bar{\omega}}(A) = \bigcup_{u \in [d]^{2k}} \left\{ v \in \text{prog}(u) : \exists u^0, \dots, u^n \in \mathbb{V}_{d,k} \text{ so that } u^0 \in A, \right. \\ \left. u^n = v \text{ and } \langle u^i, u^{i+1} \rangle \in \tilde{\mathbb{E}}_{d,k}, \bar{\omega}(\langle u^i, u^{i+1} \rangle) = 1 \forall i \right\}. \quad (4)$$

That is, $J_{\bar{\omega}}(A)$ is the set of vertices in $\bigcup_{u \in [d]^{2k}} \text{prog}(u)$ that are reachable by paths started from A and consisting only of open edges of $\tilde{\mathbb{E}}_{d,k}$. Note that in such a path, all edges have both extremities in $\bigcup_{n=0}^{2k-1} [d]^n$ except for the last one, which has only one extremity in $\bigcup_{n=0}^{2k-1} [d]^n$. In particular, $J_{\bar{\omega}}(A) \subseteq \bigcup_{n=2k}^{3k-1} [d]^n$.

Now, define the deterministic configurations $\bar{\omega}_{\mathcal{J}}^* \in \bar{\Omega}_{\mathcal{J}}$ and $\bar{\omega}_{\mathcal{L},1}^*, \bar{\omega}_{\mathcal{L},2}^* \in \bar{\Omega}_{\mathcal{L}}$ by setting

$$\bar{\omega}_{\mathcal{L},1}^* \equiv 0, \quad \bar{\omega}_{\mathcal{L},2}^* \equiv 1 \quad \text{and} \quad \bar{\omega}_{\mathcal{J}}^*(\langle u, v \rangle) = 1 \text{ if and only if } u \notin [d]^{2k-1}.$$

Let $0 < p_0 < 1$ and $0 < q < q' < 1$. By Lemma 3.1, if p and p' with $p' < p_0 < p$ are chosen sufficiently close to p_0 , then there exists a coupling of configurations

$$X = (X_\delta, X_{\ell,1}, X_{\ell,2}) \quad \text{and} \quad Y = (Y_\delta, Y_{\ell,1}, Y_{\ell,2})$$

in $\bar{\Omega}_\delta \times \bar{\Omega}_\ell \times \bar{\Omega}_\ell$ so that the following holds:

- the values of $X_\delta, X_{\ell,1}$ and $X_{\ell,2}$ in all edges are independent;
- $X_\delta, X_{\ell,1}$ and $X_{\ell,2}$ assign each edge to be open with respective probabilities p, q and $\frac{q'-q}{1-q}$;
- the values of $Y_\delta, Y_{\ell,1}$ and $Y_{\ell,2}$ in all edges are independent;
- $Y_\delta, Y_{\ell,1}$ and $Y_{\ell,2}$ assign each edge to be open with respective probabilities p', q and $\frac{q'-q}{1-q}$;
- the following event has probability one:

$$\{X = Y\} \cup \{X = (\bar{\omega}_\delta^*, \bar{\omega}_{\ell,1}^*, \bar{\omega}_{\ell,2}^*)\} \cup \{Y = (\bar{\omega}_\delta^*, \bar{\omega}_{\ell,1}^*, \bar{\omega}_{\ell,2}^*)\}. \quad (5)$$

Now take $\bar{\omega}_\delta = X_\delta, \bar{\omega}_\ell = X_{\ell,1}, \bar{\omega}'_\delta = Y_\delta, \bar{\omega}'_\ell = Y_{\ell,1} \vee Y_{\ell,2}$.

The main observation is that each of the three events in (5) implies that, for every $A \subseteq \cup_{n=0}^{k-1} [d]^n$,

$$J_{\bar{\omega}}(A) \subseteq J_{\bar{\omega}'}(A). \quad (6)$$

Indeed, on the first event we have $\bar{\omega}' \geq \bar{\omega}$, on the second event we have $J_{\bar{\omega}}(A) = \emptyset$, and on the third event $J_{\bar{\omega}'}(A)$ contains the set of sites $y \in \cup_{n=2k}^{3k-1} [d]^n$ that are in $\text{prog}(x)$ for some $x \in A$, which always contains $J_{\bar{\omega}}(A)$.

Finally, with this coupling at hand, we can sample configurations $\omega, \omega' \in \{0, 1\}^{\mathbb{E}_{d,k}}$ such that the restrictions of ω and ω' to sets of the form

$$\{ \langle u \cdot v, w \rangle \in \mathbb{E}_{d,k} : v \in \cup_{n=0}^{2k-1} [d]^n \}$$

with $u \in \cup_{m \in 2\mathbb{N}} [d]^{mk}$ are independent and sampled from the (appropriately translated) coupling measure. Then ω and ω' are distributed as $\mathbb{P}_{p,q}$ and $\mathbb{P}_{p',q'}$ respectively, and the cluster of the root in ω is a subset of the cluster of the root in ω' . This concludes the proof of (2).

We now turn to the proof of (3). As the two proofs are very similar, we now only outline the main steps of the argument.

We let $\bar{\mathbb{E}}_{d,k}^\delta, \bar{\mathbb{E}}_{d,k}^\ell, \bar{\mathbb{E}}_{d,k}, \bar{\Omega}_\delta, \bar{\Omega}_\ell$ and $J_{\bar{\omega}}(A)$ be the same as before. A special configuration $\bar{\omega}^* \in \bar{\Omega}_\delta \times \bar{\Omega}_\delta \times \bar{\Omega}_\ell$ is defined as follows:

$$\bar{\omega}_{\delta,1}^* \equiv 0, \quad \bar{\omega}_{\delta,2}^* \equiv 1, \quad \bar{\omega}_\ell^*(\langle r, s \rangle) = 1 \text{ if and only if } r \in \cup_{n=k}^{2k-1} [d]^n.$$

Using Lemma 3.1, we obtain $q' < q_0 < q$ and a coupling of $X = (X_{\delta,1}, X_{\delta,2}, X_\ell)$ and $Y = (Y_{\delta,1}, Y_{\delta,2}, Y_\ell)$ so that the following hold. The values of $X_{\delta,1}, X_{\delta,2}$ and X_ℓ in all edges are independent; $X_{\delta,1}, X_{\delta,2}$ and X_ℓ assign each edge to be open with respective probabilities $p, \frac{p'-p}{1-p}$ and q ; the values of $Y_{\delta,1}, Y_{\delta,2}$ and Y_ℓ in all edges are independent; $Y_{\delta,1}, Y_{\delta,2}$ and Y_ℓ assign each edge to be open with respective probabilities $p, \frac{p'-p}{1-p}$ and q' ; the following event has probability one:

$$\{X = Y\} \cup \{X = (\bar{\omega}_{\delta,1}^*, \bar{\omega}_{\delta,2}^*, \bar{\omega}_\ell^*)\} \cup \{Y = (\bar{\omega}_{\delta,1}^*, \bar{\omega}_{\delta,2}^*, \bar{\omega}_\ell^*)\}.$$

We then let $\bar{\omega}_\delta = X_{\delta,1}, \bar{\omega}_\ell = X_\ell, \bar{\omega}'_\delta = Y_{\delta,1} \vee Y_{\delta,2}$ and $\bar{\omega}'_\ell = Y_\ell$. This coupling then guarantees (6) as before, which concludes the proof of Theorem 2.2.

4 | COMPARISON OF DIFFERENT RANGES

In this section we prove Theorem 2.1. The general idea behind the proof is to explore short edges until reaching a dead end, then use a coupling construction to show that one has a better chance to proceed from each dead end when k is larger.

Let $u \in \mathbb{V}_{d,k}$ and $r = (r_1, \dots, r_k) \in [d]^k$, so that $e = \langle u, u \cdot r \rangle \in \mathbb{E}_{d,k}^\ell$. We define the trace of e to be the set of short edges

$$\text{trace}(e) = \{\langle u, u \cdot r_1 \rangle, \langle u \cdot r_1, u \cdot (r_1, r_2) \rangle, \dots, \langle u \cdot (r_1, \dots, r_{k-1}), u \cdot r \rangle\}.$$

Fix $\omega = (\omega_s, \omega_\ell)$, with $\omega_s \in \{0, 1\}^{\mathbb{E}_{d,k}^s}$ and $\omega_\ell \in \{0, 1\}^{\mathbb{E}_{d,k}^\ell}$, and a set $A \subseteq \mathbb{V}_{d,k}$. We let $\Pi(A)$ be the cluster of A in ω , that is, the set of vertices of $\mathbb{T}_{d,k}$ which can be reached by a path started from some vertex of A and consisting of directed edges which are open in ω (note that $\Pi(A)$ depends on A and ω but we omit ω from the notation; this will also be the case for further notation that we introduce). We also let $\pi(A)$ be the cluster of A in ω_s , that is, the set of vertices of $\mathbb{T}_{d,k}$ that can be reached by a path started from some vertex of A and consisting of *short* edges, all of which are open in ω_s . Note that $A \subseteq \pi(A) \subseteq \Pi(A)$.

We say a short edge $e = \langle u, v \rangle \in \mathbb{E}_{d,k}^s$ is a *hub* for A (in ω) if the following two conditions hold:

$$\text{prog}(v) \cap \pi(A) = \emptyset \quad \text{and} \quad \text{prog}(u) \cap \pi(A) \neq \emptyset. \quad (7)$$

We let $\sigma(A)$ denote the set of hubs for A in ω .

Lemma 4.1 *Let $\omega \in \{0, 1\}^{\mathbb{E}_{d,k}}$ and $A \subseteq \mathbb{V}_{d,k}$. Then,*

$$\text{the progenies } \text{prog}(e) \text{ for } e \in \sigma(A) \text{ are disjoint.} \quad (8)$$

Further assuming that

$$\text{there exists } w \in \mathbb{V}_{d,k} \text{ such that } A \subseteq \{w \cdot v : v \in \cup_{n=0}^k [d]^n\}, \quad (9)$$

we also have

$$\begin{aligned} &\text{for any } e = \langle u, u \cdot r \rangle \in \mathbb{E}_{d,k}^\ell \text{ such that } u \in \pi(A) \text{ and } u \cdot r \notin \pi(A), \\ &\text{there exists a unique } e' \in \text{trace}(e) \cap \sigma(A) \end{aligned} \quad (10)$$

and

$$\begin{aligned} &\Pi(A) \text{ is the disjoint union of } \pi(A) \text{ and the sets} \\ &\Pi(A) \cap \text{prog}(e) \text{ for } e \in \sigma(A). \end{aligned} \quad (11)$$

Proof To prove (8), assume that there are two distinct hubs

$$e = \langle u, v \rangle, \quad e' = \langle u', v' \rangle \in \sigma(A) : \quad \text{prog}(e) \cap \text{prog}(e') \neq \emptyset.$$

Then either $u \in \text{prog}(v')$ or $u' \in \text{prog}(v)$. Without loss of generality we assume the latter. Together with (7) applied to e' , this implies that

$$\text{prog}(v) \cap \pi(A) \supseteq \text{prog}(u') \cap \pi(A) \neq \emptyset,$$

which contradicts (7) applied to e .

Now fix an edge $e = \langle u, u \cdot r \rangle$ as in (10). Consider the k short edges in the trace of e . By the first statement, we know that at most one of these short edges is in $\sigma(A)$. In order to show that one of them is in $\sigma(A)$, it suffices to show that

$$\text{prog}(u) \cap \pi(A) \neq \emptyset \quad \text{and} \quad \text{prog}(u \cdot r) \cap \pi(A) = \emptyset. \quad (12)$$

The first claim of (12) follows from the fact that $u \in \pi(A)$; let us prove the second. We are given that $u \cdot r \notin \pi(A)$, so it suffices to prove that $\text{prog}(u \cdot r) \cap A = \emptyset$. For vertices u', v' with $v' \in \text{prog}(u')$, let $\text{dist}(u', v')$ denote the length of the unique path of short edges from u' to v' . Then, (9) gives $\text{dist}(w, v) \leq k$ for all $v \in A$. If $v \in \text{prog}(u \cdot r)$ and $v \neq u \cdot r$, then

$$\text{dist}(w, v) > \text{dist}(w, u \cdot r) = \text{dist}(w, u) + k,$$

so $v \notin A$. We also have $u \cdot r \notin A$, so the proof of (12) is complete.

Statement (11) is an immediate consequence of (8) and (10). ■

Again fix $\omega \in \{0, 1\}^{\mathbb{E}_{d,k}}$ and $A \subseteq \mathbb{V}_{d,k}$ satisfying (9). For each hub $e \in \sigma(A)$, we define

$$R(A, e) = \{e' = \langle u', v' \rangle \in \mathbb{E}_{d,k}^{\ell} : u' \in \pi(A) \text{ and } e \in \text{trace}(e')\},$$

$$\bar{S}(A, e) = \{v' \in \mathbb{V}_{d,k} : \langle u', v' \rangle \in R(A, e) \text{ for some } u' \in \mathbb{V}_{d,k}\},$$

$$S(A, e) = \{v' \in \mathbb{V}_{d,k} : \omega(\langle u', v' \rangle) = 1 \text{ for some } \langle u', v' \rangle \in R(A, e)\}.$$

Note that $S(A, e) \subseteq \bar{S}(A, e) \subseteq \text{prog}(e)$. Also note that, if $e_1, e_2 \in \sigma(A)$ are distinct, then $R(A, e_1)$ and $R(A, e_2)$ are disjoint, by (8). Finally, note that for every $e \in \sigma(A)$, we have

$$\Pi(A) \cap \text{prog}(e) = \Pi(S(A, e)),$$

so that (11) can be restated as

$$\Pi(A) = \pi(A) \cup \left(\bigcup_{e \in \sigma(A)} \Pi(S(A, e)) \right), \quad (13)$$

where the union is disjoint.

For A satisfying (9), we now let $\mathcal{C}_{p,q,k}(A)$ be the random set $\Pi(A)$ when ω is sampled from the measure $\mathbb{P}_{p,q}$ on percolation configurations on $\mathbb{T}_{d,k}$. Note that $\mathcal{C}_{p,q,k} = \mathcal{C}_{p,q,k}(\{o\})$.

We observe that, conditioning on $\pi(A)$, $\sigma(A)$ is determined and the sets $\Pi(S(A, e))$ are independent over $e \in \sigma(A)$. Indeed, $\Pi(S(A, e))$ is determined by $\pi(A)$ and $\omega(e')$ for all

$$e' = \langle u', v' \rangle \text{ with } v' \in \text{prog}(e).$$

The sets of edges displayed above are disjoint for distinct choices of $e \in \sigma(A)$. Indeed, assume $e, f \in \sigma(A)$, $e \neq f$, and $e' = \langle u', v' \rangle, f' = \langle w', x' \rangle$ are long edges with $v' \in \text{prog}(e)$, $x' \in \text{prog}(f)$. Then, since (8) gives $\text{prog}(e) \cap \text{prog}(f) = \emptyset$, we obtain $v' \neq x'$, so $e' \neq f'$.

Guided by this consideration, we now present a recursive exploration algorithm to reveal $\mathcal{C}_{p,q,k}(A)$. The algorithm starts by applying the following two steps to the set A :

Step 1. Explore $\pi(A)$ by revealing only the edges in ω_s that are necessary. More precisely, grow $\pi(A)$ progressively by starting from A and querying the open/closed-state of short edges one by one,

each time selecting a short edge $e = \langle u, v \rangle$ such that u is already included in $\pi(A)$ and v is not (and also following some lexicographic-type priority rule that guarantees that the full $\pi(A)$ is explored). Note that this also determines $\sigma(A)$, hence $\bar{S}(A, e)$ for each $e \in \sigma(A)$.

Step 2. For each $e \in \sigma(A)$, reveal $S(A, e)$. This is the same as revealing the value of $\omega_e(e')$ for each long edge $e' \in R(A, e)$.

Note that, if $e = \langle u, v \rangle \in \sigma(A)$, then $S(A, e) \subseteq \{v \cdot w : w \in \cup_{n=0}^{k-1} [d]^n\}$, so that property (9) holds with A replaced by $S(A, e)$. The algorithm then proceeds by applying Steps 1 and 2 to each of the sets $S(A, e)$, which take the role of A . That is: in Step 1 it explores $\pi(S(A, e))$, which also reveals $\sigma(S(A, e))$, and in Step 2, for each $e' \in \sigma(S(A, e))$, it reveals $S(S(A, e), e')$. The recursion then continues to further levels. By (13), this reveals the whole cluster $\mathcal{C}_{p,q,k}(A)$.

We now want to look at the distributions of $S(A, e)$ and $\Pi(S(A, e))$ for $e \in \sigma(A)$. Although these distributions are easily understood, they are somewhat clumsy to describe, so we will need some more notation.

First, fix $e = \langle u, v \rangle \in \sigma(A)$ with $v = (v_1, \dots, v_n)$. Define

$$\beta(A, e) = \{i \in \{1, \dots, k\} : (v_1, \dots, v_{n-i}) \in \pi(A)\},$$

it describes which ancestors of e have been reached from A using short edges and could reach $\text{prog}(e)$ using long edges (the ω -state of which is not looked at). Note that

$$R(A, e) = \{(\langle v_1, \dots, v_{n-i} \rangle, v \cdot w) : i \in \beta(A, e), w \in [d]^{k-i}\},$$

so that

$$\bar{S}(A, e) = \{v \cdot w : i \in \beta(A, e), w \in [d]^{k-i}\}.$$

Second, we define some shift mappings in $\mathbb{T}_{d,k}$. Given $u \in \mathbb{V}_{d,k}$, we let $\tau_u : \text{prog}(u) \rightarrow \mathbb{V}_{d,k}$ be the function

$$\tau_u(u \cdot v) = v, \quad v \in [d]_{\text{star}}.$$

If $e = \langle u, v \rangle \in \mathbb{E}_{d,k}^j$, we let $\tau_e = \tau_v$.

Third, given $b \subseteq \{1, \dots, k\}$, we let $\mathcal{A}_{q,k}(b)$ denote the distribution of the random subset of $\cup_{i \in b} [d]^{k-i}$ in which, independently, each point is included with probability q .

Let $A \subseteq \mathbb{V}_{d,k}$ satisfy (9). Conditioning on $\pi(A)$, for each $e \in \sigma(A)$ we have

$$\tau_e(S(A, e)) \stackrel{(d)}{=} \mathcal{A}_{q,k}(\beta(A, e)) \quad (14)$$

and the law of $\tau_e(\Pi(S(A, e)))$ is equal to the law of the cluster of B in $\mathbb{T}_{d,k}$, where B is chosen according to $\mathcal{A}_{q,k}(\beta(A, e))$.

We finally turn to the desired comparison between $\mathcal{C}_{p,q,k}$ for different values of the parameters. Given $A, B \subseteq \mathbb{V}_{d,k}$, let us write $A \leq B$ in case there exist $u, v \in [d]_{\text{star}}$ such that $A \subseteq \text{prog}(u)$ and $\tau_u(A) \subseteq \tau_v(B \cap \text{prog}(v))$.

Lemma 4.2 *For any $k \in \mathbb{N}$ and $q \in (0, 1)$, there exists $q' < q$ such that the following holds. Let $b' \subseteq \{1, \dots, k+1\}$ and $b = b' \cap \{1, \dots, k\}$. There exists a coupling (A, B) of random sets $A, B \subseteq [d]_{\text{star}}$ such that*

$$A \leq B, \quad A \stackrel{(d)}{=} \mathcal{A}_{q,k}(b) \quad \text{and} \quad B \stackrel{(d)}{=} \mathcal{A}_{q',k+1}(b').$$

With this lemma at hand, we are ready to conclude the proof of Theorem 2.1. Fix $p, q \in (0, 1)$ and $k \in \mathbb{N}$, and choose q' corresponding to k and q in Lemma 4.2. The idea is to compare the explorations of $\mathcal{C}_{p,q,k}$ and $\mathcal{C}_{p,q',k+1}$ using coupling. Recall that our algorithm to explore a cluster proceeds by the iterative application of two steps. Step 1 grows a portion of the cluster using only short edges, so it can be taken as the same for both explorations, since short edges have the same probability of being open in both. Step 2 inspects “exit routes”, using long edges, from the portion of cluster revealed in Step 1; Lemma 4.2 guarantees that this is better (in the sense of $\leq$ -domination) for $\mathcal{C}_{p,q',k+1}$ than for $\mathcal{C}_{p,q,k}$.

Let us now present the coupling of explorations more formally. Note that we are dealing with percolation in the two graphs $\mathbb{T}_{d,k}$ and $\mathbb{T}_{d,k+1}$ simultaneously; these graphs have the same set of vertices (namely, $[d]_{\text{star}}$) and same set of short edges, but the long edges differ. A set $A \subseteq [d]_{\text{star}}$ satisfying condition (9) for k also satisfies it when k is replaced by $k+1$. For such a set, and for $e \in \sigma(A)$, instead of $S(A, e)$ we will now write $S_k(A, e)$ and $S_{k+1}(A, e)$ to distinguish this set in the two graphs.

The coupled exploration of $\mathcal{C}_{p,q,k}$ and $\mathcal{C}_{p,q',k+1}$ starts with revealing $\pi(\{o\})$, which we can take as the same in both clusters. Thus, $\sigma(\{o\})$ is also the same in both graphs, and we enumerate

$$\sigma(\{o\}) = \{e^1, \dots, e^N\}.$$

Also write

$$A^i = S_k(\{o\}, e^i), \quad B^i = S_{k+1}(\{o\}, e^i), \quad i = 1, \dots, N.$$

By Lemma 4.2, these can be sampled with $A^i \leq B^i$, so there exist $u^i, v^i \in [d]_{\text{star}}$ and $\tilde{B}^i \subseteq B^i$ such that

$$A^i \subseteq \text{prog}(u^i), \quad \tilde{B}^i \subseteq \text{prog}(v^i), \quad \tau_{u^i}(A^i) = \tau_{v^i}(\tilde{B}^i).$$

The second level of the exploration then proceeds as follows. For each $i \in \{1, \dots, N\}$, take $\pi(\tau_{u^i}(A_i))$ and $\pi(\tau_{v^i}(\tilde{B}^i))$ as the same in both clusters, enumerate

$$\sigma(\tau_{u^i}(A^i)) = \sigma(\tau_{v^i}(\tilde{B}^i)) = \{e^{i,1}, \dots, e^{i,N_i}\},$$

and let

$$A^{i,j} = S_k(\tau_{u^i}(A^i), e^{i,j}), \quad B^{i,j} = S_{k+1}(\tau_{v^i}(\tilde{B}^i), e^{i,j}), \quad j = 1, \dots, N_i,$$

which can be sampled with $A^{i,j} \leq B^{i,j}$ for each j . Further levels are then carried out in the same way. The construction guarantees that $\mathcal{C}_{p,q,k}$ is embedded in $\mathcal{C}_{p,q',k+1}$, concluding the proof of Theorem 2.1. It remains only to prove the previous lemma.

Proof of Lemma 4.2 We can assume that $k+1 \notin b'$, so that $b = b'$. In that case, for $\hat{q} \in (0, 1)$ and $\hat{B}$ a random subset of $[d]_{\text{star}}$,

$$\hat{B} \sim \mathcal{A}_{\hat{q},k+1}(b) \quad \text{if and only if} \quad \begin{array}{l} \tau_{(1)}(\hat{B}), \dots, \tau_{(d)}(\hat{B}) \text{ i.i.d.} \\ \text{and distributed as } \mathcal{A}_{\hat{q},k}(b). \end{array} \quad (15)$$

We now define sets $S_1^*, \dots, S_d^* \subseteq [d]_{\text{star}}$ by

$$S_1^* = \emptyset, \quad S_2^*, \dots, S_d^* = \bigcup_{i \in b} [d]^{k-i}.$$

By Lemma 3.1, there exists $q' < q$ and a coupling of random sets $X_1, \dots, X_d, Y_1, \dots, Y_d \subseteq [d]_{\text{star}}$ so that $X_1, \dots, X_d$ are independent and distributed as $\mathcal{A}_{q,k}(b)$, $Y_1, \dots, Y_d$ are independent and distributed as $\mathcal{A}_{q',k}(b)$ and the following event has probability 1:

$$\begin{aligned} \{(X_1, \dots, X_d) = (Y_1, \dots, Y_d)\} \cup \{(X_1, \dots, X_d) = (S_1^*, \dots, S_d^*)\} \\ \cup \{(Y_1, \dots, Y_d) = (S_1^*, \dots, S_d^*)\}. \end{aligned} \quad (16)$$

The desired conclusion now follows by setting

$$A = X_1, \quad B = \cup_{a \in [d]} \{a \cdot u : u \in Y_a\}. \quad \blacksquare$$

ACKNOWLEDGEMENTS

The authors would like to thank Aernout van Enter for helpful discussions, and Gábor Pete for pointing out an inaccurate citation in an earlier version of this paper. B.N.B.L. would like to thank the University of Groningen and D.V. would like to thank NYU-Shanghai for support and hospitality. This project was supported by grants CNPq 309468/2014-0, FAPEMIG (Programa Pesquisador Mineiro), PIP 11220130100521CO, PICT-2015-3154, PICT-2013-2137, PICT-2012-2744, Conicet-45955 and MinCyT-BR-13/14.

REFERENCES

1. D. Ahlberg, *Asymptotics of first-passage percolation on one-dimensional graphs*, Adv. Appl. Probab. **470** (2015), no. 1, 182–209. <https://doi.org/10.1239/aap/1427814587>
2. M. Aizenman and G. Grimmett, *Strict monotonicity for critical points in percolation and ferromagnetic models*, J. Stat. Phys. **630** (1991), no. 5–6, 817–835. <https://doi.org/10.1007/BF01029985>
3. S. E. Alm and J. C. Wierman, *Inequalities for means of restricted first-passage times in percolation theory*, Combin. Probab. Comput. **80** (1999), no. 4, 307–315. <https://doi.org/10.1017/S0963548399003843>
4. E. D. Andjel and M. Sued, *An inequality for oriented 2-D percolation*. In and out of equilibrium, 2, Progr. Probab., vol. 60, Birkhäuser, Basel, 2008, pp. 21–30, http://dx.doi.org/10.1007/978-3-7643-8786-0_2.
5. A. P. F. Atman and M. Schnabel, Private communication (2010).
6. P. Balister, B. Bollobás, and O. Riordan, *Essential enhancements revisited*, Unpublished data, 2014, <http://arxiv.org/abs/1402.0834>.
7. I. Benjamini, A. Nachmias, and Y. Peres, *Is the critical percolation probability local?*, Probab. Theory Relat. Fields. **1490** (2011), no. (1–2), 261–269. <https://doi.org/10.1007/s00440-009-0251-5>
8. B. N. B. de Lima and A. Procacci, *A note on anisotropic percolation*, Lett. Math. Phys. **70** (2004), 223–229. <https://doi.org/10.1007/s11005-004-4297-0>
9. B. N. B. de Lima, R. Sanchis, and R. W. C. Silva, *Critical point and percolation probability in a long range site percolation model on $\mathbb{Z}^d$* , Stochast. Process. Appl. **1210** (2011), no. 9, 2043–2048. <https://doi.org/10.1016/j.spa.2011.05.009>
10. B. N. B. de Lima, A. Procacci, and R. Sanchis, *A remark on monotonicity in Bernoulli bond percolation*, J. Stat. Phys. **1600** (2015), no. 5, 1244–1248. <https://doi.org/10.1007/s10955-015-1284-z>
11. J.-B. Gouéré, *Monotonicity in first-passage percolation*, ALEA Lat. Am. J. Probab. Math. Stat. **110** (2014), no. 1, 565–569. <http://alea.impa.br/articles/v11/11-26.pdf>.
12. J. M. Hammersley and D. J. A. Welsh, *First-passage percolation, subadditive processes, stochastic networks, and generalized renewal theory*, Proc. Internat. Res. Semin., Statist. Lab., Univ. California, Berkeley, CA, Springer-Verlag, New York, 1965, pp. 61–110, http://dx.doi.org/10.1007/978-3-642-99884-3_7.
13. C. D. Howard, *Differentiability and monotonicity of expected passage time in Euclidean first-passage percolation*, J. Appl. Probab. **380** (2001), no. 4, 815–827. <https://projecteuclid.org/euclid.jap/1011994174url>.

14. G. K. Iliev, E. J. Janse van Rensburg, and N. Madras, *Phase diagram of inhomogeneous percolation with a defect plane*, J. Stat. Phys. **1580** (2015), no. 2, 255–299. <https://doi.org/10.1007/s10955-014-1125-5>
15. S. Martineau and V. Tassion, *Locality of percolation for Abelian Cayley graphs*, Ann. Probab. **45** (2017), 1247–1277. <https://doi.org/10.1214/15-AOP1086>
16. A. Q. Teixeira, *Comparison between oriented and directed bond percolation on $\mathbb{Z}^2$* , Oral communication, 2006.
17. J. van den Berg, *A counterexample to a conjecture of J. M. Hammersley and D. J. A. Welsh concerning first-passage percolation*, Adv. Appl. Probab. **2** (1983), 465–467. <https://doi.org/10.2307/1426447>
18. J. van den Berg, Private communication. (2007).

How to cite this article: de Lima BNB, Rolla LT, Valesin D. Monotonicity and phase diagram for multirange percolation on oriented trees. *Random Struct Alg.* 2019;55:160–172. <https://doi.org/10.1002/rsa.20805>