University of Groningen

Virtual reality experiments linking social environment and psychosis

Veling, Wim; Brinkman, Willem-Paul; Dorrestijn, Emily; van der Gaag, Mark

Published in:
Cyberpsychology, Behavior and Social Networking

DOI:
[10.1089/cyber.2012.0497](https://doi.org/10.1089/cyber.2012.0497)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2014

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Veling, W., Brinkman, W-P., Dorrestijn, E., & van der Gaag, M. (2014). Virtual reality experiments linking social environment and psychosis: a pilot study. *Cyberpsychology, Behavior and Social Networking*, 17(3), 191-195. <https://doi.org/10.1089/cyber.2012.0497>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Virtual Reality Experiments Linking Social Environment and Psychosis: A Pilot Study

Wim Veling, PhD,^{1,2} Willem-Paul Brinkman, PhD,³ Emily Dorrestijn, MSc,¹ and Mark van der Gaag, PhD^{1,4}

Abstract

Initial studies with healthy subjects and individuals with high risk for psychosis have suggested that virtual reality (VR) environments may be used to investigate social and psychological mechanisms of psychosis. One small study reported that VR can safely be used in individuals with current persecutory delusions. The present pilot study investigated the feasibility and potential negative side effects of exposure to different virtual social risk environments in patients with first episode psychosis and in healthy controls. Seventeen patients with first episode psychosis (FEP) and 24 healthy control subjects (HC) participated in four virtual experiments during which they walked for 3.5–4 minutes in a virtual café, looking for avatars with digits on their clothing. The level of paranoid thoughts, as well as psychological, physiological, and behavioral correlates of paranoid thoughts, were measured in different virtual social risk environments, manipulating two factors: population density and ethnicity of avatars. FEP and HC frequently had paranoid thoughts about avatars. Paranoia in the real world correlated strongly with paranoid thoughts about avatars in virtual environments (Spearman's $\rho = 0.67$ and 0.54 in FEP and HC respectively, $p < 0.01$). FEP kept a smaller distance to avatars than HC. In FEP, but not in HC, galvanic skin response was significantly stronger in virtual environments with avatars of other ethnicity than in the own ethnicity condition. These results suggest that VR is an acceptable and sufficiently realistic method to use in patients with first episode psychosis. VR research may help to increase our understanding of the social and psychological mechanisms of psychosis and to develop new treatment applications.

Introduction

PSYCHOTIC SYNDROMES can be understood as disorders of adaptation to social context.¹ Epidemiological studies have shown high rates of psychotic disorders in densely populated urban environments² and among ethnic minorities,^{3,4} in particular when they live in neighborhoods with relatively few others of their own ethnic group.⁵ The exact nature of these environmental exposures as well as the underlying mechanisms remain largely unknown, in part because measurement of the daily social environment of individuals, and their psychological and physiological response to it, remains a major challenge.^{1,6} Virtual reality (VR) technology offers the possibility of exposing individuals to controlled experimental social risk environments. In VR, sense data from the natural world are substituted with sense

data about an imaginary world that change in response to the user's actions. This interactive process results in a "sense of presence" in a three-dimensional virtual world.⁷ VR is relatively new in psychosis research. Freeman et al. found in small samples that exposure to a neutral VR environment could elicit paranoid ideations toward avatars in healthy individuals, individuals with at-risk mental state for psychosis, and in individuals with psychotic disorder.^{8–10} VR did not have serious side effects and was well tolerated.⁹ These findings have not yet been replicated.

The present pilot study investigated the feasibility of VR in patients with first episode psychosis and in healthy controls. We explored level of paranoid thoughts as well as psychological, physiological, and behavioral correlates of paranoid thoughts in different virtual social environments, manipulating two factors: population density and ethnicity of avatars.

¹Parnassia Psychiatric Institute, The Hague, The Netherlands.

²Department of Psychiatry and Neuropsychology, Maastricht University Medical Centre, Maastricht, The Netherlands.

³Delft University of Technology, Delft, The Netherlands.

⁴Department of Clinical Psychology, VU University and EMGO Institute for Health and Care Research, Amsterdam, The Netherlands.

Methods

Participants

Seventeen patients with first episode psychosis (FEP) were recruited from a specialist service for early psychosis in The Hague. Inclusion criteria for the study were: aged 18–40 years and a DSM IV diagnosis of schizophrenia, schizophreniform disorder, schizoaffective disorder, brief psychotic disorder, delusional disorder, or psychotic disorder not otherwise specified. Twenty-four healthy control subjects (HC) were recruited from Delft University of Technology staff and students. Written informed consent was obtained from all participants.

Virtual reality setting

The virtual environment was a café with an indoor and an outdoor part, adjacent to a shopping street, built by CleVR. Participants could navigate in the virtual environment, using a Logitech Chillstream Gamepad. They wore an Emagin Z800 3D Visor with a resolution of SVGA 800×600 24 bit, with 40° diagonal field of view, and built-in 3DOF tracker. Avatars walked in and out, chatted, and had drinks while sitting or standing at a table. Café background noises were played. The avatars were programmed to exhibit neutral behavior. They glanced up when participants approached them, and if participants blocked their way, avatars politely asked them to move aside.

Experiments

To engage the participants with the avatars, the system randomly assigned a number to five of the avatars that was visible on their clothing. Participants were instructed to find the numbered avatars and to memorize the numbers. This task had to be completed in three-and-half (HC) to four (FEP) minutes, after which the experiment ended. Patients were allowed more time because psychosis is associated with reduced cognitive performance.

The number of avatars in the café and the ethnic appearance of the avatars was manipulated. Both factors had two options. The ethnic appearance could be either mainly (95%) white European or mainly (75%) North-African (own or other ethnicity, depending on the ethnicity of the participant); the number of avatars could be either between seven and nine (low density) or between 34 and 38 (high density) avatars.

All subjects participated in four conditions: (a) own ethnicity/low population density, (b) other ethnicity/low population density, (c) own ethnicity/high population density, and (d) other ethnicity/high population density. All FEP started with own/low condition; the other experimental conditions were randomized for order of condition. A counter-balanced experimental design was used for the HC. While these designs are slightly different, both approaches were intended to prevent order effects.

Measurements

Before experiments. Baseline measures included the Green Paranoid Thoughts Scale (GPTS),¹¹ the Social Interaction Anxiety Scale (SIAS),¹² the Davos Assessment of Cognitive Biases Scale (DACOBS),¹³ the Self-Esteem Rating Scale (SERS),¹⁴ and the Simulator Sickness Questionnaire (SSQ).¹⁵ Before the experiments, neutral physiological data were col-

lected with a Mobi8 from TMSi with Xpod Oximeter. Heart rate (HR) and galvanic skin response (GSR) sensor were measured using two finger electrodes.

During and between experiments. During experiments, besides HR and GSR data, position of participants in VR relative to avatars was automatically recorded three times per second. After each experiment, paranoid ideations toward avatars were assessed with the State Social Paranoia Scale (SSPS).¹⁶ Furthermore, subjective experience of distress was measured with visual analogue scales, ranging from 1 to 10.

After experiments. The SSQ was administered again in order to measure cyber sickness. The Igroup Presence Questionnaire (IPQ)¹⁷ was used to explore sense of presence in the virtual environment.

Statistical analyses. Nonparametric tests were used for statistical analyses, or, in cases with two or more independent variables, linear mixed model analyses on aligned rank transformed data.¹⁸ Real world symptoms at baseline were compared between FEP and HC with Mann-Whitney *U* tests for independent samples. In order to investigate whether real world experiences corresponded to VR experiences, non-parametric Spearman coefficients were calculated for correlations between degree of paranoid thoughts (and correlates of paranoia, such as social anxiety and cognitive biases) at baseline and paranoid thoughts about avatars in VR. Physiological and behavioral responses to virtual environments were explored by comparing GSR, HR, and distance to avatars, averaged over all virtual experiments, between FEP and HC. The GSR in micro Siemens (μ S) values (δ) were set against the mean GSR value of the neutral (β) phase ($(\delta - \beta) / \beta$), whereby an increase indicates an increase in sweating. Finally, psychological, physiological, and behavioral responses of participants were compared across four different virtual social environments. Since the virtual experiments combined ethnicity and population density conditions (e.g., other ethnicity and high, or own ethnicity and low), results for ethnicity conditions were averaged over population density conditions (low and high), and results for population density conditions were averaged over ethnic density conditions (own and other).

Results

Sociodemographic characteristics of the participants are shown in Table 1.

TABLE 1. CHARACTERISTICS OF STUDY PARTICIPANTS

	FEP (n = 17)	HC (n = 24)
Age, mean (SD)	27.3 (5.5)	29.0 (9.2)
Male sex, n (%)	14 (82.4)	20 (83.3)
Dutch ethnicity, n (%)	6 (35.3)	24 (100.0)
DSM IV diagnosis, n (%)		
Schizophrenia	10 (58.8)	
Schizoaffective disorder	3 (17.6)	
Delusional disorder	2 (11.8)	
Psychotic disorder NOS	2 (11.8)	—

Note. FEP, first episode psychosis; HC, healthy control subjects.

TABLE 2. REAL WORLD EXPERIENCES AND PARANOID THOUGHTS ABOUT AVATARS IN VR

	FEP		HC	
	Experiences in real world, mean (SD)	Correlation with paranoia in VR ^a	Experiences in real world, mean (SD) ^b	Correlation with paranoia in VR ^a
Paranoid thoughts				
Persecution	34.2 (20.7)	0.65**	23.3 (5.1)	0.54**
Social reference	32.7 (18.4)	0.72**	18.4 (3.1)	0.32
Total	66.8 (44.4)	0.67**	41.7 (6.8)	0.54**
Social anxiety	40.2 (18.4)	0.73**	37.6 (9.9)	0.42 *
Cognitive bias				
Jumping to conclusions	20.8 (5.9)	0.42	18.6 (3.6)	0.18
Belief inflexibility	19.8 (5.8)	0.71**	14.2 (3.7)**	0.39
Attention to threat	24.8 (8.6)	0.27	19.3 (6.1)*	0.50*
External attribution	24.1 (8.5)	0.72**	15.9 (4.6)**	0.52**
Social cognition problems	21.9 (9.2)	0.91**	16.9 (4.1)*	0.16
Subjective cognitive problems	22.3 (6.5)	0.63**	17.2 (7.2)*	0.39
Safety behavior	15.2 (9.5)	0.63**	8.0 (2.4)**	0.23
Total	154.3 (44.3)	0.78**	110.0 (19.9)**	0.47*
Self-esteem				
Positive	36.9 (9.6)	-0.10	47.8 (9.7)**	-0.13
Negative	37.7 (12.5)	0.58*	23.4 (10.1)**	0.40
Total	-0.8 (14.8)	-0.46	24.4 (17.0)**	-0.23

Note. ^aSpearman nonparametric correlation coefficients between real world experiences and paranoid thoughts in VR. ^bMann-Whitney *U* test for independent samples, FEP compared to HC.

* $p < 0.05$; ** $p < 0.01$.

FEP, first episode psychosis; HC, healthy control subjects.

Before the experiments, mean degree of paranoid thoughts was 66.8 ($SD=44.4$) in FEP and 41.7 ($SD=6.8$) in HC, Mann-Whitney *U* test, $p=0.482$. FEP had significantly more cognitive biases, higher subjective distress, and lower self-esteem than HC (Table 2).

Total SSPS scores ranged from 40 to 96 in HC (40 indicates not a single paranoid thought during any of the four experiments; $M=53.1$; $SD=17.1$) and 40 to 181 in FEP ($M=67.5$; $SD=44.1$). Table 2 shows that baseline scores of paranoia, social anxiety, and cognitive biases in the real world correlated strongly and statistically significantly with degree of paranoid thoughts in VR.

Paranoid thoughts about avatars varied slightly with virtual condition (Fig. 1), but within-subject differences across conditions were not statistically significant, neither in the FEP nor in the HC condition.

Differences between FEP and HC in physiological measures, averaged over all virtual conditions, were not statistically significant (GSR: 89.5 ($SD=151.6$) vs. 66.2 ($SD=58.6$) and mean HR: 87.4 ($SD=16.2$) vs. 82.9 ($SD=11.0$); Table 3). Heart rate did not vary much across virtual conditions. With regard to GSR, there was a two-way interaction effect of ethnic density and case-control status, $df=1.32$, $F=9.824$, $p=0.004$. In FEP, GSR was significantly higher in the other ethnicity ($M=114.1$; $SD=156.8$) than in the own ethnicity condition ($M=64.9$; $SD=155.8$), whereas this was not the case in HC.

Averaged over all conditions, HC kept significantly more distance to avatars than FEP: 1.29 meters ($SD=0.06$) versus 1.25 meters ($SD=0.05$), $df=1.39$, $F=4.38$, $p=0.043$. There was an interaction effect between ethnic density and population density. Participants kept more distance to avatars in the other ethnicity environment, but only if in the low population density condition, $df=1.39$, $F=5.08$, $p=0.030$.

Symptoms of simulator sickness were low. They slightly increased during experiments but were not statistically significant between pre and post measurement. The mean SSQ score was significantly higher in FEP than in HC before but not after the experiments. Sense of presence in VR did not differ significantly between FEP and HC. The mean total IPQ score was 41.53 ($SD=12.61$) in cases and 40.75 ($SD=15.73$) in controls, Mann-Whitney *U* Test=179, $Z=-.029$, $p=0.977$.

Discussion

Main results

In this pilot study of VR experiments linking social environment to psychosis, we have found that patients with FEP as well as HC frequently had paranoid thoughts about neutral virtual characters. Symptoms and correlates of paranoia

FIG. 1. Paranoid thoughts about avatars in different virtual social environments: patients with first episode psychosis (FEP), and healthy control subjects (HC).

TABLE 3. PHYSIOLOGICAL AND BEHAVIORAL RESPONSES TO VIRTUAL SOCIAL ENVIRONMENTS^a

	FEP (n = 17)	HC (n = 24)
Galvanic skin response, % (SD) ^b	89.5 (151.6)	66.2 (58.6)
Heart rate, mean BPM (SD)	87.4 (16.2)	82.9 (11.0)
Mean distance to avatars, meters (SD)	1.25 (0.05)	1.29 (0.06) ^c

Note. ^aAveraged over all experiments. ^bMissing data galvanic skin response for six FEP (n = 11) and 1 HC (n = 23). ^cp < 0.05, Mann-Whitney U test.

FEP, first episode psychosis; HC, healthy control subjects.

in the real world strongly correlated to experiences in virtual environments, indicating that VR can be used for environmental studies of psychosis. It was feasible to expose FEP and HC to different virtual social environments. FEP kept a smaller distance to avatars than HC did. In FEP, but not in HC, galvanic skin response was significantly higher in virtual environments with other ethnicity avatars than in the own ethnicity condition.

Methodological and technical considerations

The pilot study was conducted in order to investigate the feasibility of virtual experiments in psychosis, and to explore responses of participants to different virtual environments. The main limitation is the small sample size, allowing only exploratory analyses. Based on these results, a larger study will be conducted.

Experience with use of physiological measures in VR is limited. Heart rate and skin conductance are commonly used in psychological and psychiatric research to assess an individual's emotional arousal and vulnerability to stress, as well as in studies of psychosis (liability).^{19,20} In our experiments, heart rate did not vary substantially across conditions. We used mean heart rate per condition, but variation may be detected only when smaller time windows before and after marked stressors are analyzed.

Participants were instructed to find and memorize numbers on the clothing of avatars. Because patients with psychosis often have cognitive deficits, they were given more time to complete the task. Still, since it may have taken a substantial part of their cognitive abilities to remember the digits correctly, some participants are likely to have paid less attention to the behavior of avatars and the virtual environment. This may explain why patients kept a slightly smaller distance to avatars than HC subjects.

A fully counterbalanced experimental design was used for the HC. For FEP, a crossover design was used in which the order of experiments was random except for the first. As a safety precaution, all patients started with the low population density and high ethnic density condition because this was considered the lowest social stress environment. Before the first experiment, they had some time in VR to get used to the virtual environment and the equipment, but the subjective distress and physiological arousal during the first experiment may still have been influenced by unfamiliarity with VR.

Previous research

There are few previous studies of VR in psychosis. One study exposed 20 patients with a current paranoid delusion

and 20 control subjects to a four-minute ride in a VR underground train containing neutral characters.⁹ The results were similar to our findings. A high proportion of the patients (65%) had persecutory thinking about the avatars. The VR experience did not raise levels of anxiety or symptoms of simulator sickness. Another study, using the same VR environment, reported that in 21 individuals with an at-risk mental state for psychosis, persecutory ideation in VR was predicted by higher levels of trait paranoia, anxiety, stress, immersion in VR, perseveration, and interpersonal sensitivity.¹⁰ Initial findings from South Korea suggest that VR scenarios may be useful in evaluating social skills and emotions arising from social interactions in patients with schizophrenia.^{21,22}

Conclusion

VR seems an acceptable and sufficiently realistic method to use in patients with first episode psychosis. It offers a unique way to expose individuals to controlled social environments. VR research may help to increase our understanding of the social and psychological mechanisms of psychosis and to develop new treatment applications, but larger studies are needed.

Acknowledgment

This study was supported by a grant from the Parnassia Psychiatric Institute.

Author Disclosure Statement

No competing financial interests exist.

References

1. Van Os J, Kenis G, Rutten BPF. The environment and schizophrenia. *Nature* 2010; 468:203–212.
2. Pedersen CB, Mortensen PB. Evidence of a dose-response relationship between urbanicity during upbringing and schizophrenia risk. *Archives of General Psychiatry* 2001; 58:1039–1046.
3. Bourque F, van der Ven E, Malla A. A meta-analysis of the risk for psychotic disorders among first- and second-generation immigrants. *Psychological Medicine* 2011; 41:897–910.
4. Veling W, Selten JP, Veen N, et al. Incidence of schizophrenia among ethnic minorities in the Netherlands: a four-year first-contact study. *Schizophrenia Research* 2006; 86:189–193.
5. Veling W, Susser E, van Os J, et al. Ethnic density of neighborhoods and incidence of psychotic disorders among immigrants. *American Journal of Psychiatry* 2008; 165:66–73.
6. March D, Hatch SL, Morgan C, et al. Psychosis and place. *Epidemiological Reviews* 2008; 30:84–100.
7. Freeman D. Studying and treating schizophrenia using virtual reality: a new paradigm. *Schizophrenia Bulletin* 2008; 34:605–610.
8. Freeman D, Garety PA, Bebbington P, et al. The psychology of persecutory ideation II: a virtual reality experimental study. *The Journal of Nervous & Mental Disease* 2005; 193:309–315.
9. Fornells-Ambrojo M, Barker C, Swapp D, et al. Virtual reality and persecutory delusions: safety and feasibility. *Schizophrenia Research* 2008; 104:228–236.
10. Valmaggia LR, Freeman D, Green C, et al. Virtual reality and paranoid ideations in people with an “at-risk mental state” for psychosis. *British Journal of Psychiatry* 2007; 191:s63–68.

11. Green CE, Freeman D, Kuipers E, et al. Measuring ideas of persecution and social reference: the Green et al. Paranoid Thought Scales (GPTS). *Psychological Medicine* 2008; 38:101–111.
12. Mattick RP, Clarke JC. Development and validation of measures of social phobia scrutiny fear and social interaction anxiety. *Behaviour Research & Therapy* 1998; 36:455–470.
13. Van der Gaag M, Schütz C, Ten Napel A, et al. The development of the Davos Assessment of Cognitive Biases Scale. *Schizophrenia Research* 2013; 144:63–71.
14. Lecomte T, Corbiere M, Laisne F. Investigating self-esteem in individuals with schizophrenia: relevance of the Self-Esteem Rating Scale-Short Form. *Psychiatry Research* 2006; 143:99–108.
15. Kennedy RS, Lane NE, Berbaum KS, et al. Simulator Sickness Questionnaire: an enhanced method for quantifying simulator sickness. *International Journal of Aviation Psychology* 1993; 3:203–220.
16. Freeman D, Pugh K, Green C, et al. A measure of state persecutory ideation for experimental studies. *The Journal of Nervous and Mental Disease* 2007; 195:781–784.
17. Schubert T, Friedmann F. The experience of presence: factor analytic insights. *Presence* 2001; 10:266–281.
18. Wobbrock JO, Findlater L, Gergle D, et al. (2011) The Aligned Rank Transform for nonparametric factorial analyses using only ANOVA procedures. In *Proceedings of the ACM Conference on Human Factors in Computing Systems*. New York: ACM Press; pp. 143–146.
19. Allen P, Freeman D, McGuire P. Slow habituation of arousal associated with psychosis proneness. *Psychological Medicine* 2007; 37:577–582.
20. Castro MN, Vigo DE, Weidema H, et al. Heart rate variability response to mental arithmetic stress in patients with schizophrenia: autonomic response to stress in schizophrenia. *Schizophrenia Research* 2008; 99:294–303.
21. Park IH, Kim JJ, Ku J, et al. Characteristics of social anxiety from virtual interpersonal interactions in patients with schizophrenia. *Psychiatry* 2009; 72:79–93.
22. Park KM, Ku J, Park IH, et al. Improvement in social competence in patients with schizophrenia: a pilot study using a performance-based measure using virtual reality. *Human Psychopharmacology* 2009; 24:619–627.

Address correspondence to:
 Dr. Wim Veling
 Parnassia Psychiatric Institute
 Lijnbaan 4
 2512 VA The Hague
 The Netherlands

E-mail: w.veling@parnassia.nl