University of Groningen

Communicative abilities in young children with a significant cognitive and motor developmental delay

Dhondt, Ann; Van keer, Ines; van der Putten, Annette; Maes, Bea

Published in:
Journal of Applied Research in Intellectual Disabilities

DOI:
[10.1111/jar.12695](https://doi.org/10.1111/jar.12695)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Dhondt, A., Van keer, I., van der Putten, A., & Maes, B. (2020). Communicative abilities in young children with a significant cognitive and motor developmental delay. *Journal of Applied Research in Intellectual Disabilities*, 33(3), 529-541. <https://doi.org/10.1111/jar.12695>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

ORIGINAL ARTICLE

JARID

Journal of Applied Research in Intellectual Disabilities

bild

WILEY

Communicative abilities in young children with a significant cognitive and motor developmental delay

Ann Dhondt¹ | Ines Van keer¹ | Annette van der Putten² | Bea Maes¹

¹Faculty of Psychology and Educational Sciences, Research Unit Parenting and Special Education, Catholic University of Leuven, Leuven, Belgium

²University of Groningen, Department of Special Needs Education and Youth Care, Groningen, The Netherlands

Correspondence

Ann Dhondt and Ines Van keer,
Vanderkelenstraat 32, bus 3765, 3000
Leuven, Belgium.
Emails: ann.dhondt@kuleuven.be (A. D.);
ines.vankeer@kuleuven.be (I. V.)

Funding information

Research Foundation (Flanders, Belgium),
Grant/Award Number: ZKC5734

Abstract

Background: Children with a significant cognitive and motor developmental delay are pre-symbolic communicators. The primary aim of this study was to reveal the variability within the communicative functioning of this group of children in terms of communication level, the reasons to communicate and behavioural expressions.

Methods: Twenty-six children between 14 and 58 months with a significant cognitive and motor developmental delay were recruited. The Communication Matrix of Rowland (2011, *Communication Disorders Quarterly*, 32, 190) was used to integrate different sources of information on the children's communicative functioning.

Results: These children primarily communicated at the level of pre-intentional and intentional behaviour, aimed at refusing, obtaining and, to a lesser extent, social purposes.

Conclusions: To develop or adapt early intervention strategies, and to monitor progress in communicative development, an even more nuanced view on these children's communicative utterances in terms of frequency, duration, idiosyncrasy and context relatedness is needed.

KEYWORDS

communication, communicative behaviours, communicative functions, developmental delay, intellectual disability, motor disability, profound intellectual and multiple disabilities

1 | INTRODUCTION

During early childhood, communicative skills develop rapidly from the pre-linguistic to the linguistic phase. In this latter phase, children acquire language skills starting with single words and evolving to the use of full sentences (Berk, 2007; Golinkoff, 2013). Concurrently, communication progresses from pre-intentional to intentional, while the communicative behaviours change from idiosyncratic (only understood by those who know the child well) to conventional and referential (Bates, Camaioni, & Volterra, 1975; Brady et al., 2012; Rowland, 2011; Siegel-Causey & Bashinski, 1997). In Bates et al. (1975), the most remarkable shift is described as the transition from perlocutionary to illocutionary acts. McLean and Snyder-McLean

(1987) describe perlocutionary acts as those "that produce an effect on the receiver" and illocutionary acts as those "that express a communicative intent of the speaker". Further differentiating these global shifts in communicative functioning, Rowland (2011) describes seven levels of communicative development (I-VII), where the transition from level II to level III encompasses the shift from the perlocutionary to the illocutionary acts (Bates et al., 1975). Level I refers to pre-intentional behaviour, where adults assign purpose to the child's actions, which shapes later intention. Level II is described as the level of "intentional behaviours," but these actions are not yet communicative. An example is touching an object of interest. Intentional communication emerges at Level III and is pre-symbolic and non-conventional. At Level IV, communication is still pre-symbolic, but has evolved from idiosyncratic to more conventional. An example of a behaviour at this level would be holding up the palm

Dhondt and Van keer are Joint first authors.

of the hand to request an object. Level V, often bypassed in typically developing children, refers to symbolic communication, where the child communicates through concrete tangible representations, whereas at level VI abstract symbols are used. At the highest level (VII), children combine two or three abstract symbols and eventually make use of language to communicate their messages.

In children with developmental disabilities, the communicative development is often disturbed, which results in a delayed and/or different development of communicative skills (e.g., Abbeduto, Warren, & Conners, 2007; Brady, Marquis, Fleming, & McLean, 2004; Grove, Bunning, Porter, & Olsson, 1999; Hostyn & Maes, 2009; Roberts, Price, & Malkin, 2007; Visser, Vlaskamp, Emde, Ruiter, & Timmerman, 2017). In persons with profound intellectual and multiple disabilities (PIMD; Nakken & Vlaskamp, 2007), communicative development is challenged even more due to the complex interplay between their cognitive, motor and often additional (e.g., sensory) limitations (Ogletree, Wetherby, & Westling, 1992; Olsson, 2005). The communicative abilities of persons with PIMD are described as primarily pre- or protosymbolic, including many idiosyncratic and subtle behaviours. Their movements are not always under voluntary control, which makes it difficult to correctly interpret behaviour as potentially communicative (Goldbart, 1994; Ogletree, Bartholomew, Wagaman, Genz, & Reisinger, 2012; Olsson, 2005). Therefore, communication partners consistently need to use contextual information and prior knowledge of the person to interpret their communicative utterances (Goldbart, Chadwick, & Buell, 2014; Grove et al., 1999; Hostyn, Daelman, Janssen, & Maes, 2010; Petry, Maes, & Demunyk, 2004; Vlaskamp & Oxener, 2002; Weis, 2014). However, even if the communication partner knows the person well, perlocutionary acts might not be recognized and illocutionary acts might not be interpreted as such. The lack of appropriate responses from communication partners potentially impedes further development (Olsson, 2005). Persons with PIMD also experience difficulties in exploring the environment, which, for example, hinders their ability to provide communicative signals indicating a focus of interest (Markova, 1990).

In short, limited by cognitive, motor and sensory impairments, persons with PIMD often communicate on a pre-symbolic level. This gives the impression of limited variability in this group with regard to their communicative functioning. However, pre-symbolic communicators can function on different levels of pre-symbolic communication (from pre-intentional to conventional), can be driven by different reasons to communicate and can differ in their behavioural expressions (Rowland, 2011). Furthermore, heterogeneity regarding communicative abilities is found to lead to a variety of individual growth curve outcomes (Brady et al., 2004). Therefore, in the present study, we wanted to investigate the variability in communicative abilities of young children (age < 5 years) with a significant cognitive and motor developmental delay.

The primary aim of this study was to reveal the communicative variability within this group of children in terms of communication level, reasons for communicating and behavioural expressions. At the moment, knowledge on the communicative abilities of these

children is very scarce. The secondary aim was to address the relation between communicative functioning and three child characteristics. First, we were interested in the relation with chronological age as communicative skills typically develop with age (Berk, 2007; Roberts et al., 2007). A second child characteristic of interest is the severity of the motor problems. Previous research revealed an influence of intellectual disabilities as well as motor impairments on the acquisition and occurrence of communicative skills. More specifically, it has been found that the severity of the impairments is positively related to communicative difficulties (Bhat, Galloway, & Landa, 2012; Houwen, Visser, van der Putten, & Vlaskamp, 2016; Lipscombe et al., 2016; Pennington, 2012; Voorman, Dallmeijer, Van Eck, Schuengel, & Becher, 2010). We hypothesize that children with more severe motor disabilities will have more limited communicative abilities (Cobo-Lewis, Oller, Lynch, & Levine, 1996; Gernsbacher, Stevenson, Khandakar, & Goldsmith, 2008; Iverson, 2010; Karasik, Tamis-LeMonda, & Adolph, 2011). Thirdly, the relation between communicative functioning and sensory impairments has been investigated as earlier research revealed the impact of these impairments on communicative behaviours (Bigelow, 2003; Evenhuis, Theunissen, Denkers, Verschuure, & Kemme, 2001; Tröster & Brambring, 1992)). For example, children with a significant visual impairment might have a more restricted repertoire of facial expressions (Tröster & Brambring, 1992).

Summarized, the two research questions to be answered in this manuscript are as follows:

1. Which communicative abilities are shown by young children with a significant cognitive and motor developmental delay in terms of communication level, reasons for communicating and behavioural expressions?
2. What is the relation between chronological age, motor functioning and sensory impairments on the one hand and the communicative abilities of this group of children on the other hand?

2 | METHOD

2.1 | Participants

In this study, children were included when they met the following criteria: (a) age between 6 months and 59 months; (b) severe cognitive delay characterized by children functioning at a quarter of their chronological age or lower on the Tandemlijst (Stadeus, Windey, Vermier, & Van Driessche, 1994) and (c) severe motor dysfunctions operationalized by children functioning at level IV or V (or level III for children below 2 years old) on the Gross Motor Function Classification System (GMFCS; Palisano, Rosenbaum, Bartlett, & Livingston, 2007). Children were not excluded when having additional problems (e.g., visual impairment) and/or by means of the cause of the developmental delay. The criteria that have been used are in line with the criteria of PIMD in adulthood, since the included children are expected to meet the criteria of individuals with PIMD

as described by Nakken and Vlaskamp (2007) later on in life. At this young age, however, the level of the intellectual disability has often not yet been determined and it is not easy to predict the developmental outcomes of these children when they grow older.

Children were recruited through various organizations and facilities (e.g., hospitals, diagnostic centres, early intervention teams, parent groups) in Flanders and the Netherlands. We contacted them via telephone and/or email and sent information about the project and the inclusion criteria. When children met the inclusion criteria, the researcher informed the parents or legal guardian about the project during a telephone contact and provided additional information by means of a folder and a website about the project (www.ojko.be; Dutch only). The parents or legal guardian gave written consent for participation of the children.

Twenty-six children between 14 to 58 months old with a significant cognitive and motor developmental delay participated in the study. Half of the children are male participants. Most of the children ($n = 25$) have additional visual and/or health problems. Table 1 presents detailed information about the participants.

2.2 | Procedure

Researchers visited the children at their home and/or day care facility, except for two children living in full-time residential provision. As this research was part of a broader project on the functioning of young children with a significant cognitive and motor developmental delay on different developmental domains, an extensive assessment battery was used. Therefore, the administrations were divided over two visits, which took place within a 2-week period. Observations, tests and questioning the primary caregiver were alternated to avoid children being overloaded. In case the children showed any signs of distress, observations and tests were disrupted and postponed to a later visit within 2 weeks. In this study, the information from three different observational protocols, one questionnaire and two semi-structured interviews with caregivers was integrated and used to obtain a clear overview of the children's communicative functioning (cf., by using this information to fill in the Communication Matrix). Detailed information on all of these instruments is presented in the next section ("Instruments"). This study was performed in accordance with the guidelines of the Ethical Committee of the Faculty Psychology and Educational Sciences at the University of Leuven and reported to the Privacy Commission in Belgium.

2.3 | Instruments

2.3.1 | Communication Matrix

The Communication Matrix of Rowland (2011) is an assessment tool designed to develop communicative profiles of individuals who are at the earliest stages of communicative development. The Matrix provides a clear overview of the expressive communicative abilities of

TABLE 1 Overview of participant characteristics as reported by a primary caretaker ($n = 26$)

Variable	Range (Mean)	n (%)
Age (in months)	14–58 (36.12)	
Sex		
Male		13 (50.00)
Female		13 (50.00)
Aetiology		
Acquired brain injury		2 (7.69)
Drug and alcohol abuse during pregnancy		1 (3.85)
Epilepsy		1 (3.85)
Genetic disorder		11 (42.31)
Infection during pregnancy		1 (3.85)
Lissencephaly and epilepsy		1 (3.85)
Perinatal asphyxia		2 (7.69)
Perinatal asphyxia and genetic disorder		1 (3.85)
Unknown		6 (23.08)
Visual functioning ^a		
Normal vision		13 (50.00)
Visual impairment		8 (30.77)
Blind		2 (7.69)
Unknown		3 (11.54)
Motor functioning ^b		
<0.5	0.04–1.67 (0.60)	12 (48.00)
0.5 to < 1		8 (32.00)
1 to < 1.5		3 (12.00)
≥1.5		2 (8.00)
Health problems		
Epilepsy		19 (73.08)
Gastrointestinal problems		14 (53.85)
Respiratory problems		13 (50.00)
Heart problems		2 (7.69)
Feeding tube		14 (53.85)

^aWhen defining visual functioning, the use of glasses is taken into account ($n = 3$). Caregivers were given the possible options regarding their children's visual functioning in the questionnaire by means of checkboxes. No explicit operational definitions of the categories were given.

^bMotor functioning is operationalized by the mean score (on a total of 2) on a questionnaire based on the motor questions of the Portage Program. Information on motor functioning of only 25 children is presented due to missing data.

the individual with the emphasis on what they can do (i.e., "the functional use of communication"). As all the children in this study had significant cognitive and motor developmental delays and often did not reach the level of intentional communication, the Communication

Matrix provided the most appropriate perspective to study the behavioural expressions of these children. The Communication Matrix is based on research on typically developing infants between 0 and 24 months (Rowland, 2011). It is structured around seven levels of communication development (pre-intentional behaviour, intentional behaviour, unconventional behaviour, conventional communication, concrete symbols, abstract symbols and language) and four global reasons to communicate (refuse things, obtain things, engage in social interaction and seek/provide information). Twenty-four states (at level I), functions (at level II) or intents (at level III to VII) are binary questioned throughout the Matrix, all corresponding with one of the four reasons to communicate. When answering “yes,” more specific communicative behaviours can be selected by choosing from nine behavioural categories (body movements, early sounds, facial expressions, visual behaviour, simple gestures, conventional gestures/vocalizations, concrete symbols, abstract symbols and language). An overview of the 24 states, functions, and intents by level and global reasons to communicate is provided in Table 2. A communicative state/function/intent is regarded as surpassed when a child masters the communicative state/function/intent on a subsequent level (e.g., a child has surpassed the function “A1: Expresses discomfort” when he/she masters the function “B1: Protests”; see Figure 1). The scoring system results in a one-page profile consisting of 80 cells (cf., Figure 1), representing all possible combinations of states, functions and intents within the seven levels of communication. Rowland (2011) clarifies that “where multiple behaviours are used to express a given intent at a given level, the cell is shaded according to the highest level (mastered or emerging) at which any behaviour in that cell is

coded” (p. 194). Therefore, an additional overview of the communicative behaviours used by the child to communicate the function (e.g., a child cries to express discomfort) can be composed. Also, a total score (from 0 to 160) can be computed by awarding 2 points to a cell that is checked as mastered or surpassed and by awarding 1 point to a cell that is checked as emerging. Even though the Communication Matrix is seldom used for research purposes (Rowland, 2011), previous studies have proven the added value of the Communication Matrix in research on individuals with language delays (e.g., Hategan & Talaş, 2014; Parker, 2009; Rowland & Schweigert, 2000) and with various types of disabilities (Rowland, 2011).

2.3.2 | Sources of information

Due to the idiosyncratic behaviours and fluctuating performance levels of children with a significant cognitive and motor developmental delay, it is a challenge to get an objective and representative view on their communicative functioning. Therefore, in our study, the Communication Matrices were completed by a researcher, based on multiple sources of information collected during home visits. The specific instruments (i.e., sources of information) were chosen based on their previous or possible application in this study's target group as well as on the variability in instrument type (i.e., observation, questionnaire or interview), degree of structure and freedom to support/encourage child behaviour (i.e., highly structured, semi-structured or free) and type of interaction partner (i.e., familiar caregiver or unfamiliar researcher).

Level	Reasons to communicate			
	Refuse	Obtain	Social	Information
I	A1. Expresses discomfort	A2. Expresses comfort	A3. Expresses interest in people	
II	B1. Protests	B2. Continues an action B3. Obtains more	B4. Attracts attention	
III-VII	C1. Refuses or rejects	C2. Requests more action C3. Requests new action C4. Requests more object C5. Makes choices C6. Requests new object	C8. Requests attention C9. Shows affection	
IV-VII			C10. Greets people C11. Offers/shares C12. Directs attention C13. Uses polite forms	C14. Answers Yes/No C15. Asks questions
V-VII		C7. Requests absent objects		C16. Names things C17. Comments

TABLE 2 The 24 states (A), functions (B) and intents (C) by level and reason to communicate as defined in the Communication Matrix

Note: Reprinted from Rowland (2011).

Level1 Pre-Intentional Behaviour	A1 Expresses Discomfort	A2 Expresses Comfort					A3 Expresses Interest in Other People										
Level2 Intentional Behaviour	B1 Protests	B2 Continues Action		B3 Obtains More of Something			B4 Attracts Attention										
Level3 Unconventional Communication	C1 Refuses, Rejects	C2 Requests More Action	C3 Requests New Action	C4 Requests More Object	C5 Makes Choices	C6 Requests New Object	C8 Requests Attention	C9 Shows Affection									
Level4 Conventional Communication	C1 Refuses, Rejects	C2 Requests More Action	C3 Requests New Action	C4 Requests More Object	C5 Makes Choices	C6 Requests New Object	C8 Requests Attention	C9 Shows Affection	C10 Greets People	C11 Offers, Shares	C12 Direct Your Attention	C13 Polite Social Forms	C14 Answers Yes/No Questions	C15 Asks Questions			
Level5 Concrete Symbols	C1 Refuses, Rejects	C2 Requests More Action	C3 Requests New Action	C4 Requests More Object	C5 Makes Choices	C6 Requests New Object	C7 Requests Absent Objects	C8 Requests Attention	C9 Shows Affection	C10 Greets People	C11 Offers, Shares	C12 Direct Your Attention	C13 Polite Social Forms	C14 Answers Yes/No Questions	C15 Asks Questions	C16 Names Things/ People	C17 Makes Comments
Level6 Abstract Symbols	C1 Refuses, Rejects	C2 Requests More Action	C3 Requests New Action	C4 Requests More Object	C5 Makes Choices	C6 Requests New Object	C7 Requests Absent Objects	C8 Requests Attention	C9 Shows Affection	C10 Greets People	C11 Offers, Shares	C12 Direct Your Attention	C13 Polite Social Forms	C14 Answers Yes/No Questions	C15 Asks Questions	C16 Names Things/ People	C17 Makes Comments
Level7 Language	C1 Refuses, Rejects	C2 Requests More Action	C3 Requests New Action	C4 Requests More Object	C5 Makes Choices	C6 Requests New Object	C7 Requests Absent Objects	C8 Requests Attention	C9 Shows Affection	C10 Greets People	C11 Offers, Shares	C12 Direct Your Attention	C13 Polite Social Forms	C14 Answers Yes/No Questions	C15 Asks Questions	C16 Names Things/ People	C17 Makes Comments
	Refuse	Obtain					Social					Information					

FIGURE 1 Communication Matrix (Rowland, 2018) [Colour figure can be viewed at wileyonlinelibrary.com]

First, an adapted version of the Early Social Communication Scales (ESCS; Mundy et al., 2003) was used. The ESCS is a videotaped standardized observation measurement to elicit early non-verbal communication skills, more specifically joint attention, behavioural requests and social interaction (Mundy et al., 2003). The adapted protocol is available upon request and mainly encompasses an abridgement of the original protocol, motivated by the severity of the disabilities and the limited attention span of the target group. Second, parents or professional caretakers filled in the Communicative and Symbolic Behavior Scales (CSBS; Wetherby & Prizant, 2002), a questionnaire about the communicative behaviour of the child. It consists of questions about seven clusters: emotion and eye gaze, communication, gestures, sounds, words, understanding and object use. The questions can be answered on a 3-point Likert scale indicating if a skill is not yet mastered or whether a mastered skill is used sometimes or frequently (Wetherby & Prizant, 2002). Third, a brief semi-structured conversation was conducted with the informants to gather information on how the child reacts when he/she likes/does not like something and what he/she does when he wants/does not want something. Fourth, the three communicative scales (emotional communication, receptive language and general communicative behaviour) as well as the category "additional information" of the Behavioral Appraisal Scale (BAS) provided information on the communicative functioning of the children. This instrument is a combination of observation, testing and questioning people who are familiar with the child (Vlaskamp, Van der Meulen, & Smrkovsky, 1999). Fifth, an interview about the social and emotional development was conducted by means of the revised Scale for Social

and Emotional Development (SEO-R; Claes, & Verduyn, 2012). A proxy was questioned about 13 domains in which children's social and emotional abilities are discussed (e.g., handling your own body, dealing with emotionally important persons, communication, emotion differentiation). Finally, a free-play interaction between a child and a familiar caregiver of approximately 15 min was used to observe spontaneous communicative skills in the children (based on Mahoney, 1998; 2008).

We combined information provided by persons who are very familiar with the child (i.e., parents or professional caregivers who experience the child's functioning throughout different situations and time points) as well as by researchers who were able to observe the child relatively unbiased since they have had no previous experience with the child. Next to different perspectives, information on the child was collected in three different situations by looking at a strictly structured observation protocol in which each child was given the same instructions and feedback (cf., ESCS), a semi-structured observation protocol in which the child was given the maximum possibility, encouragement and support to achieve the provided tasks (cf., BAS) and finally an unstructured play interaction. Throughout these tasks, the child was observed in interaction with at least one known (i.e., caregiver) and at least one unknown (i.e., researcher) person.

2.3.3 | Child characteristics

General information on child, family and contextual factors was collected by means of a researcher-developed questionnaire on basic

demographic and health-related information, including the variables of interest in the second research question: child's age as well as visual functioning (within three categories: normal vision, visual impairment and blind) with or without the use of glasses, as reported by a primary caretaker and presented in Table 1. Initially, the aim was to take the auditory functioning of the children also into account. As only two of the caregivers reported any degree of hearing loss in the questionnaire, this variable could not be incorporated.

The children's motor abilities were assessed by means of a separate questionnaire, based on the motor questions of the Portage Program and focused on the gross and fine motor abilities of young children (Hoekstra et al., 2011). The questionnaire consists of 145 items scored on a three-point scale: score 2 when a child masters the skill, score 1 when a child is almost mastering the skill and score 0 when a child does not master the skill. The items in this questionnaire range from questions about if the child can move his or her head, to standing or moving independently, or manipulating a toy with one or two hands. For each child, a total score was calculated by adding up the item scores and a mean score was calculated by dividing this total score by 145 (as presented in Table 1). Therefore, this mean score reflects the global motor functioning of the child, on both the gross and fine motor domain. Globally, children with the lowest mean score (0–0.5) are developing towards turning their head and obtaining some control over upper limbs (e.g., turning head or moving arm towards stimulus). Children with a mean score between 0.5 and 1 show a development towards sitting independently for a short period of time and using their upper limbs in a more controlled way (e.g., touching and holding objects). Mean scores between 1 and 1.5 are seen in children that are developing towards being able to move independently, standing with support and using their upper limbs in a more exploratory way (e.g., pushing and taking objects). A mean score of 2 would mean that the child can walk independently and uses upper limbs in a more functional way (e.g., picking up a toy and putting it in a box), which is not the case for any of the participants.

2.4 | Data processing and interrater reliability

The online version of the Communication Matrix (Rowland, 2018) was used to integrate the aforementioned sources of information (cf., "Sources of information") by means of a researcher-developed protocol (available upon request). The first part of the protocol consisted of general guidelines (e.g., score conservative when doubting) and steps that needed to be followed to fill in the Matrix (e.g., check the behaviours first and only then conclude if a state/function/intent is not used, emerging or mastered). The second part of the protocol contained the coding rules to decide whether a function is not used, emerging or mastered. A child was considered as mastering a skill when it was observed or described in minimum two of the instruments. When a communicative skill was only observed or described in one instrument or it was observed in two instruments but the skill was not fully present (e.g., a child sometimes alternates between an object and a communication

partner), it was coded as emerging. And finally, a skill was considered "not used" when it was not displayed in one of the instruments or when it was observed in one instrument but the skill was not fully present.

Two independent research assistants conducted the integration of the information into the Communication Matrix. The first author explained the coding protocol and trained the research assistants until they reached a minimum of 90% exact interrater agreement. After training, each research assistant completed the communication profiles of 13 participants. To assess the reliability of the coding protocol, the first author double-coded 40%, randomly selected, of the participants (20% per research assistant). The exact agreement was determined by dividing the number of agreements by the total number of items multiplied by 100. This resulted in an agreement of 95.1% and 97.2%, which indicated a good reliability (Cordes, 1994). Cohen's kappa for both research assistants was 0.724, which is substantial (Cohen, 1988).

2.5 | Data analysis

First, descriptive analyses were conducted to gain insight into the communicative abilities of the children. Therefore, we calculated the number of participants who showed either unused, emerging or mastered communicative states, functions or intents at the associated communicative levels. Also, we calculated the number and percentage of participants who showed specific behavioural expressions. These behaviours are predefined in the Communication Matrix as mentioned before (Rowland, 2011) and exclusively linked to specific communication levels as well as specific states, functions or intents. We did not distinguish between emerging or mastered behavioural expressions, but combined (i.e., added up) the number of participants. The denominator in calculating the percentages corresponded to the number of participants who used the specific states/functions/intents.

Second, the relationship of children's communicative functioning (total score on the Communication Matrix) with chronological age (in months) and motor functioning (total score on the motor questionnaire) was addressed using the Spearman's rho correlation test. We excluded one participant from the analysis due to missing data. We opted for a non-parametric test because of the small sample size and the presence of non-normality in the data. Additionally, bootstrapping (a resample method with replacement) was used to determine confidence intervals for these correlations (Lee & Rodgers, 1998).

The relationship with visual functioning was addressed using a Mann-Whitney *U* test, distinguishing children with normal vision ($n = 13$) and impaired vision or blindness ($n = 10$). Since visual functioning was unknown in three of the children, they were excluded from the analysis.

3 | RESULTS

Preliminary analyses revealed no significant differences between male ($n = 13$) and female ($n = 13$) participants with regard to their communicative functioning as well as their motor functioning, visual

FIGURE 2 Number of participants showing communicative states/functions/intents according to the level of mastering. Note. As is specified in Table 2, communicative states/functions/intents can be shown on different communicative levels. In this figure, the associated levels are only split up when the results vary over different levels. This is the case for C1, C2, C9, C10 and C12 [Colour figure can be viewed at wileyonlinelibrary.com]

functioning and chronological age. Therefore, data of both groups were combined in subsequent analyses.

3.1 | Descriptive analyses

Figure 2 provides an overview of the number of participants who were showing specific communicative states, functions or intents according to the level of mastering. In Table 3, an overview of specific communicative behaviours that were used by all participants is presented.

At level 1 (pre-intentional behaviour), most children mastered the three accompanying states. All 26 children “expressed discomfort” and “expressed comfort.” The state “expresses interest in other people” was mastered in 16 children and emerging in 7 children, which means it was not expressed by 3 of the participants. To express (dis)comfort, nearly all children used body movements as well as early sounds and facial expressions. However, body movements were seen in more children in relation to discomfort (96%) than comfort (73%), especially changes in posture (69% vs. 35%). Limb movements were also often observed for this purpose (i.e., by 46% and 42% of the children), while head movements were rarely observed (19% and 8%). Of the 23 children who expressed interest in other people, most used early sounds (91%) and facial expressions (87%) while body movements were less frequently used (22%).

More variation was noticed at level 2 (intentional behaviour). More specifically, “protesting” and “continuing an action” were emerging ($n = 10$ for both) or mastered ($n = 11$ resp. $n = 7$) by most participants while “obtaining more” and “attracting attention” were approximately equally distributed between “not used” ($n = 14$ and $n = 12$) and “emerging” or “mastered” ($n = 7 + 6$ and $n = 5 + 8$). Protesting was primarily expressed through body movements (71%) and early sounds (95%), but not often through facial expressions (29%). With regard to the body movements, 57% and 38% of the children used head and arm movements, respectively, while only a few children used leg movements (10%, $n = 2$) or move away from the person or object (5%, $n = 1$). Wanting an action to be continued was expressed by early sounds (82%) and facial expressions (65%), but not often by visual behaviour (35%) or movements (6%, $n = 1$). To obtain more of something, children used body

movements (75%), early sounds (67%) and visual behaviour (58%), but not often facial expressions (25%). Body movements primarily included approaching or taking the desired object (67% and 42%, respectively). When children attracted attention, they used sounds (79%) and visual behaviour (71%), while facial expressions (43%) and body movements (i.e., approaching a person, 21%) were used less often.

At levels III (unconventional communication) and IV (conventional communication), only nine of the children (35%) used one or more of the related intents. A few children refused or rejected something by using unconventional communication ($n = 7$, of which four children mastered this intent). For this purpose, all children used a simple gesture (i.e., pushing away the person or object), but in only one child this was accompanied by a facial expression and none of the children used related movements. In two of these seven children, the use of conventional communication to refuse or reject something was also emerging, which means they are learning “to shake their head deliberately for no.” Further, on level III, requesting more of an action was emerging in one child (who was learning to deliberately take the hand of the communication partner) and mastered in one other (who used leg movements and reached towards or tapped the communication partner). Showing affection through unconventional communication was present in four children (of which two children mastered this intent). Simple gestures (i.e., arm/hand movements and touching the communication partner) and facial expressions were mostly used ($n = 3$), followed by visual behaviour ($n = 2$) and early sounds ($n = 1$). Two other children showed affection through conventional communication by hugging, kissing or patting the communication partner. In one child, the ability to greet people by waving “hi” or “bye” (level IV) was emerging. Lastly, in three children, the ability to direct the communication partner’s attention to something by looking back and forth between the communication partner and an object, person or place (level IV) was emerging, with one child also learning to deliberately point at something. Intents and communicative behaviours at levels V to VII were not observed within the research group.

3.2 | Relation with child characteristics

To address the relation between the communicative abilities and child characteristics, a total score on the Communication Matrix for

TABLE 3 Number of participants using behaviours associated with specific states (A), functions (B) and intents (C)

Behaviours	States			Functions				Intents				
	A1.	A2.	A3.	B1.	B2.	B3.	B4.	C1.	C2.	C9.	C10.	C12.
	(n = 26)	(n = 26)	(n = 23)	(n = 21)	(n = 17)	(n = 12)	(n = 14)	(n = 7)	(n = 2)	(n = 6)	(n = 1)	(n = 3)
Body movements (level I-III)	25 (96%)	19 (73%)	5 (22%)	15 (71%)	1 (6%)	9 (75%)	3 (21%)	0	1 (50%)			
Approaches desired object						8 (67%)						
Approaches person							3 (21%)					
Whole-body movement								0	0			
Change in posture	18 (69%)	9 (35%)	2 (9%)									
Limb movements	12 (46%)	11 (42%)	3 (13%)									
Arm movements				8 (38%)	0	0	0	0	0			
Leg movements				2 (10%)	1 (6%)	1 (8%)	0	0	1 (50%)			
Head movements	5 (19%)	2 (8%)		12 (57%)	0	0	0	0				
Takes object						5 (42%)						
Moves away from person or object				1 (5%)								
Early sounds (level I-III) ^a	26 (100%)	26 (100%)	21 (91%)	20 (95%)	14 (82%)	8 (67%)	11 (79%)	0	0	1 (17%)		
Facial expressions (level I-III) ^b	25 (96%)	26 (100%)	20 (87%)	6 (29%)	11 (65%)	3 (25%)	6 (43%)	1 (14%)	0	3 (50%)		
Visual behaviour (level II-III) ^c					6 (35%)	7 (58%)	10 (71%)		0	2 (34%)		
Simple gestures (level III)								7 (100%)	2 (100%)	3 (50%)		
Pushes away object or person								7 (100%)				
Takes your hand									1 (50%)			
Reaches towards or taps you									1 (50%)			
Arm/hand movements										1 (17%)		
Touches you									0	2 (33%)		
Conventional gestures and vocalizations (level IV)								2 (29%)	0	2 (33%)	1 (100%)	3 (100%)
Shakes head no								2 (29%)				
Specific vocalizations								0				
Gives unwanted item to you								0				
Beckons you to come									0			
Holds hands up to you									0			
Nods head									0			
Hugs, kisses and pats you										2 (33%)		

(Continues)

TABLE 3 (Continued)

Behaviours	States			Functions				Intents				
	A1.	A2.	A3.	B1.	B2.	B3.	B4.	C1.	C2.	C9.	C10.	C12.
	(n = 26)	(n = 26)	(n = 23)	(n = 21)	(n = 17)	(n = 12)	(n = 14)	(n = 7)	(n = 2)	(n = 6)	(n = 1)	(n = 3)
Waves "hi" or "bye"	1 (100%)											
Points to something	1 (33%)											
Looks back and forth	3 (100%)											
Concrete symbols (level V)	0 0 0 0 0 0 0											
Abstract symbols (level VI)	0 0 0 0 0 0 0											
Language (level VII)	0 0 0 0 0 0 0											

Note: Specific behaviours as well as states/functions/intents can both be characterized as emerging or mastered. This table shows the number (and percentage) of participants for which the specific behaviours are either emerging or mastered (i.e., both numbers were added up). The denominator in calculating the percentages corresponds to the number of participants who used the specific states/functions/intents. Only the states/functions/intents that are used (i.e., emerging or mastered) by at least one participant are presented. Blank spaces indicate that the specific behaviours could not be chosen in combination with the specific communicative state, function or intent.

^aEarly sounds include crying, grunting, screaming, cooing, squealing, fussing and whining.

^bFacial expressions include grimacing, smiling and frowning.

^cVisual behaviour include looking at a (desired) object and looking at a person.

each child was calculated. The mean score was 10.3 ($SD = 3.83$), with scores ranging between 4 and 20 (on a total of 160). Correlational analysis (Spearman's rho) revealed a significant positive relation between total scores on the Communication Matrix and chronological age ($r_s = .391$, $p = .024$), 95% CI $[-0.035, 0.708]$. Secondly, we found that higher scores on the Communication Matrix were associated with higher scores on motor functioning ($r_s = .835$, $p = .000$, 95% CI $[0.651, 0.911]$). However, no significant correlation between chronological age and motor functioning was found ($r_s = .279$, $p = .089$). Using a Mann-Whitney U test, no significant relation between total scores on the Communication Matrix and visual functioning (normal vision vs. visual impairment/blindness) was found ($U = 52.00$, $p = .446$).

4 | DISCUSSION

4.1 | Conclusions

With regard to the first research aim, the study's results demonstrate that children with a significant cognitive and motor developmental delay primarily communicated at the level of pre-intentional and intentional behaviour, particularly aimed at refusing, obtaining and, to a lesser extent, social purposes. The low mean total score on the Communication Matrix indicates that these children scored very low on the acquisition of communicative functions that typically occur between 0 and 24 months of age (Rowland, 2011). Children used various communicative behaviours, depending on the communicated function, state or intent. This variety of communicative modalities is also found in the research of Iacono, Carter, and Hook (1998), although in older children with severe and multiple disabilities.

More specifically, at a pre-intentional level, the behaviour of all the children can be interpreted as expressing (dis)comfort, while more social communication (i.e., expressing interest in other people) was evident in most, but not all, children. All of these three states were primarily seen in early sounds and facial expressions. However, we observed that body movements were a very important way for these children to communicate comfort and especially discomfort, for which posture change was additionally important.

At the level of intentional behaviour, most children communicated to refuse something (in the form of "protesting") or, to a slightly lesser extent, to obtain something (i.e., a continuation of an action). Further, approximately half of the children (also) communicated to obtain more of something or for social purposes (i.e., attracting attention). Early sounds seemed to play a major role in communicating all of these four functions. Body movements were regularly used to protest and to obtain more of something, but seldom to continue an action or attracting attention. Facial expressions were not used by a lot of children in this regard, but did seem to play a role in trying to continue an action. In more than half of the children who showed these functions, visual behaviour was used to attract attention and obtain more of something.

By definition, children with a significant cognitive and motor developmental delay function on a pre-symbolic level, so we did not

expect them to master intents beyond level IV. However, even at the pre-symbolic level of intentional communication (level III and IV), children never communicated with the intent to give or receive information and rarely communicated to obtain something. If the children intentionally communicated on levels III and IV, protesting/refusing and social interactions were the reasons they would communicate for. Still, children with significant cognitive and motor developmental delay rely highly on their communication partners in order to “make meaning” together. They depend very much on the sensitivity and responsivity of their partners to contextualize their reactions in response to the situation (Van keer et al., 2017).

The second research question addressed the correlation between communicative functioning and three child characteristics. Correlational analyses showed that older children had a higher total score on the Communication Matrix, although this result needs to be interpreted carefully as the bootstrap 95% confidence interval included zero (Hesterberg, Monaghan, Moore, Clipson, & Epstein, 2003). Furthermore, it was found that children with better motor skills also had a higher total score on the Communication Matrix. As no significant correlation was found between chronological age and motor functioning, we can carefully conclude that both age and motor functioning have a unique relation with the children's communicative abilities. The strong relationship between the communicative and motor functioning can be partly explained by the interrelatedness of developmental domains in child development, especially in atypical populations (Diamond, 2000; Houwen et al., 2016; Roebbers & Kauer, 2009; Wang, Lekhal, Aarø, & Schjølberg, 2012). Particularly in the early stages of communicative development, severe motor impairments can impede children's communication as messages are often communicated through motor behaviour (e.g., body movements, head movements, simple gestures).

With regard to visual functioning, no significant association was found between communicative and visual functioning. It might be that visual behaviour is less decisive in (measuring) these earliest stages of communicative functioning, which is substantiated by the structure of the Matrix in which visual behaviour cannot be chosen in relation to the first communicative level. Also, it is possible that children with visual impairments express the same states/functions/intents through different behavioural modalities, and therefore, no differences in total scores can be found (Bigelow, 2003; Tröster & Brambring, 1992). Finally, for both sensory functions of vision and hearing, prevalence is known to be underestimated in persons with severe disabilities in general (Evenhuis et al., 2001; Nijs, Schouten, & Maes, 2019; Van Den Broek, Janssen, Van Ramshorst & Deen, 2006; Van Splunder, Stilma, Bernsen & Evenhuis, 2006). Therefore, the responses of the caregivers on these items in the questionnaire should be interpreted with caution.

It is important to note that all results are based on a general group analysis and that no further differentiation has been made based on the children's specific (additional) disabilities such as level of cognitive and motor functioning, and visual and hearing impairments as reported by the caregivers. We certainly acknowledge the possible influence of these characteristics, especially on the use of specific communicative behaviours. Subgroup or even individual analysis, despite

being challenged by difficulties in obtaining reliable differentiation as well as sample size issues, is an important area for future research.

4.2 | Strengths, limitations and future research

A major strength of this study is its integrative nature, by looking at different aspects of communicative functioning (i.e., levels, reasons and behavioural expressions) as well as combining several information sources. Even though previous studies already used several instruments to gain insight in the communicative abilities of children with a significant cognitive and motor developmental delay, these have rarely been integrated to obtain a comprehensive picture of their communicative functioning. In particular, information collected through proxies (questionnaires and interviews) has been alternated with observations made by the researcher. A major advantage of this type of data triangulation is that subjectivity and bias are reduced (Brady et al., 2012; Thurmond, 2001). Also, the influence of timing (e.g., an observation at a moment when the child has little attention or had an epileptic seizure) decreased because proxies give information on the daily functioning, while the researcher is more dependent on the time of the observation. To reduce the impact of the latter, the observations were made on different time points (unless this was not possible) but within a period of approximately 2 weeks.

Some limitations of the present study should be acknowledged. To start from a statistical point of view, the relatively small research sample affects the type of data analyses that could be applied and the generalizability of the results. A larger sample could make it possible to conduct other statistical analyses to reveal differences in the specific communicative behaviours of the children according to their limitations (e.g., children with and without visual impairments), and to identify subgroups of children. Nevertheless, considering the specific characteristics regarding our target group, the research sample can be assumed as rather large. Factors limiting sample size are, for example, the children's multiple disabilities, their young age, the vulnerability of the children and their caregivers, the difficulties in recruitment and the low visibility of this group. In addition, bootstrapping was conducted to cater this shortcoming and to find more robust and solid results regarding the correlation between communicative functioning and chronological age, motor functioning and visual functioning (Lee & Rodgers, 1998; Zhu, 1997).

Another limitation of the study is related to the study design. Longitudinal research (if possible with a larger sample) is needed to get a more comprehensive representation of the communicative development of these children whereby the developmental sequence (e.g., of communicative reasons, behaviours,...) and possible critical periods can be identified, which can help to give direction in individual support plans. Additionally, it would have been interesting to study other child characteristics and contextual factors as this has proven its importance for child development (e.g., Berk, 2007; Kahn, 1996; Stephenson & Dowrick, 2005). For example, the importance of sensitivity and responsivity of the primary caregiver is addressed in previous research (De Bal, 2011; Van keer et al., 2017; Warren & Brady, 2007). Also, the

development of other domains (e.g., motor development) can play an important role in the support of these children. It can be assumed that children put energy in one of the developmental domains and make a progress in that area, where the other domains stabilize or even decline. Or, as Wang et al. (2012) pointed out that the relationship between communication and motor skills is rather multifaceted instead of directional.

A final remark can be made regarding the theoretical background and structure of the Communication Matrix in relation to our specific target group. Since each profile was unique, the Matrix allowed us to identify meaningful variability in the children's communicative functioning. However, the Matrix consists of a top-down procedure in which specific communicative behaviours are predefined and exclusively linked to specific levels of communication, states/functions/intents and reasons to communicate. All of this is defined based on research in typically developing children (Rowland, 2011). Therefore, it is possible that the communicative functioning of these children is to some extent overrated (i.e., due to biased expectations based on the predefined behavioural categories), underrated (i.e., because certain meaningful behaviours are possibly not part of the predefined categories) and/or oversimplified (i.e., because a myriad of different behavioural expressions are possibly taken together in one predefined category). We know that these children often make use of idiosyncratic signals (Daelman, 2003), for example through changes in muscle tone, which are often strongly person- or context-bound and are easily misinterpreted or ignored (Grove et al., 1999). Even when recognized, the Matrix does, for example, not allow the registration of posture changes beyond level I. Further, the Matrix does not take into account differences in frequency, duration and context relatedness of communicative utterances. Also, by shading a cell according to the highest level at which any behaviour in that cell is coded, individuals with very limited means of expression "may have a similar profile to someone with an extensive repertoire of behaviours" (Rowland, 2011, p. 194). Additionally, children in our target group may not only show a delayed, but also a different developmental trajectory (Vlaskamp, 2011). This renders the use of a "surpassed"-category questionable. To conclude, the Communication Matrix is a very useful instrument within the studied target group to establish a general estimation of the children's functional use of communication; however, we suspect there is still a lot more variability that is not elucidated by this instrument as it is currently used. A more detailed view on this variability is especially necessary when adopting a longitudinal viewpoint aimed at mapping these children's (often slow) development, since their developmental steps may need to be defined in terms of "broadening" communicative skills (i.e., in terms of frequency, duration and context relatedness) rather than solely in terms of learning new skills.

4.3 | Relevance of the study

The present study supports the idea that children with a significant cognitive and motor developmental delay vary in their communicative functioning although most of them particularly function on a

pre-intentional level. The acknowledgement of this variability is important for early intervention strategies, which need to consider both the general description of the communicative abilities of these children and their uniqueness. Moreover, the detailed description of the children's communicative functioning can provide early interventionist strategies in stimulating the communicative development of this vulnerable group of children (Brady et al., 2012). This can, in its turn, result in positive outcomes on other developmental domains as communication is widely recognized as a base for general development (Bukatko & Daelher, 2004; Fogel, 1992; Stephenson & Dowrick, 2005; Vygotsky, 1978). As Brady et al. (2016) state in their Guidance for Assessment and Intervention (2016), assessment and intervention are often intertwined and co-occurring. However, assessment should preferably be dynamic, in order to identify the individual's potential to learn new skills when provided with appropriate support, which in turn should lead to setting and evaluating new goals.

ACKNOWLEDGMENTS

We gratefully thank the Research Foundation (Flanders, Belgium), Steunfonds Marguerite-Marie Delacroix (Belgium) and JKF Kinderfonds (The Netherlands) for their financial support for this project. Also, we are grateful to Gertrud Schalen, Kristien Hermans, Lien Vanderkerken, Minou Nauta, Riëtte Krijgsheld and Stephy Colla for their contribution to the broader project in which this study is embedded. We are especially grateful to Stephy Colla for initiating this study and to Lien Vanderkerken for her helpful input during both conducting and writing this manuscript. We also thank the research assistants for coding the information into the communication matrix. Lastly, we would like to thank all participating families for their time, effort and enthusiasm.

ORCID

Ann Dhondt <https://orcid.org/0000-0003-1855-8718>

Ines Van keer <https://orcid.org/0000-0002-7848-1276>

REFERENCES

- Abbeduto, L., Warren, S. F., & Conners, F. A. (2007). Language development in Down syndrome: From the prelinguistic period to the acquisition of literacy. *Mental Retardation and Developmental Disabilities Research Reviews*, 13, 247–261. <https://doi.org/10.1002/mrdd.20158>
- Bates, E., Camaioni, L., & Volterra, V. (1975). The acquisition of performatives prior to speech. *Merrill-Palmer Quarterly of Behavior and Development*, 21, 205–226.
- Berk, L. E. (2007). *Development through the lifespan* (4th ed.). Boston, MA: Pearson.
- Bhat, A. N., Galloway, J. C., & Landa, R. J. (2012). Relation between early motor delay and later communication delay in infants at risk for autism. *Infant Behavior and Development*, 35, 838–846. <https://doi.org/10.1016/j.infbeh.2012.07.019>
- Bigelow, A. E. (2003). The development of joint attention in blind infants. *Development and Psychopathology*, 15, 259–275. <https://doi.org/10.1017/S0954579403000142>
- Brady, N. C., Bruce, S., Goldman, A., Erickson, K., Mineo, B., Ogletree, B. T., ... Wilkinson, K. (2016). Communication services and supports for individuals with severe disabilities: Guidance for

- assessment and intervention. *American Journal on Intellectual and Developmental Disabilities*, 121(2), 121–138. <https://doi.org/10.1352/1944-7558-121.2.121>
- Brady, N. C., Fleming, K., Thiemann-Bourque, K., Olswang, L., Dowden, P., Saunders, M. D., & Marquis, J. (2012). Development of the communication complexity scale. *American Journal of Speech-Language Pathology*, 21, 16–28. [https://doi.org/10.1044/1058-0360\(2011/10-0099\)](https://doi.org/10.1044/1058-0360(2011/10-0099))
- Brady, N. C., Marquis, J., Fleming, K., & McLean, L. (2004). Prelinguistic predictors of language growth in children with developmental disabilities. *Journal of Speech, Language, and Hearing Research*, 47, 663–677. [https://doi.org/10.1044/1092-4388\(2004/051\)](https://doi.org/10.1044/1092-4388(2004/051))
- Bukatko, D., & Daelher, M. W. (2004). *Child Development: A thematic approach* (5th ed.). New York, NY: Houghton Mifflin Company.
- Claes, L., & Verduyn, A. (red.) (2012). *Schaal voor emotionele ontwikkeling bij mensen met een verstandelijke beperking – revised*. Antwerpen, Belgium: Garant.
- Cobo-Lewis, A. B., Oller, D. K., Lynch, M. P., & Levine, S. L. (1996). Relations of motor and vocal milestones in typically developing infants and infants with Down syndrome. *American Journal of Mental Retardation*, 100, 456–467.
- Cohen, J. (1988). *Statistical power analysis for the behavioral sciences* (2nd ed.). Hillsdale, NJ: Lawrence Erlbaum Associates.
- Cordes, A. K. (1994). The reliability of observational data: I. Theories and methods for speech-language pathology. *Journal of Speech and Hearing Research*, 37, 264–279. <https://doi.org/10.1044/jshr.3702.264>
- Daelman, M. (2003). *Een analyse van de presymbolische communicatie bij blinde kinderen met een meervoudige handicap*. Unpublished doctoral thesis, Katholieke Universiteit Universiteit Leuven, Afdeling Orthopedagogiek, Leuven.
- De Bal, C. (2011). Communicatie. In I. B. Maes, C. Vlaskamp, & A. Penne (red.), *Ondersteuning van mensen met ernstige meervoudige beperkingen: Handvatten voor een kwaliteitsvol leven* (pp. 145–164). Leuven, Belgium: Acco.
- Diamond, A. (2000). Close interrelation of motor development and cognitive development and of the cerebellum and prefrontal cortex. *Child Development*, 71, 44–56. <https://doi.org/10.1111/1467-8624.00117>
- Evenhuis, H. M., Theunissen, M., Denkers, I., Verschuure, H., & Kemme, H. (2001). Prevalence of visual and hearing impairment in a Dutch institutionalized population with intellectual disability. *Journal of Intellectual Disability Research*, 45(5), 457–464. <https://doi.org/10.1046/j.1365-2788.2001.00350.x>
- Fogel, A. (1992). Movement and communication in human infancy: The social dynamics of development. *Human Movement Science*, 11, 387–423. [https://doi.org/10.1016/0167-9457\(92\)90021-3](https://doi.org/10.1016/0167-9457(92)90021-3)
- Gernsbacher, M. A., Stevenson, J. L., Khandakar, S., & Goldsmith, H. H. (2008). Why does joint attention look atypical in autism. *Child Development Perspectives*, 2(1), 38–45. <https://doi.org/10.1111/j.1750-8606.2008.00039.x>
- Goldbart, J. (1994). Opening the communication curriculum to students with PMLDs. In J. Ware (Ed.), *Educating children with profound and multiple learning difficulties* (pp. 15–62). London, UK: David Fulton.
- Goldbart, J., Chadwick, D., & Buell, S. (2014). Speech and language therapists' approaches to communication intervention with children and adults with profound and multiple learning disability. *International Journal of Language and Communication Disorders*, 49(6), 687–701. <https://doi.org/10.1111/1460-6984.12098>
- Golinkoff, R. M. (Ed.) (2013). *The transition from prelinguistic to linguistic communication*. Delaware, DE: University of Delaware, Psychology Press.
- Grove, N., Bunning, K., Porter, J., & Olsson, C. (1999). 'See what I mean': Interpreting the meaning of communication by people with severe and profound intellectual disabilities. *Journal of Applied Research in Intellectual Disabilities*, 12, 190–203. <https://doi.org/10.1111/j.1468-3148.1999.tb00076.x>
- Hategan, C. B., & Talaş, D. (2014). Communication matrix – An assessment tool used in a case of autism spectrum disorders. *Procedia – Social and Behavioral Sciences*, 127, 169–173. <https://doi.org/10.1016/j.sbspro.2014.03.234>
- Hesterberg, T., Monaghan, S., Moore, D. S., Clipson, A., & Epstein, R. (2003). *Bootstrap methods and permutation tests: Companion chapter 18 to the practice of business statistics*. New York, NY: W.H. Freeman and Company.
- Hoekstra, A. T., van der Meulen, B. F., Oenema-Mostert, C. E., Jansen, G. G. H., Smrkovsky, M., Hoekstra, I. S. J., & Memelink, A. (2011). *Portage Programma – Nederland Revised*. Amsterdam, Nederland: Pearson.
- Hostyn, I., Daelman, M., Janssen, M. J., & Maes, B. (2010). Describing dialogue between persons with profound intellectual and multiple disabilities and direct support staff using the Scale for Dialogical Meaning Making. *Journal of Intellectual Disability Research*, 54(8), 679–690. <https://doi.org/10.1111/j.1365-2788.2010.01292.x>
- Hostyn, I., & Maes, B. (2009). Interaction between persons with profound intellectual and multiple disabilities and their partners: A literature review. *Journal of Intellectual and Developmental Disability*, 34(4), 296–312. <https://doi.org/10.3109/13668250903285648>
- Houwen, S., Visser, L., van der Putten, A., & Vlaskamp, C. (2016). The interrelationships between motor, cognitive, and language development in children with and without intellectual and developmental disabilities. *Research in Developmental Disabilities*, 53–54, 19–31. <https://doi.org/10.1016/j.ridd.2016.01.012>
- Iacono, T., Carter, M., & Hook, J. (1998). Identification of intentional communication in students with severe and multiple disabilities. *Augmentative and Alternative Communication*, 14, 102–114. <https://doi.org/10.1080/07434619812331278246>
- Iverson, J. M. (2010). Developing language in a developing body: The relationship between motor and language development. *Journal of Child Language*, 37, 229–261.
- Kahn, J. V. (1996). Cognitive skills and sign language knowledge of children with severe and profound mental retardation. *Education and Training in Mental Retardation and Developmental Disabilities*, 31, 162–168.
- Karasić, L. B., Tamis-LeMonda, C. S., & Adolph, K. E. (2011). Transition from crawling to walking and infants' actions with objects and people. *Child Development*, 82, 1199–1209. <https://doi.org/10.1111/j.1467-8624.2011.01595.x>
- Lee, W., & Rodgers, J. L. (1998). Bootstrapping correlation coefficients using univariate and bivariate sampling. *Psychological Methods*, 3, 91–103. <https://doi.org/10.1037/1082-989X.3.1.91>
- Lipscombe, B., Boyd, R. N., Coleman, A., Fahey, M., Rawicki, B., & Whittingham, K. (2016). Does early communication mediate the relationship between motor ability and social function in children with cerebral palsy? *Research in Developmental Disabilities*, 53–54, 279–286. <https://doi.org/10.1016/j.ridd.2016.02.013>
- Mahoney, G. (1998). *Child Behavior Rating Scale – Revised* (Unpublished document). Cleveland, OH: Case Western Reserve University.
- Mahoney, G. (2008). *Maternal Behavior Rating Scale – Revised* (Unpublished document). Cleveland, OH: Case Western Reserve University.
- Markova, I. (1990). Language and communication in Mental Handicap. In H. Giles, & P. W. Robinson (Eds.), *Handbook of language and social psychology* (pp. 363–380). Chichester, UK: Wiley.
- McLean, J., & Snyder-McLean, L. (1987). Form and function of communicative behavior among persons with severe developmental disabilities. *Australia and New Zealand Journal of Developmental Disabilities*, 13(2), 83–98.
- Mundy, P., Delgado, C., Block, J., Venezia, M., Hogan, A., & Seibert, J. (2003). *A manual for the Abridged Early Social Communication Scales (ESCS)*. Miami, FL: University of Miami.
- Nakken, H., & Vlaskamp, C. (2007). A need for a taxonomy for profound intellectual and multiple disabilities. *Journal of Policy*

- and Practice in Intellectual Disabilities, 4, 83–87. <https://doi.org/10.1111/j.1741-1130.2007.00104.x>
- Nijs, S., Schouten, B., & Maes, B. (2019). Visual functioning of persons with severe and profound intellectual disabilities: Observations by direct support workers and staff members and information available in personal files. *Journal of Policy and Practice in Intellectual Disabilities*, 16(4), 287–295. <https://doi.org/10.1111/jppi.12316>
- Ogletree, B. T., Bartholomew, P., Wagaman, J. C., Genz, S., & Reisinger, K. (2012). Emergent potential communicative behaviors in adults with the most severe intellectual disabilities. *Communication Disorders Quarterly*, 34(1), 56–58. <https://doi.org/10.1177/1525740112456201>
- Ogletree, B. T., Wetherby, A. M., & Westling, D. L. (1992). Profile of the prelinguistic intentional communicative behaviors of children with profound mental retardation. *American Journal on Mental Retardation*, 97, 186–196.
- Olsson, C. (2005). The use of communicative functions among pre-school children with multiple disabilities in two different setting conditions: Group versus individual patterns. *Augmentative and Alternative Communication*, 21(1), 3–18. <https://doi.org/10.1080/07434610412331270516>
- Palisano, R., Rosenbaum, P., Bartlett, D., & Livingston, M. (2007). *Gross motor function classification system expanded and revised*. CanChild Centre for Childhood Disability Research: McMaster University. Retrieved from https://www.cpqcc.org/sites/default/files/documents/HRIF_QCI_Docs/GMFCS-ER.pdf
- Parker, A. (2009). *Measuring an adapted form of Picture Exchange Communication Systems (PECS) for young children with visual impairments and developmental disabilities* (Unpublished doctoral dissertation). Lubbock, TX: Texas Tech University.
- Pennington, L. (2012). Speech and communication in cerebral palsy. *Eastern Journal of Medicine*, 17, 171–177.
- Petry, K., Maes, B., & Demuyne, J. (2004). *Geen beter leven dan een goed leven: Ouders en begeleiders over het leven van personen met ernstige meervoudige beperkingen*. Leuven, Belgium: Acco.
- Roberts, J. E., Price, J., & Malkin, C. (2007). Language and communication development in down syndrome. *Mental Retardation and Developmental Disabilities Research Reviews*, 13, 26–35. <https://doi.org/10.1002/mrdd.20136>
- Roebbers, C. M., & Kauer, M. (2009). Motor and cognitive control in a normative sample of 7-year-olds. *Developmental Science*, 12, 175–181. <https://doi.org/10.1111/j.1467-7687.2008.00755.x>
- Rowland, C. (2011). Using the communication matrix to assess expressive skills in early communicators. *Communication Disorders Quarterly*, 32, 190–201. <https://doi.org/10.1177/1525740110394651>
- Rowland, C. (2018). *Communication matrix*. Retrieved from <https://www.communicationmatrix.org/>
- Rowland, C., & Schweigert, P. (2000). Tangible symbols, tangible outcomes. *Augmentative and Alternative Communication*, 16, 61–78. <https://doi.org/10.1080/07434610012331278914>
- Siegel-Causey, E., & Bashinski, S. M. (1997). Enhancing initial communication and responsiveness of learners with multiple disabilities: A tri-focus framework for partners. *Focus on Autism and Other Developmental Disabilities*, 12, 105–120.
- Stadeus, A., Windey, K., Raman, M., Vermeir, G., & Van Driessche, S. (1994). *Tandemlijst: Voor jonge kinderen met ontwikkelingsmoeilijkheden*. Leuven, Belgium: Garant.
- Stephenson, J., & Dowrick, M. (2005). Parents' perspectives on the communication skills of their children with severe disabilities. *Journal of Intellectual and Developmental Disability*, 30, 75–85. <https://doi.org/10.1080/13668250500125031>
- Thurmond, V. A. (2001). The point of triangulation. *Journal of Nursing Scholarship*, 33, 253–258. <https://doi.org/10.1111/j.1547-5069.2001.00253.x>
- Tröster, H., & Brambling, M. (1992). Early social-emotional development in blind infants. *Child: Care Health and Development*, 18, 207–277. <https://doi.org/10.1111/j.1365-2214.1992.tb00355.x>
- Van Den Broek, E. G., Janssen, C. G. C., Van Ramshorst, T., & Deen, L. (2006). Visual impairments in people with severe and profound multiple disabilities: An inventory of visual functioning. *Journal of Intellectual Disability Research*, 50(6), 470–475. <https://doi.org/10.1111/j.1365-2788.2006.00804.x>
- Van der Meer, I., Colla, S., Van Leeuwen, K., Vlaskamp, C., Ceulemans, E., Hoppenbrouwers, K., ... Maes, B. (2017). Exploring parental behavior and child interactive engagement: A study on children with a significant cognitive and motor developmental delay. *Research in Developmental Disabilities*, 64, 131–142. <https://doi.org/10.1016/j.ridd.2017.04.002>
- Van Splunder, J. A. N. G., Stilma, J. S., Bernsen, R. M. D., & Evenhuis, H. M. (2006). Prevalence of visual impairment in adults with intellectual disabilities in the Netherlands: Cross-sectional study. *Eye*, 20(9), 1004.
- Visser, L., Vlaskamp, C., Emde, C., Ruiter, S. A. J., & Timmerman, M. E. (2017). Difference or delay? A comparison of Bayley-III Cognition item scores of young children with and without developmental disabilities. *Research in Developmental Disabilities*, 71(September), 109–119. <https://doi.org/10.1016/j.ridd.2017.09.022>
- Vlaskamp, C. (2011). Kijken naar ontwikkeling. In B. Maes, C. Vlaskamp, & A. Penne (Eds.), *Ondersteuning van mensen met ernstige meervoudige beperkingen. Handvatten voor een kwaliteitsvol leven* (pp. 59–75). Leuven, Belgium: Acco.
- Vlaskamp, C., & Oxener, G. (2002). Communicatie bij mensen met ernstige meervoudige beperkingen: Een overzicht van assessment en interventie methoden. *Nederlands Tijdschrift Voor De Zorg Aan Mensen Met Verstandelijke Beperkingen*, 28, 226–237.
- Vlaskamp, C., van der Meulen, B. F., & Smrkovsky, M. (1999). *Gedragstaxatie instrument*. Groningen, Nederland: Stichting Kinderstudies.
- Voorman, J. M., Dallmeijer, A. J., Van Eck, M., Schuengel, C., & Becher, J. G. (2010). Social functioning and communication in children with cerebral palsy: Association with disease characteristics and personal and environmental factors. *Developmental Medicine and Child Neurology*, 52, 441–447. <https://doi.org/10.1111/j.1469-8749.2009.03399.x>
- Vygotsky, L. S. (1978). *Mind in society: The development of higher psychological processes*. Cambridge, MA: Harvard University Press.
- Wang, M. V., Lekhal, R., Aarø, L. E., & Schjølberg, S. (2012). Co-occurring development of early childhood communication and motor skills: Results from a population-based longitudinal study. *Child: Care, Health and Development*, 40, 77–84.
- Warren, S. F., & Brady, N. C. (2007). The role of maternal responsivity in the development of children with intellectual disabilities. *Mental Retardation and Developmental Disabilities Research Reviews*, 13, 330–338. <https://doi.org/10.1002/mrdd.20177>
- Weis, R. (2014). *Introduction to abnormal child and adolescent psychology*. London, UK: Sage Publications.
- Wetherby, A., & Prizant, B. (2002). *Communication and symbolic behavior scales developmental profile – first, normed ed.*. Baltimore, MD: Paul H. Brookes.
- Zhu, W. (1997). Making bootstrap statistical inferences: A tutorial. *Research Quarterly for Exercise and Sport*, 68, 44–55. <https://doi.org/10.1080/02701367.1997.10608865>

How to cite this article: Dhondt A, Van der Meer I, van der Putten A, Maes B. Communicative abilities in young children with a significant cognitive and motor developmental delay. *J Appl Res Intellect Disabil*. 2020;33:529–541. <https://doi.org/10.1111/jar.12695>