University of Groningen

Searching for chemical signatures of planet formation

Rab, Ch.; Muro-Arena, G. A.; Kamp, I.; Dominik, C.; Waters, L. B. F. M.; Thi, W.-F.; Woitke, P.

Published in:
Origins

DOI:
[10.1017/S1743921319002114](https://doi.org/10.1017/S1743921319002114)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Final author's version (accepted by publisher, after peer review)

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Rab, C., Muro-Arena, G. A., Kamp, I., Dominik, C., Waters, L. B. F. M., Thi, W.-F., & Woitke, P. (2020). Searching for chemical signatures of planet formation. In *Origins: From the Protosun to the First Steps of Life* (Vol. 345, pp. 362-364). Cambridge University Press. <https://doi.org/10.1017/S1743921319002114>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Searching for chemical signatures of planet formation

Ch. Rab¹, G.A. Muro-Arena², I. Kamp¹, C. Dominik²,
 L.B.F.M. Waters^{2,3}, W-F. Thi⁴, and P. Woitke⁵

¹Kapteyn Astronomical Institute, University of Groningen, P.O. Box 800, 9700 AV Groningen, The Netherlands, email: rab@astro.rug.nl

²Anton Pannekoek Institute for Astronomy, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands

³SRON Netherlands, Sorbonnelaan 2, 3584 CA Utrecht, The Netherlands

⁴MPE, Giessenbachstrasse 1, 85748 Garching, Germany

⁵SUPA, School of Physics & Astronomy, University of St. Andrews, North Haugh, St. Andrews KY16 9SS, UK

Abstract. High spatial resolution observations with ALMA and VLT/SPHERE show gaps and rings in continuum emission of protoplanetary disks, possibly indicating ongoing planet formation. However, it is still unclear if the gas follows the dust distribution. We present radiation thermo-chemical models for the disk of HD 163296 to study the impact of dust and gas gaps on the chemistry and molecular line emission. We compare a model with only dust gaps to a model that also has gas gaps. In both models, rings and gaps are visible in (sub)mm molecular line emission. Due to chemistry, certain molecules are sensitive to dust gaps where others are more sensitive to gas depletion. Observations of multiple molecules might allow to accurately determine the degree of gas depletion within the dust gaps, information crucial to discriminate between gap formation theories (e.g. planets, ice lines).

Keywords. stars: pre-main-sequence, (stars:) planetary systems: protoplanetary disks, astrochemistry, radiative transfer, methods: numerical

1. Introduction & Methods

High spatial resolution observations with ALMA and VLT/SPHERE reveal gaps and rings in dust continuum images of protoplanetary disks (e.g. Andrews et al. 2016, Avenhaus et al. 2018). Although those gaps and rings are very prominent in dust observations, they are not yet clearly detected in the gas. An accurate measurement of the gas column densities and the possible gas depletion within the dust gaps is crucial to discriminate between different gap formation scenarios. If planets produced the gaps, one would expect strong gas depletion and gas gaps, other formation scenarios such as dust evolution, molecular ice lines or deadzones predict no or only shallow gas gaps (e.g. Isella et al. 2016, Birnstiel et al. 2015, Zhang et al. 2015, Pinilla et al. 2016).

Here we present a model for the HD 163296 protoplanetary disk to study in detail the impact of the observed dust gaps and possible gas gaps on the chemistry and on molecular line emission. We used the radiation thermo-chemical disk model PRODIMO (PROtoplanetary DIsk MOdel, e.g. Woitke et al. 2009, Kamp et al. 2017) to self-consistently calculate dust and gas temperatures, molecular abundances and synthetic observables. The models are based on the HD 163296 PRODIMO model presented in Woitke et al. (2019), but was refined by Muro-Arena et al. (2018) to fit new VLT/SPHERE and ALMA continuum observations that show prominent dust gaps and rings.

Figure 1. Two models for the HD 163296 disk, one without gas gaps (SMOOTH, *top row*) and one with gas gaps (GAPS, *bottom row*). The gray boxes mark the dust gaps present in both models. *First column:* normalized radial molecular column density profiles (H_2 traces the total gas column); *second column:* the corresponding radial intensity profiles (normalized to the peak intensity); *third column:* integrated intensity map for $C^{18}O J = 3 - 2$.

2. Results & Conclusions

In Fig. 1, we compare two models, one with a smooth gas surface density profile (SMOOTH) and one with gas gaps (GAPS). As an example, we show the column densities and observables for the molecules CO and CN. The molecular column density profiles show gaps and rings in both models. In the SMOOTH model this is solely due to the impact of dust gaps on the chemistry (e.g. temperature, radiation field). In the models CO quite nicely follows the total gas surface density (see also Facchini et al. 2018), but also in the SMOOTH model gaps are seen in the $C^{18}O$ line image. The $C^{18}O$ line is optically thick and therefore sensitive to temperature changes caused by the dust gaps (see also van der Marel et al. 2018). In the SMOOTH model, CN gaps and rings appear due to dust depletion and the column density actually peaks within the dust gaps, but in the gas GAPS model these peaks are washed out, making CN a good tracer of dust gaps without gas depletion. This example shows that self-consistent dust and gas modeling including chemistry is required to accurately infer gas column densities (gas gaps) from molecular line observations and that observations of multiple molecules will allow for a more accurate measurement of gas depletion within dust gaps.

References

- Andrews, S. M., Wilner, D. J., Zhu, Z., Birnstiel, T. et al. 2016, *ApJ* (Letters), 820, L40
 Avenhaus, H., Quanz, S. P., Garufi, A., Perez, S., Casassus, S. et al. 2018, *ApJ*, 863, 44
 Birnstiel, T., Andrews, Sean M., Pinilla, P., Kama, M. 2015, *ApJ*, 813
 Facchini, S., Pinilla, P., van Dishoeck, E. F. & de Juan Ovelar, M. 2018, *A&A*, 612
 Isella, A., Guidi, G., Testi, L., Liu, S., Li, H., Li, S. et al. 2016, *Phys. Rev. Lett.*, 117, 251101
 Kamp, I., Thi, W.-F., Woitke, P., Rab, C., Bouma, S., Ménard, F. 2017, *A&A*, 607, A41
 Muro-Arena, G. A., Dominik, C., Waters, L. B. F. M., Min, M. et al. 2018, *A&A*, 614, A24
 Pinilla, P., Flock, M., Ovelar, M. de Juan & Birnstiel, T. 2016, *A&A*, 596
 van der Marel, N., Williams, J. P., & Bruderer, S. 2018, *ApJ* (Letters), 867, L14
 Woitke, P., Kamp, I., & Thi, W.-F. 2009, *A&A*, 501, 383
 Woitke, P., Kamp, I., Antonellini, S., Anthonioz, F., et al., 2019, *PASP*, 131, 064301
 Zhang, K., Blake, G. A., & Bergin, E. A. 2015, *ApJ* (Letters), 806, L7