University of Groningen

Intraspecific niche partitioning in macrodetritivores enhances mixed leaf litter decomposition

Fontana, Simone; Berg, Matty P.; Moretti, Marco

Published in:
Functional Ecology

DOI:
[10.1111/1365-2435.13448](https://doi.org/10.1111/1365-2435.13448)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2019

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Fontana, S., Berg, M. P., & Moretti, M. (2019). Intraspecific niche partitioning in macrodetritivores enhances mixed leaf litter decomposition. *Functional Ecology*, 33(12), 2391-2401. <https://doi.org/10.1111/1365-2435.13448>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

RESEARCH ARTICLE

Functional Ecology

Intraspecific niche partitioning in macrodetritivores enhances mixed leaf litter decomposition

Simone Fontana¹ | Matty P. Berg^{2,3} | Marco Moretti¹

¹Biodiversity and Conservation
Biology, Swiss Federal Research Institute
WSL, Birmensdorf, Switzerland

²Animal Ecology Group, Department
of Ecological Science, Vrije Universiteit
Amsterdam, Amsterdam, The Netherlands

³Community and Conservation Ecology
Group, Groningen Institute of Evolutionary
Life Science, University of Groningen,
Groningen, The Netherlands

Correspondence

Simone Fontana
Email: simone.fontana@wsl.ch

Funding information

Schweizerischer Nationalfonds zur
Förderung der Wissenschaftlichen
Forschung, Grant/Award Number:
315230_170200/1

Handling Editor: Sally Power

Abstract

1. In biodiversity and ecosystem functioning research, the consideration of intraspecific trait variation and multiple trophic levels – both largely overlooked so far – has been proposed as a promising step towards a consensus about causal relationships and mechanisms.
2. Soil invertebrates contribute crucially to leaf litter decomposition and thus to the functioning of one of the most diverse ecosystem components on Earth. On the one hand, a positive effect of leaf litter diversity on decomposition was shown by several studies. On the other hand, at the interspecific level, differences in body size and feeding traits of consumers influence this process by interacting with leaf litter traits that define decomposability, such as nutrient content and physical properties. It is also known that the feeding niche of consumers can vary during their ontogeny as it depends on individual body size, metabolism and physiology. However, whether and how intraspecific trait variation of macrodetritivores contributes through niche partitioning to the positive effect of litter combinations on decomposition was never investigated.
3. Here, we tested experimentally the effect of intraspecific body size differences in the common terrestrial isopod species *Oniscus asellus* on the decomposition of the two co-occurring litter species *Acer platanoides* and *Betula pendula*. We hypothesized complementary resource use by two distinct isopod body size classes, leading to decreased competition and higher overall litter decomposition.
4. In a full factorial design with all possible combinations of litter species and isopod body size classes, litter mixtures enhanced decomposition beyond the additive effect of each single species, especially in isopod mixed-body size treatments. Our results suggest an interaction between litter and consumer diversity driving intraspecific feeding niche partitioning. Although the focus on a single detritivore species does not allow for generalizations, our results should stimulate the investigation of the mechanisms of biotic interactions among individual organisms.

KEYWORDS

body size, complementarity, decomposition, functional diversity, intraspecific trait variation, Isopoda, ontogeny, soil biota

1 | INTRODUCTION

It has been for long recognized that biodiversity can have a positive influence on ecosystem processes. However, relationships observed in a variety of systems are far from being consistent (see review by Cardinale et al., 2012) and a deep understanding of the driving mechanisms is still lacking. For this reason, a broadening of perspectives and new approaches in biodiversity and ecosystem functioning (BEF) research have been called for, such as the explicit inclusion of food web structure (e.g. Reiss, Bridle, Montoya, & Woodward, 2009; Seibold, Cadotte, MacIvor, Thorn, & Müller, 2018), the adoption of trait-based approaches (McGill, Enquist, Weiher, & Westoby, 2006) and the consideration of different levels of biological organization, especially intraspecific trait variation (Violle et al., 2012).

In community ecology, the focus has mainly been on interspecific trait variability, largely based on mean field theory, which suggests that mean trait values capture the majority of species and community dynamics (MacArthur & Levins, 1967). In the last decade, the legitimacy of this view has been strongly challenged by both theoretical work (e.g. Barabás & D'Andrea, 2016; De Laender et al., 2014; Violle et al., 2012) and empirical studies that quantify the relative contribution to trait variation of the intra- and interspecific components (Bonfanti et al., 2018; Chalmardier et al., 2017; Jung, Violle, Mondy, Hoffmann, & Muller, 2010; Rhoades, Best, & Stachowicz, 2018; Siefert et al., 2015; Volf et al., 2016). These studies showed that intraspecific trait variation can be comparable or even larger than interspecific trait variation, and trait differences among conspecific individuals can have community-wide consequences. Still, intraspecific trait variation was largely overlooked in BEF research (Bolnick et al., 2011; but see Des Roches et al., 2018; Fontana, Thomas, Moldoveanu, Spaak, & Pomati, 2018; Raffard et al., 2017; Raffard, Santoul, Cucherousset, & Blanchet, 2019), while functional traits of individuals determine their fitness in changing environments (McGill et al., 2006; Violle et al., 2007) and finally the effects of species on ecosystem processes (e.g. Bolnick et al., 2003; Violle et al., 2012). This is particularly relevant when considering biological interactions, which are by definition realized at the individual level. We urgently need to incorporate explicitly intraspecific variation (e.g. due to ontogeny, sexual dimorphism, genetic differences or phenotypic plasticity) into BEF research to link the traits of primary producers to those of consumers and thus afford a food web perspective (Coq et al., 2018).

In soil food webs, litter decomposition is a key process that sustains biogeochemical cycles across ecosystems and contributes crucially to the functioning of soils and their service delivery (Bardgett & van der Putten, 2014; Wagg, Bender, Widmer, & van der Heijden, 2014). Nevertheless, linking soil biodiversity to ecosystem processes is far from being a trivial task (Eisenhauer et al., 2017; Wall, Bardgett, & Kelly, 2010). Attempts to gain mechanistic understanding of litter decomposition should take into account the complex trait-mediated interactions between litter species and soil fauna communities, namely the co-occurrence of bottom-up and top-down effects (Hättenschwiler & Gasser, 2005). Leaf litter species traits can

influence decomposition through a variety of mechanisms (Gessner et al., 2010; Hättenschwiler, Tiunov, & Scheu, 2005), which result in non-additive effects (i.e. the observed decomposition of a litter mixture deviates from expectations derived by summing decomposition rates of litter monocultures). For example, a positive effect of leaf litter species combinations on decomposition can be driven by interspecific differences in litter quality (determined by traits such as physical and structural properties, and the concentration of nutrients and secondary metabolites), which cause the preferential consumption of one litter species when the consumer has the possibility to choose among many (e.g. Hättenschwiler et al., 2005). This so-called selection effect also depends on the identity of the consumer considered (Hättenschwiler & Gasser, 2005; Heemsbergen et al., 2004), as confirmed by results from different macrodetritivore taxa (Vos, van Ruijven, Berg, Peeters, & Berendse, 2011). Besides the selection effect, complementary effects have also been described, in which enhanced decomposition of a litter mixture is determined by the advantages of a balanced diet or by complementary resource use between microbes and macrodetritivores (Vos, van Ruijven, Berg, Peeters, & Berendse, 2013). The same authors suggested that additional consideration of macrodetritivore trait diversity might contribute to explain a greater positive effect of leaf litter trait variation (see also Heemsbergen et al., 2004; but see Bílá et al., 2014). This was already shown for two soil macrofauna classes (Diplopoda and Gastropoda), whose very distinct traits led to complementary consumption of leaf litter mixtures (De Oliveira, Hättenschwiler, & Handa, 2010).

On the contrary, the consequences of arguably smaller trait differences, such as those among individuals of the same consumer species, were not experimentally tested for their interaction with litter diversity effects so far. Animal trait-based ecology suffers from a lack of mechanistic understanding behind the link between traits and ecosystem functioning (Didham, Leather, & Basset, 2016), which hampers predictive multitrophic BEF research (but see, e.g. Coulis et al., 2015; Muñoz, Schaefer, Böhning-Gaese, & Schleuning, 2017; Schleuning, Fründ, & García, 2015; Schmitz, Buchkowski, Burghardt, & Donihue, 2015). Interestingly, in herbivores and detritivores, intraspecific trait differences can drive variation in consumption rates among individuals of a population, as experimental approaches have shown (Raffard et al., 2017; Rhoades et al., 2018; Rota, Jabiol, Chauvet, & Lecerf, 2018). However, none of these studies investigated how trait variation can determine niche partitioning within a single consumer species, and whether the resulting resource use complementarity (due to decreased competition) enhances mixed litter decomposition. During its ontogeny, a given consumer species might occupy distinct niches (e.g. Clegg, Ali, & Beckerman, 2018; Polis, 1984; Start, 2018), defined by stage-specific feeding modes due to differences in body size (Werner & Gilliam, 1984). This ontogenetic niche shift reflects changes in energetic requirements, metabolism and physiology and can result in preferences for different leaf parts (Vilisics, Szekeres, & Hornung, 2012), or litter traits or microenvironmental conditions, which depend on litter species combinations (Hättenschwiler et al., 2005).

Here, we tested experimentally whether intraspecific body size differences in the common terrestrial isopod species *Oniscus asellus* Linnaeus, 1758 (*Oniscus* hereafter) affect the decomposition of the two co-occurring litter species *Acer platanoides* (Norway maple, *Acer* hereafter) and *Betula pendula* (Silver birch, *Betula* hereafter). Isopods, as significant part of soil macrofauna in most terrestrial ecosystems, throughout their life cycle play an important role in litter decomposition (David, 2014; Hättenschwiler et al., 2005). By testing all compositions of litter species (*Acer* and *Betula* alone, as well as mixed) and *Oniscus* body size classes (large and small alone, as well as mixed) in a full factorial design, we aimed at elucidating how intraspecific body size variation of the consumers mediates the effect of litter species composition on litter mass loss. *Acer* and *Betula* have different traits (e.g. C-to-N ratio, magnesium and calcium content, physical resistance to consumption by macrodetritivores), which generally result in a slightly higher decomposition rate (=litter quality) in *Betula* (Cornelissen, 1996; Finerty et al., 2016). A combination of *Oniscus* body size classes could be expected to enhance the positive, non-additive effect of litter diversity on consumption, thanks to intraspecific feeding niche partitioning.

In this study, we tested the following specific hypotheses and related expectations:

1. Leaf litter mixtures can enhance decomposition independent of consumer identity and species composition, due to selection and/or complementarity effects driven by trait differences among litter species. Therefore, we expect positive non-additive effects of the two litter species (*Acer* and *Betula*) on total litter mass lost in all treatments of our experiment.
2. If the above pattern is enhanced by intraspecific feeding niche partitioning of the consumer, the positive effect of a litter mixture on overall decomposition is expected to be highest when the variation in consumer traits is maximized. Therefore, we expect the greatest positive difference between observed and expected total leaf litter mass loss (*Acer* + *Betula*) in the mixed-body size treatment of the consumer, assuming that body size relates to feeding traits and strategies in *Oniscus* (Vilšić et al., 2012; Werner & Gilliam, 1984).

2 | MATERIALS AND METHODS

2.1 | Litter collection and preparation

To maximize the probability of detecting non-additive effects of small- and large-sized *Oniscus* on litter decomposition, we avoided pairs of litter species with a large difference in palatability and decomposability, which would probably have caused a much higher consumption of one litter species independent of *Oniscus* body size classes' composition. Therefore, we selected two common and co-occurring tree species, that is *Acer* and *Betula*, which show moderate differences in decomposition rates determined by traits affecting resource quality for decomposers (Cornelissen, 1996; Finerty et al., 2016). Freshly fallen leaves at time of leaf abscission

without any visible sign of herbivory consumption, fungal attack or diseases were collected. In order to minimize leaf trait variation within species, we focused litter collection on a few healthy trees in a restricted area of the Canton of Zurich, Switzerland (*Acer*: 47°14'N/8°31'E in October 2016; *Betula*: 47°22'N/8°27'E in November 2016). Leaves were collected in paper bags and dried in a stove set at 40°C, subsequently put in sealed plastic bags after air was removed manually and stored in a dark aerated room. Note that during the experiment, leaf litter was dried several times (see below for details). While temperature was always set at 40°C, the duration of the drying process was adapted to initial water content and litter quantity to assure complete water removal (according to previous experience) and was always >14 hr. Prior to the experiment (September 2017), leaves' size and shape were standardized by cutting them with a metal hole punch in round pieces of 2 cm diameter.

2.2 | Animal collection and sorting

Ten and nine days before experiment start, *Oniscus* was collected in a deciduous forest in the city of Zurich, Switzerland (47°24'N/8°31'E). Animals were stored under experimental conditions (85% relative humidity; daily fluctuating temperature, day 20°C and night 15°C; 12:12-hr light-dark cycle, 8 a.m./8 p.m.) in plastic boxes (20.0 cm length; 9.5 cm width; 9.0 cm height) with a moist bottom layer of plaster of Paris. Individuals were sorted by eye into two distinct body size classes, which were defined in the attempt of maximizing differences between large and small animals given size distributions observed in field samples collected prior to the experiment (e.g. Figure S1). Body size was defined as fresh body mass, which is easy to measure on living *Oniscus* and showed a strong positive correlation with body length (Figure S1). By discarding the intermediate body sizes, we made sure fresh body mass of small (range = 10.7–45.6 mg; mean = 29.5 mg; median = 30.0 mg) and large *Oniscus* (range = 53.7–146.3 mg; mean = 80.7 mg; median = 78.0 mg) showed a non-overlapping distribution at the beginning of the experiment, with large animals being more than double as heavy as small ones on average (Figure S2). *Oniscus* with a developed marsupium (a brood pouch with eggs or embryos) were excluded, because they would probably deviate from the fresh body mass to length relationship (Figure S1) and to assure a constant number of animals per microcosm throughout the whole experiment.

2.3 | Microcosm setting

Each microcosm was composed of a transparent Perspex ring (6.3 cm diameter; 3.0 cm height), closed at the bottom with a moist layer of plaster of Paris (approx. 0.8 cm thickness) and closed on the top with a lid with a central round mesh (2.0 cm diameter; 200 µm mesh size) for gas exchange. All microcosms were put in plastic trays (Figure 1) on a moist layer of plaster of Paris (approx. 3.0 cm thickness), which allowed to keep the leaf litter in the microcosms moist without disturbance by watering the trays only (from there the water was also

FIGURE 1 Experimental setup. (a-b) Dark brown = *Acer* leaf litter discs. Light brown = *Betula* leaf litter discs. (a) For simplicity, the proportion of large and small *Oniscus* is reported here depicting only two out of four animals. (b) Control microcosms. (c) Photograph of the experimental microcosms

equally sucked by capillary action into the plaster of Paris of each microcosm).

2.4 | Experimental design

We tested litter decomposition of three *Oniscus* body size combinations (large only: 79.4 mg mean fresh body mass; small only: 29.3 mg; large and small mixed: 55.1 mg) on three litter species combinations (*Acer* only, *Betula* only, a mixture of both). This resulted in nine treatments, with 10 replicates each (total = 90 microcosms). Each

microcosm contained four animals (all large, all small, or two large and two small). In addition, 30 control microcosms with litter only were included (=3 litter species combinations × 10 replicates) to measure litter decomposition by microbes.

The experiment was conducted in a climate chamber with the same conditions as above. In order to assure ad libitum food, and so avoid the influence of food limitation on observed patterns, the experiment was carried on for 3 weeks until approximately 50% litter mass loss in the microcosms with the fastest decomposition. Although we did not expect environmental conditions to vary much among microcosms,

we subdivided them into two blocks (=trays), each one containing five replicates of each treatment in varying position (Figure 1a). Control microcosms were grouped in an additional block (Figure 1b).

2.5 | Animal acclimation

The microcosms used for *Oniscus* acclimation were identical to the ones used for the experiment. To inoculate the dried leaf litter with a natural microbial community and thus facilitate consumption by *Oniscus* (Zimmer, 2006), 15 days before experiment start a microbial wash was made by collecting 750-ml partially degraded litter in a mixed deciduous forest close to the Swiss Federal Research Institute WSL (47°21'N/8°27'E). This litter was mixed with 1.5-L tap water, shaken overnight in a 2-L bottle (150 rpm, VWR Microplate Shaker) and left 7 hr to extract microbes. The liquid fraction was then sieved in two successive steps (250 and 50- μ m metal mesh, respectively), kept hereafter in the climate chamber under experimental conditions and shaken manually from time to time to avoid anoxia. Fourteen days before experiment start, microcosms were filled with leaf litter leftovers resulting from cutting round pieces for the experiment (as described above), according to treatment (*Acer* only, *Betula* only, or a mixture of both litters in approx. equal proportion). The layer of plaster of Paris in the tray was saturated with water (and kept saturated during the following days by adding water when necessary), and each microcosm was sprayed with a standardized dose of microbial wash (approx. 2.5 ml, which was enough to reach a leaf litter water content of about 50% according to pre-tests). The same microbial treatment was applied a second time 10 days before experiment start. The day after, *Oniscus* collected in the field were transferred into microcosms, selecting individuals of similar body size within each category (small or large, single-size treatments), while maximizing body size variation in mixed-size treatments (Figure S3). After 7 days of acclimation following Moretti et al. (2017), leaf litter and faeces were carefully removed with a brush from the microcosms and *Oniscus* were starved for 2 days (under identical conditions, except for the absence of food), to make sure that they would have empty guts at the start of the experiment. The same acclimation and starving procedure as described above (but with a mixture of the two litter species only) was applied to several spare animals (divided by size class in separate boxes). During the acclimation phase, it was necessary to substitute 15 dead *Oniscus* (eight small- and seven large-sized out of a total of 360 specimens).

2.6 | Experiment preparation

The leaf litter discs were dried to remove the small amount of water that accumulated since their preparation. Immediately after that, leaf litter portions were weighted using a precision balance accurate to 0.01 mg (Mettler Toledo AE 240), and added to the microcosms (including control microcosms) according to their treatment (3 days before experiment start). A total amount of approx. 500 mg leaf litter (range = 472–548 mg) was put in each microcosm, aiming at an equal proportion of the two species in mixed litter treatments

(range = 237–275 mg each). To facilitate the identification of the two species in mixed litter treatments at the end of the experiment, leaf discs were horizontally separated in all microcosms for consistency (where both species were present, *Acer* left side and *Betula* right side of the microcosm). A standardized dose of microbial wash (approx. 2.5 ml) was also sprayed twice to each microcosm, to increase leaf water content and allow the colonization by microbes during 3 days before adding *Oniscus* (following, e.g. Bílá et al., 2014).

2.7 | Experiment and final measurements

All acclimated *Oniscus* were weighted using a precision balance accurate to 0.01 mg (Mettler Toledo AX205 DeltaRange) and then transferred into the microcosms (official start of the experiment). During the experiment, water was poured into the trays every 2–3 days to make sure the layer of plaster of Paris in the microcosms was constantly moist. As a consequence, leaf litter showed a gradient of decreasing moisture content from bottom to top of the litter layers, allowing individual *Oniscus* to find their preferred microenvironment. Every 1–2 days we checked each microcosm for dead *Oniscus* and noted dates and times when we found any. To minimize disturbance and leaf litter movements, the search was kept as short as possible and dead animals were not removed from the microcosms. The experiment was stopped after 21 days, that is when approximately half of the litter was lost in the treatment with highest decomposition. All *Oniscus* were separated from leaf litter and weighted again. Remaining leaf litter was dried and fragments were then separated according to litter species using a binocular. During litter separation, *Oniscus* faeces and extensive fungal mycelium were carefully removed. Leaf fragments were dried again and immediately weighted.

2.8 | Statistical analyses

Oniscus mortality during the experiment was very low (5%; 18 dead individuals out of 360), affecting 16 out of 90 microcosms. We decided to completely exclude from further analyses the microcosms with dead individuals. This decision was made because of the low number of microcosms affected and to avoid an estimation of partial contributions to decomposition that would have introduced uncertainties and confounding factors (e.g. unbalanced design, possible cannibalism, nutrient addition from dead animals).

Leaf litter relative consumption rate in presence of *Oniscus* (RCR) was defined by litter mass loss for each microcosm and leaf litter species separately using the following formula:

$$\text{RCR} = (\text{ML} [\text{mg}] - \text{mean ML microbial controls} [\text{mg}]) / (\text{time} [\text{d}] * \text{Oniscus mass} [\text{mg}]) \quad (1)$$

where ML is leaf litter dry mass loss, and *Oniscus mass* is the sum of four individuals' fresh body masses (averaged between the start and the end of the experiment).

To test whether the combination of two litter species given a certain *Oniscus* body size treatment resulted in additive or non-additive

FIGURE 2 Expected (=additive effect of single-litter species microcosms) versus observed leaf litter mass loss (mean values $\pm$ SE) in the presence of both litter species *Acer* and *Betula* for any given *Oniscus* body size composition. Welch's *t* test significance levels are reported (* $p < .1$; ** $p < .05$; *** $p < .01$) where appropriate (see also Table 1)

effects, we compared expected versus observed amount of leaf litter mass loss in microcosms using Welch's *t* tests (function *t.test*, R Core Team, 2018). The expected values were calculated by assuming the same *Acer* and *Betula* RCR in a litter mixture as in the corresponding monocultures (we applied the average RCR measured in single-litter treatments). In this case, equal probability of feeding on *Acer* and *Betula* was assumed; that is, expected litter decomposition was calculated by multiplying the species-specific RCR by both the body mass of *Oniscus* and half of the experiment duration (≈ 10.5 d) (see formula (1)).

To account for the variability in RCR among litter monocultures (in addition to average RCR), we also calculated 95% confidence intervals of expected consumption for single microcosms, and checked whether the observed consumption fell within this range. An overlap between observed values and the confidence intervals of expected values would indicate that the total consumption of leaf litter mixtures can be calculated from the consumption of litter monocultures (null hypothesis).

Analyses were carried out using R version 3.5.1 (R Core Team, 2018).

3 | RESULTS

The percentage of decomposed litter at the end of the experiment varied quite a lot among microcosms (range = 7.8%–54.0%, mean = 23.9%), showing that even in case of the highest decomposition there was no food limitation for *Oniscus*.

The differences between expected and observed leaf litter decomposition in the presence of both *Acer* and *Betula* simultaneously indicate non-additive effects of the two litter species. In treatments with both large- and small-sized *Oniscus*, the two litter species combined showed the strongest positive effect on total mass loss. Thus, observed total litter decomposition was on average 38% higher than expected, whereas this increase was 22% and 24% in the presence of only large- and small-sized *Oniscus*, respectively (Figure 2, Table 1). Specifically, in treatments with large-sized *Oniscus* only, mass loss was lower than expected in *Acer*, whereas it was higher for *Betula* (Figure 2, Table 1). This resulted in higher total leaf litter mass loss than expected (Figure 2, Table 1). The same general pattern of lower-than-expected *Acer* but higher-than-expected *Betula* decomposition was observed also in treatments with small-sized *Oniscus* only (Figure 2, Table 1). In this case, however, differences were weaker and non-significant.

The consistency of our results is supported by taking into account the variability in relative consumption rate among single replicates, which also showed the predominance of non-additive effects of the two litter species (Figure S4).

4 | DISCUSSION

By explicitly testing for the first time in terrestrial ecosystems the role of consumer intraspecific trait variation in mediating, through

TABLE 1 Effects of combining two leaf litter species on litter mass loss, defined as departures from additivity of single-litter relative consumption rates

	Expected versus observed leaf litter mass loss								
	<i>Acer</i>			<i>Betula</i>			Total (both species)		
	<i>t</i>	<i>n</i>	<i>p</i>	<i>t</i>	<i>n</i>	<i>p</i>	<i>t</i>	<i>n</i>	<i>p</i>
Effect of litter combination									
Large body size	2.105	16	.071	-4.350	16	.003	-2.995	16	.015
Small body size	0.343	14	.740	-2.193	14	.058	-1.692	14	.124
Mixed body sizes	-0.126	20	.902	-4.077	20	.002	-3.853	20	.002

Note: Results are reported for each *Oniscus* body size composition separately. Summary statistics refer to Welch's *t* tests performed with data reported in Figure 2. Positive *t*-values indicate 'expected is higher than observed'; vice versa for negative *t*-values.

niche partitioning, the effects of leaf litter diversity on decomposition, our experiment aimed at elucidating some fundamental mechanisms in the context of the still lively debate about BEF relationship. Our results demonstrate that intraspecific body size variation in a macrodetritivore species interacts positively with trait differences between two leaf litter species (e.g. nutrient composition and physical properties) to enhance total litter decomposition. This supports our initial hypotheses and suggests the existence of intraspecific feeding niche partitioning among small- and large-sized individuals of *Oniscus*, which was contingent upon the presence of a combination of the two litter species *Acer* and *Betula*. In fact, this resource niche partitioning increased the non-additive effects of the two litter species on decomposition and was possibly dependent on the possibility for each individual *Oniscus* to select the leaf species that had the best trait matching with consumer's metabolic and biomechanical constraints (Pawar, Dell, & Savage, 2015). Purely behavioural differences between small- and large-sized *Oniscus* might also have contributed to the observed effect. The positive influence of a litter species combination on total mass loss was also present in treatments with a single *Oniscus* body size class. However, in this case the observed effects of litter diversity were less pronounced and apparently only driven by selection effects, that is clear shifts towards the leaf litter species with highest quality or most palatable (*Betula*) (Figure 2, Table 1). In contrast, when small- and large-sized *Oniscus* were present simultaneously in combined litter samples, not only the decomposition of *Betula* increased, but also *Acer* showed the same tendency, or at least its mass loss did not decrease compared to expectations (Figure 2, Table 1).

We hypothesize that this pattern can be explained by the interaction between leaf litter and consumer traits. For instance, *Betula* litter has a higher quality (i.e. higher decomposability) compared to *Acer* which is probably mainly determined by the higher content of calcium and magnesium (Finerty et al., 2016), two fundamental components of isopods' diet and exoskeleton (Becker, Ziegler, & Epple, 2005; David & Handa, 2010; Nicolai, 1988; Zimmer, 2002). In contrast, *Betula* litter has a higher carbon-to-nitrogen ratio and tensile strength (Finerty et al., 2016). As a consequence, small-sized *Oniscus* individuals might not be able to produce enough force at the tip of their mandibles to puncture tough *Betula* leaves as

efficiently as large-sized individuals, which is consistent with the much smaller increase in *Betula* (respectively decrease in *Acer*) mass loss in small than in large specimens when litter species were combined (Figure 2, Table 1). In leaf herbivores (i.e. grasshoppers), it has been shown that biomechanical traits such as mandible strength correlate well with plant toughness (Ibanez, Lavorel, Puijalon, & Moretti, 2013). Analogously, large-sized *Oniscus* individuals could afford a clearer switch towards *Betula* to fulfil their physiological demand of calcium and magnesium, probably thanks to a higher biting strength. Alternatively, the observed switch towards *Betula* might also be caused by the lower relative consumption rates showed by large-sized *Oniscus* (data not shown); as they consumed less leaf litter per unit body mass and unit time, they needed to spend more time feeding on *Betula*. On the other hand, small-sized *Oniscus* in mixed-size treatments might have profited from the feeding activity of those large-sized *Oniscus* that selected *Betula* (the tougher of the two leaf litter species), possibly by taking advantage of resulting small leaf fragments. A similar mechanism has been proposed for interspecific interactions between litter consumers: the presence of isopods and millipedes facilitate litter consumption by earthworms (Heemsbergen et al., 2004), and the same kind of interspecific facilitation was already shown to be driven by body size differences (Tonin et al., 2018). This hypothesis is also supported by our data, which shows that the positive effect on *Betula* mass loss in mixed-body size microcosms is not intermediate between the two single-size treatments, as one would expect, but much more similar to treatments with large *Oniscus* only (Figure 2, Table 1). The complementarity between large and small *Oniscus* in the consumption of *Betula* might have in turn favoured the feeding activity on *Acer* through transfer of calcium and magnesium (and/or other nutrients) by fungal hyphae or leaching (Hättenschwiler et al., 2005). Although this mechanism remains speculative, our results confirm that *Acer* mass loss tended to be higher than expected in mixed-size treatments, while one would have predicted a clear decrease based on single-size patterns (Figure 2, Table 1). Other indirect effects might also have played a role, such as microbial community changes caused by *Oniscus* activity (e.g. Vos et al., 2011). For example, small and large *Oniscus* might selectively feed on different microbial species or strains and so modify their competitive interactions and dominance

structure, thus increasing microbial activity and/or diversity which in turn lead to a positive effect on decomposition (Gessner et al., 2010).

Previous research has already detected intraspecific differences in macrodetritivore consumption rates, but an experimental investigation of their consequences for population-level niche partitioning and process rates was lacking so far. Vilisics et al. (2012) demonstrated that intraspecific body size differences in two common isopod species influence consumption rates of a single litter species, and different size classes also showed distinct patterns of litter degradation. Significant differences in consumption rates among conspecific individuals, driven by a variety of traits, have also been shown in analogous aquatic systems (Raffard et al., 2017; Rota et al., 2018). Interestingly, Reiss, Bailey, Perkins, Pluchinotta, and Woodward (2011) specifically tested the effect of intraspecific body size variation in aquatic detritivores on ecosystem functioning. However, the absence of non-additive effects in their study might be due to the use of one single leaf litter species, which decreases the opportunities for intraspecific niche partitioning. Our results built upon this previous knowledge (the differences in consumption rates detected so far among conspecific individuals do not automatically imply complementary resource use) and allowed us to experimentally demonstrate for the first time that intraspecific trait variation matters for small-scale niche partitioning in soil fauna, as it was already shown for primary producers in different systems (Fontana, Thomas, Reyes, & Pomati, 2019; He et al., 2018), and has consequences for fundamental ecosystem processes.

Our main goal was to provide evidence of the possibility of niche partitioning among individuals of the same species, independent of the reasons why niche partitioning arose. As a consequence, we tried to maximize body size differences being aware that size correlates with many other traits. In our case, ontogeny likely played a major role in determining individual differences. However, since in natural populations generations overlap and different developmental stages coexist at very small spatial scales and compete for the same resources (Al-Dabbagh & Block, 1981; Al-Jetlawi & Nair, 1994; Paris & Pitelka, 1962), this does not diminish the relevance of our results. The same is true for the fact that total litter decomposition was highest in the presence of large-sized individuals only (Figure 2), while relative consumption rates were highest in treatments with only small-sized individuals (data not shown), in accordance with previous results (Vilisics et al., 2012) and the metabolic theory of ecology. However, our a priori hypotheses target mechanisms of niche partitioning and thus focus on departures from additivity rather than absolute decomposition rates to assess the effect of both *Oniscus* size classes and litter combinations, and their interaction. Observed mass loss generally showed a higher variability than expected mass loss (Figure 2). This is not surprising, as we calculated expected values using average relative consumption rates of the litter monocultures, whereas observed values contained sources of variability other than total *Oniscus* mass differences (e.g. individual physiological status).

Although the role of litter combinations on ecosystem processes has been investigated quite extensively (e.g. Finerty et al., 2016;

Handa et al., 2014; Hättenschwiler et al., 2005), intraspecific trait diversity of soil macrodetritivores has been largely neglected so far. Our study represents an attempt to unravel the complex interactions regulating the process of litter decomposition. We argue that scaling down our focus to the level at which ecological interactions occur – that is between litter and individual consumers – can represent a possible way forward. By demonstrating that intraspecific feeding niche partitioning plays a role in the interactions between food resource and consumer diversity, we suggest here that individual-level trait measurements are a promising way to shed light on some poorly understood aspects of the mechanisms that determine decomposition in natural ecosystems. This calls for an urgent need of standardized individual-level trait measurements, especially of invertebrates (Didham, Edwards, Leather, & Basset, 2013; Ellers et al., 2018; Moretti et al., 2017). We are aware that our focus on a single macrodetritivore species represents a limitation and makes it difficult to generalize our conclusions. Nevertheless, it is noteworthy that intraspecific niche partitioning in *Oniscus* was driven by differences in size, which is an easy-to-measure trait that determines trophic interactions across taxa. Therefore, we consider our results as a first step that should encourage increased efforts in the same direction. Further work is needed to assess to which extent the patterns we observed are general, and including a finer resolution of diversity (e.g. more litter species and size categories, or different traits) might even strengthen our findings. Our results suggest that trait variability can reduce intraspecific competition; especially in case this relationship is nonlinear, as it was showed in analogous freshwater systems for the effect of shredding macroinvertebrate density (Little, Fronhofer, & Altermatt, 2018), fine-scale processes, as the ones elucidated in our experiment, might translate into consequences for ecosystem functioning at the landscape level. Beside the possible propagation across spatial scales of the effects described here, it is crucial to consider in BEF research how intraspecific trait variation might have cascading consequences for multiple processes and services (e.g. nutrient cycles, soil fertilization, plant germination, growth and defence), especially under environmental change scenarios.

ACKNOWLEDGEMENTS

S. Fontana was supported by the Swiss National Science Foundation, Project 315230_170200/1. We thank the gardeners' team at WSL for helping with litter collection, N. Forni for contributing to animal collection in the field, P. Stucki and A. Dias for their precious information about isopod ecology and taxonomy and O. Franken for the inspiring discussions. This manuscript was improved by the helpful comments of Y. Bieri and the two anonymous reviewers.

AUTHORS' CONTRIBUTIONS

S.F., M.M. and M.P.B. conceived the ideas and designed methodology; S.F. collected and analysed the data; S.F. led the writing of the

manuscript. All authors contributed critically to the drafts and gave final approval for publication.

DATA AVAILABILITY STATEMENT

The data analysed in this paper are available in the Zenodo Repository at <https://doi.org/10.5281/zenodo.3384279> (Fontana, Berg, & Moretti, 2019).

ORCID

Simone Fontana <https://orcid.org/0000-0003-1194-1909>

Matty P. Berg <https://orcid.org/0000-0001-8442-8503>

Marco Moretti <https://orcid.org/0000-0002-5845-3198>

REFERENCES

- Al-Dabbagh, K. Y., & Block, W. (1981). Population ecology of a terrestrial isopod in two Breckland grass heaths. *Journal of Animal Ecology*, 50, 61–77. <https://doi.org/10.2307/4031>
- AlJetlawi, A. A., & Nair, G. A. (1994). Breeding and population biology of *Armadillo officinalis* (Isopoda: Oniscidea) in Benghazi, Libya. *Journal of Arid Environments*, 27, 241–255. <https://doi.org/10.1006/jare.1994.1061>
- Barabás, G., & D'Andrea, R. (2016). The effect of intraspecific variation and heritability on community pattern and robustness. *Ecology Letters*, 19, 977–986. <https://doi.org/10.1111/ele.12636>
- Bardgett, R. D., & van der Putten, W. H. (2014). Belowground biodiversity and ecosystem functioning. *Nature*, 515, 505–511. <https://doi.org/10.1038/nature13855>
- Becker, A., Ziegler, A., & Epple, M. (2005). The mineral phase in the cuticles of two species of Crustacea consists of magnesium calcite, amorphous calcium carbonate and amorphous calcium phosphate. *Dalton Transactions*, 10, 1814–1820. <https://doi.org/10.1039/b412062k>
- Bíla, K., Moretti, M., de Bello, F., Dias, A. T., Pezzatti, G. B., Van Oosten, A. R., & Berg, M. P. (2014). Disentangling community functional components in a litter-macrodetritivore model system reveals the predominance of the mass ratio hypothesis. *Ecology and Evolution*, 4, 408–416. <https://doi.org/10.1002/ece3.941>
- Bolnick, D. I., Amarasekare, P., Araújo, M. S., Bürger, R., Levine, J. M., Novak, M., ... Vasseur, D. A. (2011). Why intraspecific trait variation matters in community ecology? *Trends in Ecology and Evolution*, 26, 183–192. <https://doi.org/10.1016/j.tree.2011.01.009>
- Bolnick, D. I., Svanbäck, R., Fordyce, J. A., Yang, L. H., Davis, J. M., Hulsey, C. D., & Forister, M. L. (2003). The ecology of individuals: Incidence and implications of individual specialization. *The American Naturalist*, 161, 1–28. <https://doi.org/10.1086/343878>
- Bonfanti, J., Hedde, M., Joimel, S., Krogh, P. H., Violle, C., Nahmani, J., & Cortet, J. (2018). Intraspecific body size variability in soil organisms at a European scale: Implications for functional biogeography. *Functional Ecology*, 32, 2562–2570. <https://doi.org/10.1111/1365-2435.13194>
- Cardinale, B. J., Duffy, J. E., Gonzalez, A., Hooper, D. U., Perrings, C., Venail, P., ... Naeem, S. (2012). Biodiversity loss and its impact on humanity. *Nature*, 486, 59–67. <https://doi.org/10.1038/nature11148>
- Chalmandrier, L., Münkemüller, T., Colace, M.-P., Renaud, J., Aubert, S., Carlson, B. Z., ... Thuiller, W. (2017). Spatial scale and intraspecific trait variability mediate assembly rules in alpine grasslands. *Journal of Ecology*, 105, 277–287. <https://doi.org/10.1111/1365-2745.12658>
- Clegg, T., Ali, M., & Beckerman, A. P. (2018). The impact of intraspecific variation on food web structure. *Ecology*, 99, 2712–2720. <https://doi.org/10.1002/ecy.2523>
- Coq, S., Nahmani, J., Resmond, R., Segrestin, J., David, J. F., Schevin, P., & Kazakou, E. (2018). Intraspecific variation in litter palatability to macroarthropods in response to grazing and soil fertility. *Functional Ecology*, 32, 2615–2624. <https://doi.org/10.1111/1365-2435.13205>
- Cornelissen, J. H. C. (1996). An experimental comparison of leaf decomposition rates in a wide range of temperate plant species and types. *Journal of Ecology*, 84, 573–582. <https://doi.org/10.2307/2261479>
- Coulis, M., Fromin, N., David, J., Gavinet, J., Clet, A., Devidal, S., ... Hättenschwiler, S. (2015). Functional dissimilarity across trophic levels as a driver of soil processes in a Mediterranean decomposer system exposed to two moisture levels. *Oikos*, 124, 1304–1316. <https://doi.org/10.1111/oik.01917>
- David, J.-F. (2014). The role of litter-feeding macroarthropods in decomposition processes: A reappraisal of common views. *Soil Biology and Biochemistry*, 76, 109–118. <https://doi.org/10.1016/j.soilbio.2014.05.009>
- David, J.-F., & Handa, I. T. (2010). The ecology of saprophagous macroarthropods (millipedes, woodlice) in the context of global change. *Biological Reviews of the Cambridge Philosophical Society*, 85, 881–895. <https://doi.org/10.1111/j.1469-185X.2010.00138.x>
- De Laender, F., Melián, C. J., Bindler, R., Van den Brink, P. J., Daam, M., Roussel, H., ... Janssen, C. R. (2014). The contribution of intra- and interspecific tolerance variability to biodiversity changes along toxicity gradient. *Ecology Letters*, 17, 72–81.
- de Oliveira, T., Hättenschwiler, S., & Handa, I. T. (2010). Snail and millipede complementarity in decomposing Mediterranean forest leaf litter mixtures. *Functional Ecology*, 24, 937–946. <https://doi.org/10.1111/j.1365-2435.2010.01694.x>
- Des Roches, S., Post, D. M., Turley, N. E., Bailey, J. K., Hendry, A. P., Kinnison, M. T., ... Palkovacs, E. P. (2018). The ecological importance of intraspecific variation. *Nature Ecology and Evolution*, 2, 57–64. <https://doi.org/10.1038/s41559-017-0402-5>
- Didham, R. K., Edwards, O. R., Leather, S. R., & Basset, Y. (2013). Arthropod diversity and the future of all-taxa inventories. *Insect Conservation and Diversity*, 6, 1–4. <https://doi.org/10.1111/icad.12022>
- Didham, R. K., Leather, S. R., & Basset, Y. (2016). Circle the bandwagons – Challenges mount against the theoretical foundations of applied functional trait and ecosystem service research. *Insect Conservation and Diversity*, 9, 1–3. <https://doi.org/10.1111/icad.12150>
- Eisenhauer, N., Antunes, P. M., Bennett, A. E., Birkhofer, K., Bissett, A., Bowker, M. A., ... Powell, J. R. (2017). Priorities for research in soil ecology. *Pedobiologia*, 63, 1–7. <https://doi.org/10.1016/j.pedobi.2017.05.003>
- Ellers, J., Berg, M. P., Dias, A. T. C., Fontana, S., Ooms, A., & Moretti, M. (2018). Diversity in form and function: Vertical distribution of soil fauna mediates multidimensional trait variation. *Journal of Animal Ecology*, 87, 933–944. <https://doi.org/10.1111/1365-2656.12838>
- Finerty, G. E., de Bello, F., Bíla, K., Berg, M. P., Dias, A. T. C., Pezzatti, G. B., & Moretti, M. (2016). Exotic or not, leaf trait dissimilarity modulates the effect of dominant species on mixed litter decomposition. *Journal of Ecology*, 104, 1400–1409. <https://doi.org/10.1111/1365-2745.12602>
- Fontana, S., Berg, M. P., & Moretti, M. (2019). Data from: Intraspecific niche partitioning in macrodetritivores enhances mixed leaf litter decomposition. *Zenodo Digital Repository*, <https://doi.org/10.5281/zenodo.3384279>
- Fontana, S., Thomas, M. K., Moldoveanu, M., Spaak, P., & Pomati, F. (2018). Individual-level trait diversity predicts phytoplankton community properties better than species richness or evenness. *The ISME Journal*, 12, 356–366. <https://doi.org/10.1038/ismej.2017.160>

- Fontana, S., Thomas, M. K., Reyes, M., & Pomati, F. (2019). Light limitation increases multidimensional trait evenness in phytoplankton populations. *The ISME Journal*, 13, 1159–1167. <https://doi.org/10.1038/s41396-018-0320-9>
- Gessner, M. O., Swan, C. M., Dang, C. K., McKie, B. G., Bardgett, R. D., Wall, D. H., & Hättenschwiler, S. (2010). Diversity meets decomposition. *Trends in Ecology and Evolution*, 25, 372–380. <https://doi.org/10.1016/j.tree.2010.01.010>
- Handa, I. T., Aerts, R., Berendse, F., Berg, M. P., Bruder, A., Butenschoten, O., ... Hättenschwiler, S. (2014). Consequences of biodiversity loss for litter decomposition across biomes. *Nature*, 509, 218–221. <https://doi.org/10.1038/nature13247>
- Hättenschwiler, S., & Gasser, P. (2005). Soil animals alter plant litter diversity effects on decomposition. *Proceedings National Academy Science United States of America*, 102, 1519–1524. <https://doi.org/10.1073/pnas.0404977102>
- Hättenschwiler, S., Tiunov, A. V., & Scheu, S. (2005). Biodiversity and litter decomposition in terrestrial ecosystems. *Annual Review of Ecology, Evolution, and Systematics*, 36, 191–218. <https://doi.org/10.1146/annurev.ecolsys.36.112904.151932>
- He, P., Fontana, S., Sui, X., Gessler, A., Schaub, M., Rigling, A., ... Li, M.-H. (2018). Scale dependent responses of pine reproductive traits to experimental and natural precipitation gradients. *Environmental and Experimental Botany*, 156, 62–73. <https://doi.org/10.1016/j.envexpbot.2018.08.028>
- Heemsbergen, D. A., Berg, M. P., Loreau, M., van Hal, J. R., Faber, J. H., & Verhoef, H. A. (2004). Biodiversity effects on soil processes explained by interspecific functional dissimilarity. *Science*, 306, 1019–1020. <https://doi.org/10.1126/science.1101865>
- Ibanez, S., Lavorel, S., Puijalon, S., & Moretti, M. (2013). Herbivory mediated by coupling between biomechanical traits of plants and grasshoppers. *Functional Ecology*, 27, 479–489. <https://doi.org/10.1111/1365-2435.12058>
- Jung, V., Violle, C., Mondy, C., Hoffmann, L., & Muller, S. (2010). Intraspecific variability and trait-based community assembly. *Journal of Ecology*, 98, 1134–1140. <https://doi.org/10.1111/j.1365-2745.2010.01687.x>
- Little, C. J., Fronhofer, E. A., & Altermatt, F. (2018). Nonlinear effects of intraspecific competition alter landscape-wide upscaling of ecosystem function. *bioRxiv*, <https://doi.org/10.1101/470591>
- MacArthur, R. H., & Levins, R. (1967). The limiting similarity, convergence and divergence of coexisting species. *The American Naturalist*, 101, 377–385. <https://doi.org/10.1086/282505>
- McGill, B. J., Enquist, B. J., Weiher, E., & Westoby, M. (2006). Rebuilding community ecology from functional traits. *Trends in Ecology and Evolution*, 21, 178–185. <https://doi.org/10.1016/j.tree.2006.02.002>
- Moretti, M., Dias, A. T., de Bello, F., Altermatt, F., Chown, S. L., Azcarate, F. M., ... Berg, M. P. (2017). Handbook of protocols for standardized measurement of terrestrial invertebrate functional traits. *Functional Ecology*, 31, 558–567. <https://doi.org/10.1111/1365-2435.12776>
- Muñoz, M. C., Schaefer, H. M., Böhning-Gaese, K., & Schleuning, M. (2017). Importance of animal and plant traits for fruit removal and seedling recruitment in a tropical forest. *Oikos*, 126, 823–832. <https://doi.org/10.1111/oik.03547>
- Nicolai, V. (1988). Phenolic and mineral content of leaves influences decomposition in European forest ecosystems. *Oecologia*, 75, 575–579. <https://doi.org/10.1007/BF00776422>
- Paris, O. H., & Pitelka, F. A. (1962). Population characteristics of the terrestrial isopod *Armadillidium vulgare* in California grassland. *Ecology*, 43, 229–248. <https://doi.org/10.2307/1931979>
- Pawar, S., Dell, A. I., & Savage, V. M. (2015). From metabolic constraints on individuals to the dynamics of ecosystems. In A. Belgrano, G. Woodward, & U. Jacob (Eds.), *Aquatic functional biodiversity: An ecological and evolutionary perspective* (pp. 3–36). San Diego, CA: Academic Press.
- Polis, G. A. (1984). Age structure component of niche width and intra-specific resource partitioning: Can age groups function as ecological species? *The American Naturalist*, 123, 541–564. <https://doi.org/10.1086/284221>
- R Core Team (2018). R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from www.r-project.org
- Raffard, A., Lecerf, A., Coet, J., Buoro, M., Lassus, R., & Cucherousset, J. (2017). The functional syndrome: Linking individual trait variability to ecosystem functioning. *Proceedings of the Royal Society B: Biological Sciences*, 284(1868), 20171893–<https://doi.org/10.1098/rspb.2017.1893>
- Raffard, A., Santoul, F., Cucherousset, J., & Blanchet, S. (2019). The community and ecosystem consequences of intraspecific diversity: A meta-analysis. *Biological Reviews*, 94(2), 648–661. <https://doi.org/10.1111/brv.12472>
- Reiss, J., Bailey, R. A., Perkins, D. M., Pluchinotta, A., & Woodward, G. (2011). Testing effects of consumer richness, evenness and body size on ecosystem functioning. *Journal of Animal Ecology*, 80, 1145–1154. <https://doi.org/10.1111/j.1365-2656.2011.01857.x>
- Reiss, J., Bridle, J. R., Montoya, J. M., & Woodward, G. (2009). Emerging horizons in biodiversity and ecosystem functioning research. *Trends in Ecology and Evolution*, 24, 505–514. <https://doi.org/10.1016/j.tree.2009.03.018>
- Rhoades, O. K., Best, R. J., & Stachowicz, J. J. (2018). Assessing feeding preferences of a consumer guild: Partitioning variation among versus within species. *The American Naturalist*, 192, 287–300. <https://doi.org/10.1086/698325>
- Rota, T., Jabiol, J., Chauvet, E., & Lecerf, A. (2018). Phenotypic determinants of inter-individual variability of litter consumption rate in a detritivore population. *Oikos*, 127, 1670–1678. <https://doi.org/10.1111/oik.05228>
- Schleuning, M., Fründ, J., & García, D. (2015). Predicting ecosystem functions from biodiversity and mutualistic networks: An extension of trait-based concepts to plant-animal interactions. *Ecography*, 38, 380–392. <https://doi.org/10.1111/ecog.00983>
- Schmitz, O. J., Buchkowski, R. W., Burghardt, K. T., & Donihue, C. M. (2015). Functional traits and trait-mediated interactions: Connecting community-level interactions with ecosystem functioning. In S. Pawar, G. Woodward, & A. I. Dell (Eds.), *Trait-based ecology – From structure to function* (pp. 319–343). London, UK: Advances in Ecological Research. Elsevier Ltd.
- Seibold, S., Cadotte, M. W., MacIvor, J. S., Thorn, S., & Müller, J. (2018). The necessity of multitrophic approaches in community ecology. *Trends in Ecology and Evolution*, 33, 754–764. <https://doi.org/10.1016/j.tree.2018.07.001>
- Siefert, A., Violle, C., Chalmandrier, L., Albert, C. H., Taudiere, A., Fajardo, A., ... Wardle, D. A. (2015). A global meta-analysis of the relative extent of intraspecific trait variation in plant communities. *Ecology Letters*, 18, 1406–1419. <https://doi.org/10.1111/ele.12508>
- Start, D. (2018). Ontogeny and consistent individual differences mediate trophic interactions. *The American Naturalist*, 192, 301–310. <https://doi.org/10.1086/698693>
- Tonin, A. M., Pozo, J., Monroy, S., Basaguren, A., Pérez, J., Gonçalves, J. F. Jr, ... Boyero, L. (2018). Interactions between large and small detritivores influence how biodiversity impacts litter decomposition. *Journal of Animal Ecology*, 87, 1465–1474. <https://doi.org/10.1111/1365-2656.12876>
- Vilisics, F., Szekeres, S., & Hornung, E. (2012). Size dependent differences in litter consumption of isopods: Preliminary results. *ZooKeys*, 176, 247–259. <https://doi.org/10.3897/zookeys.176.2470>
- Violle, C., Enquist, B. J., McGill, B. J., Jiang, L., Albert, C. H., Hulshof, C., ... Messier, J. (2012). The return of the variance: Intraspecific variability in community ecology. *Trends in Ecology and Evolution*, 27, 244–252. <https://doi.org/10.1016/j.tree.2011.11.014>

- Violle, C., Navas, M.-L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I., & Garnier, E. (2007). Let the concept of trait be functional! *Oikos*, 116, 882–892. <https://doi.org/10.1111/j.2007.0030-1299.15559.x>
- Volf, M., Redmond, C., Albert, Á. J., Le Bagousse-Pinguet, Y., Biella, P., Götzenberger, L., ... de Bello, F. (2016). Effects of long and short-term management on the functional structure of meadows through species turnover and intraspecific trait variability. *Oecologia*, 180, 941–950. <https://doi.org/10.1007/s00442-016-3548-y>
- Vos, V. C. A., van Ruijven, J., Berg, M. P., Peeters, E. T. H. M., & Berendse, F. (2011). Macrodetrivore identity drives leaf litter diversity effects. *Oikos*, 120, 1092–1098. <https://doi.org/10.1111/j.1600-0706.2010.18650.x>
- Vos, V. C. A., van Ruijven, J., Berg, M. P., Peeters, E. T. H. M., & Berendse, F. (2013). Leaf litter quality drives litter mixing effects through complementary resource use among detritivores. *Oecologia*, 173, 269–280. <https://doi.org/10.1007/s00442-012-2588-1>
- Wagg, C., Bender, S. F., Widmer, F., & van der Heijden, M. G. A. (2014). Soil biodiversity and soil community composition determine ecosystem multifunctionality. *Proceedings National Academy of Science United States of America*, 111, 5266–5270. <https://doi.org/10.1073/pnas.1320054111>
- Wall, D. H., Bardgett, R. D., & Kelly, E. (2010). Biodiversity in the dark. *Nature Geoscience*, 3, 297–298. <https://doi.org/10.1038/ngeo860>
- Werner, E. E., & Gilliam, J. F. (1984). The ontogenetic niche and species interactions in size-structured populations. *Annual Review of Ecology and Systematics*, 15, 393–425. <https://doi.org/10.1146/annurev.es.15.110184.002141>
- Zimmer, M. (2002). Nutrition in terrestrial isopods (Isopoda: Oniscidea): An evolutionary-ecological approach. *Biological Reviews*, 77, 455–493. <https://doi.org/10.1017/S1464793102005912>
- Zimmer, M. (2006). The role of animal-microbe interactions in isopod ecology and evolution. *Acta Biologica Benrodis*, 13, 127–168.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Fontana S, Berg MP, Moretti M.

Intraspecific niche partitioning in macrodetritivores enhances mixed leaf litter decomposition. *Funct Ecol*. 2019;33:2391–2401. <https://doi.org/10.1111/1365-2435.13448>