University of Groningen

A homozygous variant in growth and differentiation factor 2(GDF2) may cause lymphatic dysplasia with hydrothorax and nonimmune hydrops fetalis

Aukema, Sietse M.; ten Brinke, Gerdien A.; Timens, Wim; Vos, Yvonne J.; Accord, Ryan E.; Kraft, Karianne E.; Santing, Michiel J.; Morssink, Leonard P.; Streefland, Esther; van Diemen, Cleo C.

Published in:

American Journal of Medical Genetics. Part A

DOI:

[10.1002/ajmg.a.61743](https://doi.org/10.1002/ajmg.a.61743)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Aukema, S. M., ten Brinke, G. A., Timens, W., Vos, Y. J., Accord, R. E., Kraft, K. E., Santing, M. J., Morssink, L. P., Streefland, E., van Diemen, C. C., Vrijlandt, E. J. L. E., Hulzebos, C., & Kerstjens-Frederikse, W. S. (2020). A homozygous variant in growth and differentiation factor 2(GDF2) may cause lymphatic dysplasia with hydrothorax and nonimmune hydrops fetalis. *American Journal of Medical Genetics. Part A*, 182(9), 2152-2160. <https://doi.org/10.1002/ajmg.a.61743>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

CLINICAL REPORT

A homozygous variant in growth and differentiation factor 2 (*GDF2*) may cause lymphatic dysplasia with hydrothorax and nonimmune hydrops fetalis

Sietse M. Aukema¹ | Gerdien A. ten Brinke² | Wim Timens³ | Yvonne J. Vos¹ | Ryan E. Accord⁴ | Karianne E. Kraft² | Michiel J. Santing⁵ | Leonard P. Morssink⁶ | Esther Streefland⁷ | Cleo C. van Diemen¹ | Elianne JLE Vrijlandt⁸ | Christian V. Hulzebos² | Wilhelmina S. Kerstjens-Frederikse¹

¹Department of Clinical Genetics, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands

²Beatrix Children's Hospital, University Medical Center Groningen, Groningen, The Netherlands

³Department of Pathology and Medical Biology, University of Groningen, University Medical Centre Groningen, Groningen, The Netherlands

⁴Department of Congenital Cardiothoracic Surgery, University of Groningen, University Medical Center Groningen, Beatrix Children's Hospital, Center for Congenital Heart Diseases, Groningen, The Netherlands

⁵Department of Radiology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands

⁶Department of Obstetrics and Gynaecology, Medical Center Leeuwarden, Leeuwarden, The Netherlands

⁷Department of Obstetrics and Gynecology/Prenatal diagnosis, University Medical Centre of Groningen, University of Groningen, Groningen, The Netherlands

⁸Department of Pediatric Pulmonology and Pediatric Allergy, Beatrix Children's Hospital, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands

Correspondence

Sietse M. Aukema, Department of Clinical Genetics, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands.
Email: smaukema@gmail.com

Present address

Sietse M. Aukema, Department of Clinical Genetics, Maastricht University Medical Centre, Maastricht, The Netherlands

Abstract

The etiology of nonimmune hydrops fetalis is extensive and includes genetic disorders. We describe a term-born female neonate with late onset extensive nonimmune hydrops, that is, polyhydramnios, edema, and congenital bilateral chylothorax. This newborn was successfully treated with repetitive thoracocentesis, total parenteral feeding, octreotide intravenously and finally surgical pleurodesis and corticosteroids. A genetic cause seemed plausible as the maternal history revealed a fatal nonimmune hydrops fetalis. A homozygous truncating variant in *GDF2* (c.451C>T, p.(Arg151*)) was detected with exome sequencing. Genetic analysis of tissue obtained from the deceased fetal sibling revealed the same homozygous variant. The parents and two healthy siblings were heterozygous for the *GDF2* variant. Skin and lung biopsies in the index patient, as well as the revised lung biopsy of the deceased fetal sibling, showed lymphatic dysplasia and lymphangiectasia. To the best of our knowledge, this is the first report of an association between a homozygous variant in *GDF2* with lymphatic dysplasia, hydrothorax and nonimmune hydrops fetalis.

KEYWORDS

BMP9, *GDF2*, hereditary hemorrhagic telangiectasia, lymphatic dysplasia, nonimmune hydrops fetalis, pulmonary arterial hypertension

1 | INTRODUCTION

Nonimmune hydrops fetalis is a condition characterized by excessive accumulation of fluid in at least two fetal extravascular compartments or body cavities, including ascites, pericardial effusion, pleural effusion and skin edema. Placental thickening and polyhydramnios are other frequent findings. The majority of cases of hydrops fetalis is classified as nonimmune hydrops fetalis (NIHF), that refers to hydrops due to other causes than red cell alloimmunization (Bellini et al., 2015; Norton, Chauhan, & Dashe, 2015). NIHF has a reported prevalence of 1:1700 to 1:3000 pregnancies (Norton et al., 2015) and is associated with a high mortality rate with an overall mortality up to 45% at 1 year (Fukushima et al., 2011; Nassr et al., 2018; Steurer et al., 2017). Identification of the underlying cause may guide perinatal management of contemporary cases and genetic counseling in future pregnancies (Mardy, Chetty, Norton, & Sparks, 2019; Sparks et al., 2019). NIHF has a wide array of underlying etiologies.

A recent systematic review (Bellini et al., 2015) lists 14 different etiological-categories and their relative frequencies. Cardiovascular anomalies are the most frequent cause of NIHF ($\approx 20\%$). Genetic disorders may also lead to NIHF and include a wide spectrum of diseases, including chromosomal aneuploidies (trisomies, monosomy X, triploidy), hematological diseases (α -thalassemia), monogenic syndromes (RASopathies, Kabuki syndrome), inborn errors of metabolism (lysosomal storage disease) and lymphatic dysplasia (Bellini et al., 2015; Mardy et al., 2019; McPherson, 2019; Moreno et al., 2013; Quinlan-Jones et al., 2019; Weissbach et al., 2019). Few familial cases of nonimmune hydrops fetalis have been described and in some of the cases with lymphatic dysplasia and/or lymphangiectasia bi-allelic variants in *CCBE1*, *ADAMTS3*, *FAT4*, *CALCRL*, and *PIEZO1* have been reported (Alders et al., 2009; Alders et al., 2014; Brouillard et al., 2017; Datkhaeva et al., 2018; Delabaere et al., 2008; Fotiou et al., 2015; Jacquemont, Barbarot, Boceno, Stalder, & David, 2000; Mackie et al., 2018; Njolstad, Reigstad, Westby, & Espeland, 1998; Stevenson, Pysher, Ward, & Carey, 2006; Wieacker, Muschke, Pollak, & Muller, 2005). However, even with conventional and state-of-the-art diagnostics like exome sequencing an underlying genetic defect remains undetermined in the majority of infants with hydrops (Lord et al., 2019; Yates et al., 2017). A genetic diagnosis for NIHF may affect counseling during the pregnancy, neonatal management as well as genetic counseling in future pregnancies including the possibility of preimplantation diagnostics.

In this report, we describe the familial occurrence of late onset nonimmune hydrops fetalis in two siblings in which we identified a homozygous truncating variant in *GDF2*. To the best of our knowledge, an association between *GDF2* with lymphatic dysplasia, hydrothorax, and NIHF has not been described previously. We propose an etiologic role of this gene in lymphatic dysplasia and concurrent NIHF.

2 | PATIENTS AND METHODS

2.1 | Patients

A family with two siblings with late onset (McPherson, 2019) NIHF and two healthy siblings is reported. The parents of the index patient signed a written informed consent for rapid diagnostic exome sequencing (trio analysis of the proband, mother and father), consented with targeted variant analysis of the three siblings and signed consent for publication.

2.2 | Immunohistochemistry

Tissues were fixed in neutral buffered formalin and embedded in paraffin. Immunohistochemistry was performed on fresh 3 μ m sections on a Ventana Benchmark Ultra autostainer (Ventana Medical Systems / Roche-diagnostics, Tucson, AZ). For immunohistochemical demonstration of podoplanin (PDPN) the D2-40 mouse monoclonal antibody was used (Roche Diagnostics, Catalogue Number 760-4395) in accordance with the manufacturer's instructions.

2.3 | Molecular studies

Exome sequencing and variant calling were performed as previously described (Herkert et al., 2018). In brief, the exome was captured with the Agilent SureSelect XT Human All Exon V6 kit (Agilent, Santa Clara, CA) and exome libraries were sequenced on a NextSeq500 (Illumina, San Diego, CA) with 2x 150 bp paired-end reads at an average coverage of 100x and with >90% of the exome covered >20x. Sequence reads were aligned to the human reference genome (UCSC version GRCh37/hg19) with the Burrows-Wheeler Aligner version 0.7.5a. Sambamba was used to process the aligned reads, after which we applied Genome Analysis Tool Kit (GATK) duplicate removal and performed SNP and INDEL discovery and genotyping using standard hard filtering parameters according to GATK Best Practices recommendations. Sequence variants were filtered with Cartagenia Next-Generation Sequencing-Bench Laboratory software (Agilent, Santa Clara, CA) by using an automated filtering tree. Variant classification was done according to the ACMG-guidelines (Richards et al., 2015). For data-analysis an updated version of a previously published virtual genepanel (van Diemen et al., 2017) containing approximately 3,850 genes at the time of analysis was used. In brief, this virtual gene panel contains monogenic diseases listed in the clinical genomic database (CGD) with the exception that genes associated with late-onset diseases were removed (van Diemen et al., 2017). In addition, individual genes from a standard, clinical exome-capturing panel (SureSelect Inherited Disease; Agilent, Santa Clara, CA) but not included in the CGD were added.

2.4 | Review of *GDF2* variants

We mined the published literature for patients with a germline variant in *GDF2* and summarized their genotypes and associated phenotypes (Abou Hassan et al., 2018; Eyries et al., 2019; Graf et al., 2018; Hernandez et al., 2015; Hodgson et al., 2019; G. Wang et al., 2016; X.-J. Wang et al., 2019; Wooderchak-Donahue et al., 2013; Yang et al., 2018; Zhu et al., 2019) including the present study.

3 | CLINICAL REPORT

3.1 | Case report

3.1.1 | Prenatal history

A female neonate was born via an emergency cesarean section at term to a multigravida (G5) with two healthy children. The maternal obstetric history revealed a spontaneous abortion at ~10 weeks and an intrauterine fetal death at 33 weeks of gestation. The intrauterine fetal death was due to nonimmune hydrops fetalis with hydrothorax and amniocentesis revealed a mosaic trisomy 20 by both conventional karyotyping (47,XX,+20[7]/46,XX[4]) and array analysis. The current pregnancy was again complicated by late onset NIHF with polyhydramnios and a right-sided hydrothorax first detected at 34 + 6 weeks of gestation. Prenatal evaluation showed no evidence of immune hydrops, congenital infections, or cardiac abnormalities.

3.1.2 | Post partum diagnostic evaluation and management

After birth extensive subcutaneous edema, especially in the head and neck region, without other dysmorphic features, was observed in this newborn. Apgar scores were 2, 7, and 8 after 1, 5, and 10 min, respectively. Respiratory failure developed immediately after birth requiring mechanical ventilation. A chest X-ray showed bilateral pleural effusions. Chest tubes were placed; the biochemical analyses of the fluid was consistent with chyle. Additional treatment consisted of nil per os and total parenteral nutrition. Chylous volumes averaged up to 900 ml per day, and therefore intravenous infusion of Octreotide was started. Initially there was a gradual decrease in chylous volumes, so after several weeks chest tubes could be removed and Octreotide infusion could be discontinued. Unfortunately, the production of chyle increased after a medium-chain triglyceride diet was started. Enteral feeding was discontinued and Octreotide was restarted. In the meantime, a high-resolution computed tomography (HRCT) of the chest was suggestive for interstitial lung disease with diffuse ground glass opacities, thickened inter- and intralobar septa and prominent pulmonary vessels. These findings suggested an obstructive disorder of the lymphatic outflow trajectory.

3.1.3 | Surgical management and follow-up

Surgical treatment was deemed necessary after weeks of mechanical ventilation and failure of conservative management. A video-assisted thoracic surgery (VATS) on the left side was performed with a pleurectomy and pleural drainage. Due to the presence of adhesions the right-sided VATS was converted to a thoracotomy and subsequently chemical pleurodesis was performed. A lung and skin biopsy were taken in order to provide histopathological support for the presumed diagnosis. Lung histology showed interstitial emphysema, and some lymphangiectasia, and interstitial pathology compatible with pulmonary interstitial glycogenosis (Figure 1b). Skin histology showed features of lymphatic vascular malformations (Figure 1c). Given the severity of the disease and uncertainty about the contributing role of the pulmonary interstitial glycogenosis, methylprednisolone therapy was given for 3 days. Clinical improvement occurred followed by an uneventful neonatal course. Currently, at the age of 1 year, she is doing fine on a normal diet without respiratory symptoms, without hepatosplenomegaly and without signs of peripheral lymphedema. She has a normal mental and psychomotor development.

3.1.4 | Molecular studies

A SNP-array showed no relevant chromosomal numerical abnormalities and no evidence for a (large) deletion of the *GBA* locus. In addition several large (>9 Mb) regions of homozygosity were identified (including the *GDF2* region). Targeted exome sequencing was performed on DNA from the proband and her parents. A homozygous truncating variant in *GDF2* (c.451C > T, p.(Arg151*)), and two heterozygous variants (in *WNT10A* and *GBA*) were identified. Both parents were heterozygous for the *GDF2* variant. The *WNT10A* was paternally inherited and interpreted as unrelated to the phenotype. The heterozygous *GBA* variant was maternally inherited and no second variant was found. The SNP-array was re-analyzed for the *GBA* locus and no deletion was detected. Although small deletions below the SNP-array resolution cannot be fully excluded these seem very rare in *GBA* (Pastores & Hughes, 1993). Moreover, clinically there were no signs of Gaucher's disease (normal beta-glucosidase/glucocerebrosidase activity and no increased plasma chitotriosidase activity). Therefore autosomal recessive Gaucher's disease was excluded as a cause of the fetal hydrops. Subsequent analysis of the *GDF2* variant by targeted sequencing identified homozygosity of c.451C > T / p.(Arg151*) in the affected stillbirth and heterozygosity in her unaffected siblings (Figure S4). A submission of the variant in GeneMatcher (Sobreira, Schiettecatte, Valle, & Hamosh, 2015) did not result in any matches for homozygous or compound heterozygous variants (match status last checked 04.05.2020).

4 | DISCUSSION

4.1 | General discussion

We report a family with two siblings affected by fetal hydrops carrying a homozygous truncating variant in *GDF2*. *GDF2* encodes the

FIGURE 1 Pedigree, histopathological evaluation and genotype–phenotype characteristics of (published) *GDF2* variants. (a) pedigree of the family with unaffected heterozygous depicted by half-filled symbols and affected homozygous individuals by filled symbols. The index patients is indicated by an arrow ↗. (b,c) representative histopathological analysis of lung and skin biopsies of the index patient (b) lung biopsy (x40) and (c) skin biopsy. (B) lung biopsy (bar = 1 mm) showing lymphangiectasis (long arrows) as well as interstitial emphysema (short arrows) and (c) skin biopsy (bar = 0.5 mm) showing lymphangiectasis in upper dermis as well as subcutis (arrows). The lymphatic origin was confirmed with D2-40 (PDPN) staining (data not shown). (d) schematic representation of published *GDF2*-variants. Protein domains according to Wang et al. (X.-J. Wang et al., 2019) and UniProt, <https://www.uniprot.org/uniprot/Q9UK05>. Lay-out adapted from Wagener et al. (Wagener et al., 2015) SP indicates signal peptide, AA amino acid. Splice-site variants are mapped against the amino-acid position corresponding to the position of the nucleotide change (Skoric-Milosavljevic et al., 2019). Different variants at the nucleotide level may lead to different amino-acid substitutions at the same amino-acid position. Variants reported in control cohorts as well as those classified as (likely) benign in the reviewed papers were not included in the figure

circulating bone morphogenetic protein 9 (BMP9), a protein that plays a role in angiogenesis. BMP9 is a ligand that binds with high affinity to activin receptor-like kinase 1 (ALK1) and Endoglin, members of the transforming growth factor- β family (TGF β) and is expressed in endothelial cells (Kienast et al., 2016; Lawera et al., 2019; Saito et al., 2017; Townson et al., 2012). The BMP9/ALK1/Endoglin pathway is involved in the regulation of angiogenesis by inhibition of endothelial cell proliferation and migration (David, Feige, & Bailly, 2009). In addition to its role in angiogenesis TGF β signaling is also involved in lymphatic development (James, Nalbandian, & Mukouyama, 2013; Oka et al., 2008). Indeed, analysis of the STRING database (<https://string-db.org/>) shows an enrichment of protein–protein interactions between *GDF2* and other proteins encoded by

germline predisposition genes involved in pulmonary arterial hypertension (PAH) and hereditary hemorrhagic telangiectasia (HHT) (Figure S1) as well as overlap between the genotypes and phenotypes of PAH, HHT, and hydrops (Figure S2). Variants in *GDF2* (almost exclusively heterozygous variants) have emerged as disease causing in HHT (MIM #615506) and PAH (Figure 1d) (Abou Hassan et al., 2018; Eyries et al., 2019; Graf et al., 2018; Hernandez et al., 2015; Hodgson et al., 2019; G. Wang et al., 2016; X.-J. Wang et al., 2019; Wooderchak-Donahue et al., 2013; Zhu et al., 2019). These variants are predominantly missense and protein truncating variants (nonsense and frameshift) and loss-of-function has been proposed as the pathogenic mechanism (Hodgson et al., 2019; Southgate, Machado, Graf, & Morrell, 2019). Although no clear hotspots exist, ~60% of the variants

affect the relatively small mature domain (109 AA) either directly by missense or by protein truncating variants (Figure S3). The majority of the protein truncating variants (including the Arg151* in our patients) will likely not be subjected to nonsense mediated mRNA decay (NMD) as the premature termination codons (PTC) are located in the terminal exon of the gene (exon 2) (Kurosaki, Popp, & Maquat, 2019; Lykke-Andersen & Jensen, 2015). As no functional studies were conducted it remains unclear if indeed a truncated protein is formed and whether there is any residual protein activity.

4.2 | The pulmonary arterial hypertension (PAH) phenotype

Homozygous and compound heterozygous variants in *GDF2* have been very rarely reported previously. Wang et al. (G. Wang et al., 2016) reported a child with PAH carrying a homozygous truncating variant (c.76C > T, p.[Gln26*]). In this case the early onset and severity of PAH is most likely due to the homozygosity of the variant. Incomplete penetrance of the heterozygous variant may account for the negative family history of PAH in both the parental and maternal family. Incomplete penetrance is supported by the finding that most of the reported patients with PAH and *GDF2* variants represent sporadic/idiopathic cases without a family history of PAH (Eyries et al., 2019; Graf et al., 2018; Hodgson et al., 2019; G. Wang et al., 2016; Zhu et al., 2019). Nonpenetrance of the heterozygous variant may also be the case in our family. In the series published by Abou Hassan et al. (Abou Hassan et al., 2018) one patient with PAH was compound heterozygous for two variants (c.254C>T, p.(Pro85-Leu)/c.652G>A, p.(Asp218Asn)). However, as the unaffected father of this patient carried the p.(Asp218Asn) variant in homozygous state and as this variant is frequently detected (in homozygous state) in non-PAH conditions (Hodgson et al., 2019) and has been classified as benign based on functional studies by Hodgson et al. it is unclear whether this compound heterozygous variant is comparable with other reported bi-allelic variants. Finally, a homozygous variant of undetermined clinical significance (VUS) was reported in a PAH patient by Yang et al. (Yang et al., 2018). However, the phenotype of hydrops and hydrothorax/chylothorax as seen in our family has not been reported yet. The lymphangiectasia observed in our patients are unlikely to be a manifestation of PAH as in PAH only very small angiectasies can be seen and only as component of the characteristic plexiform lesions as seen in advanced stages. In PAH teleangiectasies/lymphangiectasies are not part of the histopathological spectrum.

4.3 | The lymphatic dysplasia and hydrops phenotype

Several observations support an etiologic role for the homozygous *GDF2* variant in this NIHF phenotype. First, the two affected patients were homozygous for the c.451C > T, p.(Arg151*) while the unaffected siblings and parents were heterozygous. Second, the skin- and

lung biopsies in the index patient and affected sibling showed lymphatic dysplasia and lymphangiectasia (Figure 1b,c). The lymphangiectasia in the affected sibling was identified only after histopathological and immunohistochemical re-evaluation confirming the observation that lymphatic dysplasia is a likely underestimated etiology in the evaluation of NIHF when relying on histopathological evaluation only (Bellini et al., 2010). The third reason to suggest an etiological role for *GDF2* variants in NIHF is supplied by animal studies: *Bmp9* knock-out (KO) mice show several defects in lymphatic development and function (Levet et al., 2013; Yoshimatsu et al., 2013). As noted before, BMP9 is the ligand for the ALK1 receptor and blockade of ALK1 signaling also results in defects in lymphatic development (Niessen, Zhang, Ridgway, Chen, & Yan, 2010). Fourth, with SNP-array analysis and massive parallel sequencing other major genetic causes of NIHF were excluded. This included hematological disease, chromosomal aneuploidies, and variants in genes known to cause lymphatic dysplasia including RASopathies, (generalized) lymphedema, or inborn errors of metabolism (Bellini et al., 2015; Hakami, Dillon, Lebo, & Mason-Suares, 2016; Houweling et al., 2010; Johnston et al., 2018; Joyce et al., 2016; Mardy et al., 2019; Martin-Almedina et al., 2016; Mason-Suares et al., 2017; Meng et al., 2019; Moreno et al., 2013; Pagnamenta et al., 2019; Quinlan-Jones et al., 2019; Stuurman et al., 2019; Sudrie-Arnaud et al., 2018; Weissbach et al., 2019; Yates et al., 2017). Finally, no other causes for the NIHF, such as intra-uterine infections and cardiac abnormalities, were present. The mosaic trisomy 20 observed in cultured amnion cells from the deceased fetal sibling is unlikely to have played a role as no single case of (mosaic) trisomy 20 was identified among 1,004 cases (including 199 with chromosomal abnormalities) with NIHF in the series published by Meng et al. (Meng et al., 2019) and for reasons outlined below (see "the pathophysiological mechanism").

4.4 | The pathophysiological mechanism

Despite this specific genetic diagnosis of a homozygous variant in *GDF2*, the exact pathophysiology in these NIHF cases remains not fully elucidated for at least two reasons: First, the histological findings in the index patient were characteristic for pulmonary interstitial glycogenosis (PIG, a rare pediatric lung disease characterized by accumulation of cytoplasmic glycogen in mesenchymal cells in the alveolar interstitium, with unknown underlying etiology and clinical significance). This glycogenosis is associated with a variety of neonatal pulmonary and cardiovascular disorders and may appear to be transient with usually a favorable prognosis (Canakis, Cutz, Manson, & O'Brodovich, 2002; Cutz, Chami, Dell, Langer, & Manson, 2017; Seidl et al., 2018). The combination of PIG, pulmonary lymphangiectasia, and pleural effusion has been described previously (Cutz et al., 2017; Deutsch & Young, 2016) and supports the finding that PIG is associated with pulmonary disorders, including an abnormal lung development such as pulmonary lymphangiectasia. We presume that PIG is not responsible for the main symptoms in our case, but it is tempting to speculate that PIG is part of the phenotype with hydrops and

chylothorax or even is a consequence of impaired differentiation or maturation of the lung due to the pulmonary abnormalities. Second, the multi-treatment-strategy for our patient makes it impossible to discriminate between specific treatment effects and their role in clinical recovery. Repeating a lung HRCT and biopsy might potentially help to unravel this diagnostic issues, but is obviously potentially harmful and therefore not performed. Despite both affected siblings had similar prenatal ultrasonographic findings with hydrops and hydrothorax and both were homozygous for the *GDF2* variant the outcome was fatal in the prior pregnancy were also a mosaic trisomy 20 was detected. Mosaic trisomy 20 is a frequent finding in invasive prenatal diagnostics and (likely) in the majority of cases of extraembryonic origin (R Wallerstein et al., 2000). It is not clearly associated with a specific phenotype (Bianca et al., 2008; Robinson et al., 2005; Willis, Bird, Dell'Aquila, & Jones, 2008) but a poor outcome has been reported in individual cases (with a high-level mosaicism for trisomy 20) (Robinson et al., 2005; Robert Wallerstein et al., 2015). So the mosaic trisomy 20 might have contributed to the fatal outcome. However, in our opinion a full relation between the mosaic trisomy 20 and the hydrops phenotype seems unlikely (Robinson et al., 2005; Robert Wallerstein et al., 2015).

4.5 | Concluding remarks

In conclusion we report on a family with fatal as well as nonfatal NIHF in two siblings and propose, based on molecular evaluation, clinical- and histopathological analyses and literature review a critical role for *GDF2* in the observed phenotype. Although some uncertainty remains about the relation between this gene and the presenting symptoms, it is important to consider this specific genetic cause in case of unexplained NIHF. Hopefully other cases will be recognized after this report to confirm the causal relation. The identification of the underlying genetic defect not only provides important information for genetic counseling in future pregnancies (with a recurrence risk of 25% in the present family) but it also provides an option for preimplantation diagnostics. In addition, this report may affect perinatal counseling and management for further families. In particular because we reported not only a fatal case, but also a favorable outcome after intensive treatment of an infant with NIHF.

ACKNOWLEDGMENTS

The authors would like to thank the patient's parents for approving publication of the data.

CONFLICT OF INTEREST

The authors have no conflict of interest.

AUTHOR CONTRIBUTIONS

Designed the research: Sietse M. Aukema, Gerdien A. ten Brinke, Christian V. Hulzebos, Wilhelmina S. Kerstjens-Frederikse. Contributed clinical or pathological data: Wim Timens, Yvonne J. Vos, Ryan E. Accord, Karianne E. Kraft, Michiel J. Santing, Leonard P. Morssink,

Esther Streefland, Cleo C. van Diemen, Elianne JLE Vrijlandt. Manuscript writing: Sietse M. Aukema, Gerdien A. ten Brinke, Christian V. Hulzebos, Wilhelmina S. Kerstjens-Frederikse. Revision and approval of the manuscript: all authors.

DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

ORCID

Sietse M. Aukema <https://orcid.org/0000-0001-5824-0320>

REFERENCES

- Abou Hassan, O. K., Haidar, W., Nemer, G., Skouri, H., Haddad, F., & BouAkl, I. (2018). Clinical and genetic characteristics of pulmonary arterial hypertension in Lebanon. *BMC Medical Genetics*, 19(1), 89. <https://doi.org/10.1186/s12881-018-0608-7>
- Alders, M., Al-Gazali, L., Cordeiro, I., Dallapiccola, B., Garavelli, L., Tuysuz, B., ... Hennekam, R. C. (2014). Hennekam syndrome can be caused by *FAT4* mutations and be allelic to Van Maldergem syndrome. *Human Genetics*, 133(9), 1161–1167. <https://doi.org/10.1007/s00439-014-1456-y>
- Alders, M., Hogan, B. M., Gjini, E., Salehi, F., Al-Gazali, L., Hennekam, E. A., ... Hennekam, R. C. (2009). Mutations in *CCBE1* cause generalized lymph vessel dysplasia in humans. *Nature Genetics*, 41(12), 1272–1274. <https://doi.org/10.1038/ng.484>
- Bellini, C., Donarini, G., Paladini, D., Calevo, M. G., Bellini, T., Ramenghi, L. A., & Hennekam, R. C. (2015). Etiology of non-immune hydrops fetalis: An update. *American Journal of Medical Genetics. Part A*, 167A(5), 1082–1088. <https://doi.org/10.1002/ajmg.a.36988>
- Bellini, C., Fulcheri, E., Rutigliani, M., Calevo, M. G., Boccardo, F., Campisi, C., ... Hennekam, R. C. M. (2010). Immunohistochemistry in non-immune hydrops fetalis: A single center experience in 79 fetuses. *American Journal of Medical Genetics. Part A*, 152A(5), 1189–1196. <https://doi.org/10.1002/ajmg.a.33191>
- Bianca, S., Boemi, G., Barrano, B., Cataliotti, A., Ingegnosi, C., Indaco, L., & Ettore, G. (2008, July). Mosaic trisomy 20: Considerations for genetic counseling. *American Journal of Medical Genetics. Part A*, 146A, 1897–1898. <https://doi.org/10.1002/ajmg.a.32338>
- Brouillard, P., Dupont, L., Helaers, R., Coulie, R., Tiller, G. E., Peeden, J., ... Vikkula, M. (2017). Loss of *ADAMTS3* activity causes Hennekam lymphangiectasia-lymphedema syndrome 3. *Human Molecular Genetics*, 26(21), 4095–4104. <https://doi.org/10.1093/hmg/ddx297>
- Canakis, A.-M., Cutz, E., Manson, D., & O'Brodovich, H. (2002). Pulmonary interstitial glycogenosis: A new variant of neonatal interstitial lung disease. *American Journal of Respiratory and Critical Care Medicine*, 165(11), 1557–1565. <https://doi.org/10.1164/rccm.2105139>
- Cutz, E., Chami, R., Dell, S., Langer, J., & Manson, D. (2017). Pulmonary interstitial glycogenosis associated with a spectrum of neonatal pulmonary disorders. *Human Pathology*, 68, 154–165. <https://doi.org/10.1016/j.humpath.2017.06.026>
- Datkhaeva, I., Arboleda, V. A., Senaratne, T. N., Nikpour, G., Meyerson, C., Geng, Y., ... Janzen, C. (2018). Identification of novel *PIEZO1* variants using prenatal exome sequencing and correlation to ultrasound and autopsy findings of recurrent hydrops fetalis. *American Journal of Medical Genetics. Part A*, 176(12), 2829–2834. <https://doi.org/10.1002/ajmg.a.40533>
- David, L., Feige, J.-J., & Bailly, S. (2009). Emerging role of bone morphogenetic proteins in angiogenesis. *Cytokine & Growth Factor Reviews*, 20(3), 203–212. <https://doi.org/10.1016/j.cytogr.2009.05.001>
- Delabaere, A., Laurichesse-Delmas, H., Varlet, M. N., Clemenson, A., Dechelotte, P. J., Beaufre, A. M., ... Gallot, D. (2008, April). Recurrent

- congenital pulmonary lymphangiectasia. *Ultrasound in Obstetrics & Gynecology: The Official Journal of the International Society of Ultrasound in Obstetrics and Gynecology*, 31, 479–481. <https://doi.org/10.1002/uog.5284>
- Deutsch, G. H., & Young, L. R. (2016). Lipofibroblast phenotype in pulmonary interstitial Glycogenosis. *American Journal of Respiratory and Critical Care Medicine*, 193(6), 694–696. <https://doi.org/10.1164/rccm.201509-1809LE>
- Eyries, M., Montani, D., Nadaud, S., Girerd, B., Levy, M., Bourdin, A., ... Soubrier, F. (2019). Widening the landscape of heritable pulmonary hypertension mutations in paediatric and adult cases. *The European Respiratory Journal*, 53(3), 1801371. <https://doi.org/10.1183/13993003.01371-2018>
- Fotou, E., Martin-Almedina, S., Simpson, M. A., Lin, S., Gordon, K., Brice, G., ... Ostergaard, P. (2015). Novel mutations in PIEZO1 cause an autosomal recessive generalized lymphatic dysplasia with non-immune hydrops fetalis. *Nature Communications*, 6, 8085. <https://doi.org/10.1038/ncomms9085>
- Fukushima, K., Morokuma, S., Fujita, Y., Tsukimori, K., Satoh, S., Ochiai, M., ... Wake, N. (2011). Short-term and long-term outcomes of 214 cases of non-immune hydrops fetalis. *Early Human Development*, 87(8), 571–575. <https://doi.org/10.1016/j.earlhumdev.2011.04.015>
- Graf, S., Haimel, M., Bleda, M., Hadinnapola, C., Southgate, L., Li, W., ... Morrell, N. W. (2018). Identification of rare sequence variation underlying heritable pulmonary arterial hypertension. *Nature Communications*, 9(1), 1416. <https://doi.org/10.1038/s41467-018-03672-4>
- Hakami, F., Dillon, M. W., Lebo, M., & Mason-Suares, H. (2016). Retrospective study of prenatal ultrasound findings in newborns with a Noonan spectrum disorder. *Prenatal Diagnosis*, 36(5), 418–423. <https://doi.org/10.1002/pd.4797>
- Herkert, J. C., Abbott, K. M., Birnie, E., Meems-Veldhuis, M. T., Boven, L. G., Benjamins, M., ... Jongbloed, J. D. H. (2018). Toward an effective exome-based genetic testing strategy in pediatric dilated cardiomyopathy. *Genetics in Medicine: Official Journal of the American College of Medical Genetics*, 20(11), 1374–1386. <https://doi.org/10.1038/gim.2018.9>
- Hernandez, F., Huether, R., Carter, L., Johnston, T., Thompson, J., Gossage, J. R., ... Elliott, A. M. (2015). Mutations in RASA1 and GDF2 identified in patients with clinical features of hereditary hemorrhagic telangiectasia. *Human Genome Variation*, 2, 15040. <https://doi.org/10.1038/hgv.2015.40>
- Hodgson, J., Swietlik, E. M., Salmon, R. M., Hadinnapola, C., Nikolic, I., Wharton, J., ... Morrell, N. W. (2019). Characterization of GDF2 mutations and levels of BMP9 and BMP10 in pulmonary arterial hypertension. *American Journal of Respiratory and Critical Care Medicine*, 201(5), 575–585. <https://doi.org/10.1164/rccm.201906-1141OC>
- Houweling, A. C., de Mooij, Y. M., van der Burgt, I., Yntema, H. G., Lachmeijer, A. M. A., & Go, A. T. J. I. (2010, March). Prenatal detection of Noonan syndrome by mutation analysis of the PTPN11 and the KRAS genes. *Prenatal Diagnosis*, 30, 284–286. <https://doi.org/10.1002/pd.2458>
- Jacquemont, S., Barbarot, S., Boceno, M., Stalder, J. F., & David, A. (2000). Familial congenital pulmonary lymphangiectasia, non-immune hydrops fetalis, facial and lower limb lymphedema: Confirmation of Njolstad's report. *American Journal of Medical Genetics*, 93(4), 264–268.
- James, J. M., Nalbandian, A., & Mukoyama, Y. (2013). TGFbeta signaling is required for sprouting lymphangiogenesis during lymphatic network development in the skin. *Development (Cambridge, England)*, 140(18), 3903–3914. <https://doi.org/10.1242/dev.095026>
- Johnston, J. J., van der Smagt, J. J., Rosenfeld, J. A., Pagnamenta, A. T., Alsward, A., Baker, E. H., ... Biesecker, L. G. (2018). Autosomal recessive Noonan syndrome associated with biallelic LZTR1 variants. *Genetics in Medicine: Official Journal of the American College of Medical Genetics*, 20(10), 1175–1185. <https://doi.org/10.1038/gim.2017.249>
- Joyce, S., Gordon, K., Brice, G., Ostergaard, P., Nagaraja, R., Short, J., ... Mansour, S. (2016). The lymphatic phenotype in Noonan and Cardiofaciocutaneous syndrome. *European Journal of Human Genetics: EJHG*, 24(5), 690–696. <https://doi.org/10.1038/ejhg.2015.175>
- Kienast, Y., Jucknischke, U., Scheiblich, S., Thier, M., de Wouters, M., Haas, A., ... Lorenz, S. (2016). Rapid activation of bone Morphogenic protein 9 by receptor-mediated displacement of pro-domains. *The Journal of Biological Chemistry*, 291(7), 3395–3410. <https://doi.org/10.1074/jbc.M115.680009>
- Kurosaki, T., Popp, M. W., & Maquat, L. E. (2019). Quality and quantity control of gene expression by nonsense-mediated mRNA decay. *Nature Reviews. Molecular Cell Biology*, 20(7), 406–420. <https://doi.org/10.1038/s41580-019-0126-2>
- Lawera, A., Tong, Z., Thorikay, M., Redgrave, R. E., Cai, J., van Dinther, M., ... Li, W. (2019). Role of soluble endoglin in BMP9 signaling. *Proceedings of the National Academy of Sciences of the United States of America*, 116(36), 17800–17808. <https://doi.org/10.1073/pnas.1816661116>
- Levet, S., Ciais, D., Merdzhanova, G., Mallet, C., Zimmers, T. A., Lee, S.-J., ... Vittet, D. (2013). Bone morphogenetic protein 9 (BMP9) controls lymphatic vessel maturation and valve formation. *Blood*, 122(4), 598–607. <https://doi.org/10.1182/blood-2012-12-472142>
- Lord, J., McMullan, D. J., Eberhardt, R. Y., Rinck, G., Hamilton, S. J., Quinlan-Jones, E., ... Maher, E. R. (2019). Prenatal exome sequencing analysis in fetal structural anomalies detected by ultrasonography (PAGE): A cohort study. *Lancet (London, England)*, 393(10173), 747–757. [https://doi.org/10.1016/S0140-6736\(18\)31940-8](https://doi.org/10.1016/S0140-6736(18)31940-8)
- Lykke-Andersen, S., & Jensen, T. H. (2015). Nonsense-mediated mRNA decay: An intricate machinery that shapes transcriptomes. *Nature Reviews. Molecular Cell Biology*, 16(11), 665–677. <https://doi.org/10.1038/nrm4063>
- Mackie, D. I., al Mutairi, F., Davis, R. B., Kechele, D. O., Nielsen, N. R., Snyder, J. C., ... Caron, K. M. (2018). hCALCRL mutation causes autosomal recessive nonimmune hydrops fetalis with lymphatic dysplasia. *The Journal of Experimental Medicine*, 215(9), 2339–2353. <https://doi.org/10.1084/jem.20180528>
- Mardy, A. H., Chetty, S. P., Norton, M. E., & Sparks, T. N. (2019). A system-based approach to the genetic etiologies of non-immune hydrops fetalis. *Prenatal Diagnosis*, 39(9), 732–750. <https://doi.org/10.1002/pd.5479>
- Martin-Almedina, S., Martinez-Corral, I., Holdhus, R., Vicente, A., Fotou, E., Lin, S., ... Ostergaard, P. (2016). EPHB4 kinase-inactivating mutations cause autosomal dominant lymphatic-related hydrops fetalis. *The Journal of Clinical Investigation*, 126(8), 3080–3088. <https://doi.org/10.1172/JCI85794>
- Mason-Suares, H., Toledo, D., Gekas, J., Lafferty, K. A., Meeks, N., Pacheco, M. C., ... Lebo, M. S. (2017). Juvenile myelomonocytic leukemia-associated variants are associated with neo-natal lethal Noonan syndrome. *European Journal of Human Genetics: EJHG*, 25(4), 509–511. <https://doi.org/10.1038/ejhg.2016.202>
- McPherson, E. (2019). Hydrops fetalis in a cohort of 3,137 stillbirths and second trimester miscarriages. *American Journal of Medical Genetics. Part A*, 179(12), 2338–2342. <https://doi.org/10.1002/ajmg.a.61340>
- Meng, D., Li, Q., Hu, X., Wang, L., Tan, S., Su, J., ... Fu, C. (2019). Etiology and outcome of non-immune Hydrops Fetalis in southern China: Report of 1004 cases. *Scientific Reports*, 9(1), 10726. <https://doi.org/10.1038/s41598-019-47050-6>
- Moreno, C. A., Kanazawa, T., Barini, R., Nomura, M. L., Andrade, K. C., Gomes, C. P., ... Cavalcanti, D. P. (2013). Non-immune hydrops fetalis: A prospective study of 53 cases. *American Journal of Medical Genetics. Part A*, 161A(12), 3078–3086. <https://doi.org/10.1002/ajmg.a.36171>
- Nassr, A. A., Ness, A., Hosseinzadeh, P., Salmanian, B., Espinoza, J., Berger, V., ... Shamshirsaz, A. A. (2018). Outcome and treatment of Antenatally diagnosed nonimmune Hydrops Fetalis. *Fetal Diagnosis and Therapy*, 43(2), 123–128. <https://doi.org/10.1159/000475990>

- Niessen, K., Zhang, G., Ridgway, J. B., Chen, H., & Yan, M. (2010). ALK1 signaling regulates early postnatal lymphatic vessel development. *Blood*, 115(8), 1654–1661. <https://doi.org/10.1182/blood-2009-07-235655>
- Njolstad, P. R., Reigstad, H., Westby, J., & Espeland, A. (1998). Familial non-immune hydrops fetalis and congenital pulmonary lymphangiectasia. *European Journal of Pediatrics*, 157(6), 498–501. <https://doi.org/10.1007/s004310050862>
- Norton, M. E., Chauhan, S. P., & Dashe, J. S. (2015). Society for maternal-fetal medicine (SMFM) clinical guideline #7: Nonimmune hydrops fetalis. *American Journal of Obstetrics and Gynecology*, 212(2), 127–139. <https://doi.org/10.1016/j.ajog.2014.12.018>
- Oka, M., Iwata, C., Suzuki, H. I., Kiyono, K., Morishita, Y., Watabe, T., ... Miyazono, K. (2008). Inhibition of endogenous TGF-beta signaling enhances lymphangiogenesis. *Blood*, 111(9), 4571–4579. <https://doi.org/10.1182/blood-2007-10-120337>
- Pagnamenta, A. T., Kaisaki, P. J., Bennett, F., Burkitt-Wright, E., Martin, H. C., Ferla, M. P., ... Stewart, H. (2019). Delineation of dominant and recessive forms of LZTR1-associated Noonan syndrome. *Clinical Genetics*, 95(6), 693–703. <https://doi.org/10.1111/cge.13533>
- Pastores, G. M., & Hughes, D. A. (1993). In M. P. Adam, H. H. Ardinger, R. A. Pagon, S. E. Wallace, L. J. H. Bean, K. Stephens, & A. Amemiya (Eds.), *Gaucher disease*. Seattle (WA): University of Washington.
- Quinlan-Jones, E., Lord, J., Williams, D., Hamilton, S., Marton, T., Eberhardt, R. Y., ... Kilby, M. D. (2019). Molecular autopsy by trio exome sequencing (ES) and postmortem examination in fetuses and neonates with prenatally identified structural anomalies. *Genetics in Medicine: Official Journal of the American College of Medical Genetics*, 21(5), 1065–1073. <https://doi.org/10.1038/s41436-018-0298-8>
- Richards, S., Aziz, N., Bale, S., Bick, D., Das, S., Gastier-Foster, J., ... Rehms, H. L. (2015). Standards and guidelines for the interpretation of sequence variants: A joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genetics in Medicine: Official Journal of the American College of Medical Genetics*, 17(5), 405–424. <https://doi.org/10.1038/gim.2015.30>
- Robinson, W. P., McGillivray, B., Lewis, M. E. S., Arbour, L., Barrett, I., & Kalousek, D. K. (2005). Prenatally detected trisomy 20 mosaicism. *Prenatal Diagnosis*, 25(3), 239–244. <https://doi.org/10.1002/pd.1121>
- Saito, T., Bokhove, M., Croci, R., Zamora-Caballero, S., Han, L., Letarte, M., ... Jovine, L. (2017). Structural basis of the human Endoglin-BMP9 interaction: Insights into BMP signaling and HHT1. *Cell Reports*, 19(9), 1917–1928. <https://doi.org/10.1016/j.celrep.2017.05.011>
- Seidl, E., Carls, J., Reu, S., Wetzke, M., Ley-Zaporozhan, J., Brasch, F., ... Griesse, M. (2018). Pulmonary interstitial glycogenosis - a systematic analysis of new cases. *Respiratory Medicine*, 140, 11–20. <https://doi.org/10.1016/j.rmed.2018.05.009>
- Skoric-Milosavljevic, D., Tjong, F. V. Y., Barc, J., Backx, A. P. C. M., Clur, S.-A. B., van Spaendonck-Zwarts, K., ... Postma, A. v. (2019). GATA6 mutations: Characterization of two novel patients and a comprehensive overview of the GATA6 genotypic and phenotypic spectrum. *American Journal of Medical Genetics. Part A*, 179(9), 1836–1845. <https://doi.org/10.1002/ajmg.a.61294>
- Sobreira, N., Schiettecatte, F., Valle, D., & Hamosh, A. (2015). GeneMatcher: A matching tool for connecting investigators with an interest in the same gene. *Human Mutation*, 36(10), 928–930. <https://doi.org/10.1002/humu.22844>
- Southgate, L., Machado, R. D., Graf, S., & Morrell, N. W. (2019). Molecular genetic framework underlying pulmonary arterial hypertension. *Nature Reviews Cardiology*, 17(2), 85–95. <https://doi.org/10.1038/s41569-019-0242-x>
- Sparks, T. N., Thao, K., Lianoglou, B. R., Boe, N. M., Bruce, K. G., Datkhaeva, I., ... Norton, M. E. (2019). Nonimmune hydrops fetalis: Identifying the underlying genetic etiology. *Genetics in Medicine: Official Journal of the American College of Medical Genetics*, 21(6), 1339–1344. <https://doi.org/10.1038/s41436-018-0352-6>
- Steurer, M. A., Peyvandi, S., Baer, R. J., MacKenzie, T., Li, B. C., Norton, M. E., ... Moon-Grady, A. J. (2017). Epidemiology of live born infants with nonimmune Hydrops Fetalis—insights from a population-based dataset. *The Journal of Pediatrics*, 187, 182–188.e3. <https://doi.org/10.1016/j.jpeds.2017.04.025>
- Stevenson, D. A., Pysker, T. J., Ward, R. M., & Carey, J. C. (2006). Familial congenital non-immune hydrops, chylothorax, and pulmonary lymphangiectasia. *American Journal of Medical Genetics. Part A*, 140(4), 368–372. <https://doi.org/10.1002/ajmg.a.31093>
- Stuurman, K. E., Joosten, M., van der Burg, I., Elting, M., Yntema, H. G., Meijers-Heijboer, H., & Rinne, T. (2019). Prenatal ultrasound findings of rasopathies in a cohort of 424 fetuses: Update on genetic testing in the NGS era. *Journal of Medical Genetics*, 56(10), 654–661. <https://doi.org/10.1136/jmedgenet-2018-105746>
- Sudrie-Arnaud, B., Marguet, F., Patrier, S., Martinovic, J., Louillet, F., Broux, F., ... Bekri, S. (2018). Metabolic causes of nonimmune hydrops fetalis: A next-generation sequencing panel as a first-line investigation. *Clinica Chimica Acta. International Journal of Clinical Chemistry*, 481, 1–8. <https://doi.org/10.1016/j.cca.2018.02.023>
- Townson, S. A., Martinez-Hackert, E., Greppi, C., Lowden, P., Sako, D., Liu, J., ... Grinberg, A. v. (2012). Specificity and structure of a high affinity activin receptor-like kinase 1 (ALK1) signaling complex. *The Journal of Biological Chemistry*, 287(33), 27313–27325. <https://doi.org/10.1074/jbc.M112.377960>
- van Diemen, C. C., Kerstjens-Frederikse, W. S., Bergman, K. A., de Koning, T. J., Sikkema-Raddatz, B., van der Velde, J. K., ... Wijnga, C. (2017). Rapid targeted genomics in critically ill newborns. *Pediatrics*, 140(4), e20162854. <https://doi.org/10.1542/peds.2016-2854>
- Wagener, R., Aukema, S. M., Schlesner, M., Haake, A., Burkhardt, B., Claviez, A., ... Molecular Mechanisms in Malignant Lymphomas Network Project of the Deutsche Krebshilfe. (2015). The PCBP1 gene encoding poly(rc) binding protein 1 is recurrently mutated in Burkitt lymphoma. *Genes, Chromosomes and Cancer*, 54(9), 555–564. <https://doi.org/10.1002/gcc.22268>
- Wallerstein, R., Yu, M. T., Neu, R. L., Benn, P., Lee Bowen, C., Crandall, B., ... Wang, H. (2000). Common trisomy mosaicism diagnosed in amniocytes involving chromosomes 13, 18, 20 and 21: Karyotype-phenotype correlations. *Prenatal Diagnosis*, 20(2), 103–122.
- Wallerstein, R., Misra, S., Dugar, R. B., Alem, M., Mazzoni, R., & Garabedian, M. J. (2015). Current knowledge of prenatal diagnosis of mosaic autosomal trisomy in amniocytes: Karyotype/phenotype correlations. *Prenatal Diagnosis*, 35(9), 841–847. <https://doi.org/10.1002/pd.4620>
- Wang, G., Fan, R., Ji, R., Zou, W., Penny, D. J., Varghese, N. P., & Fan, Y. (2016). Novel homozygous BMP9 nonsense mutation causes pulmonary arterial hypertension: A case report. *BMC Pulmonary Medicine*, 16, 17. <https://doi.org/10.1186/s12890-016-0183-7>
- Wang, X.-J., Lian, T.-Y., Jiang, X., Liu, S.-F., Li, S.-Q., Jiang, R., ... Jing, Z.-C. (2019). Germline BMP9 mutation causes idiopathic pulmonary arterial hypertension. *The European Respiratory Journal*, 53(3), 1801609. <https://doi.org/10.1183/13993003.01609-2018>
- Weissbach, T., Kushnir, A., Rasslan, R., Rosenblatt, O., Yinon, Y., Berkenstadt, M., ... Kassif, E. (2019). Fetal pleural effusion: Contemporary methods of genetic evaluation. *Prenatal Diagnosis*, 39(9), 751–757. <https://doi.org/10.1002/pd.5497>
- Wieacker, P., Muschke, P., Pollak, K.-H., & Muller, R. (2005, January). Autosomal recessive non-immune hydrops fetalis caused by systemic lymphangiectasia. *American Journal of Medical Genetics. Part A*, 132A, 318–319. <https://doi.org/10.1002/ajmg.a.30392>
- Willis, M. J. H., Bird, L. M., Dell'Aquila, M., & Jones, M. C. (2008). Expanding the phenotype of mosaic trisomy 20. *American Journal of Medical Genetics. Part A*, 146A(3), 330–336. <https://doi.org/10.1002/ajmg.a.32126>

- Wooderchak-Donahue, W. L., McDonald, J., O'Fallon, B., Upton, P. D., Li, W., Roman, B. L., ... Bayrak-Toydemir, P. (2013). BMP9 mutations cause a vascular-anomaly syndrome with phenotypic overlap with hereditary hemorrhagic telangiectasia. *American Journal of Human Genetics*, 93(3), 530–537. <https://doi.org/10.1016/j.ajhg.2013.07.004>
- Yang, H., Zeng, Q., Ma, Y., Liu, B., Chen, Q., Li, W., ... Zhou, Z. (2018). Genetic analyses in a cohort of 191 pulmonary arterial hypertension patients. *Respiratory Research*, 19(1), 87. <https://doi.org/10.1186/s12931-018-0789-9>
- Yates, C. L., Monaghan, K. G., Copenheaver, D., Retterer, K., Scuffins, J., Kucera, C. R., ... Juusola, J. (2017). Whole-exome sequencing on deceased fetuses with ultrasound anomalies: Expanding our knowledge of genetic disease during fetal development. *Genetics in Medicine: Official Journal of the American College of Medical Genetics*, 19(10), 1171–1178. <https://doi.org/10.1038/gim.2017.31>
- Yoshimatsu, Y., Lee, Y. G., Akatsu, Y., Taguchi, L., Suzuki, H. I., Cunha, S. I., ... Watabe, T. (2013). Bone morphogenetic protein-9 inhibits lymphatic vessel formation via activin receptor-like kinase 1 during development and cancer progression. *Proceedings of the National Academy of Sciences of the United States of America*, 110(47), 18940–18945. <https://doi.org/10.1073/pnas.1310479110>
- Zhu, N., Pauciulo, M. W., Welch, C. L., Lutz, K. A., Coleman, A. W., Gonzaga-Jauregui, C., ... Nichols, W. C. (2019). Novel risk genes and mechanisms implicated by exome sequencing of 2572 individuals with pulmonary arterial hypertension. *Genome Medicine*, 11(1), 69. <https://doi.org/10.1186/s13073-019-0685-z>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.

How to cite this article: Aukema SM, ten Brinke GA, Timens W, et al. A homozygous variant in growth and differentiation factor 2 (*GDF2*) may cause lymphatic dysplasia with hydrothorax and nonimmune hydrops fetalis. *Am J Med Genet Part A*. 2020;1–9. <https://doi.org/10.1002/ajmg.a.61743>