University of Groningen

Core Outcome Domains for Trials in Autosomal Dominant Polycystic Kidney Disease

Cho, Yeoungjee; Rangan, Gopala; Logeman, Charlotte; Ryu, Hyunjin; Sautenet, Benedicte; Perrone, Ronald D; Nadeau-Fredette, Annie-Claire; Mustafa, Reem A; Htay, Htay; Chonchol, Michel

Published in:
American Journal of Kidney Diseases

DOI:
[10.1053/j.ajkd.2020.01.005](https://doi.org/10.1053/j.ajkd.2020.01.005)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Cho, Y., Rangan, G., Logeman, C., Ryu, H., Sautenet, B., Perrone, R. D., Nadeau-Fredette, A-C., Mustafa, R. A., Htay, H., Chonchol, M., Harris, T., Gutman, T., Craig, J. C., Ong, A. C. M., Chapman, A., Ahn, C., Coolican, H., Kao, J. T-W., Gansevoort, R. T., ... Tong, A. (2020). Core Outcome Domains for Trials in Autosomal Dominant Polycystic Kidney Disease: An International Delphi Survey. *American Journal of Kidney Diseases*, 76(3), 361-373. <https://doi.org/10.1053/j.ajkd.2020.01.005>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Core Outcome Domains for Trials in Autosomal Dominant Polycystic Kidney Disease: An International Delphi Survey

Yeoungjee Cho, Gopala Rangan, Charlotte Logeman, Hyunjin Ryu, Benedicte Sautenet, Ronald D. Perrone, Annie-Claire Nadeau-Fredette, Reem A. Mustafa, Htay Htay, Michel Chonchol, Tess Harris, Talia Gutman, Jonathan C. Craig, Albert C.M. Ong, Arlene Chapman, Curie Ahn, Helen Coolican, Juliana Tze-Wah Kao, Ron T. Gansevoort, Vicente Torres, York Pei, David W. Johnson, Andrea K. Viecelli, Armando Teixeira-Pinto, Martin Howell, Angela Ju, Karine E. Manera, and Allison Tong

Rationale & Objective: Outcomes reported in trials involving patients with autosomal dominant polycystic kidney disease (ADPKD) are heterogeneous and rarely include patient-reported outcomes. We aimed to identify critically important consensus-based core outcome domains to be reported in trials in ADPKD.

Study Design: An international 2-round online Delphi survey was conducted in English, French, and Korean languages.

Setting & Participants: Patients/caregivers and health professionals completed a 9-point Likert scale (7-9 indicating critical importance) and a Best-Worst Scale.

Analytical Approach: The absolute and relative importance of outcomes were assessed. Comments were analyzed thematically.

Results: 1,014 participants (603 [60%] patients/caregivers, 411 [40%] health professionals) from 56 countries completed round 1, and 713 (70%) completed round 2. The prioritized outcomes were kidney function (importance

score, 8.6), end-stage kidney disease (8.6), death (7.9), blood pressure (7.9), kidney cyst size/growth (7.8), and cerebral aneurysm (7.7). Kidney cyst-related pain was the highest rated patient-reported outcome by both stakeholder groups. Seven themes explained the prioritization of outcomes: protecting life and health, directly encountering life-threatening and debilitating consequences, specificity to ADPKD, optimizing and extending quality of life, hidden suffering, destroying self-confidence, and lost opportunities.

Limitations: Study design precluded involvement from those without access to internet or limited computer literacy.

Conclusions: Kidney function, end-stage kidney disease, and death were the most important outcomes to patients, caregivers, and health professionals. Kidney cyst-related pain was the highest rated patient-reported outcome. Consistent reporting of these top prioritized outcomes may strengthen the value of trials in ADPKD for decision making.

Complete author and article information provided before references.

Correspondence to Y. Cho (yeoungjee.cho@health.qld.gov.au)

Am J Kidney Dis. 76(3):361-373. Published online April 28, 2020.

doi: 10.1053/j.ajkd.2020.01.005

© 2020 by the National Kidney Foundation, Inc.

Autosomal dominant polycystic kidney disease (ADPKD) is the most common genetic cause of chronic kidney disease (CKD). Up to 70% of patients with ADPKD progress to end-stage kidney disease (ESKD) by the age of 65 years.^{1,2} Although kidney function often remains stable for many years after diagnosis, patients with ADPKD often experience debilitating symptoms such as pain that are related to kidney cyst growth and enlargement,³⁻⁶ and they are at risk for extrarenal complications, including stroke from ruptured intracranial aneurysm and severe polycystic liver disease.⁷

Although there is some evidence to support lifestyle interventions (eg, salt restriction⁸) and therapeutic agents (eg, tolvaptan) in improving kidney function and total kidney volume,⁹⁻¹³ outcomes important to patients with ADPKD and their caregivers, such as kidney pain, fatigue,¹⁴ and anxiety/psychosocial distress,¹⁵ are infrequently reported in trials in ADPKD.^{7,16-18} Trials in ADPKD most frequently report surrogate outcomes, including kidney function, kidney/cyst volume, and blood pressure.¹⁹ As part of the Standardized Outcomes in Nephrology–Polycystic Kidney Disease (SONG-PKD) initiative,²⁰ the aim of this study was

to generate consensus among patients, caregivers, and health professionals on critically important outcomes to be reported in trials in ADPKD. This will inform the development of a core outcome set, defined as an “agreed minimum set of outcomes to be reported in all trials,”^{21,22} which can lead to improved consistency in reporting outcomes important to patients and clinicians across trials in ADPKD to better enable shared decision making.

Methods

Study Design

The Delphi technique²³ is an internationally accepted approach used to establish consensus on core outcome sets for trials in nephrology and other disciplines.²⁴⁻²⁸ Participants’ anonymities are maintained, and they are able to complete the survey independently and allow for widespread international participation through online dissemination.²⁹ The survey was conducted online in 3 languages (English, French, and Korean). The English survey was translated into French and Korean by a bilingual health professional and cross-checked by a second bilingual

professional to ensure accuracy. The survey was pilot tested by members of the SONG-PKD Steering Group including 3 patients with ADPKD. There were 2 iterative rounds completed by a panel of participants with experience or expertise in ADPKD. Due to stability in results from rounds 1 and 2, we did not proceed with an additional round. In the second round, participants could review their own score from round 1, the distribution of scores (overall, patients/caregivers, and health professionals), and comments provided by participants. The SONG-PKD Delphi process is shown in Figure S1.

Participant Selection and Recruitment

Patients, caregivers, and health professionals with experience in ADPKD were eligible to participate. Patients/caregivers (aged ≥ 18 years) included patients with ADPKD across all stages of CKD (stages 1-5; dialysis [hemodialysis and peritoneal dialysis], and transplantation) and family members and friends. Health professionals included physicians (nephrologists, hepatologists, surgeons, and geneticists), nurses, allied health professionals, researchers, policy makers, industry, and regulators.

To include a diverse range of participants, we used multiple recruitment strategies. Patients were recruited from hospitals, patient/consumer organizations, the SONG database, and social media. Health professionals were recruited through the SONG database, investigator networks, and professional organizations (Table S1). Participants received an e-mail invitation including a link to the survey after registering their e-mail on the SONG website (www.songinitiative.org). Ethics approval was provided by the University of Sydney (2015-228) and participating institutions (Table S1). Informed consent was obtained prior to survey completion from each participant.

Data Collection

Selection of Outcome Domains

The 41 outcome domains included in the survey were identified from a systematic review of ADPKD trials and a study of patient/caregiver priorities for outcomes in ADPKD.¹⁴ The order in which outcome domains appeared in the survey was random and each outcome was accompanied by a plain language definition (Table S2). The SONG-PKD Steering Group and investigators reviewed the list of outcomes. The survey was administered online using Qualtrics (Qualtrics software) from June 2018 to February 2019.

Round 1

Participants scored the importance of each of the 41 outcome domains using a 9-point Likert scale. Scores 1 to 3 indicated "limited importance," 4 to 6 indicated "important but not critical," and 7 to 9 indicated "critical importance" (Fig S1), based on the Grading of Recommendations Assessment, Development and Evaluation (GRADE) process.³⁰ For each outcome, there was an option to enter

comments in free-text boxes. Participants also had an "unsure" option and could suggest new outcomes. New outcomes suggested by $>10\%$ of participants were considered for inclusion in the second round. Outcomes with a mean score less than 6 or median score less than 7 in both groups (patients/caregivers and health professionals) were excluded in round 2. Comments from each outcome in round 1 were separately evaluated to determine any overlap between outcome domains and the need to revise the outcome domains for round 2.

Round 2

A total of 17 outcomes were included in round 2. Participants could review their own scores from round 1, as well as the distribution of scores (overall and divided according to status of patients/caregivers and health professionals) displayed as percentage of participants in a column graph. Deidentified comments from round 1 were provided. After reviewing these results, participants were asked to re-rate the outcomes using the 9-point Likert scale following the same methods used for round 1. On completion of rating all outcomes, participants were asked to complete a Best-Worst Scale (BWS) survey to examine the relative importance of each outcome.²⁴

The BWS consists of choice tasks in which a participant is asked to indicate the best and worst items/options, with the overall aim to obtain a ranking of items in the order of preference.^{31,32} Five Best-Worst choice sets, each containing 6 of 17 outcomes selected using a balanced incomplete block design, were presented to each participant. For each Best-Worst choice set, participants had to select the most important and least important outcome.

Data Analysis

Quantitative Analysis

For every outcome in each round, mean score, median score, and proportion of participants who rated the outcomes as critically important (from 7 to 9) were calculated. Scores were separately calculated for patients/caregivers and health professionals, and their mean differences in scores and changes between 2 survey rounds were analyzed using *t* test. Results from the BWS survey were incorporated in a multinomial logistic regression model to determine the relative importance. Utility functions containing all outcomes and interaction terms for participant characteristics were constructed. Subsequently, the mean regression coefficients of these functions provided the relative importance scores for each outcome,³² in which a scale of 1 represented the least important and 9 represented the most important.

Statistical analyses were performed using Stata (version 14.0; StataCorp LP), SPSS (IBM SPSS Statistics for Windows, version 25.0), Excel (Microsoft Corp, version 16.0), and NLOGIT V6 (Econometric Software Inc) for the BWS.

Definition of Consensus for Core Outcomes

Due to unknown distribution of scores, the threshold for consensus for the core outcomes domains could not be

Table 1. Characteristics of Patients/Caregivers

Characteristic	Round 1 (n = 603)	Round 2 (n = 406)
Participant type ^a		
Patient	536 (89%)	372 (92%)
Caregiver/family member	107 (18%)	65 (16%)
Sex ^b		
Female	388 (65%)	252 (62%)
Male	212 (35%)	154 (38%)
Age group		
18-30 y	49 (8%)	27 (7%)
31-40 y	117 (19%)	72 (18%)
41-50 y	139 (23%)	84 (21%)
51-60 y	175 (29%)	125 (31%)
61-70 y	102 (17%)	82 (20%)
71-80 y	21 (3%)	16 (4%)
Marital status		
Married/de facto	476 (79%)	336 (83%)
Partner	18 (3%)	14 (3%)
Single	71 (12%)	34 (8%)
Divorced/separated/widowed	38 (6%)	22 (5%)
Employment status		
Employed full-time	299 (50%)	200 (49%)
Employed part-time/casual	86 (14%)	62 (15%)
Unemployed	39 (6%)	21 (5%)
Retired	97 (16%)	81 (20%)
Student or other	82 (14%)	42 (10%)
Education		
Did not complete high school	39 (6%)	27 (7%)
High school graduate	83 (14%)	45 (11%)
Professional certificate	97 (16%)	65 (16%)
Undergraduate degree	242 (40%)	159 (39%)
Postgraduate degree	142 (24%)	110 (27%)
Current type of treatment (patients only)		
No kidney replacement therapy	401 (67%)	275 (74%)
Peritoneal dialysis	9 (1%)	7 (2%)
Hemodialysis	37 (6%)	22 (6%)
Kidney transplant	89 (15%)	68 (18%)
Age group at diagnosis of ADPKD (patients only) ^c		
<10 y	11 (3%)	8 (3%)
11-20 y	51 (14%)	32 (13%)
21-30 y	101 (28%)	75 (30%)
31-40 y	117 (32%)	70 (28%)
41-50 y	51 (14%)	38 (15%)
51-60 y	24 (7%)	19 (8%)
>60 y	6 (2%)	6 (2%)
Family history of ADPKD (patients only)		
Yes	393 (73%)	276 (74%)
No	96 (18%)	68 (18%)
Unsure	47 (9%)	28 (8%)
Country		
Republic of Korea	182 (30%)	94 (23%)
United States	134 (22%)	86 (21%)
United Kingdom	104 (17%)	90 (22%)
Australia	75 (12%)	58 (14%)

(Continued)

Table 1 (Cont'd). Characteristics of Patients/Caregivers

Characteristic	Round 1 (n = 603)	Round 2 (n = 406)
Canada	30 (5%)	28 (7%)
Italy	20 (3%)	9 (2%)
Other ^d	58 (10%)	41 (10%)

Abbreviation: ADPKD, autosomal dominant polycystic kidney disease.

^aSome have multiple roles.^bN = 603 for round 1 due to missing data.^cN = 361 and 248 for rounds 1 and 2, respectively, due to missing data.^dOther includes 22 countries (in descending order of number of participants): Norway, France, Singapore, Spain, Netherlands, New Zealand, South Africa, China, Germany, Pakistan, Poland, Switzerland, Belarus, Denmark, Hong Kong, India, Ireland, Japan, Malaysia, Nepal, Nigeria, Turkey.

prespecified. The Delphi survey aimed to identify 3 to 5 outcome domains that were critically important to both stakeholder groups. “Consensus” for the critical outcome domains was defined based on both patient/caregiver and health professional groups yielding median scores of 7 or higher and mean scores of 7 or higher, as well as the proportions of both stakeholder groups rating the outcome as “critically important (defined as scores 7-9)” being >75% in round 2. These thresholds were discussed and approved by the SONG-PKD Steering Group. Scores obtained from the BWS were used to examine the relative differences in preference scores between patients/caregivers and health professionals.

Qualitative Analysis

Comments from the survey were imported into Hyper-RESEARCH (version 3.7) software for data analysis. Using thematic analysis, investigators coded the text (in English, Korean [Y.C.], and French [B.S.]) and inductively identified themes focusing on reasons for ratings, differences between stakeholder groups and changes in scores across rounds. A third investigator (A.T.) read the qualitative data and reviewed the preliminary analysis to ensure that the themes captured all the data.

Results

Participant Characteristics

In the first round, 1,014 participants from 56 countries completed the survey, of whom 603 (60%) were patients/caregivers and 411 (40%) were health professionals. In the second round, 713 participants (70% overall retention rate) from 47 countries completed the survey, which included 406 (57%) patients/caregivers and 307 (63%) health professionals (Tables 1 and 2).

Of the 406 patients/caregivers who completed both rounds, 275 (74%) were patients not receiving kidney replacement therapy, 7 (2%) were receiving peritoneal dialysis, 22 (6%) were receiving hemodialysis, and 68 (18%) were kidney transplant recipients. Overall, 65 caregivers/family members (total N > 406 due to multiple roles, eg, patients who were also caregivers to other family

Table 2. Characteristics of Health Professionals

Characteristic	Round 1 (n = 411)	Round 2 (n = 307)
Participant role^a		
Nephrologist	272 (66%)	214 (70%)
Researcher	72 (18%)	36 (12%)
Nurse	62 (15%)	32 (10%)
Industry	11 (3%)	6 (2%)
Dietician	6 (1%)	5 (2%)
Policy maker	6 (1%)	5 (2%)
Surgeon	5 (1%)	4 (1%)
Social worker	3 (1%)	2 (<1%)
Other	28 (7%)	26 (8%)
Sex^b		
Female	217 (53%)	166 (54%)
Male	185 (46%)	140 (46%)
Age group		
18-30 y	22 (5%)	15 (5%)
31-40 y	121 (29%)	84 (27%)
41-50 y	106 (26%)	76 (25%)
51-60 y	108 (26%)	85 (28%)
61-70 y	49 (12%)	42 (14%)
71-80 y	5 (1%)	5 (2%)
Experience in ADPKD		
≤10 y	141 (34%)	100 (33%)
11-20 y	110 (27%)	82 (27%)
21-30 y	82 (20%)	68 (22%)
>30 y	56 (14%)	47 (15%)
Not applicable	22 (5%)	10 (3%)
No. of ADPKD trials as investigators		
0	229 (56%)	163 (53%)
1-5	111 (27%)	97 (32%)
6-10	16 (4%)	10 (3%)
11-15	2 (<1%)	2 (<1%)
>15	4 (1%)	4 (1%)
Not applicable	49 (12%)	31 (10%)
Country		
Australia	64 (16%)	55 (18%)
France	49 (12%)	39 (13%)
Republic of Korea	49 (12%)	25 (8%)
United States	45 (11%)	36 (12%)
Canada	31 (8%)	24 (8%)
United Kingdom	28 (7%)	26 (8%)
Singapore	17 (4%)	1 (<1%)
Hong Kong	14 (3%)	9 (3%)
Other ^c	114 (27%)	92 (30%)

Abbreviation: ADPKD, autosomal dominant polycystic kidney disease.

^aSome have multiple roles.^bN ≠ 411 for round 1, due to missing data.^cOther includes 42 countries (in descending order): Italy, New Zealand, Japan, Spain, China, India, Portugal, Belgium, Argentina, Netherlands, Egypt, Germany, Greece, Lithuania, Malaysia, Pakistan, Poland, Thailand, Brazil, Finland, Nigeria, Philippines, Vietnam, Albania, Algeria, Austria, Belarus, Bosnia and Herzegovina, Bulgaria, Croatia, Denmark, Ecuador, El Salvador, Libya, Montenegro, Romania, Russia Federation, Slovakia, Sweden, Syrian Arab Republic, Ukraine, United Arab Emirates.

members affected by ADPKD) from 23 countries participated, including from the Republic of Korea (23%), United Kingdom (22%), United States (21%), Australia (14%), and

Canada (7%). Of the 307 health professionals who also completed both rounds, 214 (70%) were nephrologists, 36 (12%) were researchers, and 32 (10%) were nurses. Dietitians, policy makers, surgeons, a geneticist, a hepatologist, industry representatives, and a psychologist also participated. Health professionals were from 41 countries, including Australia (18%), France (13%), United States (12%), United Kingdom (8%), Republic of Korea (8%), and Canada (8%).

Rating Scores

Round 1

Mean and median scores and proportions of participants scoring the outcomes from 7 to 9 separated by patients/caregivers and health professionals for each of the 41 outcome domains in round 1 are provided in [Table S3](#). The top 5 outcomes with the highest mean scores for patients/caregivers were kidney function (8.5; defined for participants as the ability of the kidney to remove waste from the body and balance fluids), ESKD (8.4), cerebral aneurysm (8.0), kidney cyst size/growth (8.0), and blood pressure (7.9). For health professionals, the top 5 outcomes were kidney function (8.4), ESKD (8.4), death (7.8), cerebral aneurysm (7.5), and blood pressure (7.5). Twenty-four outcomes had mean scores less than 6.0 or median scores less than 7.0 across both groups and were excluded from round 2. Although fatigue did not meet the criteria based on mean and median scores, it was included in round 2 to ensure the inclusion of at least 5 patient-reported outcomes in round 2. Kidney cyst infection and cyst-related pain/bleeding were combined as kidney cyst pain/bleeding/infection due to overlap based on comments (ie, prioritized due to pain caused by bleeding or infection) and similarity in scores. None of the new outcomes were suggested by >10% of participants ([Table S4](#)) and were therefore not included in the next round.

Round 2

For each of the 17 outcome domains in round 2, mean and median scores and proportion of participants scoring the outcome as critically important are shown in [Table S5](#). The top 5 prioritized outcomes from all participants were kidney function (8.6), ESKD (8.6), death (7.9), blood pressure (7.9), and kidney cyst size/growth (7.8). However, the top 5 outcomes list for patients/caregivers included cerebral aneurysm (7.8) instead of death. In general, top 5 outcomes according to language subtypes were comparable, except inclusion of cerebral aneurysm instead of kidney cyst size/growth and death in French and Korean, respectively ([Table S6](#)). Mean scores of outcomes were generally higher for female patients/caregivers compared with male participants for all outcomes ([Table S7](#)).

Changes in Scores From Round 1 to 2

Changes in mean scores from the first and second rounds are presented in [Figures 1](#) (patients/caregivers) and [2](#) (health professionals). Between the 2 rounds, mean

scores for each outcome were generally stable, with the top 10 outcomes consistent across both rounds (Tables S3 and S5). Patient/caregiver mean scores increased between rounds for the following 4 outcomes: life participation (mean score difference, 0.27; $P = 0.006$), ESKD (0.19; $P = 0.005$), kidney function (0.14; $P = 0.02$), and kidney cyst size/growth (0.15; $P = 0.05$). For health professionals, mean scores increased for 5 outcomes from round 1 to 2: death (0.33; $P = 0.002$), cardiovascular disease (0.30; $P = 0.003$), chronic pain (0.21; $P = 0.05$), ESKD (0.20; $P = 0.002$), and blood pressure (0.19; $P = 0.04$).

In round 2, more health professionals rated blood pressure (round 1 vs 2: 82% vs 89%; $P = 0.02$), cardiovascular disease (81 vs 88%; $P = 0.007$), and fatigue (50% vs 55%; $P = 0.04$) as critically important compared with round 1. Kidney function (95% vs 98%; $P = 0.007$) and life participation (73% vs 79%; $P = 0.002$) were the only outcomes with a higher proportion of patients/caregivers

who rated it to be critically important compared with round 1 (Fig 3).

Differences Between Stakeholder Groups

Differences in mean scores between patients/caregivers and health professionals for rounds 1 and 2 are shown in Figure 4. Of the 17 outcomes in round 2, patients/caregivers rated 14 outcomes higher than health professionals on the Likert scale, with the greatest difference in scores for liver cyst (absolute mean difference, 0.81; $P < 0.001$), fatigue (0.74; $P < 0.001$), and kidney cyst size/growth (0.68; $P < 0.001$). Death was the only outcome that was rated higher on the Likert scale by health professionals (0.37; $P = 0.001$). Differences in mean scores were comparable between the 2 participant groups for hospitalization (0.05; $P = 0.7$) and ESKD (0.07; $P = 0.3$). Similar differences were observed when results were analyzed by comparing proportions of critically important

Figure 1. Mean scores of patients/caregivers in rounds 1 and 2. Abbreviation: ESKD, end-stage kidney disease.

Figure 2. Mean scores of health professionals in rounds 1 and 2. Abbreviation: ESKD, end-stage kidney disease.

outcomes between patients/caregivers and health professionals (Fig 3).

Best-Worst Scale

In the BWS survey, both stakeholder groups identified ESKD as the most important outcome, but there were notable differences in the subsequent order of outcomes (Figs 5 and S2-S4). Patients and caregivers prioritized ESKD, kidney function, cerebral aneurysm, cardiovascular disease, and blood pressure in descending order, whereas health professionals considered death to be the second most important outcome, followed by kidney function and cardiovascular disease. Patients and caregivers identified chronic pain, kidney cyst size/growth, and kidney cyst pain/bleeding/infection to be as important as death. Results from the BWS survey were comparable for all participants when outcomes were analyzed according to language, except that kidney cyst size/growth was given higher priority by participants who completed the survey in Korean language. Patients and

caregivers also highly rated chronic pain, which was the most important outcome among participants who completed the French-language survey.

Thematic Analysis

Seven themes reflecting the reasons for and changes and differences in the rating of outcomes were identified: protecting life and health, directly encountering life-threatening and debilitating consequences, specificity to ADPKD, optimizing and extending quality of life, hidden suffering, destroying self-confidence, and lost opportunities. The themes reflected the perspectives of all stakeholder groups unless otherwise specified. Illustrative quotations supporting each theme are provided in Box 1.

Protecting Life and Health

Outcomes such as blood pressure, kidney function, and kidney cyst size/growth were rated highly because they were considered important “biomarkers” to “keep

healthy” and “to prevent damage,” to delay progression to ESKD or the development of cardiovascular disease. Cerebral aneurysm was noted to be uncommon and not relevant to all patients with ADPKD but was rated highly due to its life-threatening consequences.

Directly Encountering Life-Threatening and Debilitating Consequences

Some patients witnessed life-threatening consequences, such as premature death or severe disability involving their family members caused by cardiovascular disease and aneurysm/stroke, and these were thus rated critically important: “Heart problems is what killed my father, who had PKD. His transplanted kidney was still working but his heart failed.” Other outcomes, such as ability to do usual activities (ie, life participation), were prioritized based on having direct experience with outcomes that caused fear and frustration, “Near end-stage, lots of activities are impossible. Very frustrating.”

Specificity to ADPKD

Outcomes such as cardiovascular disease were considered “important but not specific to ADPKD” and similar views were held for blood pressure, which was “not any more important in PKD than in any other kidney disease” by some health professionals.

Optimizing and Extending Quality of Life

Living with ADPKD was “long lasting” and patients were “more concerned with quality of life than survival,” with lower scores for death among patients/caregivers

compared with results from health professionals. Outcomes directly related to day-to-day symptom burden, such as kidney cyst size/growth, kidney cyst–related pain/bleeding/infection, and chronic pain, were rated highly because they caused a “huge amount of morbidity,” which “has a big impact on day-to-day life.”

Hidden Suffering

Outcomes including chronic pain, kidney cyst–related pain/bleeding/infection, and mood were rated highly by patients/caregivers because they were often “overlooked,” “minimized,” and “misunderstood” by others. Moreover, these outcomes often led to “debilitating” symptom burden, which could be present “despite not being on dialysis (or anywhere near it).”

Destroying Self-confidence

For some patients, change in appearance or weight from ADPKD was critically important. It led to shattering “self-confidence—as the cysts/kidneys grew, it brought up issues other than just physical health. It’s harder to find well-fitting clothes and...can contribute to an existing feeling of sadness, anger, and despair,” as well as “embarrassment and fear of social stigma.” Often patients felt they had “no control” over these outcomes.

Lost Opportunities

Financial impact was rated highly by some participants who lost employment opportunities due to illnesses caused by ADPKD. Even if it was not directly experienced, participants recognized this to be an important concern: “many PKD

Figure 3. Proportion of outcomes rated critically important (scores 7-9 using a 9-point Likert scale), important (scores 4-6), and limited importance (scores 1-3) for patients/caregivers and health professionals in rounds 1 and 2.

Figure 4. Difference in mean scores between patients/caregivers and health professionals for rounds 1 and 2. Error bars refer to 95% confidence interval.

Figure 5. Mean relative importance scores of patients/caregivers and health professionals based on the Best-Worst Scale. Ordered by the mean importance scores of patients/caregivers (bars with 95% confidence interval). Abbreviation: ESKD, end-stage kidney disease.

Box 1. Themes and Selected Illustrative Quotations**Protecting life and health**

"I like to know how my kidney is going so I am able to keep healthy." (kidney function; Patient)

"Not common, but very life threatening." (aneurysm/stroke; HP)

"BP to be controlled to prevent damage" (BP; Patient)

"Cyst growth is associated with decline in kidney function and is an important surrogate outcome measure." (cyst size/growth; HP)

"Reducing blood pressure can be beneficial for extending the loss of function of the kidney." (BP; Patient)

"Kidney size is the most feared aspect of the disease for me. Very debilitating and absolutely ruins the way you look as well." (cyst size/growth; Patient)

"This is very important to me when my kidneys will start to fail." (ESKD; Patient)

Directly encountering life-threatening and debilitating consequences

"Heart problems is actually what killed my father, who had PKD; his transplanted kidney was still working but his heart failed." (CVD; PCG)

"My father died at 52 and had heart attack which caused transplant to fail and required bilateral [amputation]." (CVD; PCG)

"My dad died of a heart attack, probably from uncontrolled BP." (BP; PCG)

"When I was diagnosed (more than 40 years ago), I read in a medical dictionary that average age of death with PKD was 57. My dad died at 57. My sister died at 60. Death is critical outcome for me." (death; PCG)

"I am in a wheelchair due to a brain bleed after going into cardiac arrest after having an aneurysm coiled." (ability to do usual activities; Patient)

"Near end stage, lots of activities are impossible. Very frustrating." (life participation; Patient)

"Father had aneurysm at age 46 and he lived but was disabled. This impacted our family of 8 tremendously." (aneurysm/stroke; PCG)

Specificity to ADPKD

"CVD is important but not specific to ADPKD so although should be reported in all trials in CKD not a specific outcome for ADPKD." (CVD; HP)

Optimizing and extending quality of life

"PKD is usually long lasting. More concerned with quality of life than survival." (death; Patient)

"Increased size can lead to decreased function and pain." (cyst size/growth; Patient)

"Ability to function with enlarged kidneys." (cyst size/growth; Patient)

"I suffer from constant chronic pain and it impacts on my quality of life so much." (chronic pain; Patient)

"It has a big impact on day-to-day life." (chronic pain; Patient)

Hidden suffering

"Hidden symptom, feeling of being heavily pregnant, even though I'm slim." (chronic pain; Patient)

"Despite not being on dialysis (or anywhere near it just yet), I still get kidney pain which fluctuates." (chronic pain; Patient)

"I have been disability status based on my pain." (chronic pain; Patient)

"Full abdomen feeling makes me think I'm full all the time." (weight change; Patient)

"Mental and emotional health are so very important in getting through life with PKD." (depression; Patient)

"I've had a couple of very serious cyst infections which caused agonizing pain." (cyst-related pain/bleeding; Patient)

"In my experience this complication was overlooked and became quite painful and made me very sick." (cyst infection; Patient)

"These are debilitating, painful and sometimes make people (patients and carers) feel hopeless." (cyst infection; Patient)

"I think pain in PKD patients is hugely misunderstood by doctors and often overlooked." (chronic pain; Patient)

"When mentioning pain physician tends to minimize this complaint." (chronic pain; Patient)

"A lot of people do not understand how PKD can affect someone long term." (impact on family/friends; Patient)

"Dealing with a disease for which there is no cure can really get to a person especially if you're not getting support from family and friends. It's hard to explain how you feel to someone who doesn't have this disease and it gets frustrating." (mood; Patient)

Destroying self-confidence

"For a patient, the importance of appearance is much underestimated. It's all about self-confidence, as the cysts/kidneys grow it brings up issues other than just physical health. It's harder to find well-fitting clothes and lack of confidence can contribute to an existing feeling of sadness, anger, despair." (weight change; Patient)

"Huge belly is incredibly depressing and debilitating." (appearance; Patient)

"I feel like I have no control over how people see me." (appearance; Patient)

Lost opportunities

"Lost my job, was too unwell to return." (financial impact; Patient)

"Many PKDers lose their jobs or face employment disciplinary procedures because of illness and hospitals. People don't get promoted I'm sure, or have to cut short their careers because of physical limitations." (financial impact; Patient)

Abbreviations: ADPKD, autosomal dominant polycystic disease; BP, blood pressure; CKD, chronic kidney disease; CVD, cardiovascular disease; ESKD, end-stage kidney disease; HP, health professional; PCG, patient/caregiver; PKD, polycystic kidney disease.

patients lose their jobs or face employment disciplinary procedures because of illness and hospitalizations.”

Discussion

Kidney function, ESKD, death, blood pressure, and kidney cyst size/growth were the highest prioritized outcomes in ADPKD among patients/caregivers and health professionals. Kidney cyst size/growth was in the top 5 outcomes for patients/caregivers only, whereas death was in the top 5 for health professionals only. Kidney cyst-related pain/bleeding/infection was the highest rated patient-reported outcome by both stakeholder groups, whereas other patient-reported outcomes, such as anxiety, muscle pain, itch/skin, and sexual function, were deemed less important. Kidney pain due to cyst size/growth and infection/bleeding were highly prioritized due to their relentless symptom burden limiting life participation and as an indicator of disease progression toward ESKD. Similarly, other outcomes affecting quality of life, including liver cysts, financial impact, and fatigue, were rated higher by patients/caregivers because these outcomes disrupted daily living and restricted their ability to fulfil their social roles and goals. In contrast, health professionals placed greater emphasis on death.

Both patients/caregivers and health professionals consistently prioritized kidney function and ESKD as the most important outcomes in ADPKD. ESKD (ie, kidney failure requiring kidney replacement therapy) was a feared consequence to be avoided because it threatened quality of life, and kidney function was used to monitor progression to ESKD through its trajectory. The paramount importance attributed to kidney function and ESKD is in line with prior work on identifying outcomes important to patients with ADPKD and their caregivers.¹⁴ Other outcomes with life-threatening or life-changing consequences, such as cerebral aneurysm and cardiovascular disease, and those believed to promote their occurrence, such as blood pressure, were also highly prioritized by both groups. Blood pressure may also have been perceived as critically important by patients and clinicians because hypertension is reported to affect up to 70% of patients and often diagnosed around 30 years of age, usually before any apparent decrease in kidney function.^{33–35} In contrast, surrogate outcomes perceived not to impose a similar risk for harm, such as anemia, proteinuria, or lipid levels, were considered less important, even though these are frequently reported in ADPKD trials.¹⁹

In general, health professionals gave lower priority to most outcomes compared with patients/caregivers except for 3 clinical outcomes: ESKD, death, and hospitalization. Having direct experience of ADPKD, patients placed greater emphasis on life participation, indicating that they were limited by day-to-day symptom burden (eg, cyst-related pain and fatigue) and its impact on

practical aspects of life (eg, financial impact). They rated these similarly to death in terms of relative importance. This finding also reinforces the profound impacts of ADPKD on lifestyle and well-being, which have been identified in previous studies.^{14,15,36} Often these impacts have been perceived by patients to be underestimated by their physicians, and their symptom burden can result in dissatisfaction with care and treatment.³⁶ The high priority given to lifestyle-related outcomes has also been repeatedly shown in studies conducted in hemodialysis, peritoneal dialysis, and kidney transplant populations.^{27,28,37,38}

Outcome domains related to cysts were prioritized highly by both stakeholder groups for different reasons. Health professionals, particularly from the Republic of Korea, rated kidney cyst size/growth highly due to its importance as a biomarker of disease progression.³⁹ In contrast, patients/caregivers who completed the survey in French placed highest importance on chronic pain due to its detrimental impact on quality of life. Patients from the Republic of Korea were also concerned about changes in physical appearance from enlarged cysts, which led to embarrassment and social stigma limiting their employment opportunities. Cyst-related outcomes were perceived to be underrecognized and inadequately managed, which in part explained the higher prioritization of these outcomes. Other studies have also shown that cyst-related symptom burden is associated with worse quality of life and is a barrier to achieving long-term life goals.^{17,18} An increase in economic burden from health care use due to cyst complications in ADPKD before clinically evident decreases in kidney function is now widely acknowledged,^{15,40,41} and tools are being developed to better capture the cyst-related symptom burden of ADPKD (eg, ADPKD-Impact Scale and genetic psychosocial risk instrument for PKD [GPRI-PKD]).^{15,42}

We have shown that kidney cyst-related pain/bleeding/infection was the most important patient-reported outcome. This is in contrast to other patient CKD populations, those receiving hemodialysis⁴³ and transplant recipients⁴⁴, in which fatigue and life participation were identified as core patient-reported outcomes, respectively. This highlights the need for a core outcome set that is specific to ADPKD.

This Delphi survey involved a large and diverse range of participants from 56 countries with a high retention rate of 70% from round 1 to 2. The survey was available in multiple languages to enable wider engagement. The qualitative data elucidated reasons that explain the prioritization of outcomes.

However, there are some potential limitations. The survey did not measure details including ethnicity or stage of CKD (eg, CKD stage 1) to allow subgroup analysis of outcomes. Although the stages of treatment (eg, dialysis and transplantation) were measured, small numbers precluded

subgroup analysis. Moreover, participants were asked to respond to the survey in the overall context of ADPKD rather than relating to their current treatment stage. The number of surveys completed in French ($n = 47$) was relatively low compared with Korean ($n = 96$) and English ($n = 461$). Furthermore, the online mode of administration used to ensure wider participation precluded involvement from those without access to the internet or with limited computer literacy. However, the top prioritized outcomes in this study were similar to those identified in prior studies of patients with ADPKD elicited through other methods, such as focus groups and workshops.^{14,17,18}

In conclusion, the most important outcomes to patients/caregivers and health professionals were ESKD, kidney function, cerebral aneurysm, and blood pressure. Kidney cyst pain and life participation were the most highly prioritized patient-reported outcomes by patients/caregivers. Before finalizing the core set of outcome domains, public consultation will be sought through a consensus workshop involving patients, caregivers, and health professionals, and any members of the public will be able to access the proposed core outcomes over a 2-week time frame and provide feedback through the SONG website. All input will be reviewed and considered by the SONG-PKD Steering Group to establish a core outcome set. Although the elements discussed as end points in this survey have been studied previously, they have generally been inconsistently reported. Therefore, when a core outcome set has been identified, outcomes measures will be developed through a systematic process (systematic review and workshop) before its implementation in trials. Establishing and implementing a core outcome set will help improve the relevance and consistency of evidence to better inform shared decision making for patients with ADPKD.

Supplementary Material

Supplementary File (PDF)

Figure S1: SONG-PKD Delphi process.

Figure S2: Mean relative importance scores of all participants according to survey completion language based on the BWS.

Figure S3: Mean relative importance scores of health professionals according to survey completion language based on the BWS.

Figure S4: Mean relative importance scores of patients and caregivers according to survey completion language based on the Best-Worst Scale.

Table S1: Collaborating organizations and institutions.

Table S2: Outcomes assessed during round 1 and their definition.

Table S3: Round 1 mean scores, median scores, and proportion rating 7-9 of patients/caregivers and health professionals for 41 outcomes.

Table S4: Outcomes suggested by participants in round 1.

Table S5: Round 2 means, medians, proportions rating 7-9, and BWS scores of patients/caregivers and health professionals for 17 outcomes.

Table S6: Round 2 mean scores and median scores of participants according to survey completion language.

Table S7: Round 2 mean scores of patients/caregivers according to sex.

Article Information

Authors' Full Names and Academic Degrees: Yeoungjee Cho, PhD, Gopala Rangan, PhD, Charlotte Logeman, MPH, Hyunjin Ryu, MD, Benedicte Sautenet, PhD, Ronald D. Perrone, MD, Annie-Claire Nadeau-Fredette, MSc, Reem A. Mustafa, PhD, Htay Htay, MD, Michel Chonchol, MD, Tess Harris, MA, Talia Gutman, MPH, Jonathan C. Craig, PhD, Albert C.M. Ong, DM, Arlene Chapman, PhD, Curie Ahn, PhD, Helen Coolican, MA, Juliana Tze-Wah Kao, PhD, Ron T. Gansevoort, PhD, Vicente Torres, PhD, York Pei, PhD, David W. Johnson, PhD, Andrea K. Viecelli, PhD, Armando Teixeira-Pinto, PhD, Martin Howell, PhD, Angela Ju, PhD, Karine E. Manera, MIPH, and Allison Tong, PhD.

Authors' Affiliations: Department of Nephrology, Princess Alexandra Hospital (YC, DWJ, AKV); Australasian Kidney Trials Network, University of Queensland (YC, DWJ, AKV); Translational Research Institute, Brisbane (YC, DWJ); Centre for Transplant and Renal Research, Westmead Institute for Medical Research, The University of Sydney (GR); Department of Renal Medicine, Westmead Hospital, Western Sydney Local Health District (GR); Sydney School of Public Health, The University of Sydney (CL, TG, AT-P, MH, AJ, KEM, AT); Centre for Kidney Research, The Children's Hospital at Westmead, Sydney, Australia (CL, TG, AT-P, MH, AJ, KEM, AT); Department of Internal Medicine, Seoul National University Hospital, Seoul, South Korea (HR, CA); Department of Nephrology Hypertension, Dialysis, Kidney Transplantation, Tours Hospital, SPHERE – INSERM 1246, University of Tours and Nantes, Tours, France (BS); Division of Nephrology, Tufts Medical Center, Tufts University School of Medicine, Boston, MA (RDP); Department of Nephrology, Hopital Maisonneuve-Rosemont, Montreal, Canada (A-CN-F); Department of Internal Medicine, Division of Nephrology and Hypertension, University of Kansas Medical Center, Kansas City, KS (RAM); Department of Renal Medicine, Singapore General Hospital, Bukit Merah, Singapore (HH); Department of Nephrology, University of Colorado, Denver, CO (MC); Polycystic Kidney Disease International, London, United Kingdom (TH); College of Medicine and Public Health, Flinders University, Adelaide, Australia (JCC); Academic Nephrology Unit, Department of Infection Immunity & Cardiovascular Disease, University of Sheffield, Sheffield, United Kingdom (ACMO); Department of Medicine, The University of Chicago, Chicago, IL (AC); Polycystic Kidney Disease Foundation of Australia, Roseville, NSW, Australia (HC); School of Medicine, Fu Jen Catholic University and Fu Jen Catholic University Hospital (JT-WK); Department of Internal Medicine, National Taiwan University Hospital, Taiwan (JT-WK); Faculty of Medical Sciences, University Medical Center Groningen, Groningen, the Netherlands (RTG); Department of Nephrology and Hypertension, Mayo Clinic, Rochester, MN (VT); Division of Nephrology and Division of Genomic Medicine, University of Toronto, Toronto, Canada (YP); and Department of Nephrology, Mater Hospital, Brisbane, Australia (AKV).

Address for Correspondence: Yeoungjee Cho, PhD, Department of Nephrology, Princess Alexandra Hospital, Ipswich Road, Woolloongabba, Brisbane, QLD 4102, Australia. E-mail: yeoungjee.cho@health.qld.gov.au

Authors' Contributions: Research idea and study design: YC, GR, TG, CL, BS, JCC; data acquisition: YC, BS, CL, HR, TG, AT, A-CN-F, RAM, HH, MC, AKV; data analysis/interpretation: CL, YC, BS, HR, TG, AT, KEM, AJ; statistical analysis: YC, AT-P, BS, MH, KEM, AT; supervision or mentorship: GR, RDP, TH, JCC, ACMO, AC, CA, HC, JT-WK, RTG, VT, YP, DWJ, AT. Each author contributed important intellectual content during manuscript drafting or revision, accepts personal accountability for the author's own

contributions, and agrees to ensure that questions pertaining to the accuracy or integrity of any portion of the work are appropriately investigated and resolved.

Support: Dr Cho is supported by a National Health and Medical Research Council (NHMRC) Early Career Fellowship (APP1126256). Dr Tong is supported by a NHMRC Fellowship (APP11067716). Ms Manera is supported by an NHMRC Postgraduate Scholarship (APP1151343). Dr Johnson is supported by an NHMRC Practitioner Fellowship (APP1117534). The study was financially supported by a grant from Polycystic Kidney Disease Foundation of Australia. The funders or affiliated institutions played no role in study design; collection, analysis, and interpretation of data; writing the report; and the decision to submit the report for publication.

Financial Disclosure: The authors declare that they have no relevant financial interests related to current study.

Acknowledgements: We thank all the patients, caregivers, and health professionals who gave their time to participate in this study.

Peer Review: Received July 14, 2019. Evaluated by 3 external peer reviewers and a methods reviewer, with direct editorial input from an Associate Editor and the Editor-in-Chief. Accepted in revised form January 5, 2020.

References

- ANZDATA. 39th Annual Report: Chapter 12—End stage kidney disease among indigenous peoples of Australia and New Zealand. Adelaide, South Australia: ANZDATA; 2016.
- Levy M, Feingold J. Estimating prevalence in single-gene kidney diseases progressing to renal failure. *Kidney Int*. 2000;58(3):925-943.
- Miskulin DC, Abebe KZ, Chapman AB, et al. Health-related quality of life in patients with autosomal dominant polycystic kidney disease and CKD stages 1-4: a cross-sectional study. *Am J Kidney Dis*. 2014;63(2):214-226.
- Schrier RW, Abebe KZ, Perrone RD, et al. Blood pressure in early autosomal dominant polycystic kidney disease. *N Engl J Med*. 2014;371(24):2255-2266.
- Torres VE, Chapman AB, Perrone RD, et al. Analysis of baseline parameters in the HALT polycystic kidney disease trials. *Kidney Int*. 2012;81(6):577-585.
- Saran R, Robinson B, Abbott KC, et al. US Renal Data System 2017 Annual Data Report: epidemiology of kidney disease in the United States. *Am J Kidney Dis*. 2018;71(3)(suppl 1):S1-S676.
- Chapman AB, Devuyst O, Eckardt KU, et al. Autosomal-dominant polycystic kidney disease (ADPKD): executive summary from a Kidney Disease: Improving Global Outcomes (KDIGO) Controversies Conference. *Kidney Int*. 2015;88(1):17-27.
- Torres VE, Abebe KZ, Schrier RW, et al. Dietary salt restriction is beneficial to the management of autosomal dominant polycystic kidney disease. *Kidney Int*. 2017;91(2):493-500.
- Gansevoort RT, Meijer E, Chapman AB, et al. Albuminuria and tolvaptan in autosomal-dominant polycystic kidney disease: results of the TEMPO 3:4 Trial. *Nephrol Dial Transplant*. 2016;31(11):1887-1894.
- Torres VE, Chapman AB, Devuyst O, et al. Tolvaptan in patients with autosomal dominant polycystic kidney disease. *N Engl J Med*. 2012;367(25):2407-2418.
- Torres VE, Chapman AB, Devuyst O, et al. Multicenter, open-label, extension trial to evaluate the long-term efficacy and safety of early versus delayed treatment with tolvaptan in autosomal dominant polycystic kidney disease: the TEMPO 4:4 Trial. *Nephrol Dial Transplant*. 2018;33(3):477-489.
- Torres VE, Harris PC. Autosomal dominant polycystic kidney disease: the last 3 years. *Kidney Int*. 2009;76(2):149-168.
- Torres VE, Wang X, Qian Q, Somlo S, Harris PC, Gattone VH 2nd. Effective treatment of an orthologous model of autosomal dominant polycystic kidney disease. *Nat Med*. 2004;10(4):363-364.
- Cho Y, Sautenet B, Gutman T, et al. Identifying patient-important outcomes in polycystic kidney disease: an international nominal group technique study. *Nephrology (Carlton)*. 2019;24(12):1214-1224.
- Simms RJ, Thong KM, Dworschak GC, Ong AC. Increased psychosocial risk, depression and reduced quality of life living with autosomal dominant polycystic kidney disease. *Nephrol Dial Transplant*. 2016;31(7):1130-1140.
- Smith KA, Thompson AM, Baron DA, Broadbent ST, Lundstrom GH, Perrone RD. Addressing the need for clinical trial end points in autosomal dominant polycystic kidney disease: a report from the Polycystic Kidney Disease Outcomes Consortium (PKDOC). *Am J Kidney Dis*. 2019;73(4):533-541.
- Tong A, Rangan GK, Ruospo M, et al. A painful inheritance—patient perspectives on living with polycystic kidney disease: thematic synthesis of qualitative research. *Nephrol Dial Transplant*. 2015;30(5):790-800.
- Tong A, Tunnicliffe DJ, Lopez-Vargas P, et al. Identifying and integrating consumer perspectives in clinical practice guidelines on autosomal-dominant polycystic kidney disease. *Nephrology (Carlton)*. 2016;21(2):122-132.
- Sautenet B. *Range and Variability of Outcomes Reported in randomized Trials Conducted in Polycystic Kidney Disease: A Systematic Review*. Melbourne, Australia: World Congress of Nephrology; 2019.
- Cho Y, Sautenet B, Rangan G, et al. Standardised Outcomes in Nephrology-Polycystic Kidney Disease (SONG-PKD): study protocol for establishing a core outcome set in polycystic kidney disease. *Trials*. 2017;18(1):560.
- Tong A, Manns B, Wang AYM, et al. Implementing core outcomes in kidney disease: report of the Standardized Outcomes in Nephrology (SONG) implementation workshop. *Kidney Int*. 2018;94(6):1053-1068.
- Williamson PR, Altman DG, Blazeby JM, et al. Developing core outcome sets for clinical trials: issues to consider. *Trials*. 2012;13:132.
- Boukdedid R, Abdoul H, Loustau M, Sibony O, Alberti C. Using and reporting the Delphi method for selecting healthcare quality indicators: a systematic review. *PLoS One*. 2011;6(6):e20476.
- Blackwood B, Ringrow S, Clarke M, et al. Core Outcomes in Ventilation Trials (COVenT): protocol for a core outcome set using a Delphi survey with a nested randomised trial and observational cohort study. *Trials*. 2015;16:368.
- Evangelidis N, Tong A, Manns B, et al. Developing a set of core outcomes for trials in hemodialysis: an international Delphi survey. *Am J Kidney Dis*. 2017;70(4):464-475.
- MacLennan S, Bekema HJ, Williamson PR, et al. A core outcome set for localised prostate cancer effectiveness trials: protocol for a systematic review of the literature and stakeholder involvement through interviews and a Delphi survey. *Trials*. 2015;16:76.
- Sautenet B, Tong A, Manera KE, et al. Developing consensus-based priority outcome domains for trials in kidney transplantation: a multinational Delphi survey with patients, caregivers and health professionals. *Transplantation*. 2017;101(8):1875-1886.

28. Manera KE, Tong A, Craig J, et al. Developing consensus-based outcome domains for trials in peritoneal dialysis: an international Delphi survey. *Kidney Int.* 2019;96(3):699-710.
29. Sinha IP, Smyth RL, Williamson PR. Using the Delphi technique to determine which outcomes to measure in clinical trials: recommendations for the future based on a systematic review of existing studies. *PLoS Med.* 2011;8(1):e1000393.
30. Schunemann H, Brozek J, Oxman AD. *GRADE Handbook for Grading Quality of Evidence and Strength of Recommendation.* 2009.
31. Louviere JJ, Flynn TN. Using best-worst scaling choice experiments to measure public perceptions and preferences for healthcare reform in Australia. *Patient.* 2010;3(4):275-283.
32. Flynn TN, Louviere JJ, Peters TJ, Coast J. Best-worst scaling: what it can do for health care research and how to do it. *J Health Econ.* 2007;26(1):171-189.
33. Chapman AB, Schrier RW. Pathogenesis of hypertension in autosomal dominant polycystic kidney disease. *Semin Nephrol.* 1991;11(6):653-660.
34. Eccer T, Schrier RW. Hypertension in autosomal-dominant polycystic kidney disease: early occurrence and unique aspects. *J Am Soc Nephrol.* 2001;12(1):194-200.
35. Schrier RW, Johnson AM, McFann K, Chapman AB. The role of parental hypertension in the frequency and age of diagnosis of hypertension in offspring with autosomal-dominant polycystic kidney disease. *Kidney Int.* 2003;64(5):1792-1799.
36. Baker A, King D, Marsh J, et al. Understanding the physical and emotional impact of early-stage ADPKD: experiences and perspectives of patients and physicians. *Clin Kidney J.* 2015;8(5):531-537.
37. Evangelidis N, Tong A, Manns B, et al. Developing a set of core outcomes for trials in hemodialysis: an international Delphi survey. *Am J Kid Med.* 2017;70(4):464-475.
38. Tong A, Winkelmayer WC, Wheeler DC, et al. Nephrologists' perspectives on defining and applying patient-centered outcomes in hemodialysis. *Clin J Am Soc Nephrol.* 2017;12(3):454-466.
39. Chebib FT, Torres VE. Recent advances in the management of autosomal dominant polycystic kidney disease. *Clin J Am Soc Nephrol.* 2018;13(11):1765-1776.
40. Knight T, Schaefer C, Krasa H, Oberdhan D, Chapman A, Perrone RD. Medical resource utilization and costs associated with autosomal dominant polycystic kidney disease in the USA: a retrospective matched cohort analysis of private insurer data. *Clinicoecon Outcomes Res.* 2015;7:123-132.
41. Barnawi RA, Attar RZ, Alfaer SS, Safdar OY. Is the light at the end of the tunnel nigh? A review of ADPKD focusing on the burden of disease and tolvaptan as a new treatment. *Int J Nephrol Renovasc Dis.* 2018;11:53-67.
42. Oberdhan D, Cole JC, Krasa HB, et al. Development of the Autosomal Dominant Polycystic Kidney Disease Impact Scale: a new health-related quality-of-life instrument. *Am J Kidney Dis.* 2018;71(2):225-235.
43. Tong A, Manns B, Hemmelgarn B, et al. Establishing core outcome domains in hemodialysis: report of the Standardized Outcomes in Nephrology-Hemodialysis (SONG-HD) Consensus Workshop. *Am J Kidney Dis.* 2017;69(1):97-107.
44. Tong A, Gill J, Budde K, et al. Toward establishing core outcome domains for trials in kidney transplantation: report of the Standardized Outcomes in Nephrology-Kidney Transplantation Consensus Workshops. *Transplantation.* 2017;101(8):1887-1896.