University of Groningen

Impulsive waves in ghost-free infinite derivative gravity in anti-de Sitter spacetime

Dengiz, Suat; Kilicarslan, Ercan; Kolar, Ivan; Mazumdar, Anupam

Published in:
Physical Review D

DOI:
[10.1103/PhysRevD.102.044016](https://doi.org/10.1103/PhysRevD.102.044016)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Dengiz, S., Kilicarslan, E., Kolar, I., & Mazumdar, A. (2020). Impulsive waves in ghost-free infinite derivative gravity in anti-de Sitter spacetime. *Physical Review D*, 102(4), [044016].
<https://doi.org/10.1103/PhysRevD.102.044016>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Impulsive waves in ghost-free infinite derivative gravity in anti-de Sitter spacetime

Suat Dengiz^{1,*}, Ercan Kilicarslan^{2,†}, Ivan Kolář^{3,‡} and Anupam Mazumdar^{3,§}

¹*Department of Mechanical Engineering, University of Turkish Aeronautical Association, 06790 Ankara, Turkey*

²*Department of Physics, Usak University, 64200 Usak, Turkey*

³*Van Swinderen Institute, University of Groningen, 9747 AG Groningen, The Netherlands*

(Received 22 June 2020; accepted 22 July 2020; published 7 August 2020)

We study exact impulsive gravitational waves propagating in anti-de Sitter spacetime in the context of the ghost-free infinite derivative gravity. We show that the source-free theory does not admit any anti-de Sitter wave solutions other than that of Einstein's general relativity. The situation is significantly different in the presence of sources. We construct impulsive-wave solutions of the infinite derivative gravity generated by massless particles and linear sources in four and three dimensions. The singularities corresponding to distributional curvature at the locations of the sources get smeared by the nonlocalities. The obtained solutions are regular everywhere. They reduce to the corresponding solutions of general relativity in the infrared regime and in the local limit.

DOI: [10.1103/PhysRevD.102.044016](https://doi.org/10.1103/PhysRevD.102.044016)

I. INTRODUCTION

Einstein's general relativity (GR) has surpassed all observations from Solar System tests to gravitational waves so far [1]. However, it is not well constrained at short distances, i.e., in the ultraviolet regime (UV). Newton's $1/r$ -potential was experimentally tested up to approximately $5 \mu\text{m}$ [2], which corresponds to 0.001 eV . Beyond these scales, gravitational interaction has not been constrained by direct experiments. Furthermore, as one approaches the short distances, GR has several problems. From the classical point of view, it suffers from the presence of spacetime singularities [3]; at the quantum level, it fails to be perturbatively renormalizable.

It has been known for a while that nonlocal terms actions can improve UV behavior. Nonlocal theories containing *form factors* with an infinite number of derivatives have brought considerable interest in the context of quantum field theories [4–11] and quantum gravity [12–16]. In particular, it was shown that infinite derivative gravity (IDG) may resolve cosmological [17] and black-hole singularities [18]. In order to avoid introducing ghostlike instabilities, the form factors are chosen as analytic functions with no roots in the complex plane (i.e., exponential of entire functions); see [12,13,18]. Moreover, the form factor of such a nonlocal action emerges from the world line approximation of one-loop amplitude in string

theory [19,20]. There were also first attempts in studying initial value problem of IDG using diffusion equation method [21,22] and constructing perturbative Hamiltonian [23] using nonlocal Hamiltonian formalism of [24,25].

Recently, there has been further progress in finding solutions of linearized IDG. It was shown that IDG may avoid not only black-hole type singularities [26–33], but also topological defects such as p branes [34], cosmic strings [35], and NUT-like singularities [36]. The exact pp-wave solutions have been studied in [37].

In this paper, we study the nonexpanding gravitational waves of the Siklos type in anti-de Sitter universe, the anti-de Sitter (*AdS*) waves, which are generalizations of the so-called pp waves in flat space in the context of the ghost-free infinite derivative gravity presented in [38,39]. The main focus of this work are the *impulsive waves*, which have been studied extensively in GR with a cosmological constant [40–46]. These solutions are generated by null sources with Dirac-delta stress-energy tensor and belong to the class of almost universal spacetimes [47,48]. The impulsive-wave solution of IDG corresponding to a massless point particle was obtained in [37]. Here, we follow-up by extending the analysis to the AdS spacetime in four and three dimensions. We illustrate how the nonlocality affects the gravitational waves in AdS if the sources are absent or present.

The layout of the paper is as follows: In Sec. II, we briefly review the ghost-free infinite derivative gravity. In Sec. III, we study the AdS wave solutions in the source-free case. Sections IV and V are dedicated to the constructions of the impulsive gravitational waves of IDG in $3+1$ and

*sdengiz@thk.edu.tr

†ercan.kilicarslan@usak.edu.tr

‡i.kolar@rug.nl

§anupam.mazumdar@rug.nl

2 + 1 dimensions, respectively. In Sec. VI, we conclude with a brief discussion of our results. The Supplemental Material is attached to the Appendixes.

II. INFINITE DERIVATIVE GRAVITY

The most general quadratic in curvature (parity-invariant and torsion-free) theory of IDG in four dimensions with a cosmological constant Λ [17,18,38,39] is given by the Lagrangian density¹

$$\mathcal{L} = \frac{\sqrt{-g}}{16\pi G} [R - 2\Lambda + \alpha_c (R\mathcal{F}_1(\Box_s)R + R_{\mu\nu}\mathcal{F}_2(\Box_s)R^{\mu\nu} + C_{\mu\nu\rho\sigma}\mathcal{F}_3(\Box_s)C^{\mu\nu\rho\sigma})], \quad (2.1)$$

where $G = M_p^{-2}$ is Newton's gravitational constant, $\Box_s \equiv \Box/M_s^2$, and $\alpha_c = 1/M_s^2$. The dimensionful constant M_s is the *scale of nonlocality* at which nonlocal interactions become manifest. In the *local limit*, $M_s \rightarrow \infty$, the theory reproduces Einstein's general relativity. The form factors $\mathcal{F}_i(\Box_s)$ are analytic functions of d'Alembert operator $\Box \equiv g_{\mu\nu}\nabla^\mu\nabla^\nu$,

$$\mathcal{F}_i(\Box_s) \equiv \sum_{n=0}^{\infty} f_{i,n} \frac{\Box^n}{M_s^{2n}}, \quad (2.2)$$

where $f_{i,n}$ are dimensionless coefficients. The form factors give rise to nonlocal gravitational interactions. They are crucial to make the theory ghost-free, and the analyticity is required for obtaining the low energy limit similar to that of GR. The equations of motion for the action (2.1) are given in Appendix A.

III. ADS WAVE SPACETIMES IN IDG

The field equations of the infinite derivative gravity are very complicated [49], so a mere attempt of finding exact solutions to the theory is an extremely daunting task. To handle the situation, we focus on the AdS wave metric ansatz, which can be written in the Kerr-Schild form,²

$$g_{\mu\nu} = \bar{g}_{\mu\nu} + 2H\lambda_\mu\lambda_\nu, \quad (3.1)$$

where $\bar{g}_{\mu\nu}$ denotes the AdS background metric, and H is a scalar function that satisfies $\lambda^\mu\partial_\mu H = 0$. Here, λ_μ is a nonexpanding, nontwisting, and shear-free null vector satisfying

$$\lambda^\mu\lambda_\mu = 0, \quad \nabla_\mu\lambda_\nu = \xi_{(\mu}\lambda_{\nu)}, \quad \xi_\mu\lambda^\mu = 0, \quad (3.2)$$

where ξ_μ is a vector in the transverse direction. Due to the fact that the curvature scalar R is constant, there is no

contribution from the nonlocal form factor $R\mathcal{F}_1(\Box_s)R$ to the field equations except a constant term. In addition, the Ricci tensor becomes [53–55]

$$R_{\mu\nu} = -\frac{3}{\ell^2}g_{\mu\nu} + \lambda_\mu\lambda_\nu\mathcal{O}H, \quad (3.3)$$

where ℓ is the AdS radius and $\mathcal{O}$ denotes the operator

$$\mathcal{O} \equiv -\left(\Box + 2\xi^\mu\partial_\mu + \frac{1}{2}\xi^\mu\xi_\mu - \frac{4}{\ell^2}\right). \quad (3.4)$$

Furthermore, one should note that the traceless Ricci tensor takes the form

$$S_{\mu\nu} = \lambda_\mu\lambda_\nu\mathcal{O}H, \quad (3.5)$$

which is of the type N in the aspect of null alignment classification [56,57]. Moreover, one can derive the following formulas for the (repeated) action of the d'Alembert operator [54]:

$$\begin{aligned} \Box(\lambda_\mu\lambda_\nu H) &= \bar{\Box}(\lambda_\mu\lambda_\nu H) = -\lambda_\mu\lambda_\nu\left(\mathcal{O} + \frac{2}{\ell^2}\right)H \\ \Box^n S_{\mu\nu} &= \bar{\Box}^n S_{\mu\nu} = (-1)^n\lambda_\mu\lambda_\nu\left(\mathcal{O} + \frac{2}{\ell^2}\right)^n\mathcal{O}H, \end{aligned} \quad (3.6)$$

where $\bar{\Box} = \bar{g}^{\mu\nu}\bar{\nabla}_\mu\bar{\nabla}_\nu$ is the AdS background d'Alembert operator. Throughout the calculations, one needs to use the following identity of higher-order derivative of the Weyl tensor:

$$\nabla_\mu\nabla_\nu\Box^n C^{\mu\alpha\nu\beta} = \frac{1}{2}\left(\Box + \frac{R}{3}\right)^n\left(\Box - \frac{R}{3}\right)S^{\alpha\beta}. \quad (3.7)$$

By using the recursive relations above, one can easily convert the field equations of the IDG for the AdS wave metric to a rather simple form,

$$\begin{aligned} &\left(\Lambda + \frac{3}{\ell^2}\right)g_{\mu\nu} + \left[1 + \alpha_c\left[\left(2f_{1,0} + \frac{f_{2,0}}{2}\right)R + \left(\bar{\Box} + \frac{2}{\ell^2}\right)\mathcal{F}_2(\bar{\Box}_s) + 2\mathcal{F}_3\left(\bar{\Box}_s - \frac{4}{M_s^2\ell^2}\right)\left(\bar{\Box} + \frac{4}{\ell^2}\right)\right]\right]S_{\mu\nu} = 0. \end{aligned} \quad (3.8)$$

The trace part of the equation determines the cosmological constant in terms of the AdS radius,

$$\Lambda = -\frac{3}{\ell^2}. \quad (3.9)$$

Note that (3.8) reduces to the field equations for pp waves on Minkowski background [37] in the limit $\ell \rightarrow \infty$

¹We use mostly positive metric signature, $(-, +, +, +)$.

²For detailed properties of AdS waves and the Kerr-Schild metrics, we refer the reader to [50–53].

(i.e., $\Lambda \rightarrow 0$). The traceless part of the field equations yields nonlocal equations,

$$\left[1 + \alpha_c \left[-\frac{12}{\ell^2} \left(2f_{1,0} + \frac{f_{2,0}}{2} \right) + \left(\bar{\square} + \frac{2}{\ell^2} \right) \mathcal{F}_2(\bar{\square}_s) + 2\mathcal{F}_3 \left(\bar{\square}_s - \frac{4}{M_s^2 \ell^2} \right) \left(\bar{\square} + \frac{4}{\ell^2} \right) \right] \right] \times \left(\bar{\square} + \frac{2}{\ell^2} \right) \lambda_\mu \lambda_\nu H = 0. \quad (3.10)$$

It is important to stress here that the full equations for AdS waves (3.10) are equivalent to the linearized field equations for the Kerr-Schild perturbations $h_{\mu\nu} = g_{\mu\nu} - \bar{g}_{\mu\nu} = 2H\lambda_\mu\lambda_\nu$. Therefore, the solutions of the full equations that we obtain below are also solutions of the linearized equations for the transverse-traceless fluctuations around AdS background.

To ensure that the theory has no extra degrees of freedom and no ghosts on the AdS background, we choose the form factors³ [39],

$$\begin{aligned} \mathcal{F}_1(\square_s) &= \mathcal{F}_2(\square_s) = 0, \\ \mathcal{F}_3(\square_s) &= \frac{1}{2} \frac{e^{-(\square_s + \frac{6}{\ell^2 M_s^2})} - 1}{\square_s + \frac{8}{\ell^2 M_s^2}}. \end{aligned} \quad (3.11)$$

The AdS wave equation (3.10) then turns into

$$e^{-(\bar{\square}_s + \frac{2}{M_s^2 \ell^2})} \left(\bar{\square} + \frac{2}{\ell^2} \right) \lambda_\mu \lambda_\nu H = 0. \quad (3.12)$$

Let us write AdS wave metric [59] using the null coordinates, $u = (x - t)/\sqrt{2}$ and $v = (x + t)/\sqrt{2}$,

$$ds^2 = \frac{\ell^2}{z^2} (2dudv + dy^2 + dz^2) + 2H(u, y, z)du^2, \quad (3.13)$$

where $z = 0$ corresponds to the conformal infinity of AdS spacetime [60]. In these coordinates, $\xi_\mu = 2z^{-1}\delta_\mu^z$, and, thus,

$$\mathcal{O} = -\left(\bar{\square} + \frac{4z}{\ell^2} \partial_z - \frac{2}{\ell^2} \right), \quad \bar{\square} = \frac{z^2}{\ell^2} \partial^2 - \frac{2z}{\ell^2} \partial_z - \frac{4z^2}{\ell^2} \partial_u \partial_v, \quad (3.14)$$

where we introduced $\partial^2 \equiv \partial_y^2 + \partial_z^2$.

Employing the first formula of (3.6), the field equations (3.12) reduce to

³Let us remark that this choice of $\mathcal{F}_3(\square_s)$ is non-analytic. An alternative analytic choice $\mathcal{F}_3(\square_s) = \frac{1}{2} (e^{-(\square_s + \frac{8}{\ell^2 M_s^2})} - 1) / (\square_s + \frac{8}{\ell^2 M_s^2})$ (discussed in [58]) would only affect the overall constant L and L_4 by the factor $e^{2/\ell^2 M_s^2}$ without changing our conclusions.

$$e^{\frac{z^2 \partial^2 + 2z \partial_z - 2}{M_s^2 \ell^2}} (z^2 \partial^2 + 2z \partial_z - 2) H = 0. \quad (3.15)$$

This equation can be solved using the *eigenvalue method* described in [61]. Let us consider the eigenvalue problem of the operator in the round brackets,

$$(z^2 \partial^2 + 2z \partial_z - 2) H_w = -w^2 H_w, \quad (3.16)$$

where H_w are eigenfunctions and w are the corresponding eigenvalues. By acting with the full nonlocal operator on H_w , we obtain

$$e^{\frac{z^2 \partial^2 + 2z \partial_z - 2}{M_s^2 \ell^2}} (z^2 \partial^2 + 2z \partial_z - 2) H_w = -e^{\frac{w^2}{M_s^2 \ell^2}} w^2 H_w. \quad (3.17)$$

The general solution H of the linear equation (3.15) is a superposition of such functions H_w for which $e^{w^2/M_s^2 \ell^2} w^2 = 0$. Since the exponential has no roots in the complex plane, the only solution is the function H_0 (the eigenvalue $w = 0$). Therefore, the original equation effectively reduces just to the equation

$$(z^2 \partial^2 + 2z \partial_z - 2) H = 0. \quad (3.18)$$

In other words, the only AdS wave solutions of the source-free theory are those of the Einstein's general relativity.⁴ The solutions of (3.18) are well known [52,62],

$$H(u, y, z) = z^{-\frac{1}{2}} [c_1 I_{\frac{3}{2}}(\zeta z) + c_2 K_{\frac{3}{2}}(\zeta z)] \sin(\zeta y + c_3), \quad (3.19)$$

where ζ and c_i are functions of the null coordinate u . The functions $I_{3/2}$ and $K_{3/2}$ are modified Bessel functions of the first and second kinds, respectively.

In fact, this result is expected since the source-free theory is not affected by nonlocalities if the equations of motion are linear, which is exactly the case of the field equations with the AdS wave metric ansatz. In order to see the nonlocal effects, we need to consider the field equations with a nonzero source. The equations derived above remain intact in the presence of nonzero null sources $T_{\mu\nu} dx^\mu dx^\nu = T_{uu} du^2$.⁵

IV. IMPULSIVE WAVES IN 3+1 DIMENSIONS

In this section, we will search for impulsive gravitational waves⁶ that are generated by massless sources in IDG. Since we put a nonzero stress energy on the right-hand side of equations of motion, we can expect that the resulting

⁴This is true also in the Minkowski background, which was studied in [37]. The source-free solution presented in [37] is incorrect because of a mistake in the Fourier transform.

⁵A feasible way of such a source is to consider a nonminimally coupled scalar field with a certain potential [63,64].

⁶For the details on the impulsive gravitational waves in GR with a cosmological constant, see [40–46].

solutions will be affected by the presence of nonlocal form factors with infinite derivatives.

A. Massless pointlike source

Let us begin with the impulsive AdS wave metric,

$$ds^2 = \frac{\ell^2}{z^2} (2dudv + dy^2 + dz^2) + 2\delta(u)H(y, z)du^2, \quad (4.1)$$

and consider a massless point particle traveling in the positive x direction with momentum $p^\mu = E(\delta_t^\mu + \delta_x^\mu)$. Such a particle is described by a source with stress-energy tensor $T_{uu} = Ez_0^2 \ell^{-2} \delta(u) \delta(y) \delta(z - z_0)$. The AdS wave equation then reads

$$e^{\frac{z^2 \partial_z^2 + 2z \partial_z - 2}{M_s^2 \ell^2}} (z^2 \partial_z^2 + 2z \partial_z - 2) H(y, z) = -L \delta(y) \delta(z - z_0), \quad (4.2)$$

where we introduced the constant $L = 16\pi G E z_0^2$. Let us recall that the homogeneous solution is given by (3.19). Since it is the same for the local as well as nonlocal theory, we will focus on finding a particular solution only.

In order to solve (4.2), we first take the Fourier transform in coordinate y ,

$$e^{\frac{z^2 \partial_z^2 + 2z \partial_z - k^2 z^2 - 2}{M_s^2 \ell^2}} (z^2 \partial_z^2 + 2z \partial_z - k^2 z^2 - 2) \hat{H}(k, z) = -\frac{L}{\sqrt{2\pi}} \delta(z - z_0). \quad (4.3)$$

Using the substitution $\hat{H}(k, z) = V(k, z)/\sqrt{z}$, we can rewrite this equation as

$$e^{-\mathcal{A}(k)/M_s^2 \ell^2} \mathcal{A}(k) V(k, z) = -\frac{L\sqrt{z_0}}{\sqrt{2\pi}} \delta(z - z_0), \quad (4.4)$$

where we introduced the k -dependent operator

$$\mathcal{A}(k) \equiv z^2 \partial_z^2 + 2z \partial_z - k^2 z^2 - 2. \quad (4.5)$$

Similar to the homogeneous case, we will first study the eigenvalue problem for this operator. Assuming $k > 0$, one can show that

$$\mathcal{A}(k) K_{i\beta}(kz) = -(\beta^2 + 9/4) K_{i\beta}(kz), \quad (4.6)$$

where $K_{i\beta}$ are modified Bessel functions of imaginary order. In order to make further progress, it is essential to

⁷Our convention for the Fourier transform is

$$\hat{f}(k) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} dx f(y) e^{-iky}, \quad f(y) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} dk \hat{f}(k) e^{iky}.$$

FIG. 1. The function $H(y, z)$ for $z_0 = 1$, $G = 1$, $E = 1$, $\ell = 1$, and $M_s = 4$. The meshed red surface represents the solution of IDG, and the gray surface depicts the corresponding solution of GR.

express the right-hand side of (4.4) in terms of the eigenfunctions $K_{i\beta}(kz)$. Fortunately, this is possible thanks to the identity presented in [65],

$$\delta(z - z_0) = \frac{2}{\pi^2 z_0} \int_0^\infty d\beta \beta \operatorname{sh}(\pi\beta) K_{i\beta}(kz_0) K_{i\beta}(kz), \quad (4.7)$$

for arbitrary $k > 0$. Thus, we can write

$$\begin{aligned} V(k, z) &= -\frac{L\sqrt{z_0}}{\sqrt{2\pi}} \frac{e^{\mathcal{A}(k)/M_s^2 \ell^2}}{\mathcal{A}(k)} \delta(z - z_0) \\ &= \frac{\sqrt{2}L}{\pi^2 \sqrt{z_0}} \int_0^\infty d\beta \frac{e^{-(\beta^2 + 9/4)/M_s^2 \ell^2}}{\beta^2 + 9/4} \beta \operatorname{sh}(\pi\beta) \\ &\quad \times K_{i\beta}(kz_0) K_{i\beta}(kz). \end{aligned} \quad (4.8)$$

After taking the inverse Fourier transform, the particular solution of (4.2) takes the form of the integral

$$\begin{aligned} H(y, z) &= \frac{16GEz_0^{\frac{3}{2}}}{\pi^2 \sqrt{z}} \int_{\mathbb{R}} dk \int_0^\infty d\beta \frac{e^{-(\beta^2 + 9/4)/M_s^2 \ell^2}}{\beta^2 + 9/4} \beta \operatorname{sh}(\pi\beta) \\ &\quad \times K_{i\beta}(|k|z_0) K_{i\beta}(|k|z) e^{iky}, \end{aligned} \quad (4.9)$$

where we also employed the fact that $H(y, z) = H(-y, z)$, as it follows from (4.2). This integral does not seem to have a closed form, but we can evaluate it numerically as shown in Fig. 1.

The GR solution can be obtained by taking the local limit $M_s \rightarrow \infty$ of the integrand in (4.9). Using the identity [66],

$$\int_0^\infty d\beta \frac{\beta \operatorname{sh}(\pi\beta)}{\beta^2 + 9/4} K_{i\beta}(|k|z_0) K_{i\beta} = \begin{cases} \frac{\pi^2}{2} I_{\frac{3}{2}}(|k|z) K_{\frac{3}{2}}(|k|z_0), \\ \frac{\pi^2}{2} I_{\frac{3}{2}}(|k|z_0) K_{\frac{3}{2}}(|k|z), \end{cases} \quad (4.10)$$

which holds for $z < z_0$ and $z > z_0$, respectively, we arrive at the function

$$H_{\text{GR}} = \frac{2GE}{z^2} \left[(y^2 + z^2 + z_0^2) \log \left(1 + \frac{4zz_0}{y^2 + (z - z_0)^2} \right) - 4zz_0 \right]. \quad (4.11)$$

This GR solution represents an impulsive gravitational wave that is generated by a massless particle; see, for example, [40,42,67].

It is clear that the impulsive-wave solution of GR diverges at the location of the particle, where it has distributional curvature. On the other hand, the nonlocal impulsive-wave solution of IDG is regular everywhere due to the improved behavior of the propagator in the UV scale. Let us remark that we could replace $\delta(u)$ by a more realistic smooth regularization of Dirac-delta $\delta_\epsilon(u)$ thanks to the linearity of equations and the independence of the coordinate v [derivative ∂_u in (3.14) never applies]. In this sense, all curvature tensors can be considered as regular. Near the conformal infinity $z = 0$, the nonlocal solution approaches GR.

B. Massless linear source

Let us consider a specific example of a null matter distribution, $T_{uu} = Ez_0 \ell^{-2} \delta(u) \delta(z - z_0)$, for which one can find an impulsive-wave solution in a closed form. This particular stress-energy tensor describes a linear null source that moves in x direction with momentum $p^\mu = E(\delta_t^\mu + \delta_x^\mu)$ and extends to infinity in y direction. The trajectory of this surface is visualized in the Poincaré spherical model of Lobachevsky space in Fig. 2. Details of this representation are reviewed in Appendix B.

This choice of the source allows the profile function H to be independent of y . Thus, the field equation takes a simpler form

$$e^{\frac{z^2 \partial_z^2 + 2z \partial_z - 2}{M_s^2 \ell^2}} (z^2 \partial_z^2 + 2z \partial_z - 2) H(z) = -L_4 \delta(z - z_0), \quad (4.12)$$

where $L_4 = 16\pi GEz_0$. Thanks to the absence of ∂_y , this equation can be solved directly using the *heat-kernel method* [26]. After transforming the equation to the coordinate $w = \log z$ and defining $\tilde{H}(w) = H(e^w)$, we can write

FIG. 2. The trajectory of the source at $z = z_0$ represented in Poincaré spherical model of the Lobachevsky space. The solid lines correspond to the location of the source at a given time t . They extend from $y = 0$ (at the dashed line) toward $y = \pm\infty$ (at the conformal infinity).

$$\begin{aligned} \tilde{H}(w) &= -L_4 e^{-w_0} \frac{e^{(\partial_w^2 + \partial_w - 2)/M_s^2 \ell^2}}{\partial_w^2 + \partial_w - 2} \delta(w - w_0) \\ &= L_4 e^{-w_0} \int_{1/M_s^2 \ell^2}^\infty ds e^{s(\partial_w^2 + \partial_w - 2)} \delta(w - w_0) \\ &= L_4 e^{-w_0} \int_{1/M_s^2 \ell^2}^\infty ds e^{-2s} e^{s\partial_w^2} \delta(w - w_0 + s) \\ &= L_4 e^{-w_0} \int_{1/M_s^2 \ell^2}^\infty ds e^{-2s} \int_{\mathbb{R}} d\tilde{w} \frac{e^{\frac{(w - \tilde{w})^2}{4s}}}{\sqrt{4\pi s}} \delta(\tilde{w} - w_0 + s), \end{aligned} \quad (4.13)$$

where we applied the shift operator $e^{s\partial_w}$ on the third line and expressed the action of $e^{s\partial_w^2}$ using the heat kernel on the fourth line. This integral can be easily found. Returning back to the variable z , we obtain the particular solution of (4.12),

$$\begin{aligned} H(z) &= \frac{8\pi GE}{3z^2 z_0} \left[z_0^3 \operatorname{erfc} \left(\frac{3}{2M_s \ell} - \frac{M_s \ell}{2} \log \left(\frac{z}{z_0} \right) \right) \right. \\ &\quad \left. + z^3 \operatorname{erfc} \left(\frac{3}{2M_s \ell} + \frac{M_s \ell}{2} \log \left(\frac{z}{z_0} \right) \right) \right], \end{aligned} \quad (4.14)$$

which is plotted in Fig. 3.

By taking the local limit $M_s \rightarrow \infty$, we can recover the GR solution,

$$H_{\text{GR}} = \frac{8\pi GEz}{3z_0} \left(1 + \frac{z_0^3}{z^3} - \left| 1 - \frac{z_0^3}{z^3} \right| \right). \quad (4.15)$$

As can be easily seen, this GR solution has a discontinuity and distributional curvature at the location of the source

FIG. 3. The function $H(z)$ for $z_0 = 1$, $G = 1$, $E = 1$, $\ell = 1$, and $M_s = 4$. The dashed red curve denotes the solution of IDG, and the solid black curve represents the corresponding solution of GR.

$z = z_0$, while the IDG solution is completely smooth everywhere. This is again caused by the fact that the form factor with infinite number of derivatives effectively smears the delta-like distributions in the stress-energy tensor. As before, the non-local solution approaches the GR solution near the conformal infinity $z = 0$.

V. IMPULSIVE WAVES IN 2+1 DIMENSIONS

Now that we have discussed gravitational waves in 3+1 dimensions, let us study the solutions in 2+1 dimensions. In this section, we will not repeat details that remain almost the same, but focus on the important differences from the four-dimensional case.

Since the Weyl tensor is identically zero in 2+1 dimensions, the IDG action contains only the form factors of $\mathcal{F}_1(\square_s)$ and $\mathcal{F}_2(\square_s)$. Traceless part of the source-free field equations in three dimensions is reduced to

$$\left[1 + \alpha_c \left[-\frac{12}{\ell^2} \left(f_{1,0} + \frac{f_{2,0}}{3} \right) + \left(\bar{\square} + \frac{2}{\ell^2} \right) \mathcal{F}_2(\bar{\square}_s) \right] \right] \times \left(\bar{\square} + \frac{2}{\ell^2} \right) \lambda_\mu \lambda_\nu H = 0. \quad (5.1)$$

Furthermore, one needs to set the form factor $\mathcal{F}_2(\square_s)$ to be in the following form in order to avoid ghostlike degrees of freedom [68]:

$$\mathcal{F}_2(\square_s) = C \frac{e^{-\left(\square_s + \frac{2}{M_s^2 \ell^2}\right)} - 1}{\square_s + \frac{2}{M_s^2 \ell^2}}, \quad (5.2)$$

where we denoted $C = 1 + \text{th}(M_s^{-2} \ell^{-2})$. It is also important to note that the field equation is independent of the form factor $\mathcal{F}_1(\square_s)$. We refer the reader to [68] for the explicit form of $\mathcal{F}_1(\square_s)$.

The AdS wave metric in 2+1 dimensions is

$$ds^2 = \frac{\ell^2}{z^2} (2dudv + dz^2) + 2H(u, z)du^2. \quad (5.3)$$

A similar arguments to those in Sec. III could be used to show that there are no new solutions of the homogeneous equation. In the next section, we focus on particular solutions in the presence of the nonzero source.

A. Massless pointlike source

Consider a pointlike particle moving in the positive x direction with the momentum $p^\mu = E(\delta_t^\mu + \delta_x^\mu)$ with the stress-energy tensor $T_{uu} = Ez_0^2 \ell^{-2} \delta(u) \delta(z - z_0)$. This source together with the impulsive-wave profile $H = \delta(u)H(z)$ leads to the equation

$$e^{-\frac{z^2 \partial_z^2 + 3z \partial_z}{M_s^2 \ell^2}} (z^2 \partial_z^2 + 3z \partial_z) H(z) = -L_3 \delta(z - z_0), \quad (5.4)$$

where $L_3 = 16\pi G_3 E z_0^2 / C$.

By introducing $w = \log z$, $\tilde{H}(w) = H(e^w)$, and employing the heat-kernel method, we find

$$\begin{aligned} \tilde{H}(w) &= -L_3 e^{-w_0} \frac{e^{(\partial_w^2 + 2\partial_w)/M_s^2 \ell^2}}{\partial_w^2 + 2\partial_w} \delta(w - w_0) \\ &= L_3 e^{-w_0} \int_{1/M_s^2 \ell^2}^{\infty} ds \int_{\mathbb{R}} d\tilde{w} \frac{e^{-\frac{(w-\tilde{w})^2}{4s}}}{\sqrt{4\pi s}} \delta(\tilde{w} - w_0 + 2s), \end{aligned} \quad (5.5)$$

which can be easily calculated. The resulting particular solution of (5.4) is

$$\begin{aligned} H(z) &= \frac{4\pi G_3 E z_0}{C z^2} \left[z_0^2 \text{erfc} \left(\frac{1}{M_s \ell} - \frac{M_s \ell}{2} \log \left(\frac{z}{z_0} \right) \right) \right. \\ &\quad \left. + z^2 \text{erfc} \left(\frac{1}{M_s \ell} + \frac{M_s \ell}{2} \log \left(\frac{z}{z_0} \right) \right) \right]. \end{aligned} \quad (5.6)$$

This function is depicted in Fig. 4.

By calculating the local limit $M_s \rightarrow \infty$, we can arrive at the GR solution,

$$H_{\text{GR}} = 4\pi G_3 E z_0 \left(1 + \frac{z_0^2}{z^2} - \left| 1 - \frac{z_0^2}{z^2} \right| \right). \quad (5.7)$$

Unlike the four-dimensional GR solution of a pointlike massless particle, which diverges, this three-dimensional GR solution is regular but has a discontinuity at $z = z_0$ [46]. This discontinuity is again cured by infinite derivatives. The IDG impulsive-wave solution is smooth everywhere. The full solution (with the homogeneous part $c_1/z^2 + c_2$) approaches the GR solution at the conformal infinity $z = 0$. Note that the metric of the GR solution is actually just the AdS metric. This is a consequence of the

FIG. 4. The function $H(z)$ for $z_0 = 1$, $G_3 = 1$, $E = 1$, $\ell = 1$, and $M_s = 4$. The dashed red curve represents the solution of IDG, and the solid black curve depicts the corresponding solution of GR.

fact that the three-dimensional GR has no local degrees of freedom [69,70]. The spacetime, however, differs from an empty AdS by the presence of nontrivial (global) topological defects that causes distributional curvature, which is not present in the nonlocal case.

VI. CONCLUSIONS

In this paper, we studied the nonexpanding gravitational waves of the Siklos type solutions of the ghost-free infinite derivative gravity in anti-de Sitter spacetime with the main focus on the impulsive waves which are generated by Dirac-delta source. We argued that the source-free infinite derivative gravity does not admit any new AdS wave

solutions other than that of Einstein's general relativity. It was demonstrated that the nonlocality described by form factors with the infinite number of derivatives plays a role only in the presence of a nonzero source.

We found the exact impulsive waves corresponding to massless pointlike and linear sources propagating in four- and three-dimensional anti-de Sitter spacetimes. It turned out that the nonlocalities smear all the divergences and discontinuities (corresponding to distributional curvature) that are present in the local impulsive-wave solutions. The obtained solutions of the infinite derivative gravity are regular everywhere. They reduce to the impulsive-waves solutions of general relativity in the local limit $M_s \rightarrow \infty$ and in the infrared regime (near the conformal infinity of AdS). Simply put, the solutions get modified due to the nonlocal effects only in the ultraviolet regime, but not in the infrared regime.

ACKNOWLEDGMENTS

We thank Bayram Tekin, Tahsin C. Sisman, Ayse K. Karasu, Pavel Krtouš, and Eloy Ayon-Beato for useful discussions and suggestions. We also thank an anonymous referee for constructive suggestions. The works of S. D., E. K., and A. M. are supported by the TUBITAK Grant No. 119F241. I. K. and A. M. are supported by Netherlands Organization for Scientific Research Grant No. 680-91-119.

APPENDIX A: EQUATIONS OF MOTION OF IDG

The equations of motion following from the action given (2.1) were found in [49]. Using a common notation for a power of d'Alembert operator, $\square^n X_{\beta\dots}^{\alpha\dots} \equiv X_{\beta\dots}^{\alpha\dots(n)}$, they can be written as

$$\begin{aligned}
 G^{\alpha\beta} + \Lambda g^{\alpha\beta} + \frac{\alpha_c}{2} [4G^{\alpha\beta} \mathcal{F}_1(\square)R + g^{\alpha\beta} R \mathcal{F}_1(\square)R - 4(\nabla^\alpha \nabla^\beta - g^{\alpha\beta} \square) \mathcal{F}_1(\square)R - 2\Omega_1^{\alpha\beta} + g^{\alpha\beta} (\Omega_{1\rho}^\rho + \bar{\Omega}_1) + 4R^\alpha{}_\nu \mathcal{F}_2(\square)R^{\nu\beta} \\
 - g^{\alpha\beta} R_\nu{}^\mu \mathcal{F}_2(\square)R_\mu{}^\nu - 4\nabla_\nu \nabla^\beta (\mathcal{F}_2(\square)R^{\nu\alpha}) + 2\square(\mathcal{F}_2(\square)R^{\alpha\beta}) + 2g^{\alpha\beta} \nabla_\mu \nabla_\nu (\mathcal{F}_2(\square)R^{\mu\nu}) - 2\Omega_2^{\alpha\beta} + g^{\alpha\beta} (\Omega_{2\rho}^\rho + \bar{\Omega}_2) - 4\Delta_2^{\alpha\beta} \\
 - g^{\alpha\beta} C^{\mu\nu\rho\sigma} \mathcal{F}_3(\square)C_{\mu\nu\rho\sigma} + 4C_{\mu\nu\sigma}^\alpha \mathcal{F}_3(\square)C^{\beta\mu\nu\sigma} - 4(R_{\mu\nu} + 2\nabla_\mu \nabla_\nu) (\mathcal{F}_3(\square)C^{\beta\mu\nu\alpha}) - 2\Omega_3^{\alpha\beta} + g^{\alpha\beta} (\Omega_{3\gamma}^\gamma + \bar{\Omega}_3) - 8\Delta_3^{\alpha\beta}] = 0, \quad (A1)
 \end{aligned}$$

where the symmetric tensors are

$$\begin{aligned}
 \Omega_1^{\alpha\beta} &= \sum_{n=1}^{\infty} f_{1,n} \sum_{l=0}^{n-1} \nabla^\alpha R^{(l)} \nabla^\beta R^{(n-l-1)}, & \bar{\Omega}_1 &= \sum_{n=1}^{\infty} f_{1,n} \sum_{l=0}^{n-1} R^{(l)} R^{(n-l)}, \\
 \Omega_2^{\alpha\beta} &= \sum_{n=1}^{\infty} f_{2,n} \sum_{l=0}^{n-1} R_{\nu}{}^{\mu;\alpha(l)} R_{\mu}{}^{\nu;\beta(n-l-1)}, & \bar{\Omega}_2 &= \sum_{n=1}^{\infty} f_{2,n} \sum_{l=0}^{n-1} R_{\nu}{}^{\mu(l)} R_{\mu}{}^{\nu(n-l)}, \\
 \Omega_3^{\alpha\beta} &= \sum_{n=1}^{\infty} f_{3,n} \sum_{l=0}^{n-1} C_{\nu\rho\sigma}^{\mu;\alpha(l)} C_{\mu}{}^{\nu\rho\sigma;\beta(n-l-1)}, & \bar{\Omega}_3 &= \sum_{n=1}^{\infty} f_{3,n} \sum_{l=0}^{n-1} C_{\nu\rho\sigma}^{\mu(l)} C_{\mu}{}^{\nu\rho\sigma(n-l)}, \\
 \Delta_2^{\alpha\beta} &= \frac{1}{2} \sum_{n=1}^{\infty} f_{2,n} \sum_{l=0}^{n-1} [R_{\sigma}{}^{\nu(l)} R^{(\beta|\sigma|\alpha)(n-l-1)} - R_{\sigma}{}^{\nu;\alpha(l)} R^{(\beta)\sigma(n-l-1)}]_{;\nu}, \\
 \Delta_3^{\alpha\beta} &= \frac{1}{2} \sum_{n=1}^{\infty} f_{3,n} \sum_{l=0}^{n-1} [C^{\rho\nu(l)}_{\sigma\mu} C_{\rho}{}^{(\beta|\sigma\mu|\alpha)(n-l-1)} - C^{\rho\nu}{}_{\sigma\mu}{}^{;\alpha(l)} C_{\rho}{}^{\beta)\sigma\mu(n-l-1)}]_{;\nu}. \quad (A2)
 \end{aligned}$$

FIG. 5. Surfaces of constant x , y , and z coordinates of the Lobachevsky space depicted in the Poincaré spherical model.

APPENDIX B: POINCARÉ SPHERICAL MODEL

Poincaré spherical model is a compactified representation of the Lobachevsky space,

$$ds^2 = \frac{\ell^2}{z^2} (dx^2 + dy^2 + dz^2), \quad (\text{B1})$$

which is a spatial part of the AdS metric. The surface of the sphere is the conformal infinity. The standard conformally flat coordinates x , y , and z are visualized in Fig. 5.

-
- [1] C. M. Will, The confrontation between general relativity and experiment, *Living Rev. Relativity* **17**, 4 (2014).
 - [2] D. J. Kapner, T. S. Cook, E. G. Adelberger, J. H. Gundlach, B. R. Heckel, C. D. Hoyle, and H. E. Swanson, Tests of the Gravitational Inverse-Square Law Below the Dark-Energy Length Scale, *Phys. Rev. Lett.* **98**, 021101 (2007).
 - [3] S. Hawking and G. Ellis, The large scale structure of space-time, <https://doi.org/10.1017/CBO9780511524646>.
 - [4] G. V. Efimov, Non-local quantum theory of the scalar field, *Commun. Math. Phys.* **5**, 42 (1967).
 - [5] G. V. Efimov and S. Z. Seltser, Gauge invariant nonlocal theory of the weak interactions, *Ann. Phys. (N.Y.)* **67**, 124 (1971).
 - [6] G. V. Efimov, On the construction of nonlocal quantum electrodynamics, *Ann. Phys. (N.Y.)* **71**, 466 (1972).
 - [7] N. V. Krasnikov, Nonlocal gauge theories, *Theor. Math. Phys.* **73**, 1184 (1987).
 - [8] J. W. Moffat, Finite nonlocal gauge field theory, *Phys. Rev. D* **41**, 1177 (1990).
 - [9] D. Evens, J. W. Moffat, G. Kleppe, and R. P. Woodard, Nonlocal regularizations of gauge theories, *Phys. Rev. D* **43**, 499 (1991).
 - [10] E. T. Tomboulis, Nonlocal and quasilocal field theories, *Phys. Rev. D* **92**, 125037 (2015).
 - [11] L. Buoninfante, G. Lambiase, and A. Mazumdar, Ghost-free infinite derivative quantum field theory, *Nucl. Phys. B* **944**, 114646 (2019).
 - [12] E. Tomboulis, Renormalizability and asymptotic freedom in quantum gravity, *Phys. Lett.* **97B**, 77 (1980).
 - [13] E. Tomboulis, Superrenormalizable gauge and gravitational theories, [arXiv:hep-th/9702146](https://arxiv.org/abs/hep-th/9702146).
 - [14] L. Modesto, Super-renormalizable quantum gravity, *Phys. Rev. D* **86**, 044005 (2012).
 - [15] A. A. Tseytlin, On singularities of spherically symmetric backgrounds in string theory, *Phys. Lett. B* **363**, 223 (1995).
 - [16] W. Siegel, Stringy gravity at short distances, [arXiv:hep-th/0309093](https://arxiv.org/abs/hep-th/0309093).
 - [17] T. Biswas, A. Mazumdar, and W. Siegel, Bouncing universes in string-inspired gravity, *J. Cosmol. Astropart. Phys.* **03** (2006) 009.
 - [18] T. Biswas, E. Gerwick, T. Koivisto, and A. Mazumdar, Towards Singularity and Ghost Free Theories of Gravity, *Phys. Rev. Lett.* **108**, 031101 (2012).
 - [19] S. Abel and N. A. Dondi, UV completion on the worldline, *J. High Energy Phys.* **07** (2019) 090.
 - [20] S. Abel, L. Buoninfante, and A. Mazumdar, Nonlocal gravity with worldline inversion symmetry, *J. High Energy Phys.* **01** (2020) 003.
 - [21] G. Calcagni, L. Modesto, and G. Nardelli, Non-perturbative spectrum of non-local gravity, *Phys. Lett. B* **795**, 391 (2019).
 - [22] G. Calcagni, L. Modesto, and G. Nardelli, Initial conditions and degrees of freedom of non-local gravity, *J. High Energy Phys.* **05** (2018) 087.
 - [23] I. Kolar and A. Mazumdar, Hamiltonian for scalar field model of infinite derivative gravity, *Phys. Rev. D* **101**, 124028 (2020).
 - [24] J. Llosa and J. Vives, Nonlocal Lagrangians and Hamiltonian formalism, *Int. J. Mod. Phys. D* **03**, 211 (1994).
 - [25] J. Gomis, K. Kamimura, and J. Llosa, Hamiltonian formalism for space-time noncommutative theories, *Phys. Rev. D* **63**, 045003 (2001).

- [26] V. P. Frolov, A. Zelnikov, and T. de Paula Netto, Spherical collapse of small masses in the ghost-free gravity, *J. High Energy Phys.* **06** (2015) 107.
- [27] V. P. Frolov and A. Zelnikov, Head-on collision of ultra-relativistic particles in ghost-free theories of gravity, *Phys. Rev. D* **93**, 064048 (2016).
- [28] J. Edholm, A. S. Koshelev, and A. Mazumdar, Behavior of the Newtonian potential for ghost-free gravity and singularity-free gravity, *Phys. Rev. D* **94**, 104033 (2016).
- [29] E. Kilicarslan, Weak field limit of infinite derivative gravity, *Phys. Rev. D* **98**, 064048 (2018).
- [30] L. Buoninfante, G. Harmsen, S. Maheshwari, and A. Mazumdar, Nonsingular metric for an electrically charged point-source in ghost-free infinite derivative gravity, *Phys. Rev. D* **98**, 084009 (2018).
- [31] L. Buoninfante, A. S. Koshelev, G. Lambiase, and A. Mazumdar, Classical properties of non-local, ghost- and singularity-free gravity, *J. Cosmol. Astropart. Phys.* **09** (2018) 034.
- [32] L. Buoninfante, A. S. Cornell, G. Harmsen, A. S. Koshelev, G. Lambiase, J. Marto, and A. Mazumdar, Towards non-singular rotating compact object in ghost-free infinite derivative gravity, *Phys. Rev. D* **98**, 084041 (2018).
- [33] J. Boos, J. P. Soto, and V. P. Frolov, Ultrarelativistic spinning objects (gyratons) in non-local ghost-free gravity, *Phys. Rev. D* **101**, 124065 (2020).
- [34] J. Boos, V. P. Frolov, and A. Zelnikov, Gravitational field of static p-branes in linearized ghost-free gravity, *Phys. Rev. D* **97**, 084021 (2018).
- [35] J. Boos, Angle deficit & non-local gravitoelectromagnetism around a slowly spinning cosmic string, [arXiv:2003.13847](https://arxiv.org/abs/2003.13847).
- [36] I. Kolar and A. Mazumdar, NUT charge in linearized infinite derivative gravity, *Phys. Rev. D* **101**, 124005 (2020).
- [37] E. Kilicarslan, pp -waves as exact solutions to ghost-free infinite derivative gravity, *Phys. Rev. D* **99**, 124048 (2019).
- [38] T. Biswas, A. S. Koshelev, and A. Mazumdar, Gravitational theories with stable (anti-)de Sitter backgrounds, *Fundam. Theor. Phys.* **183**, 97 (2016).
- [39] T. Biswas, A. S. Koshelev, and A. Mazumdar, Consistent higher derivative gravitational theories with stable de Sitter and anti-de Sitter backgrounds, *Phys. Rev. D* **95**, 043533 (2017).
- [40] M. Hotta and M. Tanaka, Shock wave geometry with nonvanishing cosmological constant, *Classical Quantum Gravity* **10**, 307 (1993).
- [41] J. Podolsky and J. B. Griffiths, Impulsive gravitational waves generated by null particles in de Sitter and anti-de Sitter backgrounds, *Phys. Rev. D* **56**, 4756 (1997).
- [42] J. Podolsky and J. B. Griffiths, Impulsive waves in de Sitter and anti-de Sitter space-times generated by null particles with an arbitrary multipole structure, *Classical Quantum Gravity* **15**, 453 (1998).
- [43] J. Podolsky, Exact impulsive gravitational waves in space-times of constant curvature, in *Gravitation: Following the Prague Inspiration (A Volume in Celebration of the 60th Birthday of Jiri Bičák)*, edited by O. Semerák, J. Podolský, and M. Zofka (World Scientific, Singapore, 2002), pp. 205–246.
- [44] J. Podolsky and J. B. Griffiths, Nonexpanding impulsive gravitational waves with an arbitrary cosmological constant, *Phys. Lett. A* **261**, 1 (1999).
- [45] J. Bicak and J. Podolsky, Gravitational waves in vacuum space-times with cosmological constant. 1. Classification and geometrical properties of nontwisting type N solutions, *J. Math. Phys. (N.Y.)* **40**, 4495 (1999).
- [46] R. G. Cai and J. B. Griffiths, Null particle solutions in three-dimensional (anti-)de Sitter spaces, *J. Math. Phys. (N.Y.)* **40**, 3465 (1999).
- [47] S. Hervik, V. Pravda, and A. Pravdova, Type III and N universal spacetimes, *Classical Quantum Gravity* **31**, 215005 (2014).
- [48] M. Kuchynka, T. Malek, V. Pravda, and A. Pravdova, Almost universal spacetimes in higher-order gravity theories, *Phys. Rev. D* **99**, 024043 (2019).
- [49] T. Biswas, A. Conroy, A. S. Koshelev, and A. Mazumdar, Generalized ghost-free quadratic curvature gravity, *Classical Quantum Gravity* **31**, 015022 (2014).
- [50] R. P. Kerr and A. Schild, Some algebraically degenerate solutions of Einstein's gravitational field equations, *Proc. Symp. Appl. Math.* **17**, 199 (1965).
- [51] T. Malek and V. Pravda, Kerr-Schild spacetimes with (A)dS background, *Classical Quantum Gravity* **28**, 125011 (2011).
- [52] I. Gullu, M. Gurses, T. C. Sisman, and B. Tekin, AdS waves as exact solutions to quadratic gravity, *Phys. Rev. D* **83**, 084015 (2011).
- [53] M. Gurses, T. C. Sisman, and B. Tekin, New exact solutions of quadratic curvature gravity, *Phys. Rev. D* **86**, 024009 (2012).
- [54] M. Gurses, T. C. Sisman, and B. Tekin, AdS-plane wave and pp -wave solutions of generic gravity theories, *Phys. Rev. D* **90**, 124005 (2014).
- [55] M. Gurses, T. C. Sisman, B. Tekin, and S. Hervik, AdS-Wave Solutions of F(Riemann) Theories, *Phys. Rev. Lett.* **111**, 101101 (2013).
- [56] R. Milson, A. Coley, V. Pravda, and A. Pravdova, Alignment and algebraically special tensors in Lorentzian geometry, *Int. J. Geom. Methods Mod. Phys.* **02**, 41 (2005).
- [57] A. Coley, S. Hervik, G. O. Papadopoulos, and N. Pelavas, Kundt spacetimes, *Classical Quantum Gravity* **26**, 105016 (2009).
- [58] K. Sravan Kumar, S. Maheshwari, and A. Mazumdar, Perturbations in higher derivative gravity beyond maximally symmetric spacetimes, *Phys. Rev. D* **100**, 064022 (2019).
- [59] S. T. C. Siklos, in *Galaxies, Axisymmetric Systems and Relativity*, edited by M. A. H. MacCallum (Cambridge University Press, Cambridge, United Kingdom, 1985).
- [60] J. Podolsky, Interpretation of the Siklos solutions as exact gravitational waves in the anti-de Sitter universe, *Classical Quantum Gravity* **15**, 719 (1998).
- [61] N. Barnaby and N. Kamran, Dynamics with infinitely many derivatives: Variable coefficient equations, *J. High Energy Phys.* **12** (2008) 022.
- [62] A. Chamblin and G. Gibbons, Supergravity on the Brane, *Phys. Rev. Lett.* **84**, 1090 (2000).
- [63] E. Ayon-Beato and M. Hassaine, Exploring AdS waves via nonminimal coupling, *Phys. Rev. D* **73**, 104001 (2006).
- [64] E. Ayon-Beato and M. Hassaine, Higher-dimensional AdS waves and pp -waves with conformally related sources, *Phys. Rev. D* **75**, 064025 (2007).
- [65] S. Murashima and T. Kiyono, Modified Bessel functions of purely imaginary order $K_{is}(x)$, $I_{is}(x)$ and their related functions, *Res. Rep. Kagoshima Univ. Faculty Eng.* **15**, 91 (1973).

- [66] I. S. Gradshteyn and I. M. Ryzhik, in *Table of Integrals, Series, and Products*, 7th ed. edited by A. Jeffrey and D. Zwillinger, (Academic Press, New York, 2007).
- [67] X. O. Camanho, J. D. Edelstein, J. Maldacena, and A. Zhiboedov, Causality constraints on corrections to the graviton three-point coupling, *J. High Energy Phys.* **02** (2016) 020.
- [68] A. Mazumdar and G. Stettinger, New massless and massive infinite derivative gravity in three dimensions and perturbations around Minkowski and in (A)dS, *Nucl. Phys.* **B956**, 115024 (2020).
- [69] S. Deser, R. Jackiw, and G. 't Hooft, Three-dimensional Einstein gravity: Dynamics of flat space, *Ann. Phys. (N.Y.)* **152**, 220 (1984).
- [70] S. Deser and R. Jackiw, Three-dimensional cosmological gravity: Dynamics of constant curvature, *Ann. Phys. (N.Y.)* **153**, 405 (1984).