University of Groningen

The absence of the peroxiredoxin Pmp20 causes permeabilisation of the peroxisomal membrane and necrotic cell death

Bener Aksam, Eda; Jungwirth, Helmut; Kohlwein, Sepp D.; Ring, Julia; Madeo, Frank; Veenhuis, Marten; Klei, Ida J. van der

Published in:
Chemistry and physics of lipids

DOI:
[10.1016/j.chemphyslip.2008.05.171](https://doi.org/10.1016/j.chemphyslip.2008.05.171)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2008

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Bener Aksam, E., Jungwirth, H., Kohlwein, S. D., Ring, J., Madeo, F., Veenhuis, M., & Klei, I. J. V. D. (2008). The absence of the peroxiredoxin Pmp20 causes permeabilisation of the peroxisomal membrane and necrotic cell death. *Chemistry and physics of lipids*, 154, S62.
<https://doi.org/10.1016/j.chemphyslip.2008.05.171>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

PO 110

Conformational change of apolipoprotein A-I and promotion of HDL formation at acidic conditions

Masakazu Fukuda, Minoru Nakano, Tetsurou Handa

Graduate School of Pharmaceutical Sciences, Kyoto University, Japan

The molecular mechanism by which nascent high-density lipoprotein (HDL) forms via the interaction of apolipoprotein A-I (apoA-I) and transmembrane ABCA1 is poorly understood. Here, as ABCA1 has been reported to localize to acidic intracellular compartments including the Golgi and endosome, we studied the interaction of apoA-I with model membranes in acidic conditions. Pure phosphatidylcholine (PC) liposomes were persistent against apoA-I at pH levels above 5.0, but were progressively transformed into reconstituted HDLs (rHDLs) by apoA-I at lower pH. CD and ANS fluorescence measurements of lipid-free apoA-I indicated that the accelerated formation of rHDLs was caused by the formation of α -helical structure and the increased hydrophobicity of apoA-I in acidic conditions. The addition of phosphatidylserine (PS) increased the acidity at bilayer's surface and enabled the formation of discoidal rHDLs even at the pH of the endosome and slightly lower pH of the Golgi. These results suggest a following new scenario of the nascent HDL formation; ABCA1 that colocalizes with apoA-I in the acidic intracellular compartments including the Golgi and endosome increases the acidity at the membrane's surface in the luminal side by its PS translocase activity and causes apoA-I to form nascent HDL.

doi:10.1016/j.chemphyslip.2008.05.169

PO 111

The accumulation of two atypical sphingolipids cause hereditary sensory neuropathy type 1 (HSAN1)

T. Hornemann, A. Penno, A. von Eckardstein

University Hospital Zurich, Rämistrasse 100, CH-8091 Zürich, Switzerland

Hereditary sensory neuropathy I (HSAN1) is an autosomal dominant inherited neuropathy that primarily affects peripheral sensory neurons. Patients suffer from a severe sensory loss leading to painless injuries and chronic skin ulcers. The disease is caused by several missense mutations in the *SPTLC1* gene of serine palmitoyltransferase (SPT). SPT catalyses the condensation of serine with palmitoyl-CoA—the first step in the de-novo synthesis pathway of sphingolipids.

We discovered recently that the HSN1 mutations in SPT lead to a shift in the substrate specificity of this enzyme. The mutant SPT can also metabolise alanine and glycine instead of serine as alternative substrates. The conjugation of palmitoyl-CoA with alanine or glycine results in the formation of the two atypical sphingolipids—Deoxy-sphinganine (DoxSA, m18:0) and 1-amino-2-deoxy-*n*-heptadecane (ADHD, m17:0). Hek293 cells which express the mutant form of SPTLC1 show a pronounced accumulation of these two metabolites. The absence of the C₁-OH group in DoxSA and ADHD blocks the further transformation towards complex sphingolipids (e.g. sphingomyeline or glycosphingolipids) but also prevents the degradation via the formation of Sphingosine-1P. Consequently, those “dead end” metabolites accumulate in the cells of HSN1 patient. This was confirmed by analyzing EBV lymphoblast lines from 12 HSN1 patients. The HSN1 lymphoblasts show 5–10-fold higher levels of DoxSA and ADHD compared to controls.

In concordance with this we find, furthermore, highly elevated levels of DoxSA and ADHD in the blood of HSN1 patients.

We therefore conclude that the toxic accumulation of these atypical sphingolipids provides the pathophysiological background for HSN1.

doi:10.1016/j.chemphyslip.2008.05.170

PO 112

The absence of the peroxiredoxin Pmp20 causes permeabilisation of the peroxisomal membrane and necrotic cell deathEda Bener Aksam¹, Helmut Jungwirth², Sepp D. Kohlwein², Julia Ring², Frank Madeo², Marten Veenhuis^{1,3}, Ida J. van der Klei^{1,3}¹ Molecular Cell Biology, Groningen Biomolecular Sciences and Biotechnology Institute, University of Groningen, Haren, The Netherlands² Institute of Molecular Biosciences, University of Graz, 8010 Graz, Austria³ Kluyver Centre for Genomics of Industrial Fermentation, Julianalaan 67, 2628 BC Delft, The Netherlands

Peroxisomes are important cellular organelles, which contain H₂O₂ producing oxidases together with catalase, which degrades H₂O₂. The presence of catalase in peroxisomes is generally assumed to prevent release of H₂O₂ into the cytoplasm. Peroxisomes, however, also contain other anti-oxidant enzymes, among others peroxiredoxins (Prx's). The physiological function of peroxisomal Prx's is still speculative.

Prx's are involved in the degradation of H₂O₂ and organic hydroperoxides. Prx's have been localized to the cytosol, the endoplasmic reticulum, mitochondria, nuclei and peroxisomes.

The first peroxisomal Prx, Pmp20, was identified in *Candida boidinii* and has glutathione peroxidase activity towards alkyl hydroperoxides and H₂O₂ (Horiguchi et al., 2001).

We identified the Pmp20 homologue of the yeast *Hansenula polymorpha* and analyzed its function *in vivo*. During growth of *H. polymorpha* on methanol massive amounts of H₂O₂ are produced in peroxisomes. We show that cells of a *H. polymorpha* PMP20 disruption strain (*pmp20*) have a severe growth defect on methanol, which is paralleled by permeabilisation of the peroxisomal membrane and leakage of peroxisomal matrix proteins into the cytosol.

Methanol-induced *pmp20* cells accumulated enhanced levels of lipid peroxidation products. Moreover, the fatty acid composition of methanol induced *pmp20* cells differed relative to WT controls, suggesting an effect on fatty acid homeostasis. Plating assays and FACS-based analysis of cell death markers revealed that *pmp20* cells show loss of clonogenic efficiency and membrane integrity, when cultured on methanol.

We conclude that the absence of the peroxisomal peroxiredoxin leads to loss of peroxisome membrane integrity and necrotic cell death.

Reference

Horiguchi, et al., 2001. J. Biol. Chem. 276, 14279–14288.

doi:10.1016/j.chemphyslip.2008.05.171