University of Groningen

Lung allograft rejection in the rat

Prop, Jochum

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

1984

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Prop, J. (1984). *Lung allograft rejection in the rat*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

SUMMARY

Although the first lung transplantation in a patient was performed over 20 years ago, it is only during the last 3 years that some of these operations have been successful. One of the major obstacles, impeding success for so many years, was the strong rejection response induced by lung grafts. Based on these poor results it was suggested, that rejection of lung grafts was more vigorous and more resistant to immunosuppressive treatment than rejection of other organ grafts. The mechanism underlying this suggested difference between rejection of lungs and other organs was however obscure.

The question whether lungs are rejected more vigorously than other organs is treated in this thesis. Particular attention is paid to the possible role of the bronchus-associated lymphoid tissue (BALT) transplanted within the lung graft, and the influence of inhaled stimuli reaching the lung graft via the airways.

In *paper 1* it is shown that in the absence of rejection the function of lung grafts is slightly affected by various factors of the transplantation procedure, in particular by a disruption of the hilar structures and ischemia. The grafts largely recover from this reimplantation response, and their function then remains constant for the rest of the observation period.

In *paper 2* it is shown that on the other hand during rejection the function of lung grafts drops sharply. The pace of rejection in the various donor-recipient combinations studied depends on the degree of incompatibility of donor and recipient. This functional study confirms that lungs are more rapidly rejected than other organs, when transplanted across the major histocompatibility barrier.

Paper 3 focuses on the induction of the rejection response against lung allografts. The first features of rejection are found in the graft's BALT which is heavily infiltrated by recipient lymphocytes, while simultaneously lymphocytes from the graft leave the BALT and home into the recipient's lymphoid tissues. It is concluded that the graft's BALT intensifies the contact between graft and recipient, so as to accelerate the rejection response in lung grafts.

In *paper 4*, the histological changes in lung grafts during rejection, investigated appear to form a universal pattern in all donor-recipient combinations studied. This pattern consists of four consecutive rejection phases, each with distinct morphological characteristics. Previous reports on the histological changes in lung grafts described more isolated pictures, but these

are found to generally fit in the rejection pattern as defined in this paper.

In *paper 5*, the influence of BALT upon lung allograft rejection is eliminated either by roentgen irradiation of the donor, or by retransplantation of the graft. The prolonged survival of irradiated grafts confirms that BALT lymphocytes of the graft accelerate the rejection response. The much longer survival of retransplanted grafts however indicates that cells other than lymphocytes, probably dendritic cells, are the major stimulus for rejection.

Paper 6 demonstrates the immunological potency of the lymphoid tissue of lung grafts which is capable of mounting a lethal Graft-vs-Host disease when lungs are transplanted from a parental strain into a F1 recipient. It is conceivable that Graft-vs-Host responses interfere with the rejection response in fully allogeneic combinations.

In *paper 7*, the influence of inhaled stimuli on the subsequent rejection of lung grafts is investigated. Therefore grafted rats are exposed to various degrees of stimuli via the airways. In this way it is shown that inhaled stimuli aggravate the rejection response against the graft.

In *paper 8* is described how rejection of lung grafts can be prevented by treatment with cyclosporine, a new and very potent immunosuppressive drug. Surprisingly, one dose of cyclosporine injected three days after transplantation suffices to prolong survival of most grafts indefinitely.

In the concluding chapter, it is stated that the extraordinarily rapid rejection of lung grafts might well be explained by the role of BALT in lung grafts in combination with the effect of inhaled stimuli. The implications of our studies for some aspects of lung transplantation which might be relevant for clinical lung transplantation (such as graft preservation, donor-recipient matching, diagnosis of rejection, and immunosuppression) are discussed here in more detail.