University of Groningen

Monte Carlo Simulation of a Kinetic Ising Model of the Glass Transition

Leutheusser, E.; Raedt, Hans De

Published in:
Solid State Communications

DOI:
[10.1016/0038-1098\(86\)90491-6](https://doi.org/10.1016/0038-1098(86)90491-6)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1986

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Leutheusser, E., & Raedt, H. D. (1986). Monte Carlo Simulation of a Kinetic Ising Model of the Glass Transition. *Solid State Communications*, 57(6). [https://doi.org/10.1016/0038-1098\(86\)90491-6](https://doi.org/10.1016/0038-1098(86)90491-6)

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

MONTE CARLO SIMULATION OF A KINETIC ISING MODEL OF THE GLASS TRANSITION

E. Leutheusser

Technische Universität München, D-8046 Garching, West Germany

and

Hans De Raedt*

Max-Planck-Institut für Physik und Astrophysik, D-8000 München, West Germany

(Received 3 September 1985 by B. Mühlischlegel)

Results of Monte Carlo experiments for the two-spin facilitated kinetic Ising model on a cubic lattice are presented and compared with a theoretical prediction.

IN A RECENT LETTER Fredrickson and Andersen [1] introduced a special kinetic Ising model as a model for structural relaxation in dense liquids and glasses. The Hamiltonian is the usual Ising one with ferromagnetic coupling J between nearest neighbor spins in a magnetic field H which tends to align the spins downwards. As in the Glauber model [2] the dynamics is described by a master equation with transition probabilities satisfying the detailed balance principle. In a special version, the two-spin facilitated model, the down-flip rate of a particular spin σ_j is chosen as

$$w_{j,\text{down}}(\sigma_j) = \alpha m_j(m_j - 1)/2,$$

where m_j is the number of nearest neighbor up-spins of spin σ_j , α sets the time scale and the up-flip rate $w_{j,\text{up}}(\sigma_j)$ is chosen to satisfy the detailed balance condition. Thus the spin flip rate is zero unless σ_j has at least two neighboring up-spins which can facilitate the flipping of σ_j . Fredrickson and Andersen [4] presented arguments, but could not prove, that the Markov chain corresponding to the master equation dynamics is irreducible [3]. If the Markov process is irreducible this model has the same equilibrium properties as the Ising model so that any phase transition in a nonzero magnetic field must be of dynamical origin [1].

Fredrickson and Anderson [1, 4] developed a diagrammatic perturbation theory for this model and by resuming diagrams of leading order in the small concentration c of up-spins, they derived a self-consistent equation for the equilibrium single-spin autocorrelation function $C(t) = [\langle \sigma_i(t)\sigma_i(0) \rangle - \langle \sigma_i \rangle^2] / [1 - \langle \sigma_i \rangle^2]$. In particular they predict that by lowering the concentration c of up-spins the relaxation time τ increases and diverges

at a critical value c^* below which spin fluctuations are frozen as is manifested in $C(t)$ decaying to a nonzero value f ($f^* < f < 1$ for $c^* > c > 0$) in the infinite time limit.

In order to test these theoretical predictions we have performed Monte Carlo experiments [5] for the two-spin facilitated kinetic Ising model with $N = 16^3$ spins on a cubic lattice. To test the size dependence runs were made also for $N = 32^3$, but no significant changes were observed. We chose $J = 0$ since for non-interacting spins equilibrium configurations at a given temperature T and magnetic field H with a concentration of up-spins $c = [1 + \exp(2H/k_B T)]^{-1}$ can be established efficiently. This means that the initial configuration of up- and down spins is random with given concentration c . Furthermore some approximations of the theory [4] become exact in this limit.

The Monte Carlo runs were performed up to 10^4

Fig. 1. Relaxation time τ and exponent β versus fraction of up-spins c determined by Monte Carlo simulation compared with theory [1, 4] (full line).

* Permanent address: Physics Department, University of Antwerp, B-2610 Wilrijk, Belgium.

Monte Carlo steps (MCS). We find that $C(t)$ can reasonably well be approximated by a Kohlrausch law [6] $\Phi(t) = \exp(-(t/\tau)^\beta)$ in the time regime $6 < t < 60$ (α is chosen so that 100 MCS correspond to 6 time units of [1, 4]) with $\beta = \beta(c)$ as shown in Fig. 1. The fitted τ is expected to show the same qualitative behavior as the one defined by the time integral of $C(t)$. In qualitative agreement with the theory we find that τ increases when the temperature is lowered, as is shown in Fig. 1. However, the relaxation time does not diverge at the predicted value $c^* = 0.0904$, where $f^* = 2/3$. It is interesting to note that an improvement of the theory [4] yields a lower value $c^* = 0.0681$, where $f^* = 0.71$. Our results, however, show no indication of a divergency at this concentration either. In order to locate the experimental value of c^* runs were performed for $c = 0.0474$ and $c = 0.0266$ for times up to $t = 6000$, but even in this time regime $C(t)$ did not settle to a constant nonzero value f .

Summarizing our results we find that the relaxation time increases with decreasing c but we do not find a

diverging τ in the regime predicted by theory [1, 4], a divergency at lower values of c cannot be ruled out, however. If a nonzero value of c^* exists, it is at least three times smaller than predicted by the theory [1, 4].

Acknowledgements — We are pleased to thank Fredrickson and Andersen for sending us preprints prior to publication. One of us (H.D.R.) thanks the National Fund of Scientific Research, Belgium, for financial support.

REFERENCES

1. G.H. Fredrickson & H.C. Andersen, *Phys. Rev. Lett.* **53**, 1244 (1984).
2. R.J. Glauber, *J. Math. Phys.* (N.Y.) **4**, 294 (1963).
3. J.M. Hammersley & D.C. Handscomb, *Monte Carlo Methods*, Methuen, London, (1964).
4. G.H. Fredrickson & H.C. Andersen, preprints.
5. K. Binder & D. Stauffer, *Applications of the Monte Carlo Method in Statistical Physics*, (Edited by K. Binder) Springer, Berlin, (1984).
6. R. Kohlrausch, *Ann. Phys.* (Leipzig), **12**, 393 (1847).