University of Groningen

Cloning, characterization and disruption of a (p)ppGpp synthetase gene (relA) of *Streptomyces coelicolor* A3(2)

Chakraborty, Rekha; White, Janet; Takano, Eriko; Bibb, Mervyn

Published in:
Molecular Microbiology

DOI:
[10.1046/j.1365-2958.1996.390919.x](https://doi.org/10.1046/j.1365-2958.1996.390919.x)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1996

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Chakraborty, R., White, J., Takano, E., & Bibb, M. (1996). Cloning, characterization and disruption of a (p)ppGpp synthetase gene (relA) of *Streptomyces coelicolor* A3(2). *Molecular Microbiology*, 19(2). <https://doi.org/10.1046/j.1365-2958.1996.390919.x>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Cloning, characterization and disruption of a (p)ppGpp synthetase gene (*relA*) of *Streptomyces coelicolor* A3(2)

Rekha Chakraborty, Janet White, Eriko Takano and Mervyn Bibb*

Department of Genetics, John Innes Centre, Norwich Research Park, Colney, Norwich NR4 7UH, UK.

Summary

An internal segment of the (p)ppGpp synthetase gene, *relA*, of *Streptomyces coelicolor* A3(2) was amplified from genomic DNA using the polymerase chain reaction and used as a hybridization probe to isolate the complete gene from a cosmid library. *relA* lies downstream of a gene (*apt*) that apparently encodes adenine phosphoribosyltransferase and is transcribed from two promoters, *relAp1* and *relAp2*, and by transcriptional readthrough from *apt*. While the level of *relAp2* transcripts remained relatively constant, *relAp1* activity apparently peaked during transition phase, following a decline in readthrough transcription from *apt*. Disruption of *relA* using an *att*[−] derivative of the temperate phage ϕ C31 abolished ppGpp synthesis on amino acid depletion. When grown on agar, the disruptants grew more slowly than a control lysogen made with an *att*⁺ phage vector and gave smaller colonies that sporulated normally. The *relA* mutation had no consistent or marked effect on actinorhodin production in either liquid- or agar-grown cultures, indicating that elevated levels of (p)ppGpp are not essential for triggering the onset of antibiotic production.

Introduction

Streptomycetes are Gram-positive mycelial soil bacteria that undergo a complex process of morphological differentiation leading to sporulation (Chater and Hopwood, 1993). They also produce a plethora of secondary metabolites, many with important applications as antibiotics in human medicine and in agriculture. Antibiotic production in streptomycetes is generally confined to stationary phase in liquid cultures, and usually coincides with the onset of morphological differentiation in agar-grown cultures. While little is understood of the mechanisms that regulate antibiotic production (Champness and Chater, 1994), most of the published data are consistent with a role for growth rate,

or the cessation of growth, in determining the onset of secondary metabolism (Chater and Bibb, 1996).

In *Escherichia coli*, the highly phosphorylated guanine nucleotides ppGpp and pppGpp have been implicated in growth-rate control of gene expression (Sarubbi *et al.*, 1988; Hernandez and Bremer, 1990; 1993; Schreiber *et al.*, 1991), and in regulating stationary-phase gene expression, mediating at least some of their effects through activation of the stationary-phase sigma factor, σ^S (Gentry *et al.*, 1993). Does (p)ppGpp play a role in growth-phase-dependent gene expression in streptomycetes and, more specifically, in triggering the onset of antibiotic production? Positive correlations were first observed between (p)ppGpp and antibiotic biosynthesis in *Streptomyces aureofaciens* (Simuth *et al.*, 1979) and *Streptomyces galilaeus* (Hamagishi *et al.*, 1980), producers of chlortetracycline and aclacinomycin, respectively. Later, Ochi (1986) observed an eightfold increase in formycin production by *Streptomyces lavendulae* after provoking (p)ppGpp accumulation by amino acid depletion. Moreover, a *relC* mutant of *S. lavendulae* defective in ribosomal protein L11, and consequently deficient in (p)ppGpp synthesis on amino acid starvation, was impaired in formycin production (Ochi 1986; 1990a). Similar *relC* mutants were isolated from *Streptomyces antibioticus* (Ochi, 1987), *Streptomyces griseoflavus* (Ochi, 1988), *Streptomyces griseus* (Ochi, 1990a), and *Streptomyces coelicolor* (Ochi, 1990b). The mutants accumulated only low levels of (p)ppGpp after nutritional shiftdown (on average c. 15% of wild type) and were deficient in antibiotic production, leading to the suggestion that (p)ppGpp plays a central role in triggering antibiotic biosynthesis (Ochi, 1990b). Consistent with this idea, reduced levels of actinomycin biosynthetic enzymes and mRNA were found in a *relC* mutant of *S. antibioticus* (Kelly *et al.*, 1991), and (p)ppGpp accumulation was noted to coincide with transcription of the pathway-specific activator gene, *brpA*, for bialaphos production by *Streptomyces hygroscopicus* (Holt *et al.*, 1992). However, Bascaran *et al.* (1991) concluded that (p)ppGpp did not play a role in antibiotic production in *Streptomyces clavuligerus*. Production of cephalosporins (mainly cephamycin C) occurred during a phase of slow exponential growth (doubling times of c. 7 h and 18 h in rich and minimal media, respectively), and increased in stationary phase, while ppGpp levels remained constant from the beginning of growth. *relC*-like mutants accumulated lower amounts of ppGpp than the wild-type strain after nutritional shiftdown

Received 26 July, 1995; revised 8 September, 1995; accepted 13 September, 1995. *For correspondence. E-mail bibb@bbsrc.ac.uk; Tel. (01603) 452571; Fax (01603) 456884.

(varying between 8% and 85% of the wild-type levels), but there was no simple correlation between the levels of ppGpp and cephalosporin production (some produced more, and others less, cephalosporin than the wild-type strain).

We have examined the relationship between (p)ppGpp synthesis and the production of undecylprodigiosin and actinorhodin in *S. coelicolor* (Strauch *et al.*, 1991; Takano *et al.*, 1992; Takano and Bibb, 1994). Transcription of pathway-specific activator genes for each antibiotic (*redD* for undecylprodigiosin and *actII-ORF4* for actinorhodin) conspicuously increased as the culture entered stationary phase, coinciding with a peak in the level of (p)ppGpp; this was followed by transcription of representative *red* and *act* biosynthetic structural genes, and production of the antibiotics. In contrast, transcription of the *rrnD* rRNA gene set decreased markedly (E. Takano and M. J. Bibb, unpublished). (p)ppGpp synthesis was also induced by amino acid depletion during rapid exponential growth, and was followed by an immediate and marked decrease in *rrnD* transcription (Strauch *et al.*, 1991). Transcription of *actII-ORF4* was detected within 30 min of the nutritional shift-down and transcription of the *act* biosynthetic genes after a further 30 min (Takano and Bibb, 1994).

Although correlations between (p)ppGpp synthesis and the onset of antibiotic production clearly exist, they do not establish a role for the highly phosphorylated nucleotides in triggering antibiotic biosynthesis. Although the *relC* mutants isolated by Ochi (1986; 1987; 1988; 1990a,b) show a marked reduction in antibiotic production, they grow at about half the maximal rate of the parent strain, presumably reflecting impaired protein synthesis. Thus it is difficult to assess whether the effect on antibiotic production is a direct consequence of reduced levels of (p)ppGpp, or an indirect effect of the L11 mutation on protein synthesis.

In *E. coli*, (p)ppGpp is synthesized predominantly by the ribosome-bound RelA, which is activated when an uncharged tRNA binds to the acceptor site of translating ribosomes (Cashel and Rudd, 1987). Therefore (p)ppGpp synthesis can be invoked by amino acid depletion or by addition of tRNA synthetase inhibitors, such as serine hydroxamate (SHX) (Tosa and Pizer, 1971). Interestingly, the enzyme responsible for (p)ppGpp degradation in *E. coli*, SpoT, is a homologue of RelA, the two proteins showing 31% amino acid sequence identity (Metzger *et al.*, 1989). Although the mechanism is not understood, starvation for carbon and energy reduces SpoT activity and results in elevated levels of (p)ppGpp (Cashel and Rudd, 1987). Furthermore, the residual level of (p)ppGpp synthetic activity observed in *relA* null mutants probably reflects a synthetic role for SpoT in addition to its degradative function (Hernandez and Bremer, 1991; Xiao, *et al.*, 1991), although this has not been demonstrated *in*

vitro. A *relA* homologue from *Vibrio* sp. strain S14 has been cloned, sequenced and disrupted (Flårdh *et al.*, 1994); the mutant exhibited relaxed control of RNA synthesis, and failed to accumulate (p)ppGpp upon addition of SHX. *relA*-dependent (p)ppGpp synthesis was also invoked by carbon-source depletion, which is not observed in *E. coli*; presumably glucose deprivation in the vibrio results in a rapid decrease in the level of aminoacylation of at least some tRNAs. *relA* homologues have also been identified in *Streptococcus equisimilis* H46A (Mechold *et al.*, 1993), *Mycobacterium leprae* (EMBL Accession Number U00011) and *Mycoplasma genitalium* (Peterson *et al.*, 1995); genetic and/or biochemical studies that might differentiate between synthetic and degradative roles for these genes have not been published. The isolation of *relC* mutants of *S. coelicolor* (Ochi, 1990b), and the ability of the wild-type strain to produce (p)ppGpp on amino acid depletion or on addition of SHX (Strauch *et al.*, 1991), suggests that *S. coelicolor* possesses a ribosome-dependent (p)ppGpp synthetase. In an attempt to establish if (p)ppGpp plays an essential role in triggering antibiotic production in *S. coelicolor*, we set out to isolate and disrupt its *relA* homologue.

Results

Isolation of a relA homologue from S. coelicolor using the polymerase chain reaction

Alignment of the amino acid sequences of RelA homologues from *E. coli* (RelA (Metzger *et al.*, 1988) and SpoT (Metzger *et al.*, 1989)), *M. leprae* (EMBL U00011), *S. equisimilis* H46A (Mechold *et al.*, 1993) and *Vibrio* sp. strain S14 (Flårdh *et al.*, 1994) revealed several blocks of conserved amino acids (Fig. 1). Those corresponding to amino acid residues 317–324 and 426–433 of *E. coli* RelA were chosen for the design of 16-fold and 64-fold degenerate oligonucleotides for use as primers in the polymerase chain reaction (PCR), using the preferred codons from 64 streptomycete genes (Wright and Bibb, 1992); *Bam*HI sites were incorporated at the 5' end of each primer to facilitate subsequent cloning. An amplified product of approximately 400 bp was obtained, consistent with the size expected for a *relA* homologue. The PCR product was digested with *Bam*HI and cloned in *Bam*HI-cleaved pBluescript SK+ in *E. coli* DH5 α , yielding pJ6034. Sequencing from both ends of the pJ6034 insert using universal and reverse primers revealed the expected internal segment of a *relA* homologue.

The pJ6034 insert was isolated as a *Bam*HI fragment, labelled with ³²P by random oligonucleotide priming, and used as a probe to isolate six hybridizing cosmids from a library of *S. coelicolor* M145 DNA. Digestion of each of the six cosmid DNAs with *Bgl*II, *Pvu*II, *Sac*I and

EcoRI + *Bam*HI revealed common restriction fragments, and Southern analysis using the same probe identified hybridizing *Pst*I fragments of approximately 4 kb and 6 kb. Cloning of these gel-purified *Pst*I fragments in *Pst*I-cleaved pBluescript SK+ yielded pJ6031 (4 kb insert) and pJ6037 (6 kb insert). A restriction map of the contiguous 10 kb region is shown in Fig. 2. Hybridization of the probe to cosmids L2 and L7 (H. M. Kieser, personal communication) localized the *relA* homologue on *Asel* fragment L and, more precisely, to a position between *bldD* and *argA* in the 12 o'clock region of the combined physical and genetic map of the *S. coelicolor* chromosome (Kieser *et al.*, 1992).

RelA of *S. coelicolor* has an N-terminal extension that is absent from all known homologues

The sequence of a 4515-nucleotide (nt) region that extends 2058 nt upstream and 2457 nt downstream of the *Pst*I site within *relA* was determined (EMBL X87267). Analysis of the sequence using the FRAME program (Bibb *et al.*, 1983) revealed three open reading frames (ORFs) with protein-coding character (Fig. 2). The largest of these (*relA*; nt positions 796–3336) would encode a protein of 847 amino acids (94.2 kDa) with a high degree of similarity to all of the known RelA homologues (Fig. 1). *S. coelicolor* RelA is most similar to the homologue from *M. leprae* (66% amino acid sequence identity), and least similar to RelA of *Vibrio* sp. strain 14 (38% identity), and has an N-terminal extension of approximately 90 amino acids that is absent from all of the other homologues. Unusually, translation of the *relA* transcript apparently initiates at a UUG codon (Fig. 3). Five nucleotides upstream of the UUG codon there is a sequence (AGGAG) that is complementary to a region close to the 3' end of the 16S rRNA of *S. coelicolor* (Baylis and Bibb, 1987) that is likely to serve as a ribosome-binding site.

A second ORF of 546 nt (nt positions 65 to 610) occurs upstream of, and reading in the same direction as, *relA* and is preceded 7 nt upstream by a potential ribosome-binding site (GGA). The ORF would encode a protein of 182 amino acids (18.6 kDa) with a high degree of identity to adenine phosphoribosyltransferases (APTs) from a variety of organisms. The most similar homologue found in searches of the databases was the APT of *E. coli* (51% amino acid sequence identity; Hershey and Taylor, 1986). APTs catalyse the conversion of adenine and 5-phosphoribosyl pyrophosphate to AMP in a salvage reaction that is energetically less costly than *de novo* synthesis.

The third ORF of >1095 nt is located 3' of *relA* and is oriented in the opposite direction (nt positions 3421 to 4515; Fig. 2). It terminates 86 nt from the end of the *relA* coding region; its N-terminus appears to occur outside of the sequenced region. Searches of the databases

revealed similarity (46% amino acid sequence identity in an overlap of 175 amino acids) to an ORF of unknown function from *M. leprae* (EMBL U00019) that is not located adjacent to the mycobacterial *relA* homologue. A 22 nt inverted repeat occurs in the non-coding region between *relA* and the convergent ORF (nt positions 3358–3404), and may function as a factor-independent bidirectional transcriptional terminator (reviewed in Lewin, 1994).

Transcription of relA occurs from multiple promoters in a growth-phase-dependent manner

To map the transcriptional start site of *relA*, S1 nuclease protection experiments were carried out using RNA from *S. coelicolor* strain M145 at different stages of growth. A 458 bp *Stu*I–*Sal*I fragment (nt positions 474–931) that includes the non-coding region between *apt* and *relA* (Fig. 2) was cloned in pBluescript SK+ that had been cleaved with *Sma*I and *Sal*I, yielding pJ6068. The 498 bp *Sac*I–*Sal*I fragment of pJ6068, uniquely labelled at the 5' end of the *Sal*I site (which lies 136 nt downstream of the UUG start codon) was used as probe; the presence of 40 nt of polylinker at the unlabelled end of the probe served to distinguish between probe reannealing and transcriptional readthrough from *apt*. Putative transcriptional start sites were identified at nt positions 682 and 751–752 (*relAp*1 and *relAp*2, respectively, in Figs 3 and 4b); a protected fragment corresponding in size to the *Stu*I–*Sal*I segment of the probe (Fig. 4b; FLP) indicated additional transcription of *relA* from a promoter upstream of, or within, *apt*. While the level of transcripts initiated at *relAp*2 remained constant, transcription from *relAp*1 appeared to peak during transition phase, following a marked decrease in the level of transcriptional readthrough from *apt* (Fig. 4b). An inverted repeat of nine nucleotides occurs between the *apt* and *relA* coding regions (nt positions 716–738), and overlaps the –35 region of *relAp*2 (Fig. 3).

Since the changes in the levels of the *relAp*1 and readthrough transcripts coincided with elevated levels of ppGpp during transition phase, the effect of nutritional shiftdown on *relA* transcription was assessed (Fig. 5). While there was a clear decrease in the level of the readthrough transcript 30 min after shiftdown, there was a marked increase in the amount of *relAp*1 transcript.

To confirm that the RNA-protected fragments assigned to *relAp*1 and *relAp*2 represented transcriptional start sites, rather than artefacts of S1 nuclease digestion, the 498 bp *Sac*I–*Sal*I and 466 bp *Bam*HI–*Sal*I fragments of pJ6068 were used as templates for *in vitro* transcription assays using RNA polymerase isolated from *S. coelicolor* strain M145. Both fragments gave run-off transcripts of approximately 250 nt and 180 nt (Fig. 6) that corresponded to the expected sizes predicted from the S1

Ecrela							0				
Vsrela							0				
Mrel							0				
Screla	MPDEAQPLTA	AKPESASASA	AKPAPSAPQA	KNDTHGPIQH	APAAPVDKPA	EQQPRPKPLP	60				
Ecspot							0				
Serel							0				
Ecrela				..MVAVR	SAH	INKAGFDPE	KWIASLGLITS	28			
Vsrela				..MVAVR	GAH	LKENTTFELV	SWVESLRLQDA	28			
Mrel				..MTAQR	S.T	ISPVLEPLVA	VHKEFYDPKAN	27			
Screla	AERPQNAPVV	RAPAGQPARS	GSSNRVRARL	ARLGVQEFANP		YNPVLEPLLR	IVRGNDPKIE	120			
Ecspot						..MYLFFESLNO	LIQTYLPEDQ	19			
Serel					MAKE	INLTGSEVVA	LAAKYMNETD	24			
Ecrela	KKSCECLAET	WAYC..LQQT	QGHPPDASLL	WRGVEMVEIL	STLSMDADTL	RANILFPLAD	86				
Vsrela	KVSSRIEGT	YQRC..IELA	AEQENSGPL	WRGRELVEIL	VTLSMDADTL	IAADLYPLV	85				
Mrel	LSI..VQRA	FEVAQDRHAS	QLKRSGDPYI	THPLAVANIL	AEELGMDITL	VAADLHDTVE	84				
Screla	TSTLRQIERA	YQVAERDAHHRG	QKRKSGDPYI	THPLAVANTIL	AEELGMDITL	MAGLHDTVE	180				
Ecspot	IKR...LRQA	YLVARDAHEG	QTRSSGDPYI	THPLAVACIL	AEEMKLDYET	MEALHDTVE	76				
Serel	AAF...VKK	LDYATAAHFY	QVHRKSGEPI	VHPIQVAGIL	ADHLDAVTV	ACGEIHDVTE	81				
Ecrela	ANVVSSEDVLR	ESVGRKSVN	LHGV	RDMAAI	RQLKATHTDS	VSSSEQVDN	VF	146			
Vsrela	GGCYSTDAL	EEYSSTIL	LHL	VQGV	EQMC	AAQAAQVDN	VF	145			
Mrel	DTGYTLEALS	EEFGDEVGH	L	VDGV	TFLDRV	..AAAEGETI	VF	136			
Screla	DTGYTLEALS	RDFFCDVVT	L	VDGV	TFLDRV	..AAAEGETI	VF	232			
Ecspot	DIPATYQDME	QLFGKSV	VAEL	VEGV	SKLDKL	KKEAAQAE	VF	130			
Serel	DITITLDNIE	FDFFGKD	VRDI	VDGV	TFLGKV	E...YKS	HEEQLAENHE	135			
Ecrela	RCVVVVKLAER	IAHLFEVKDA	PEDERVLAAK	ECTNIYAPLA	NFLGIGQLKH	ELEDYCFRY	206				
Vsrela	RCVVVVKLAER	ICNLFVKDQ	PDEVRRAAQ	ECANIIYAPLA	NFLGIGQLKH	ELEDYCFRY	205				
Mrel	RVLDVVKLAER	LHNHETMR	YI	ETLEVIAPLA	HFLGMASVK	ELEDYCFRY	196				
Screla	RVLDVVKLAER	LHNHETMR	YI	ETLEVIAPLA	HFLGMASVK	ELEDYCFRY	292				
Ecspot	RVLDVVKLAER	THNHE	TGLS	ETLEVIAPLA	HFLGIIHINT	ELEDYCFRY	190				
Serel	RVLDVVKLAER	LHNHE	TGLS	ETLEVIAPLA	HFLGIIHINT	ELEDYCFRY	195				
Ecrela	HTEVKRKIAK	LLHEFFLDRE	HYIEEFVGH	L	RAEMKAEGVK	AEVYGFPEKH	I	266			
Vsrela	HPDKTKQIAK	QLSEFPIDRE	DHITHFVDD	L	SDAMKASNIR	AEVYGFPEKH	I	265			
Mrel	HPDKTKQIAK	LVAGAPSRD	THLAKVR	AEI	ISDLGASKIK	AEVYGFPEKH	I	256			
Screla	YPKMYDEIVR	LVAFAPKRD	EMIAQVTD	DEV	QQDLRAARIK	AEVYGFPEKH	I	352			
Ecspot	YBNRFRVKE	VVKAFFGRNK	EMIQVILSEI	EGRLQEAAG	TP	AEVYGFPEKH	I	250			
Serel	NETEFYKISH	MMNFKERERE	ALVDDIVTKI	KSYTTEQGLF	G	AEVYGFPEKH	I	255			
Ecrela	NLAFFDELFDV	RAVRIRIVAE	RL	QDCYAAALG	IV	HTHYRHL	PDE	326			
Vsrela	NLAFFDELFDV	RAVRIRIVAE	EEL	QDCYAAALG	IV	HTHYRHL	PDE	325			
Mrel	GRDFDTHIDL	VGIRILCDEI	E	RDCYAAALG	IV	HSLLQPM	AGR	316			
Screla	GRDFDTHIDL	VGIRILCDEI	E	RDCYAAALG	IV	HSLLQPM	AGR	412			
Ecspot	EQRFHSDIMDI	YAFRVIVNDS	D	TCYRV	LGQM	HSLLQPM	AGR	310			
Serel	KKRFFDQITFDL	IATRCVME	ETQ	SDVYAMV	GYI	HELRW	PMPGR	315			
Ecrela	LGPFGGKTV	EI	QIRTKQM	HED	AEELGVAAAH	K	YKEGAAAGGA	373			
Vsrela	LGPFGGKTV	EI	QIRTKQM	HED	AEELGVAAAH	K	YKEGAAAGGA	372			
Mrel	VGPFGGKTV	EI	QIRTKQM	HED	AEELGVAAAH	K	YKEGAAAGGA	367			
Screla	IGPFGGKTV	EI	QIRTKQM	HED	AEELGVAAAH	K	YKEGAAAGGA	472			
Ecspot	IGPFGGKTV	EI	QIRTKQM	HED	AEELGVAAAH	K	YKEGAAAGGA	355			
Serel	YGPFGGKTV	EI	QIRTKQM	HED	AEELGVAAAH	K	YKEGAAAGGA	361			
Ecrela	IAHLRKLIAW	QBEEMADSG	EB	MLDEV	RSQV	VE	DDR	VYVFTPK	432		
Vsrela	IAHLRKLIAW	QBEEMADSG	EB	MLDEV	RSQV	VE	DDR	VYVFTPK	431		
Mrel	MAHMRQLLDH	QBEEMADSG	EB	MLDEV	RSQV	VE	DDR	VYVFTPK	426		
Screla	MAHMRQLLDH	QBEEMADSG	EB	MLDEV	RSQV	VE	DDR	VYVFTPK	531		
Ecspot	QRWQMSLLEL	QBEEMADSG	EB	MLDEV	RSQV	VE	DDR	VYVFTPK	414		
Serel	MNRIKELVLEL	QBEEMADSG	EB	MLDEV	RSQV	VE	DDR	VYVFTPK	421		
Ecrela	SDVGHRCIGA	KIGGRIVP	PFT	YQLQMG	GDQV	IE	ITITQKQPN	..	490		
Vsrela	SEVGHRCIGA	KIGGRIVP	PFT	YQLQMG	GDQV	IE	ITITQKQPN	..	489		
Mrel	TEVGHRCIGA	RVNGFLV	PALE	YQLQMG	GDQV	IE	ITITQKQPN	..	482		
Screla	TEVGHRCIGA	RVNGFLV	PALE	YQLQMG	GDQV	IE	ITITQKQPN	..	587		
Ecspot	TDIGHACVGA	RVDROPY	PPLS	QPLTSG	QOTVE	ITITQKQPN	..	468			
Serel	TOVGEKALGA	RVNGFLV	PALE	YQLQMG	GDQV	IE	ITITQKQPN	..	476		
Ecrela	KIHAHFRKQD	RDKNILAG	RQ	ILDELE	HLG	ISLK	..	..EAEKHL	LP	542	
Vsrela	KIHAHFRKQD	RDKNILAG	RQ	ILDELE	HLG	ISLK	..	..EAEKHL	LP	541	
Mrel	KIHAHFRKQD	RDKNILAG	RQ	ILDELE	HLG	ISLK	..	..EAEKHL	LP	539	
Screla	KIHAHFRKQD	RDKNILAG	RQ	ILDELE	HLG	ISLK	..	..EAEKHL	LP	644	
Ecspot	KIHAHFRKQD	RDKNILAG	RQ	ILDELE	HLG	ISLK	..	..EAEKHL	LP	523	
Serel	KIHAHFRKQD	RDKNILAG	RQ	ILDELE	HLG	ISLK	..	..EAEKHL	LP	533	
Ecrela	LAATGGGDIR	LNQMVN	FL	..	..QSQFNKPSA	EEQDAAAL	KQ	LQOKSYTPQN	RSKDNGR	..V	597
Vsrela	LAATGGGDIR	LNQMVN	FL	..	..QSQFNKPSA	EEQDAAAL	KQ	LQOKSYTPQN	RSKDNGR	..V	598
Mrel	LAATGGGDIR	LNQMVN	FL	..	..QSQFNKPSA	EEQDAAAL	KQ	LQOKSYTPQN	RSKDNGR	..V	592
Screla	LAATGGGDIR	LNQMVN	FL	..	..QSQFNKPSA	EEQDAAAL	KQ	LQOKSYTPQN	RSKDNGR	..V	698
Ecspot	LAATGGGDIR	LNQMVN	FL	..	..QSQFNKPSA	EEQDAAAL	KQ	LQOKSYTPQN	RSKDNGR	..V	558
Serel	LAATGGGDIR	LNQMVN	FL	..	..QSQFNKPSA	EEQDAAAL	KQ	LQOKSYTPQN	RSKDNGR	..V	593

Fig. 1. Alignment of the amino acid sequence of *S. coelicolor* RelA (Screla) with *E. coli* RelA (Ecrela), *E. coli* SpoT (Ecspot), *Vibrio* sp. strain 14 RelA (Vsrela), and RelA homologues from *M. leprae* (Mrel) and *S. equisimilis* (Sere). The sequences were aligned and displayed using the PILEUP and PRETTYBOX programs, respectively, of the UWGCG package (Devereux *et al.*, 1984). The locations of the conserved amino acid sequences that were used to design the PCR primers are indicated by arrows under the alignment. Black boxes indicate amino acids present in at least four of the six sequences, and shaded boxes represent conservative substitutions at these positions.

nuclease protection data for *relAp1* and *relAp2*, respectively.

Disruption of relA results in abolition of ppGpp synthesis on amino acid depletion, but has no marked effect on actinorhodin production

To assess the role of *relA* in (p)ppGpp synthesis and antibiotic production in *S. coelicolor*, the gene was disrupted using an *att⁻* derivative of the temperate *Streptomyces* phage ϕ C31 (Bruton and Chater, 1983). An internal 934 bp segment of the *relA* coding region (Fig. 2) that extended from a *Bam*HI site (nt position 1290) located 494 nt downstream of the translational start codon to a *Pvu*II site (nt position 2227) located 1108 nt upstream of the translation stop codon was cloned in the *E. coli* vector pMT3000 (Paget *et al.*, 1994) that had been cleaved with *Bam*HI and *Sma*I, yielding pIJ6070. The internal *relA* fragment was excised from pIJ6070 as a 942 bp *Bam*HI–*Sst*I fragment, and ligated with the *att⁻* *c⁺* ϕ C31 phage vector KC516 (Rodicio *et al.*, 1985) that had been cleaved with *Bgl*II and *Sst*I; KC516 carries the *tsr* gene that confers resistance to the antibiotic

thiostrepton (Thio). The ligation mixture was used to transfect protoplasts of *S. lividans* strain 1326, and likely recombinant clones were identified by plaque hybridization using the 934 bp *Bam*HI-*relA*-*Pvu*II fragment labelled with 32 P by random oligonucleotide priming as a probe. Restriction analysis of phage DNA derived from eight positive plaques revealed that six contained the required construct, KC936. After single-plaque purification, high-titre phage stocks of KC936, and of the *att*⁺ *c*⁺ phage vector KC301 (Hopwood *et al.*, 1985), were prepared from *S. lividans* 1326 top layers and spotted onto lawns of *S. coelicolor* strain J1929, a Pgl⁻ derivative of *S. coelicolor* M145 (Bedford *et al.*, 1995). After sporulation, the plates were replicated to the rich medium R5 supplemented with 5 μ g ml⁻¹ Thio to obtain lysogens. After two rounds of single-colony purification on the same selective medium, seven independent Thio^R isolates derived from KC936 and one from the control phage KC301 were replicated to lawns of *S. lividans* 1326 spores and shown to release phages. Southern analysis of *Bam*HI-digested total DNA isolated from each of the KC936 lysogens using the purified *Bam*HI-*Pvu*II fragment as a 32 P-labelled probe confirmed disruption of *relA* (Fig. 7). The 3.2 kb *Bam*HI fragment in J1929 was

Fig. 2. Restriction map of the *apt-relA* region of *S. coelicolor*. The positions of *apt*, *relA* and the convergent downstream ORF are indicated by arrows. The locations of the PCR product (PCR), the internal fragment used to disrupt *relA* (KC936), the end-labelled fragment used in the S1 nuclease protection experiments (S1), and the templates used in the *in vitro* transcription assays (IVT) are shown above the restriction map; the black circle indicates the labelled end of the probe used for S1 nuclease mapping and the thickened line the extent of the sequenced region.

F L G G R A R L E P A L A G A P L E A L
 CTTCTCGGCGGGCGGCGGCTGGAGCCGGCGCTGGCCGGGGCACCGCTGGAGGCACT 600
 L T V *
 GCTCACGGTCTGAACACGCGAACCGCACGGCTGGGGCGGGCCGGAGGAATCCGGCGCC 660
 CGCCCAGGCGTTGCAGGGGTTGACGGCCGTACAGGCCCTCACATTGGCCAGAAGGCGAT 720
 CCTGGAGGGCAGGATCGCTACCATGGGGTATCCGGAGCCTGACCGGGGGACCCGGATCGC 780
 M P D E A Q P L T A A K P E S
 GCACGAGGAGTCTCTTGCCAGACGAGGCCAGCCACTGACCGCCGCAAGCCCGAGTCC 840

replaced by fragments of 5.1 kb and >40 kb, as predicted from the restriction maps of the *relA* region and of KC516. Southern analysis of the same DNA samples digested with *Pst*I again gave the pattern expected from disruption of *relA*.

All seven independent J1929::KC936 *relA* disruptants were streaked out for single colonies on R5 and minimal medium, each supplemented with $5 \mu\text{g ml}^{-1}$ Thio, together with the control lysogen J1929::KC301. All of the disruptant colonies grew more slowly than those containing KC301. All of the J1929::KC936 derivatives appeared to sporulate normally and produced the purple-pigmented antibiotic actinorhodin, but their mature colonies were significantly smaller than those of J1929::KC301. The difference in growth rate and in the size of mature colonies was also observed on R2, SFM and SMMS agar media, both in the presence and absence of $5 \mu\text{g ml}^{-1}$ Thio.

To assess the effect of disrupting *relA* on (p)ppGpp synthesis, J1929::KC936 and J1929::KC301 were grown

in SMM liquid medium containing $5 \mu\text{g ml}^{-1}$ Thio until mid-exponential phase ($\text{OD}_{450}=0.5$) and subjected to amino acid depletion by rapidly transferring the cultures to SMM lacking amino acids. Nucleotides were extracted with 1 M formic acid 0, 7.5 and 15 min after nutritional shift-down and analysed by high-performance liquid chromatography (HPLC) (Fig. 8). While the levels of ppGpp in J1929::KC301 7.5 min after amino acid depletion (542 and 292 pmol mg^{-1} dry weight in duplicate experiments) were comparable to earlier estimates for *S. coelicolor* strain M145 (200 pmol mg^{-1} dry weight; Strauch *et al.*, 1991), there was no increase in the level of ppGpp in the *relA* disruptant J1929::KC936 (7.3 and 7.8 pmol mg^{-1} dry weight in cultures with pre-shift-down ppGpp levels of 8.3 and 10.6 pmol mg^{-1} dry weight, respectively). While the level of GTP in the *relA*⁺ J1929::KC301 dropped markedly on shiftdown, there was no reduction in J1929::KC936 (Fig. 8).

Growth of *S. coelicolor* in SMM is limited by the availability

Fig. 4. A. Growth curve of *S. coelicolor* M145 with arrows showing the times at which RNA was isolated for S1 nuclease mapping. B. S1 nuclease mapping of the transcription start sites of *relA*. Asterisks indicate the probable start points of transcription; the sequences given are those of the template strand. Lanes A>C, G+A and G are sequence ladders derived from the probe using the method of Maxam and Gilbert (1980). Probe, the end-labelled fragment used as a probe; tRNA, the results obtained using 50 μg of *E. coli* tRNA; FLP, full-length protection of probe sequences homologous to the *relA* region.

Fig. 5. A. Growth curve of *S. coelicolor* M145 after nutritional shiftdown with arrows showing the times at which RNA was isolated for S1 nuclease mapping.

B. The effect of nutritional shiftdown on *relA* transcription. See the legend to Fig. 4 for abbreviations.

of nitrogen (E. Takano and M. J. Bibb, unpublished), and (p)ppGpp synthesis occurs in *S. coelicolor* strain M145 during the transition between exponential growth and stationary phase (Strauch *et al.*, 1991). To assess the effect of the *relA* mutation on ppGpp synthesis during this transition period, ppGpp levels were monitored by HPLC. While the level of ppGpp peaked at 14.3 pmol mg⁻¹ dry weight in J1929::KC301 (similar to the 16 pmol mg⁻¹ dry weight observed by Strauch *et al.*, 1991), it remained at a constant level in J1929::KC936.

Disruption of *relA* appeared to have no deleterious effect on actinorhodin production in liquid culture, either after nutritional shiftdown or upon entry into stationary

phase. On the contrary, in both cases actinorhodin production by J1929::KC936 preceded that by J1929::KC301 by several hours. Surprisingly, in contrast to the agar-grown cultures, J1929::KC936 did not show a reduction in growth rate in liquid SMM when compared with J1929::KC301. With agar-grown cultures, although actinorhodin production in the mutant was delayed by a few hours on SMMS, there was no perceptible difference on minimal medium, R2 and R5, and on SFM it occurred several hours earlier than in the *relA*⁺ strain. The same results were obtained both in the presence and absence of Thio.

Discussion

Earlier attempts to isolate a (p)ppGpp synthetase gene from *S. coelicolor* using cloned *relA* homologues from *E. coli* (54 mol% G + C), *Serratia marcescens* (59%) and *M. leprae* (59%) as hybridization probes failed (Takano, 1993; R. Chakraborty, unpublished), presumably reflecting the considerable differences in base composition between genera (the GC content of the *S. coelicolor* *relA* homologue is 70.4%). In contrast, the PCR-based strategy reported here readily allowed for the isolation of a *relA* homologue from *S. coelicolor*; comparison of the degenerate primer sequences with the sequence of *relA* revealed 20/23 identities for the 5' primer and 23/23 for the 3' primer. Indeed, the same primers and amplification conditions gave PCR products of a similar size from *Mycobacterium smegmatis* and *Myxococcus xanthus* genomic DNAs (with GC contents deduced from the nucleotide sequence databases of 65% and 68%, respectively; R. Chakraborty, unpublished), suggesting that they may be useful for the isolation of *relA* homologues from other bacterial species with a relatively high genomic GC content.

Although the PCR product might have represented a

Fig. 6. *In vitro* transcription from *relAp1* and *relAp2*. The *relA* restriction fragments used as templates are shown at the top of the Figure, and the positions of run-off and end-to-end (ETE) transcripts are indicated on the right; *ctc*, a control *in vitro* transcription assay using a *ctc* promoter fragment from *Bacillus subtilis* that gives a run-off transcript of 155 nt (Buttner *et al.*, 1988). SM, size markers derived from the ³²P end-labelled 100 bp ladder and the *ctc* run-off transcript.

Fig. 7. Southern analysis of *relA* disruptants of *S. coelicolor* J1929. One microgram of *Bam*HI-digested genomic DNA from seven independent J1929::KC936 isolates, and from the parental strain J1929 was probed with the purified *Bam*HI-*relA*-*Pvu*II fragment that had been labelled with 32 P by random oligonucleotide priming. SM, size markers derived from phage λ DNA that had been digested with *Pst*I (the filter was also probed with phage λ DNA that had been labelled by random oligonucleotide priming).

functional homologue of *spoT* involved in (p)ppGpp degradation rather than synthesis, disruption using the *att* $^{-}$ ϕ C31 derivative KC516 established that the *S. coelicolor* gene encodes a (p)ppGpp synthetase. ppGpp synthesis invoked by amino acid depletion in mid-exponential phase, and during the transition between exponential growth and stationary phase under conditions of nitrogen limitation, was abolished in the *relA* mutant. While ATP levels after shutdown were similar in both J1929::KC936

Fig. 8. HPLC analysis of ppGpp and NTP levels after amino acid depletion of J1929::KC301 (*relA* $^{+}$) and J1929::KC936 (*relA* $^{-}$). The results are the average values from two independent experiments. NTP levels were standardized to those observed immediately before amino acid depletion in each culture (100%); ppGpp levels were standardized to the levels observed 7.5 min after shutdown in the J1929::KC301 cultures. Nucleotide concentrations (in $\mu\text{mol mg}^{-1}$ dry weight) immediately before shutdown were: J1929::KC301 - ATP, 5668 and 5495; CTP, 490 and 647; UTP, 696 and 969; GTP, 1063 and 2056; ppGpp, >1 and 1.7. J1929::KC936 - ATP, 6347 and 4641; CTP, 520 and 752; UTP, 1468 and 1685; GTP, 1800 and 1022; ppGpp, 8.3 and 10.6.

and J1929::KC301 (perhaps reflecting the higher intracellular concentrations and wide-ranging metabolic role of ATP), the concentrations of CTP and UTP were higher in J1929::KC301 than in the mutant, presumably indicating stringent control of stable RNA synthesis in the *relA* $^{+}$ strain. While GTP levels in the *relA* mutant after shutdown were similar to those of the other NTPs, they fell markedly in the *relA* $^{+}$ strain (as previously noted by Strauch *et al.*, 1991), presumably reflecting, at least in part, ppGpp synthesis.

Since the level of ppGpp observed in the *relA* mutant (on average 1.7% of that in the *relA* $^{+}$ strain after shutdown and close to the limit of detection of the HPLC assay) did not increase upon amino acid depletion, it seems unlikely that it reflects reconstitution of a functional *relA* gene by excision of KC936 (Southern analysis failed to reveal any prophage excision in cultures grown in the presence of Thio; Fig. 7). It may reflect *relA*-independent ppGpp synthesis mediated either by a functional homologue of *spoT* or by a homologue of a novel (p)ppGpp synthetase activity recently identified in *S. antibioticus* (Jones, 1994a,b; G. H. Jones, personal communication). Alternatively, the N-terminal segment of RelA that remains after disruption may possess synthetic activity that does not require ribosome activation; consequently, the level of ppGpp would not be expected to increase upon amino acid depletion, consistent with the experimental observations.

The results presented here suggest that elevated levels of (p)ppGpp are not essential for triggering actinorhodin production in *S. coelicolor*. However, since differences in the timing of actinorhodin synthesis were observed between J1929::KC936 and J1929::KC301 with some media, it is possible that variations in the level of ppGpp can influence the onset of actinorhodin production. Moreover, since the *relA* mutant still makes ppGpp, we cannot exclude the possibility that the phosphorylated nucleotide is required for transcription of antibiotic biosynthetic genes. In this context, it may be pertinent to note the effect of disrupting *relA* of *S. coelicolor* J1501 with KC936 (data not shown). J1501 (*hisA1 uraA1 strA1 Pgl* $^{-}$ *SCP1* $^{-}$ *SCP2* $^{-}$) is a commonly used strain, but for reasons that are not understood it produces much less actinorhodin and undecylprodigiosin than prototrophic plasmid-free strains. It also differs from the wild-type strain in the regulation of carbon-source utilization (S. Angell, personal communication) and its genome has undergone several large deletions and duplications (Kieser *et al.*, 1992). Disruption of *relA* with KC936 reproducibly gave J1501::KC936 lysogens that were either unaffected in antibiotic production or that produced undecylprodigiosin but not actinorhodin; the two phenotypes occurred in approximately equal numbers. While the basis for this heterogeneity and for the difference from J1929 is not clear, it is possible that the metabolism of ppGpp in J1501 differs from the wild-type

strain, and that the deficiency in actinorhodin production reflects either lower basal levels or the complete absence of ppGpp in J1501::KC936 (disruptants unaffected in actinorhodin production might have undergone secondary mutations that compensate for the loss of (p)ppGpp synthesis). The isolation of further mutants of J1929 that are completely devoid of (p)ppGpp should determine whether the highly phosphorylated nucleotides play an obligate role in actinorhodin production in the wild-type strain. If they do not, then the frequently observed correlation between ppGpp synthesis and the onset of antibiotic production might simply reflect the coincidental growth-phase-dependent initiation of antibiotic synthesis and the role of ppGpp in the growth-rate control of primary metabolic genes.

Of the six fully sequenced *relA* homologues, only for *spoT* of *E. coli* is there convincing evidence for ppGpp degradative activity. Comparison of all six amino acid sequences (Fig. 1) reveals several conserved regions, but fails to identify motifs that might be used to discriminate readily between enzymes with synthetic or degradative activities, and there seems to be little conservation of gene organization around the *relA* homologues studied so far that might give a clue to function. The *apt* and *relA* genes of *S. coelicolor* are preceded by ORFs (EMBL X85969) that are homologous to *secDF* of *E. coli*; SecD and SecF appear to act to improve the efficiency of protein export, but are not absolutely required (Pogliano and Beckwith, 1994). Both *relA* and *spoT* of *E. coli* are contained within polycistronic transcription units, and neither maps close to *apt* or *secDF* (Rudd and Cashel, 1987); the *relA* homologue of *Vibrio* sp. strain S14 is preceded by a homologue of the gene found immediately upstream of *E. coli relA* (Flårdh *et al.*, 1994). The *relA* homologue of *M. leprae* is preceded by an ORF reading in the same direction (EMBL U00011), and is followed by a converging ORF, but neither encodes a protein similar to those flanking *S. coelicolor relA*. In *S. equisimilis*, the *relA* homologue is followed by a small ORF of unknown function that is transcribed in the same orientation as the putative (p)ppGpp synthetase, and then by the converging streptokinase gene, *spc* (Mechold *et al.*, 1993). Only in *M. genitalium* is there a resemblance of the gene organization found in *S. coelicolor*; the *relA* homologue of *M. genitalium* (designated *spoT* by Peterson *et al.*, 1995) is preceded by a small ORF of unknown function, and then by a homologue of *apt*.

S. coelicolor relA is transcribed from multiple promoters in a growth-phase-dependent manner. The start site for the *relAp2* transcript, which persists at a constant level throughout growth, is preceded by a -10 region (TACCAT) that is similar to the consensus sequence (TAGYYT, where Y=G or A) deduced by Strohl (1992) for streptomycete promoters that are probably transcribed by the major RNA polymerase holoenzyme (Brown *et al.*, 1992), but lacks a recognizable -35 region. Perhaps the inverted

repeat (Fig. 3) extending from -13 to -33 binds a transcriptional activator that would obviate the need for a conventional -35 sequence. However, since transcription from *relAp2* occurred *in vitro* with purified RNA polymerase, the requirement for any such activator is presumably not absolute. Interestingly, transcription from *relAp1* peaked during transition phase, and after nutritional shiftdown, raising the possibility that *relAp1* is itself subject to positive stringent control. Although the level of the RNA-protected fragment corresponding to *relAp1* was noticeably lower than that derived from *relAp2*, the longer transcript may be translated more efficiently, and while differential transcription of *relA* in *E. coli* appears to play no role in regulating ppGpp synthesis, it is possible that increased levels of *relA* transcription contribute, at least in part, to the elevated levels of ppGpp in *S. coelicolor*. Alternatively, increased levels of transcription from *relAp1* may simply compensate for the decrease in readthrough transcription from *apt*, which is presumably subject to negative stringent control, and may serve to maintain constant levels of RelA. While the sequences upstream of the *relAp1* transcription start site do not resemble any other known streptomycete promoter (Strohl, 1992), the sequence TTGCA, which starts 10-11 bases upstream of the start site, matches perfectly the consensus -12 sequence of promoters recognized by σ^{54} (Gross *et al.*, 1992); further analysis will be required to identify the RNA polymerase holoenzyme responsible for initiating transcription at *relAp1*. The significance of the transcriptional readthrough from *apt*, and its growth-phase dependence, remains to be determined.

Experimental procedures

Bacterial strains, and culture and transformation conditions

S. coelicolor strains J1929 (Bedford *et al.*, 1995) and M145 (Hopwood *et al.*, 1985), and *S. lividans* strain 1326 (Hopwood *et al.*, 1985), were manipulated as previously described (Hopwood *et al.*, 1985). SMM is the modified minimal medium of Takano *et al.* (1992); SMMS is a solidified version of SMM containing 1.5% (w/v) agar, but lacks antifoam and PEG 6000. SFM contains 2% (w/v) mannitol, 2% (w/v) soya flour and 2% (w/v) agar in tap water, and was autoclaved twice. *E. coli* K-12 DH5 α (Hanahan, 1983) was grown and transformed according to Sambrook *et al.* (1989), and transformants were selected with carbenicillin at a final concentration of 200 $\mu\text{g ml}^{-1}$.

Isolation of plasmid DNA from *E. coli*

Plasmid DNA was isolated from *E. coli* strains using the following procedure (C. Thomas, personal communication), which reproducibly gave DNA of high purity that could be used directly for sequencing. A 10 ml culture grown overnight at 37°C in L broth containing 200 $\mu\text{g ml}^{-1}$ carbenicillin was harvested by centrifugation, resuspended in 1 ml of TEG (25 mM Tris pH 8, 10 mM EDTA, 50 mM glucose), and left at room

temperature for 5 min; 2 ml of 0.2 M NaOH/1% (w/v) SDS was added and the mixture was left at room temperature for 5 min. Sodium acetate (pH 4.8) (1.5 ml of 3 M) was added, and the mixture was vortexed, left on ice for 10 min, and microcentrifuged for 10 min. The supernatant was extracted with an equal volume of phenol:chloroform (1:1, v/v), and the nucleic acid precipitated by adding an equal volume of isopropanol and leaving on ice for 10 min. The pellet was harvested by centrifugation, washed in 80% (v/v) ethanol, dried, and redissolved in 1 ml of water. LiCl (333 µl of 8 M) was added and the solution was left at -20°C for at least 20 min. After centrifugation, the supernatant was transferred to a clean tube, and 70 µl of 3 M sodium acetate (pH 6) and 3.5 ml of ethanol were added. After storage at -20°C for at least 20 min, the DNA was harvested by centrifugation, washed with 70% (v/v) ethanol, and the pellet resuspended in 200 µl.

PCR

The synthetic oligonucleotides 5'-CATGGATCCAACGG-(CG)TACGAG(AT)(CG)(CG)(AC)T(CG)CACAC and 5'-CATGGATCCGTGTG(CG)A(CT)(CG)GCGTA(CG)GCGAAGTC (bracketed nt indicate positions of degeneracy) were used in the DNA PCR (Erich, 1989) to amplify an internal segment of the *relA* homologue from *S. coelicolor* M145 DNA. The reaction mixture contained 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 1.5 mM MgCl_2 , 0.1% (w/v) gelatin, 200 µM of each of the four dNTPs, 2.5 U of *Taq* polymerase, 40 pmol of each primer, and 50 ng of chromosomal DNA in a final volume of 100 µl. After denaturation at 95°C for 5 min, the samples were subjected to 25 cycles of denaturation (94°C , 1 min), annealing (55°C , 40 s), and extension (72°C , 50 s) and then incubated for 10 min at 72°C . PCR products were analysed by agarose (1.2% w/v) gel electrophoresis.

Nucleotide sequencing

The nucleotide sequence of the *relA* region was determined by dideoxy-sequencing (Sanger *et al.*, 1977) using the Promega TaqTrack sequencing kit and a mixture of double- and single-stranded DNA templates derived by subcloning from pIJ6031 and pIJ6037. Synthetic oligonucleotides were used to close gaps in the sequence. The region encoding *relA* and *apt* submitted to the databases (EMBL X87267) was sequenced on both strands; parts of the sequence downstream of *relA* (nt positions 3351 to 4515) were determined on only one strand and the data are available from the authors on request.

S1 nuclease protection studies

pIJ6068, pBluescript SK+ with the 458 bp *StuI*-*relA*-*SalI* fragment (nt positions 474-931) cloned between the *SmaI* and *SalI* sites of the vector, was digested with *SalI* and calf-intestinal alkaline phosphatase, and the DNA extracted with phenol:chloroform (1:1, v/v), ethanol-precipitated and digested with *SstI*. The gel-purified 498 bp *SstI*-*SalI* fragment (10 pmol) was labelled using polynucleotide kinase and 30 pmol of [γ - ^{32}P]-ATP (3000 Ci mmol $^{-1}$, Amersham) (Sambrook *et al.*, 1989). Unincorporated label was removed from the probe by passage through a Pharmacia Nick column. Fifty micrograms

of RNA, isolated from cultures of *S. coelicolor* M145 grown in SMM liquid medium (Takano *et al.*, 1992), were used in each S1 nuclease protection experiment with about 0.2 pmol (approximately 10^5 Cerenkov c.p.m.) of the probe. Hybridizations were carried out in NaTCA buffer (Murray, 1986) at 45°C overnight after denaturation at 65°C for 15 min. S1 nuclease digestions and analysis of RNA-protected fragments were carried out as previously described (Janssen *et al.*, 1989). The endpoints of the RNA-protected fragments were determined by sequencing a sample of the same end-labelled fragment that was used as probe using the method of Maxam and Gilbert (1980).

In vitro transcription

RNA polymerase was isolated from a transition-phase culture of *S. coelicolor* M145, as described by Buttner and Brown (1985), and *in vitro* run-off transcription assays were performed according to Buttner *et al.* (1987). The 498 bp *SstI*-*SalI* and 466 bp *BamHI*-*SalI* fragments of pIJ6068 were purified by gel electrophoresis and used as templates. Transcripts were analysed on denaturing 6% polyacrylamide gels using the Gibco BRL 100 bp ladder that had been end-labelled with ^{32}P as size marker.

Plaque lifts and Southern hybridization

Plaque lifts and Southern blots (Southern, 1975) were carried out as described by Hopwood *et al.* (1985), using gel-purified DNA fragments containing internal segments of *relA* labelled with ^{32}P by random oligonucleotide priming (Sambrook *et al.*, 1989).

Amino acid depletion and (p)ppGpp assays

Exponentially growing cultures ($\text{OD}_{450\text{ nm}}$ values of 0.5-0.6) were harvested by filtration (Whatman nitrocellulose filters, 0.45 µm pore size) from SMM medium containing $5\text{ }\mu\text{g ml}^{-1}$ Thio, rapidly washed with either SMM or SMM lacking cas-amino acids, resuspended in the same medium that was used for washing, and quickly returned to the shaking incubator; the 15 mM $(\text{NH}_4)_2\text{SO}_4$ present in SMM provides a source of inorganic nitrogen for the continued growth of cultures depleted of amino acids. (p)ppGpp assays were carried out as described by Strauch *et al.* (1991).

Note added in proof

We have made a *relA* null mutant of the prototrophic *S. coelicolor* strain M600 in which most of the *relA* coding region (corresponding to amino acid residues 168-686) has been deleted. On SMMS medium the mutant, which must lack *relA*-dependent ppGpp synthetase activity, does not produce γ -actinorhodin (the blue diffusible lactone form of actinorhodin; L. Bystrykh, personal communication) or actinorhodin (the purple mycelial form); on the richer R5 medium, γ -actinorhodin production is abolished, although some actinorhodin is made, while on R2 medium (R5 lacking yeast extract) there is no apparent effect on antibiotic production. We believe that the N-terminal segment of RelA that remains after the disruption reported in this paper possesses ribosome-independent

ppGpp synthetase activity, and that the elevated basal level of ppGpp observed in the disruptant (8.3 and 10.6 pmol mg⁻¹ dry weight compared with >1 and 1.7 pmol mg⁻¹ dry weight for the *relA*⁺ congenic strain) represents the constitutive synthesis of ppGpp, and accounts for the reduction in growth rate observed on solid media. These recent results also suggest that under some nutritional conditions ppGpp plays an essential role in antibiotic production in *S. coelicolor*.

Acknowledgements

We thank Mark Buttner, Belén Floriano, Gabriela Kelemen, Mark Paget, Gilles van Wezel and Andreas Wietzorrek for enthusiastic support, Mike Cashel and Dan Gentry for advice and encouragement, and Mark Buttner, Keith Chater, David Hopwood and Tobias Kieser for comments on the manuscript. This work was supported by a grant to the John Innes Centre from the Biotechnology and Biological Sciences Research Council (BBSRC) and the John Innes Foundation, and by a BBSRC Studentship to R.C.

References

- Bascaran, V., Sanchez, L., Hardisson, C., and Braña, A.F. (1991) Stringent response and initiation of secondary metabolism in *Streptomyces clavuligerus*. *J Gen Microbiol* **137**: 1625–1634.
- Baylis, H.A., and Bibb, M.J. (1987) The nucleotide sequence of a 16S rRNA gene from *Streptomyces coelicolor* A3(2). *Nucl Acids Res* **15**: 1716.
- Bedford, D.J., Laity, C., and Buttner, M.J. (1995) Two genes involved in the phase-variable ϕ C31-resistance mechanism of *Streptomyces coelicolor* A3(2). *J Bacteriol* **177**: 4681–4689.
- Bibb, M.J., Findlay, P.R., and Johnson, M.W. (1984) The relationship between base composition and codon usage in bacterial genes and its use in the simple and reliable identification of protein coding sequences. *Gene* **30**: 157–166.
- Brown, K., Wood, S., and Buttner, M.J. (1992) Isolation and characterization of the major vegetative RNA polymerase of *Streptomyces coelicolor* A3(2); renaturation of a sigma subunit using GroEL. *Mol Microbiol* **6**: 1133–1139.
- Bruton, C.J., and Chater, K.F. (1983) Mutational cloning in *Streptomyces* and the isolation of antibiotic production genes. *Gene* **26**: 67–78.
- Buttner, M.J., and Brown, N.L. (1985) RNA polymerase–DNA interactions in *Streptomyces*. *In vitro* studies of a *S. lividans* plasmid promoter with *S. coelicolor* RNA polymerase. *J Mol Biol* **185**: 177–188.
- Buttner, M.J., Fearnley, I.M., and Bibb, M.J. (1987) The agarase gene (*dagA*) of *Streptomyces coelicolor* A3(2): nucleotide sequence and transcriptional analysis. *Mol Gen Genet* **209**: 101–109.
- Buttner, M.J., Smith, A.J., and Bibb, M.J. (1988) At least three different RNA polymerase holoenzymes direct transcription of the agarase gene (*dagA*) of *Streptomyces coelicolor* A3(2). *Cell* **52**: 599–607.
- Cashel, M., and Rudd, K.E. (1987) The stringent response. In *Escherichia coli and Salmonella typhimurium: Cellular and Molecular Biology*. Neidhardt, F., Ingraham, J.L., Low, K.B., Magasanik, B., Schaechter, M., and Umberger, H.E. (eds). Washington DC: American Society for Microbiology, pp. 1410–1438.
- Chamness, W.C., and Chater, K.F. (1994) Regulation and integration of antibiotic production and morphological differentiation in *Streptomyces* spp. In *Regulation of Bacterial Differentiation*. Piggot, P., Moran, C.P., and Youngman, P. (eds). Washington DC: American Society for Microbiology, pp. 61–93.
- Chater, K.F., and Bibb, M.J. (1996) Regulation of bacterial antibiotic production. In *Biotechnology, Volume 7: Products of Secondary Metabolism*. Kleinkauf, H., and von Döhren, H. (eds). Weinheim: VCH, in press.
- Chater, K.F., and Hopwood, D.A. (1993) *Streptomyces*. In *Bacillus subtilis and Other Gram-Positive Bacteria: Biochemistry, Physiology, and Molecular Genetics*. Sonenshein, A.L., Hoch, J.A., and Losick, R. (eds). Washington, D.C.: American Society for Microbiology, pp. 83–89.
- Devereux, J., Haeberli, P., and Smithies, O. (1984) A comprehensive set of sequence analysis programs for the VAX. *Nucl Acids Res* **12**: 387–395.
- Erich, H.A. (1989) *PCR Technology*. New York: Stockton Press.
- Flårdh, K., Axberg, T., Albertson, N.H., and Kjelleberg, S. (1994) Stringent control during carbon starvation of marine *Vibrio* sp. strain 14: molecular cloning, nucleotide sequence and deletion of the *relA* gene. *J Bacteriol* **176**: 5949–5957.
- Gentry, D.R., Hernandez, V.J., Nguyen, L.H., Jensen, D.B., and Cashel, M. (1993) Synthesis of the stationary-phase sigma factor σ^S is positively regulated by ppGpp. *J Bacteriol* **175**: 7982–7989.
- Gross, C.A., Lonetto, M., and Losick, R. (1992) Bacterial sigma factors. In *Transcriptional Regulation*. Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press, pp. 129–176.
- Hamagishi, Y., Yoshimoto, A., Oki, T., and Inui, T. (1980) Occurrence of guanosine-5'-diphosphate-3'-diphosphate and adenosine-5'-diphosphate-3'-diphosphate in *Streptomyces galilaeus*. *Agric Biol Chem* **44**: 1003–1007.
- Hanahan, D. (1983) Studies on transformation of *Escherichia coli* with plasmids. *J Mol Biol* **166**: 557–580.
- Hernandez, V.J., and Bremer, H. (1990) Guanosine tetraphosphate (ppGpp) dependence of the growth rate control of *rrnB* P1 promoter activity in *Escherichia coli*. *J Biol Chem* **265**: 11605–11614.
- Hernandez, V.J., and Bremer, H. (1991) *Escherichia coli* ppGpp synthetase II activity requires *spoT*. *J Biol Chem* **266**: 5991–5999.
- Hernandez, V.J., and Bremer, H. (1993) Characterization of RNA and DNA synthesis in *Escherichia coli* strains devoid of ppGpp. *J Biol Chem* **268**: 10851–10862.
- Hershey, H.V., and Taylor, M.W. (1986) Nucleotide sequence and deduced amino acid sequence of *Escherichia coli* adenine phosphoribosyltransferase and comparison with other analogous enzymes. *Gene* **43**: 287–293.
- Holt, T.G., Chang, C., Laurentwinter, C., Murakami, T., Garrels, J.I., Davies, J.E., and Thompson, C.J. (1992) Global changes in gene expression related to antibiotic synthesis in *Streptomyces hygroscopicus*. *Mol Microbiol* **6**: 969–980.
- Hopwood, D.A., Bibb, M.J., Chater, K.F., Kieser, T., Bruton,

- C.J., Kieser, H.M., Lydiate, D.J., Smith, C.P., Ward, J.M., and Schrempf, H. (1985) *Genetic Manipulation of Streptomyces: A Laboratory Manual*. Norwich: John Innes Foundation.
- Janssen, G.R., Ward, J.M., and Bibb, M.J. (1989) Unusual transcriptional and translational features of the aminoglycoside phosphotransferase gene (*aph*) from *Streptomyces fradiae*. *Genes Dev* **3**: 415–429.
- Jones, G.H. (1994a) Purification and properties of ATP-GTP 3'-pyrophosphotransferase (guanosine pentaphosphate synthetase) from *Streptomyces antibioticus*. *J Bacteriol* **176**: 1475–1481.
- Jones, G.H. (1994b) Activation of ATP-GTP 3'-pyrophosphotransferase (guanosine pentaphosphate synthetase) in *Streptomyces antibioticus*. *J Bacteriol* **176**: 1482–1487.
- Kelly, K.S., Ochi, K., and Jones, G.H. (1991) Pleiotropic effects of a *relC* mutation in *Streptomyces antibioticus*. *J Bacteriol* **173**: 2297–2300.
- Kieser, H.M., Kieser, T., and Hopwood, D.A. (1992) A combined genetic and physical map of the *Streptomyces coelicolor* A3(2) chromosome. *J Bacteriol* **174**: 5496–5507.
- Lewin, B. (1994) *Genes V*. New York: Oxford University Press and Cell Press.
- Maxam, A.M., and Gilbert, W. (1980) Sequencing end-labelled DNA with base specific chemical cleavages. *Meth Enzymol* **65**: 449–560.
- Mechold, U., Steiner, K., Vettermann, S., and Malke, H. (1993) Genetic organisation of the streptokinase region of the *Streptococcus equisimilis* H64A chromosome. *Mol Gen Genet* **241**: 129–140.
- Metzger, S., Dror, I.B., Aizenman, E., Schreiber, G., Toone, M., Friesen, J.D., Cashel, M., and Glaser, G. (1988) The nucleotide sequence and characterization of the *relA* gene of *Escherichia coli*. *J Biol Chem* **263**: 15699–15704.
- Metzger, S., Sarubbi, E., Glaser, G., and Cashel, M. (1989) Protein sequences encoded by the *relA* and the *spoT* genes of *Escherichia coli* are interrelated. *J Biol Chem* **264**: 9122–9125.
- Murray, M.G. (1986) Use of sodium trichloroacetate and mung bean nuclease to increase sensitivity and precision during transcript mapping. *Anal Biochem* **158**: 165–170.
- Ochi, K. (1986) Occurrence of the stringent response in *Streptomyces* sp. and its significance for the inhibition of morphological and physiological differentiation. *J Gen Microbiol* **132**: 2621–2631.
- Ochi, K. (1987) A *rel* mutation abolishes the enzyme induction needed for actinomycin synthesis by *Streptomyces antibioticus*. *Agric Biol Chem* **51**: 829–835.
- Ochi, K. (1988) Nucleotide pools and stringent response in regulation of *Streptomyces* differentiation. In *Biology of Actinomycetes '88*. Okami, Y., Beppu, T., and Ogawara, H. (eds). Tokyo: Scientific Societies Press, pp. 330–337.
- Ochi, K. (1990a) *Streptomyces relC* mutants with an altered ribosomal protein ST-L11 and genetic analysis of a *Streptomyces griseus relC* mutant. *J Bacteriol* **172**: 4008–4016.
- Ochi, K. (1990b) A relaxed (*rel*) mutant of *Streptomyces coelicolor* A3(2) with a missing ribosomal protein lacks the ability to accumulate ppGpp, A-factor and prodigiosin. *J Gen Microbiol* **136**: 2405–2412.
- Paget, M.S.B., Hintermann, G., and Smith, C.P. (1994) Construction and application of streptomycete promoter probe vectors which employ the *Streptomyces glaucescens* tyrosinase-encoding gene as reporter. *Gene* **146**: 105–110.
- Peterson, S.N., Lucier, T., Heitzman, K., Smith, E.A., Bott, K.F., Hu, P.-C., and Hutchison, III, C.A. (1995) Genetic map of the *Mycoplasma genitalium* chromosome. *J Bacteriol* **177**: 3199–3204.
- Pogliano, K.J., and Beckwith, J. (1994) Genetic and molecular characterization of the *Escherichia coli* *secD* operon and its products. *J Bacteriol* **176**: 804–814.
- Rodicio, M.R., Bruton, C.J., and Chater, K.F. (1985) New derivatives of the *Streptomyces* phage ϕ C31 useful for the cloning and functional analysis of *Streptomyces* DNA. *Gene* **34**: 283–292.
- Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989) *Molecular Cloning. A Laboratory Manual*. Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press.
- Sanger, F., Nicklen, S., and Coulson, A.R. (1977) DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci USA* **74**: 5463–5467.
- Sarubbi, E., Rudd, K.E., and Cashel, M. (1988) Basal ppGpp level adjustment shown by new *spoT* mutants affect steady state growth rates and *rrnA* ribosomal promoter regulation in *Escherichia coli*. *Mol Gen Genet* **213**: 214–222.
- Schreiber, G., Metzger, S., Aizenman, E., Roza, S., Cashel, M., and Glaser, G. (1991) Overexpression of the *relA* gene in *Escherichia coli*. *J Biol Chem* **266**: 3760–3767.
- Simuth, J., Hudec, J., Chan, H.T., Danyi, O., and Zelinka, J. (1979) The synthesis of highly phosphorylated nucleotides, RNA and protein by *Streptomyces aureofaciens*. *J Antibiot* **32**: 53–58.
- Southern, E.M. (1975) Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J Mol Biol* **98**: 503–517.
- Strauch, E., Takano, E., Baylis, H.A., and Bibb, M.J. (1991) The stringent response in *Streptomyces coelicolor* A3(2). *Mol Microbiol* **5**: 289–298.
- Strohl, W.R. (1992) Compilation and analysis of DNA sequences associated with apparent *Streptomyces* promoters. *Nucl Acids Res* **20**: 961–974.
- Takano, E. (1993) ppGpp and antibiotic production in *Streptomyces coelicolor* A3(2). PhD Thesis, University of East Anglia, UK.
- Takano, E., and Bibb, M.J. (1994) The stringent response, ppGpp and antibiotic production in *Streptomyces coelicolor* A3(2). *Actinomycetologica* **8**: 1–10.
- Takano, E., Gramajo, H.C., Strauch, E., Andres, N., White, J., and Bibb, M.J. (1992) Transcriptional regulation of the *redD* transcriptional activator gene accounts for growth-phase-dependent production of the antibiotic undecylprodigiosin in *Streptomyces coelicolor* A3(2). *Mol Microbiol* **6**: 2797–2804.
- Tosa, T., and Pizer, L.I. (1971) Biochemical basis for the antimetabolite action of L-serine hydroxamate. *J Bacteriol* **106**: 972–982.
- Wright, F., and Bibb, M.J. (1992) Codon usage in the G + C-rich *Streptomyces* genome. *Gene* **113**: 55–65.
- Xiao, H., Kalman, M., Ikehara, K., Zemel, S., Glaser, G., and Cashel, M. (1991) Residual guanosine 3',5'-bispyrophosphate synthetic activity of *relA* null mutants can be eliminated by *spoT* null mutations. *J Biol Chem* **266**: 5980–5990.