University of Groningen

Novel monopentamethylcyclopentadienyl alkoxides of La and Ce; X-ray crystal structure of (C₅Me₅Ce(OCMe₃)₂)₂

Heeres, Hero J.; Teuben, Jan H.; Rogers, Robin D.

Published in:
Journal of Organometallic Chemistry

DOI:
[10.1016/0022-328X\(89\)85334-3](https://doi.org/10.1016/0022-328X(89)85334-3)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1989

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Heeres, H. J., Teuben, J. H., & Rogers, R. D. (1989). Novel monopentamethylcyclopentadienyl alkoxides of La and Ce; X-ray crystal structure of (C₅Me₅Ce(OCMe₃)₂)₂. *Journal of Organometallic Chemistry*, 364(1). [https://doi.org/10.1016/0022-328X\(89\)85334-3](https://doi.org/10.1016/0022-328X(89)85334-3)

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Journal of Organometallic Chemistry, 364 (1989) 87-96
 Elsevier Sequoia S.A., Lausanne - Printed in The Netherlands
 JOM 09540

Novel monopentamethylcyclopentadienyl alkoxides of La and Ce; X-ray crystal structure of $(C_5Me_5Ce(OCMe_3)_2)_2$

Hero J. Heeres, Jan H. Teuben *,

Department of Chemistry, University of Groningen, 9747 AG Groningen (The Netherlands)

and Robin D. Rogers *

Department of Chemistry, Northern Illinois University, DeKalb, Illinois, (U.S.A.)

(Received September 19th, 1988)

Abstract

Treatment of monomeric $(C_5Me_5)_2LnCH(SiMe_3)_2$ ($Ln = La$ and Ce) compounds with $HOCMe_3$ gives the novel compounds $(C_5Me_5Ln(OCMe_3)_2)_2$. An X-ray diffraction study of the cerium analogue reveals a dimeric structure with two symmetrically bridging and two terminal alkoxide ligands in a *cis* configuration. Crystal data: monoclinic space group $P2_1/c$, a 13.163(5), b 17.429(4), c 18.988(4) Å, β 100.75(2)°, V 4279.7 Å³ and $Z = 4$. Least-squares refinement leads to a final R (based on F) value of 0.044 for 4860 observed reflections. Treatment of $(C_5Me_5)_2CeCH(SiMe_3)_2$ with an excess of 2,6-di-*t*-butylphenol at room temperature gives $(C_5Me_5)_2(2,6\text{-di-}t\text{-butylphenoxo})$ cerium. Tris(2,6-di-*t*-butylphenoxo)cerium and C_5Me_5H are formed when the mixture is heated to 80 °C. The intermediate (C_5Me_5) bis(2,6-di-*t*-butylphenoxo)cerium is not observed under these conditions.

Introduction

The organometallic chemistry of the early lanthanides (La-Nd) is developing rapidly. Most investigations have involved complexes containing bridged dicyclopentadienyl [1] or substituted cyclopentadienyl ligands [2]. The pentamethylcyclopentadienyl group in particular seems to be a very suitable co-ligand, and a number of bis(pentamethylcyclopentadienyl) early lanthanide halide, carbyl, and amido complexes have been reported [3]. However, only a few stable mono-cyclopentadienyl or substituted cyclopentadienyl complexes are known viz: $C_5Me_5LnI_2 \cdot 3THF$ ($Ln = La$ and Ce) [4], $(C_5Me_5NdCl_3)(Na(OEt_2)_2)$ [3b], $C_5Me_5Nd(N(SiMe_3)_2)_2$ [3b], $(Li(THF)_2)_2(\mu-Cl)_4(C_5H_5)Ln \cdot THF$ ($Ln = La$ and Nd) [5] and $HmiLnCl_2$ ($Ln = La$ and Nd , $Hmi =$ heptamethylindenyl) [6]. As a part of a

program aimed at developing general methods for the synthesis of mono- C_5Me_5 early-lanthanide complexes we have begun to examine the removal of C_5Me_5 ligands from well-defined bis- C_5Me_5 complexes by use of protic acids. This approach has been used successfully to synthesize mixed-ligand ytterbium organometallics involving cyclopentadienyl and chelating ligands (e.g. β -diketonates) starting from $(\text{C}_5\text{H}_5)_3\text{Yb}$ [7]. We describe here the reactions of $(\text{C}_5\text{Me}_5)_2\text{LnCH}(\text{SiMe}_3)_2$ ($\text{Ln} = \text{La}$ and Ce) with HOCMe_3 and 2,6-di-*t*-butylphenol. These alcohols were chosen because bulky alkoxide ligands are attractive in respect of their high solubility and large size, which offer the possibility of obtaining stable sterically-saturated complexes. The CMe_3 groups also simplify NMR spectroscopy, which is especially useful for the identification of paramagnetic cerium(III) complexes.

Experimental

General considerations

All the compounds are extremely air sensitive and manipulations were carried out by Schlenk or glovebox techniques under nitrogen or argon. Solvents were distilled from Na/K alloy prior to use. HOCMe_3 (Merck) and 2,6-di-*t*-butylphenol (Janssen) were stored under nitrogen over molecular sieves (4 Å) at 50 °C. $\text{C}_5\text{Me}_5\text{H}$ was made by Burger's method [8] and converted into $\text{C}_5\text{Me}_5\text{Li}$ as described in ref. 9. $\text{LiCH}(\text{SiMe}_3)_2$ was prepared by a published procedure [10] and sublimed (90 °C/0.01 mmHg) before use. $(\text{C}_5\text{Me}_5)_2\text{CeCH}(\text{SiMe}_3)_2$ was prepared as described in ref. 11 and the lanthanum analogue similarly. Tris(2,6-di-*t*-butylphenoxo)cerium* was prepared in 65% yield by a slightly modification of the procedure described for the corresponding yttrium complex [12], the reaction mixture being stirred under reflux for 4 d instead of 8 h. IR spectra were recorded on a Pye-Unicam SP3-300 spectrophotometer with Nujol mulls between KBr discs. NMR spectra were recorded on a Bruker WH-90 or Varian VXR-300 spectrometer. Chemical shifts are reported in parts per million and were referenced to residual protons in deuterated solvents (benzene- d_6 : δ 7.15; toluene- d_8 : δ 6.98). ^{13}C chemical shifts were referenced to benzene- d_6 (δ 127.96). Elemental analyses were performed at the Micro-analytical Department of the University of Groningen under supervision of Mr A.F. Hamminga.

1. The synthesis of $(\text{C}_5\text{Me}_5)_2\text{Ce}(\text{OCMe}_3)_2$ (**1**)

HOCMe_3 (0.35 ml; 3.6 mmol) was added from a syringe to a stirred solution of 1.02 g (1.8 mmol) $(\text{C}_5\text{Me}_5)_2\text{CeCH}(\text{SiMe}_3)_2$ in pentane (20 ml) at room temperature. The color of the solution immediately changed from red to light-green. The solution was stirred for another 15 min and then concentrated to ca. 5 ml. Crystallization at -80 °C gave 0.42 g (0.50 mmol, 56%) of **1** as yellow crystals. IR (cm^{-1}): 2710(w), 2130(w), 1380(m), 1350(m), 1220(m), 1195(s), 1170(s), 1020(m), 990(vs), 920(vs), 800(m), 770(m), 750(m), 520(m), 500(s), 490(m). ^1H NMR (benzene- d_6 , 25 °C): δ 11.80 (s, *lw* 15 Hz, 9H); δ 6.87 (s, *lw* 19 Hz, 15H); δ -35.45 (s, *lw* 33 Hz, 9H). ^{13}C NMR (benzene- d_6 , 25 °C): δ 162.3 (s, C_5Me_5); δ 124.1 (s, OCMe_3); δ 54.6 (q,

* The synthesis of this compound by treatment of $\text{Ce}[\text{N}(\text{SiMe}_3)_2]_3$ with 2,6-di-*t*-butylphenol in pentane was reported recently [13].

$J(\text{C-H})$ 123 Hz, OCMe_3); δ 11.2 (q, $J(\text{C-H})$ 125 Hz, C_5Me_5); δ -5.6 (s, OCMe_3); δ -18.6 (q, $J(\text{C-H})$ 120 Hz, OCMe_3). Anal. Found: C, 51.27; H, 7.79; Ce, 33.23; mol wt (C_6H_6), 760. $\text{C}_{18}\text{H}_{33}\text{CeO}_2$ calcd.: C, 51.28; H, 7.89; Ce, 33.24%; mol wt (C_6H_6), 843.

2. The synthesis of $(\text{C}_5\text{Me}_5)_2\text{La}(\text{OCMe}_3)_2$ (**2**)

Complex **2** was prepared from $(\text{C}_5\text{Me}_5)_2\text{LaCH}(\text{SiMe}_3)_2$ in 30% yield by a procedure similar to that described for **1**. IR (cm^{-1}): 2720(w), 1380(m), 1355(s), 1220(s), 1190(s), 1170(s), 1020(m), 985(vs), 920(s), 800(w), 770(s), 750(m), 510(s), 490(s), 460(m). ^1H NMR (benzene- d_6 , 20 ° C); δ 2.20 (s, 15H); δ 1.33 (s, 9H); δ 1.27 (s, 9H). ^{13}C NMR (benzene- d_6 , 20 ° C); δ 119.1 (s, C_5Me_5); δ 73.6 (s, OCMe_3); δ 71.0 (s, OCMe_3); δ 34.4 (q, $J(\text{C-H})$ 124 Hz, OCMe_3); δ 33.5 (q, $J(\text{C-H})$ 125 Hz, OCMe_3); δ 11.6 (q, $J(\text{C-H})$ 125 Hz, C_5Me_5). Anal. Found: C, 51.49, H, 7.88; La, 32.79. $\text{C}_{18}\text{H}_{33}\text{LaO}_2$ calcd.: C, 51.43; H, 7.91; La, 33.04%.

3. The synthesis of $(\text{C}_5\text{Me}_5)_2(2,6\text{-di-}t\text{-butylphenoxo})\text{cerium}$ (**3**)

2,6-Di-*t*-butylphenol (0.327 g, 1.6 mmol) was added to a solution of 0.903 g (1.6 mmol) of $(\text{C}_5\text{Me}_5)_2\text{CeCH}(\text{SiMe}_3)_2$ in pentane (50 ml). The deep red solution was stirred for 30 min at room temperature and then concentrated to ca. 20 ml. Crystallization at -80 ° C gave 0.647 g (1.1 mmol, 69%) of **3** as deep red crystals. IR (cm^{-1}): 3040(w), 2720(w), 2110(w), 1575(m), 1400(s), 1350(m), 1250(s), 1195(w), 1090(m), 1010(m), 850(s), 810(m), 790(m), 735(s), 630(m). ^1H NMR (benzene- d_6 , 20 ° C): δ 7.93 (d, $lw < 10$ Hz, 2H); δ 7.28 (t, $lw < 10$ Hz, 1H); δ 2.79 (s, lw 12 Hz, 30H); δ -7.5 (s, lw 24 Hz, 18H). ^{13}C NMR (benzene- d_6 , 20 ° C); δ 194.3 (s, OC_6H_3); δ 165.6 (s, C_5Me_5); δ 149.8 (s, OC_6H_3); δ 126.2 (d, $J(\text{C-H})$ 154 Hz, OC_6H_3); δ 120.0 (d, $J(\text{C-H})$ 158 Hz, OC_6H_3); δ 35.3 (s, CMe_3); δ 20.0 (q, $J(\text{C-H})$ 125 Hz, CMe_3); δ 5.6 (q, $J(\text{C-H})$ 125 Hz, C_5Me_5). Anal. Found: C, 66.42; H, 8.35; Ce, 22.87. $\text{C}_{34}\text{H}_{51}\text{CeO}$ calcd.: C, 66.30; H, 8.35; Ce, 22.75%.

X-ray data collection, structure determination and refinement for **1**

A yellow crystal of the title compound, obtained by slowly cooling a saturated pentane solution to -30 ° C, was mounted in a glass-capillary under argon. The space group was shown to be the centric $P2_1/c$ from the systematic absences. A summary of the data collection parameters is given in Table 1.

The cerium atom positions were determined by use of the direct program Multan [14]. Difference Fourier maps based on the cerium atoms showed the positions of the C_5Me_5 and oxygen ligands, and revealed the presence of rotational disorder in the CMe_3 groups. Three orientations of equal occupancy (1/3) were resolved for each CMe_3 group and refined in alternate cycles. High thermal motion (but no resolvable disorder) was observed for the C_5Me_5 ligands. The hydrogen atoms were not included in the final refinement. Refinement of the nonhydrogen atoms with anisotropic temperature factors and of the disordered carbon atoms with isotropic thermal parameters in alternate cycles led to final values of $R = 0.044$ and $R_w = 0.047$. The final positional parameters are given in Table 2.

Results and discussion

As a start to this investigation we devised a general route to $(\text{C}_5\text{Me}_5)_2\text{LnX}$ derivatives (X = carbyl, amido, hydride; Ln = La and Ce) starting from

Table 1

Crystal data, summary of intensity data collection and structure refinement for **1**

Color/Shape	Yellow/parallelepiped
Mol. wt	843.2
Space group	P2 ₁ /c
Temp., °C	20
Cell Constants ^a	
<i>a</i> , Å	13.163(5)
<i>b</i> , Å	17.429(4)
<i>c</i> , Å	18.988(4)
β , deg	100.75(2)
Cell Vol., Å ³	4279.7
Formula units/unit cell	4
<i>D</i> _{calc} , gcm ⁻³	1.31
μ (calc), cm ⁻¹	20.0
Diffractometer/Scan	Enraf-Nonius CAD4/ θ -2 θ
Range of rel. transm. factors, %	85/100
Radiation, graphite monochromator	Mo- <i>K</i> α (λ 0.71073 Å)
Crystal dimensions, mm	0.35 × 0.45 × 0.75
Scan width	0.80 + 0.35 tan θ
Standard reflections	(10,0,0), (0, 12,0), (0,0,18)
Decay of standards	± 2%
Reflections measured	8132
2 θ range, deg	2 ≤ 2 θ ≤ 50
Range of <i>h</i> , <i>k</i> , <i>l</i>	+ 15, + 20, ± 22
Reflections observed ($F_o \geq 5\sigma(F_o)$) ^b	4860
Computer programs ^c	SHELX [16]
Structure solution	MULTAN [14]
No. of parameters varied	208
Weights	($\sigma(F_o)^2$) ⁻¹
GOF	1.97
$R = \Sigma F_o - F_c / \Sigma F_o $	0.044
<i>R</i> _w	0.047
Largest feature final diff. map	0.6e ⁻ /Å

^a Least-squares refinement of $((\sin \theta)/\lambda)^2$ values for 25 refl. $\theta > 19^\circ$. ^b Corrections: Lorentz-polarization and absorption (empirical, psi scan). ^c Neutral atom scattering factors and anomalous dispersion corrections from ref. 17.

$((C_5Me_5)_2LnCl)_n$ [11]. Monomeric salt-free complexes were obtained when bulky X groups, e.g. CH(SiMe₃)₂ and N(SiMe₃)₂ were used. The reaction of (C₅Me₅)₂CeCH(SiMe₃)₂ with 2 equivalents of HOCEMe₃ in pentane gave C₅Me₅H and CH₂(SiMe₃)₂ (¹H NMR) and a single organocerium complex, which analyzed as C₅Me₅Ce(OCMe₃)₂ (**1**) (eq. 1).

Complex **1** was isolated in 55% yield as intense yellow, very air-sensitive crystals, which are readily soluble in aromatic and aliphatic hydrocarbons. Molecular weight determinations (cryoscopy in benzene) indicate that **1** is dimeric in solution. ¹H NMR and ¹³C NMR spectra (benzene-*d*₆) both show a single C₅Me₅ ligand and the

Table 2

Final fractional coordinates for $[(\eta^5\text{-C}_5\text{Me}_5)\text{Ce}(\text{O-t-Bu})_2]_2$

Atom	<i>x/a</i>	<i>y/b</i>	<i>z/c</i>
Ce(1)	0.14231(4)	0.62648(3)	0.83094(3)
Ce(2)	0.34239(4)	0.56063(3)	0.72157(3)
O(1)	0.2267(5)	0.6625(3)	0.7341(3)
O(2)	0.2541(5)	0.5212(3)	0.8173(3)
O(3)	-0.0137(5)	0.6022(4)	0.7856(3)
O(4)	0.2794(5)	0.4945(4)	0.6298(3)
C(1)	0.1884(9)	0.7165(6)	0.6758(6)
C(2)a ^a	0.081(2)	0.693(2)	0.638(1)
C(3)a	0.262(2)	0.698(1)	0.618(1)
C(4)a	0.203(2)	0.799(1)	0.709(1)
C(2)b	0.104(3)	0.663(2)	0.619(2)
C(3)b	0.272(2)	0.732(2)	0.636(1)
C(4)b	0.128(2)	0.775(2)	0.699(2)
C(2)c	0.072(3)	0.727(2)	0.669(2)
C(3)c	0.205(3)	0.680(2)	0.605(2)
C(4)c	0.259(2)	0.788(2)	0.692(2)
C(5)	0.2384(9)	0.4431(6)	0.8376(5)
C(6)a	0.332(2)	0.434(1)	0.916(1)
C(7)a	0.130(2)	0.436(1)	0.853(1)
C(8)a	0.261(2)	0.390(1)	0.775(1)
C(6)b	0.156(2)	0.448(2)	0.889(2)
C(7)b	0.188(3)	0.398(2)	0.773(2)
C(8)b	0.333(3)	0.415(2)	0.885(2)
C(6)c	0.116(3)	0.423(2)	0.812(2)
C(7)c	0.301(2)	0.391(2)	0.800(1)
C(8)c	0.267(3)	0.432(2)	0.913(2)
C(9)	-0.1233(9)	0.5885(8)	0.7569(6)
C(10)a	-0.167(2)	0.542(2)	0.813(2)
C(11)a	-0.173(2)	0.673(2)	0.741(2)
C(12)a	-0.112(2)	0.535(2)	0.683(1)
C(10)b	-0.167(3)	0.652(2)	0.704(2)
C(11)b	-0.123(3)	0.514(2)	0.709(2)
C(12)b	-0.178(3)	0.572(2)	0.815(2)
C(10)c	-0.186(3)	0.662(2)	0.768(2)
C(11)c	-0.138(3)	0.574(2)	0.682(2)
C(12)c	-0.149(3)	0.515(2)	0.800(2)
C(13)	0.237(1)	0.4533(9)	0.5644(6)
C(14)a	0.114(2)	0.458(2)	0.571(1)
C(15)a	0.290(2)	0.506(2)	0.506(1)
C(16)a	0.296(3)	0.389(2)	0.554(2)
C(14)b	0.235(2)	0.520(2)	0.501(2)
C(15)b	0.304(3)	0.362(2)	0.582(2)
C(16)b	0.132(3)	0.432(2)	0.555(2)
C(14)c	0.290(3)	0.463(2)	0.509(2)
C(15)c	0.121(3)	0.480(2)	0.539(2)
C(16)c	0.244(3)	0.364(2)	0.580(2)
C(17)	0.131(1)	0.7574(7)	0.9174(6)
C(18)	0.076(1)	0.698(1)	0.9487(7)
C(19)	0.148(1)	0.6455(8)	0.9791(6)
C(20)	0.249(1)	0.6680(7)	0.9684(5)
C(21)	0.237(1)	0.7374(6)	0.9296(5)
C(22)	0.086(1)	0.8291(7)	0.8841(7)
C(23)	-0.037(1)	0.696(1)	0.9559(8)
C(24)	0.126(1)	0.5755(8)	1.0231(6)

Table 2 (continued)

Atom	<i>x/a</i>	<i>y/b</i>	<i>z/c</i>
C(25)	0.354(1)	0.6323(8)	0.9983(6)
C(26)	0.327(1)	0.7850(7)	0.9119(7)
C(27)	0.5475(9)	0.5159(7)	0.7645(8)
C(28)	0.5415(8)	0.5839(9)	0.8005(7)
C(29)	0.5283(9)	0.6452(7)	0.7529(8)
C(30)	0.5282(9)	0.615(1)	0.6838(8)
C(31)	0.5407(9)	0.534(1)	0.6927(8)
C(32)	0.571(1)	0.4351(8)	0.798(1)
C(33)	0.558(1)	0.590(1)	0.8837(7)
C(34)	0.529(1)	0.7285(7)	0.770(1)
C(35)	0.524(1)	0.658(1)	0.6143(9)
C(36)	0.553(1)	0.476(1)	0.635(1)

^a Designations a, b, c, refer to the threefold disorder observed for the ^tBu groups. Occupancy factors were fixed at 0.3333.

presence of two different OCMe₃ groups. The solid state structure, as determined by X-ray diffraction, confirms the dimeric nature of **1** (vide infra). The temperature dependence of the ¹H NMR resonances of **1** (20 to 105 °C, Fig. 1) are as expected for a paramagnetic compound, showing the linear Curie-Weiss relation between chemical shift and the reciprocal of the temperature. These data also indicate that exchange between bridging and terminal OCMe₃ groups is slow on ¹H NMR time scale even at 105 °C.

The reaction of **1** with another equivalent of HOCMe₃ was monitored by ¹H NMR spectroscopy (toluene-*d*₈, 20 °C). Instead of C₅Me₅H formation, exchange between free HOCMe₃ and one of the alkoxide ligands took place. The OCMe₃ resonance at 11.61 ppm disappeared, and a new broad resonance at ca. 6 ppm was observed, and sharpened considerably upon heating to 80 °C (*lw*(80 °C) 650 Hz). Attempts to obtain the activation parameters for this fluctuational process were

Fig. 1. Plot of paramagnetic shifts against the reciprocal of the temperature for the ¹H NMR resonances of (C₅Me₅Ce(OCMe₃)₂)₂ (**1**).

hampered owing to slow decomposition of **1** under the conditions used, to give free C_5Me_5H and unidentified cerium complexes. From these observations we tentatively assign the resonance at 11.61 ppm to the more accessible terminal $OCMe_3$ ligands.

White crystalline $(C_5Me_5)_2La(OCMe_3)_2$ (**2**) was obtained by a procedure similar to that used for **1**. The isolated yield (30%) is rather low owing to the extremely high solubility of the complex in hydrocarbons, which makes crystallization rather cumbersome. The IR and NMR spectral characteristics of **2** also suggest a dimeric structure. The 1H NMR spectrum contains two distinct $OCMe_3$ groups, at 1.33 and 1.27 ppm. The ^{13}C NMR spectra in benzene- d_6 also show signals from two chemically inequivalent $OCMe_3$ groups, with quarternary carbon signals at 73.6 and 71.0 ppm and the methyl carbon resonances at 34.4 and 33.5 ppm.

In an attempt to obtain monomeric complexes of the $C_5Me_5Ce(OR)_2$ type, we examined the reaction of $(C_5Me_5)_2CeCH(SiMe_3)_2$ with 2,6-di-*t*-butylphenol. Treatment of $(C_5Me_5)_2CeCH(SiMe_3)_2$ with 2,6-di-*t*-butylphenol at room temperature gave $CH_2(SiMe_3)_2$ (1H NMR) and intensely-red crystalline $(C_5Me_5)_2(2,6\text{-di-}t\text{-butylphenoxo})\text{cerium}$ (**3**, eq. 2).

The reaction of $(C_5Me_5)_2CeCH(SiMe_3)_2$ with 2 equivalents of 2,6-di-*t*-butylphenol was monitored by 1H NMR spectroscopy (benzene- d_6). Fast reaction took place at room temperature, with quantitative formation of **3** and $CH_2(SiMe_3)_2$. C_5Me_5H was liberated when the mixture was heated to 80 °C, giving a mixture of tris(2,6-di-*t*-butylphenoxo)cerium (**4**) and **3** (eq. 3).

The outcome of the reaction can in principle be accounted for by assuming disproportionation of initially formed $(C_5Me_5)_2(2,6\text{-di-}t\text{-butylphenoxo})\text{cerium}$ (**5**) to **3** and tris(2,6-di-*t*-butylphenoxo)cerium. However monomeric **5** has been prepared independently, and is stable towards disproportionation under these conditions [15], so it is more likely that the rate of C_5Me_5H abstraction from **5** exceeds that of the first C_5Me_5H abstraction from **3**.

From this result it is clear that **5** is less stable towards the excess of alcohol than **1**. These differences can be rationalized on steric grounds. $C_5Me_5Ce(OCMe_3)_2$ is sterically unsaturated and dimerizes to **1**. In contrast, monomeric **5** is more open, and this results in enhanced reactivity towards alcohols.

The molecular structure of 1

To confirm the stoichiometry and assign the geometry of **1** and **2**, a crystallographic study was undertaken on the cerium analogue. The solid state structure of the molecule is depicted in Fig. 2. The molecule is dimeric with a cerium(III) ion coordinated to one $\eta^5\text{-}C_5Me_5$ ligand, one terminal $OCMe_3$, and two bridging $OCMe_3$ groups, giving the metal a formal coordination number of 6. The geometry

Fig. 2. X-Ray structure of $(C_5Me_5Ce(OCMe_3)_2)_2$, showing the numbering scheme. For clarity, only one orientation of each disordered $OCMe_3$ group is shown.

Table 3

Bond distances (Å) and angles (deg) for $[(\eta^5-C_5Me_5)Ce(O-t-Bu)_2]_2$

Atoms	Distance	Atoms	Distance
Ce(1)-O(1)	2.401(6)	Ce(1)-O(2)	2.396(6)
Ce(1)-O(3)	2.116(6)	Ce(1)-C(17)	2.83(1)
Ce(1)-C(18)	2.84(2)	Ce(1)-C(19)	2.82(1)
Ce(1)-C(20)	2.82(1)	Ce(1)-C(21)	2.82(1)
Ce(2)-O(1)	2.380(6)	Ce(2)-O(2)	2.431(6)
Ce(2)-O(4)	2.124(6)	Ce(2)-C(27)	2.78(1)
Ce(2)-C(28)	2.80(1)	Ce(2)-C(29)	2.82(1)
Ce(2)-C(30)	2.84(1)	Ce(2)-C(31)	2.80(1)
O(1)-C(1)	1.47(1)	O(2)-C(5)	1.44(1)
O(3)-C(9)	1.46(1)	O(4)-C(13)	1.45(1)
C-Cav.	1.6(1)	C-Me(Cp)	1.52(3)
C-C(Cp)	1.41(2)	Cent2-Ce(2)	2.54
Cent1 ^a -Ce(1)	2.56		
Atoms	Angle	Atoms	Angle
O(1)-Ce(1)-O(2)	74.7(2)	O(1)-Ce(1)-O(3)	107.0(2)
O(2)-Ce(1)-O(3)	111.8(2)	O(1)-Ce(2)-O(3)	74.4(2)
O(1)-Ce(2)-O(4)	109.1(2)	O(2)-Ce(2)-O(4)	107.0(2)
Ce(1)-O(1)-Ce(2)	106.1(2)	Ce(1)-O(1)-C(1)	127.0(6)
Ce(2)-O(1)-C(1)	122.6(6)	Ce(1)-O(2)-Ce(2)	104.7(2)
Ce(1)-O(2)-C(5)	124.9(6)	Ce(2)-O(2)-C(5)	125.4(6)
Ce(1)-O(3)-C(9)	176.9(7)	Ce(2)-O(4)-C(13)	176.6(7)
O-C-Cav.	107 (4)	C-C-Cav.	112 (8)
C-C-C(Cp)	108 (1)	C-C-Me(Cp)	126 (2)
Cent1-Ce(1)-O(1)	122.1	Cent2-Ce(2)-O(1)	122.9
Cent1-Ce(1)-O(2)	120.1	Cent2-Ce(2)-O(2)	123.3
Cent1-Ce(1)-O(3)	114.8	Cent(2)-Ce(2)-O(4)	113.8

^a Cent1 refers to the centroid of the Cp ring containing C(17)-C(21); Cent2, C(27)-C(31).

around the cerium (III) ion is best described as a distorted tetrahedron. Relevant bond lengths and angles, with estimated standard deviations, are given in Table 3.

The C_5Me_5 groups are η^5 -coordinated, with endocyclic C-Ce bond distances between 2.78(1) and 2.84(2) Å. These values are in the range observed for $\text{C}_5\text{Me}_5\text{CeI}_2 \cdot 3\text{THF}$ [4] and $(\text{C}_5\text{Me}_5)_2\text{CeCH}(\text{SiMe}_3)_2$ [11]. The oxygens of the bridging OCMe_3 group and the two cerium atoms are coplanar within 0.017 Å, with a Ce-Ce distance of 3.822(1) Å. The sum of the internal angles is 359.9°.

The observed *cis*-configuration of the terminal OCMe_3 and C_5Me_5 ligands seems to be favoured because it minimizes the steric repulsions between the C_5Me_5 ring and the bridging OCMe_3 ligands. The angles O(1)-O(2)-C(5) and O(2)-O(1)-C(1) are 159.3(5) and 163.3(5)°, respectively, so the CMe_3 group of the bridging alkoxide ligands are pointed away from the bulky C_5Me_5 ring. The Ce-O terminal OCMe_3 distances (av. 2.120(4) Å) are at the short end of the range previously observed for a number of cerium complexes; viz. 2.14 Å (mean) for tris(2,6-di-*t*-butylphenoxo)cerium [13] and 2.23 Å (mean) for tris(2,6-di-*t*-butylphenoxo)bis(NCCMe₃)cerium [13]. The Ce-O bridging OCMe_3 bond lengths (mean 2.40 Å) are rather long.

Supplementary material: Tables of atom coordinates, thermal parameters, complete tables of bond distances and angles and structure factors for **1** have been deposited at the National Auxiliary Publications Service no. 04644 P.O. Box 3513, Grand Central Station, New York, N.Y. 10017, U.S.A.

Conclusions

Alcoholysis of the Ce-C bonds of $(\text{C}_5\text{Me}_5)_2\text{CeCH}(\text{SiMe}_3)_2$ provides an attractive route to cerium alkoxides/aryloxides. The Ce-C σ -bond is easily cleaved, and the nature of the alcohol used and the reaction conditions determine whether the C_5Me_5 ligands will be liberated.

Acknowledgement

We thank Shell Research B.V. for financial support of this research. The U.S. National Science Foundation Chemical Instrumentation Program provided funds used to purchase the diffractometer (NIU).

References

- 1 C. Qian, C. Ye, H. Lu, Y. Li, Y. Huang, J. Organomet. Chem., 263 (1984) 333.
- 2 (a) M.F. Lappert, A. Singh, J.L. Atwood, W.E. Hunter, J. Chem. Soc., Chem. Commun., (1981) 1190; (b) M.F. Lappert, A. Singh, J.L. Atwood, W.E. Hunter, J. Chem. Soc., Chem. Commun., (1981) 1191; (c) M.F. Lappert, A. Singh, J.L. Atwood, W.E. Hunter, H.M. Zhang, J. Chem. Soc. Chem. Commun., (1983) 69.
- 3 (a) A.L. Wayda, W.J. Evans, Inorg. Chem., 19 (1980) 2190. (b) T.D. Tilley, R.A. Andersen, Inorg. Chem., 20 (1981) 3267; (c) G. Jeske, H. Lauke, H. Mauermann, P.N. Swebston, H. Schumann, T.J. Marks, J. Am. Chem. Soc., 107 (1985) 8091; (d) M.D. Rausch, K.J. Moriarty, J.L. Atwood, J.A. Weeks, W.E. Hunter, H.G. Brittain, Organometallics, 5 (1986) 1281; (e) H. Schumann, I. Albrecht, J. Loebel, E. Hahn, M.B. Hossain, D. Helm, *ibid.*, 5 (1986) 1297.
- 4 P.N. Hazin, J.C. Huffman, J.W. Bruno, Organometallics, 6 (1987) 23.
- 5 J. Zhongsheng, H. Ninghai, L. Yi, X. Xialong, L. Guozli, Inorg. Chim. Acta, 142 (1988) 333.
- 6 M. Tsutsui, L.B. Chen, D.E. Bergbreiter, T.K. Miyamoto, J. Am. Chem. Soc., 104 (1982) 855.

- 7 (a) G. Bielang, R.D. Fischer, *Inorg. Chim. Acta*, 36 (1987) L389; (b) H. Ma, Z. Ye, *J. Organomet. Chem.*, 326 (1987) 369.
- 8 U. Burger, A. Delay, F. Maznot, *Helv. Chim. Acta*, 57 (1974) 2106.
- 9 R.S. Threlkel, J.E. Bercaw, *J. Organomet. Chem.*, 136 (1977) 1.
- 10 D.J. Davidson, D.H. Harris, M.F. Lappert, *J. Chem. Soc. Dalton Trans.*, (1976) 2268.
- 11 H.J. Heeres, J. Renkema, M. Booiij, A. Meetsma, J.H. Teuben, *Organometallics*, in press.
- 12 P.B. Hitchcock, M.F. Lappert, R.G. Smith, *Inorg. Chim. Acta*, 139 (1987) 183.
- 13 H.A. Stecker, A. Sen, A.L. Rheingold, *Inorg. Chem.*, 27 (1988) 1130.
- 14 G. Germain, P. Main, M.M. Woolfson, *Acta, Cryst., A*, 27 (1971) 368.
- 15 H.J. Heeres, A. Meetsma, J.H. Teuben, *J. Chem. Soc., Chem. Commun.*, (1988) 962.
- 16 G.M. Sheldrick, SHELX, a system of computer programs for X-ray structure determination as locally modified, 1976.
- 17 International Tables For X-Ray Crystallography, Kynock Press, Birmingham, England, Vol IV, 1972, 72, 99, 149.