University of Groningen

Controlling the Morphology of Aggregates of an Amphiphilic Synthetic Receptor through Host-Guest Interactions

Suksai, Chomchai; Figueiras Gómez, Sergio; Chhabra, Anjali; Liu, Jingyuan; Skepper, Jeremy N.; Tuntulani, Thawatchai; Otto, Sijbren

Published in:
Langmuir

DOI:
[10.1021/la0609470](https://doi.org/10.1021/la0609470)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2006

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Suksai, C., Figueiras Gómez, S., Chhabra, A., Liu, J., Skepper, J. N., Tuntulani, T., & Otto, S. (2006). Controlling the Morphology of Aggregates of an Amphiphilic Synthetic Receptor through Host-Guest Interactions. *Langmuir*, 22(14). <https://doi.org/10.1021/la0609470>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Supplementary Material

Controlling the Morphology of Aggregates of an Amphiphilic Synthetic Receptor through Host-Guest Interactions

Chomchai Suksai,^{a,e} Sergio Figueiras Gómez,^b Anjali Chhabra,^c Jingyuan Liu,^a Jeremy N. Skepper,^d Thawatchai Tuntulani^e and Sijbren Otto^{*a}

^a Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge, CB2 1EW, UK
Fax: +44 1223 336017; Tel: +44 1223 336509; E-mail: so230@cam.ac.uk

^b Current address: Departamento de Química Orgánica y Unidad Asociada al CSIC, Universidad de Santiago de Compostela, 15706 Santiago de Compostela, Spain.

^c Current address: Department of Chemistry, Indian Institute of Technology Bombay, India.

^d Multi-Imaging Center, Department of Anatomy, University of Cambridge, Downing Street, Cambridge, CB2 3DY, UK.

^e Department of Chemistry, Faculty of Science, Chulalongkorn University, Bangkok, 10330, Thailand.

Contents

S2	General procedures
S2	Synthesis of 3
S4	Synthesis and purification of 1b
S5	Characterisation of receptor 1b by LC-MS
S7	TOCSY-NMR spectrum of receptor 1b <i>rac</i>
S8	¹ H NMR titration of MeOD- <i>d</i> ₄ and VT- ¹ H NMR spectra of receptor 1b <i>rac</i>
S9	Re-equilibration experiments
S11	Isothermal titration calorimetry
S11	Dynamic light scattering studies
S12	Transmission electron microscopy
S12	Nile Red solubilisation
S13	References

General Procedures. Receptor **1a**,¹ building block **2**,¹ and guests **4**² and **5**³ were prepared following literature procedures. ¹H-NMR and 2D-NMR spectra were recorded on a Bruker ATM 500 MHz Spectrometer at room temperature. VT ¹H-NMR spectra were recorded on a Bruker DPX 500 Spectrometer.

Analytical HPLC was carried out on a Hewlett Packard 1100 instrument coupled to a UV analyser, set to 250 nm. The data was processed using HP Chemstation software. Separations were performed using a Nucleodur C₁₈ column (4.6 cm × 250 mm, 100 Å, 5 µm), flow rate 1 mL/min, 75:25:0.1 acetonitrile:water:trifluoroacetic acid as mobile phase at 45⁰ C.

LC-MS Method and Parameters. LC was performed using an Agilent 1100 series HPLC equipped with an online degasser, binary pump, autosampler, heated column compartment and diode array detector. MS was performed using an Agilent XCT ion trap MSD mass spectrometer. The mixture of **2**, **3** and **4** (see main text) was analyzed by injecting 10 µl of the solution onto a Nucleodur C₁₈ column (4.6 cm × 250 mm, 100 Å, 5 µm) using a 75:25:0.1 acetonitrile:water:trifluoroacetic acid mixture as the mobile phase at a flow rate of 1 mL/min. Mass spectra (negative ion mode) were acquired in ultra scan mode using a drying temperature of 350⁰C, a nebuliser pressure of 55.00 psi, a drying gas flow of 121 mL/min, a capillary voltage of 4000 V and an ICC target of 200,000 ions. We tuned for a target mass of 1200.

Synthetic Procedures for building block 3 are outlined in Scheme S1.

Scheme S1. Synthesis of building block **3**. Chemicals and reagents: (a) DMF, *N,N*-dimethylthiocarbamoyl chloride, DABCO, 15 hours,¹ (b) diphenyl ether, 230⁰C, 3 hours,¹ (c) Me₃SiI, reflux 100⁰C, 15 hours, (d) oxalyl chloride, DCM, 4 hours, (e) dodecylamine, NEt₃, DCM, 12 hours and (f) diethylene glycol, KOH, 105⁰C, 4 hours.

3,5-Bis(dimethylcarbamoylsulfanyl)benzoic acid (iv)

A mixture of 3,5-bis(dimethylcarbamoylsulfanyl)benzoic acid methyl ester (iii)¹ (2.8 g, 8.18 mmol) and Me₃SiI (6.22 mL, 45 mmol) was refluxed at 105°C for 15 hours. Diethyl ether (200 mL) was added and the solution was washed with 10% NaOH and dried over Na₂SO₄. A white solid (2.42 g, 7.36 mmol) was obtained in 90% yield after removing the diethyl ether. ¹H NMR (CDCl₃, 400 MHz, (ppm)): δ = 8.19 (s, 2H, *ArH*), 7.85 (s, 1H, *ArH*), 3.09 (m, 12H, -N(CH₃)₂); ¹³C NMR (CDCl₃, 100 MHz, ppm); δ = 166.68, 164.12, 144.85, 135.73, 128.90, 128.05, 34.98. Exact mass: calcd: 329.0630; found: 329.0630 (M+H⁺).

3,5-Bis(dimethylcarbamoylsulfanyl)benzoyl chloride (v)

Oxalyl chloride (1.3 mL, 15mmol) was added to a solution of (iv) (0.98 g; 3 mmol) in 8 mL of dry dichloromethane with 4 drops of DMF under a nitrogen atmosphere. The solution was stirred at room temperature for 3 hours. The solvent was removed under vacuum and the crude product was used in the next step without further purification.

N-dodecyl-bis(3,5-dimethylcarbamoylsulfanyl)benzamide (vi)

To a solution of dodecylamine (0.64g, 3.5 mmol) and triethylamine (0.55 mL, 4 mmol) in dry dichloromethane (50 mL) was added a solution of acid chloride v (1.09 g; 3 mmol) in dry dichloromethane (15 mL). The mixture was stirred at room temperature for 15 hours. Then the solution was washed with 10% HCl and water, respectively. After drying over MgSO₄ and removal of the solvent, the residue was recrystallised from MeOH, giving 1.33 g (2.7 mmol; 90%) of a white solid. ¹H NMR (CDCl₃, 400 MHz, ppm): δ = 7.86 (s, 2H, *ArH*), 7.73 (s, 1H, *ArH*), 6.22 (m, 1H, -NH-), 3.39 (q, J = 6 Hz, 2H, -NHCH₂-), 3.05 (m, 12H, -N(CH₃)₄), 1.59 (m, J = 7.6 Hz, 2H, -NHCH₂CH₂-), 1.25 (m, 18H, *alkyl chain* CH₂), 0.87 (t, J = 6.8 Hz, 3H, -CH₃); ¹³C NMR (CDCl₃, 100 MHz, ppm): δ = 164.89, 143.42, 135.19, 133.51, 129.12, 44.78, 39.25, 35.98, 30.90, 28.61, 28.53, 28.32, 25.99, 21.67, 13.10. Exact mass: calcd: 518.2487; found: 518.2487(M+Na⁺).

N-dodecyl-3,5-dimercaptobenzamide (vii)

Compound vi (0.3 g; 0.60 mmol) was dissolved in 35 mL of a degassed solution of KOH (1.75 M) in diethylene glycol and heated at 105°C for 30 min under a nitrogen atmosphere. The solution was cooled to room temperature and 250 mL of degassed H₂O

was added rapidly, followed by 10% HCl (30 mL). The white precipitate (0.20 g; 0.58 mmol; 97%) was filtered and extensively washed with water. ^1H NMR (CDCl_3 , 400 MHz, ppm): δ 7.39 (s, 1H, *ArH*), 7.26 (d, $J = 7.2$ Hz, 2H, *ArH*), 5.9 (m, 1H, $-\text{NH}-$), 3.42 (q, $J = 7.2$ Hz, 2H, $-\text{NHCH}_2-$), 1.55 (m, 2H, $-\text{NHCH}_2\text{CH}_2-$), 1.25 (m, 18H, $-\text{CH}_2-$), 0.87 (t, $J = 7.2$ Hz, 3H, $-\text{CH}_3$); ^{13}C NMR (CDCl_3 , 100 MHz, ppm): δ 166.99, 135.66, 132.14, 130.65, 123.89, 39.41, 31.05, 28.77, 28.73, 28.68, 28.46, 26.13, 21.82, 13.25. Exact mass: calcd: 354.1925; found: 354.1925($\text{M}+\text{H}^+$).

Synthesis and Purification of Receptor 1b

Receptor **1b** was prepared by stirring a mixture of building blocks **2** (6.67 mM), **3** (3.33 mM) and template **4** (10 mM) in DMSO for 24 h at room temperature. The receptor was isolated by using a Gilson preparative HPLC. Aliquots of 2 mL of the solution were chromatographed using a Nucleodur C_{18} preparative column (25.0 cm $\times$ 2.1 cm, 100 Å, 5 μm) with a Nucleodur C_{18} guard column (5.0 cm $\times$ 2.1 cm, 100 Å, 5 μm), at a flow rate of 20 mL/min, using 75:25:0.1 acetonitrile:water:trifluoroacetic acid as mobile phase. The chromatography was performed at 45 $^\circ$ C by immersing the column, the guard column and a length of HPLC tubing (to ensure heat exchange) into a water bath. Retention times were 35-40 minutes for the major *rac* diastereomer. The collected fractions of 10 runs were combined and dried *in vacuo*, and redissolved in 10 mL of DMSO and rechromatographed in 5 injections. The collected product was dried *in vacuo*, filtered and washed extensively with water (250 mL).

^1H NMR of **1b** *rac* ($\text{D}_2\text{O}:\text{CD}_3\text{CN}$ 1:1, 500 MHz, ppm) δ 8.59 (s, 1H, *ArH*), 8.10 (m, 6H, *ArH*), 7.78 (d, $J = 7.5$ Hz, 2H, *ArH*), 7.69 (m, 4H, *ArH*), 7.55 (m, 2H, *ArH*), 6.37 (s, 2H, $-\text{CH}-$), 6.29 (s, 2H, $-\text{CH}-$), 3.78 (q, $J = 6.6$ Hz, 2H, $-\text{NHCH}_2-$), 2.02 (t, $J = 6.1$ Hz, 2H, $-\text{NHCH}_2\text{CH}_2-$), 1.63 (m, 18H, $-\text{CH}_2-$), 1.21 (t, $J = 7.1$ Hz, 3H, $-\text{CH}_3$).

^1H NMR of **1b** *meso* ($\text{D}_2\text{O}:\text{CD}_3\text{CN}$ 1:1, 500 MHz, ppm) δ 8.28 (s, 2H, *ArH*), 8.06 (s, 2H, *ArH*), 8.02 (s, 2H, *ArH*), 7.98 (s, 1H, *ArH*), 7.74 (d, $J = 7.5$ Hz, 2H, *ArH*), 7.66 (m, 4H, *ArH*), 7.55 (d, $J = 7.5$ Hz, 2H, *ArH*), 6.33 (s, 2H, $-\text{CH}-$), 6.27 (s, 2H, $-\text{CH}-$), 3.82 (q, $J = 6.5$ Hz, 2H, $-\text{NHCH}_2-$), 2.05 (t, $J = 7.0$ Hz, 2H, $-\text{NHCH}_2\text{CH}_2-$), 1.75 (m, 18H, $-\text{CH}_2-$), 1.11 (t, $J = 7.5$ Hz, 3H, $-\text{CH}_3$).

Characterization of the Solution of Receptors **1b** by LC-MS

Scheme S2 Thiol oxidation to form receptor **1b** as a mixture of stereoisomers.

Figure S1. LC-MS analysis (negative ion mode) of the solution of receptors **1b** in DMSO.

Figure S2. A mass spectrum retention time 31 min (top) and 36.5 min (bottom) for receptor **1b meso** and **1b rac**, respectively.

Characterization of Receptor **1b rac** after Purification by Preparative HPLC

Figure S3. LC-MS analysis (negative ion mode) of the solution of receptor **1b rac** after purification by preparative HPLC.

Figure S4. A mass spectrum $t = 37$ min of receptor **1b rac** after purification by preparative HPLC.

Figure S5. Part of the TOCSY NMR spectrum of receptor **1b** *rac* (0.6 mM in D₂O (pD 8.7) : MeOD-*d*₄ 2:1).

Figure S6. Part of the ^1H NMR titration of $\text{MeOD-}d_4$ into a solution of receptor **1b** *rac* (1 mM in D_2O (pD 8.7)). The volume ratio of D_2O (pD 8.7) to $\text{MeOD-}d_4$ was: (a) 1:0; (b) 1:0.16; (c) 1:0.32; (d) 1:0.5; (e) 1:0.64; (f) 1:0.8; (g) 1:1; and (h) 1:1.16.

Figure S7. Part of the VT- ^1H NMR spectrum of receptor **1b** *rac* at 1 mM in D_2O (pD 8.7).

Re-Equilibration Experiments

To assign the diastereomers of receptor **1b** (*rac RR/RR* and *SS/SS*, *meso RR/SS*), we have studied the early stages of the re-equilibration of **1b** and its transformation into receptor **1a** of which we have previously assigned the diastereomers.⁴ Each diastereomer of receptor **1b** was isolated by preparative HPLC and dissolved in 1:1 solvent mixture of 10 mM borate buffer pH 9: acetonitrile (0.1 mg/mL of major or minor isomer). To this solution was added 1 equivalent of dithiol building block **viii**. The build up of small amounts of the expected **1a** isomers (Scheme S3) was monitored with time by HPLC. Separations were achieved by injecting 5 μ L of the mixture onto a Nucleodur C₁₈ column (4.6 cm $\times$ 250 mm, 100 Å, 5 μ m) at 45⁰C using a flow rate 1 mL/min. The following gradient was used:

Time (minutes)	Acetronitrile + 0.1%	H ₂ O + 0.1%
	Trifluoroacetic acid	Trifluoroacetic acid
0	50	50
20	50	50
25	75	25
60	75	25
65	50	50
75	50	50

The results were compared with the HPLC analysis of an independently generated mixture made from building blocks **2rac** and **viii**, which contains all stereoisomers of macrocycles **1a** which we have assigned previously (Scheme S3 and Figure S8).⁴

(A) Receptor **1b *rac***

(B) Receptor **1b *meso***

Scheme S3 Correlation between the diastereomers of **1b** with those of **1a** that are formed in the early stages of re-equilibration of receptor **1b** with building block **viii**.

Figure S8. HPLC chromatograms of the product mixture obtained upon re-equilibration of **(A)** receptor **1b** (major *rac* isomer; 0.1 mM) and **(B)** receptor **1b** (minor *meso* isomer; 0.1 mM) in the presence of building block **viii** (1 equivalent) in H₂O (pH 9):CH₃CN 1:1. The top traces represent the HPLC analysis of an independently generated mixture made from building blocks **2rac** and **viii**, which contains all stereoisomers of macrocycles **1a** which we have assigned previously.

Isothermal Titration Calorimetry Isothermal titration calorimetry was performed at 25°C using a MCS isothermal titration calorimeter (MicroCal, Northampton, MA, USA). All solutions were degassed before titrations. Titrations were performed in 10 mM borate buffer pH 9. The host concentration in the cell was 0.075 mM and the guest (0.75 mM) was titrated in 10 µL steps using 30 injections spaced at intervals of 200 seconds with a syringe stirring speed at 300 rpm. The binding constants were obtained using the one-site binding model provided in the ORIGIN software (version 2.9).

Dynamic Light Scattering Studies Dynamic light scattering measurements were performed on a Malvern Instruments Zetasizer Nano ZS. Samples were filtered through a membrane filter (0.45 µm) prior to measurements to remove any dust particles – control experiments without filtration gave essentially identical results confirming that this procedure does not remove parts of the sample.

Dynamic Light Scattering Data

Figure S9. Dynamic Light Scattering data for a solution of **1b** *rac* (0.1 mg/mL in 10 mM borate buffer pH 9). The graph represents the volume average taken over 14 measurements.

Figure S10. Dynamic Light Scattering data for a solution of **1b•5** (0.1 mg/mL of **1b** *rac* and 4 equivalent of **5**) in 10 mM borate buffer pH 9. The graph represents the volume average taken over 14 measurements.

Transmission Electron Microscopy TEM was conducted on a Philips CM100 electron microscope operating at 80 kV. A 300 mesh carbon-coated copper grid was placed on top of a drop of sample solution (0.1 mg/mL) for 30 minutes and then gently blotted with filter paper. The specimen was negatively stained using a drop of 2% PTA (pH 7) solution, incubated for 5 minutes, finally gently blotted and dried at room temperature.

Nile Red Solubilisation Nile red (0.10 mg) was added to 3mL of a solution of **1b•5** and equilibrated for 2 hours. The solution was filtered before measurements. UV-Vis experiments were carried out on a Hewlett Packard 8452A spectrophotometer. Fluorescence experiments were measured on an Aminco-Bowman spectrometer.

Figure S11. The emission spectrum of Nile Red in a solution of **1b•5** (0.1 mg/mL of **1b** *rac* and 4 equivalent of **5**) in 10 mM borate buffer pH 9 ($\lambda_{\text{ex}} = 570$ nm).

References

1. S. Otto, R. L. E. Furlan, and J. K. M. Sanders, *Science*, 2002, **297**, 590.
2. M. A. Petti, T. J. Sheppard, R. E. Barrans and D. A. Dougherty, *J. Am. Chem. Soc.*, 1988, **110**, 6825,
3. G. K. van der Wel, J. W. Wijnen and J. F. B. N. Engberts, *J. Org. Chem.*, 1996, **61**, 9001.
4. P. T. Corbett, manuscript in preparation