University of Groningen

Stereoselective Addition of Benzophenone to (C₅Me₅)Hf(2,3-dimethyl-1,3-butadiene)Cl

Ausema, Johannes B.; Hessen, Bart; Teuben, Jan H.

Published in:
 Organometallics

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
 Publisher's PDF, also known as Version of record

Publication date:
 1987

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Ausema, J. B., Hessen, B., & Teuben, J. H. (1987). Stereoselective Addition of Benzophenone to (C₅Me₅)Hf(2,3-dimethyl-1,3-butadiene)Cl. *Organometallics*, 6(6).

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Preliminary Communication

Recl. Trav. Chim. Pays-Bas **106**, 465-466 (1987)

0165-0513/87/08465-02\$1.00

STEREOSELECTIVE ADDITION OF BENZOPHENONE TO $(C_5Me_5)Hf(2,3\text{-DIMETHYL-1,3-BUTADIENE})Cl$

Johannes B. Ausema, Bart Hessen and Jan H. Teuben

Department of Inorganic Chemistry, University of Groningen, Nijenborgh 16, 9747 AG Groningen, The Netherlands

Abstract. The reaction of $(C_5Me_5)Hf(s\text{-}cis\text{-}C_6H_{10})Cl$ with benzophenone results in sequential insertion of two ketone molecules in the Hf-diene bond through adduct intermediates. The final 1,3-dioxa-2-hafnacyclonon-6-ene has a *cis*-configuration around the double bond, in contrast with related products from $(C_5H_5)_2Zr(diene)$.

In recent years, 1,3-butadiene complexes of the early transition metals (Group 4 and 5) have been the subject of much interest. Especially the $Cp_2M(diene)$ ($M = Zr, Hf$) 18-electron system has been investigated [1], and it became clear that the *s-cis* diene complexes behaved like σ^2, π -metallacyclopentene species, with pronounced nucleophilic character of the diene methylene carbon atoms. Thus insertion of polar unsaturated molecules like ketones and nitriles into the metal-diene bond is observed, with large regio- and stereoselectivity [1]. Here we wish to report the reactivity of the 14-electron *s-cis* 2,3-dimethyl-1,3-butadiene (C_6H_{10}) complex $Cp^*Hf(C_6H_{10})Cl$ (**1**, $Cp^* = \eta^5\text{-}C_5Me_5$) [2] with benzophenone, where the steric and electronic unsaturation of the mono(pentamethylcyclopentadienyl) system leads to a different stereoselectivity.

Yellow **1** reacts with one mole of Ph_2CO , initially to give a green 16 electron adduct, **1**. $OCPh_2$ [3], which in solution (hexane, $-15^\circ C$) rapidly forms the pale yellow singly inserted product $Cp^*Hf(OCPh_2-CH_2-CH_2-C(Me)=C(Me)CH_2)Cl$ (**2**) [4]. This compound is analogous to the products found in the metallocene Zr system [5]. **2** can react with a second equivalent of Ph_2CO , again initially forming an adduct, **2**. $OCPh_2$. This adduct is less stable than **1**. $OCPh_2$, and can only be detected by its transient, intense green colour. It rapidly reacts to form the colourless doubly inserted product $Cp^*Hf(OCPh_2-CH_2-CH_2-C(Me)=C(Me)-CH_2-CPh_2-O)Cl$ (**3**) [6].

Apart from the possibility to detect and even isolate intermediate adducts, the product formation in this system shows several differences with that in the Cp_2Zr -system. Firstly, in the zirconocene system doubly inserted products could only be obtained with dienes without substituents on the 2- and/or 3-position. Secondly, whereas the doubly inserted products in the zirconocene system have a *trans*-configuration around the double bond in the 1,3-dioxa-2-zirconacyclonon-6-ene, the configuration around the double bond in **3** is *cis*. This can be seen from the NMR-spectra of **3**, as the two methyl groups of the former 2,3-dimethyl-1,3-butadiene ligand are identical, indicating a mirror plane in the molecule, and thus a *cis*-configuration around the double bond. These differences in product formation are probably due to a different reaction mechanism for the second insertion step. The Cp_2Zr -2-oxacyclohept-5-ene is not accessible for the ketone, and reaction only takes place after rearrangement, through a bicyclic transition state leading to a *trans*-configuration in the final product (Scheme 1). This transition state is too severely hindered for products from 2- or 2,3-substituted dienes [1a]. The mono(pentamethylcyclopentadienyl) Hf-system is sterically hindered on one side of the molecule only, and an adduct can be formed. Through a 4-center transition state the second insertion can then take place without rearrangement or significant influence from the substituents on the 2,3-

Scheme 1

Scheme 2

-positions of the original diene (Scheme 2). This yields a product with a *cis*-arrangement around the double bond.

Preliminary results for the reaction of **1** with 2,6-xylylnitrile include formation of an adduct of much greater stability than the benzophenone adduct, and, upon heating this adduct in toluene at 60°C, disproportionation to yield equimolar amounts of **1** and a doubly inserted product. Our further attention will be directed to a more complete understanding of the selectivity in this kind of reactions, and also towards the reactivity of the Ti-analogue of **1** [2a], as the reactivity of Ti-butadiene complexes is virtually unexplored.

References and Notes

- [1] a. H. Yasuda, K. Tatsumi, and A. Nakamura, *Acc. Chem. Res.* **18**, 120 (1985) and references cited therein.
b. G. Erker, C. Krüger and G. Müller, *Adv. in Organometal. Chem.* **24**, 1 (1985) and references cited therein.
- [2] a. J. Blenkins, H.J. de Liefde Meijer, and J.H. Teuben, *Organometallics*, **2**, 1483 (1983).
b. J. Blenkins, B. Hessen, F. van Bolhuis, A.J. Wagner, and J.H. Teuben, *Organometallics* **6**, 459 (1987).
- [3] The adduct **1**.OCPPh₂ precipitates from pentane at -30°C and can be handled as a solid at 0°C. IR (KBr/silicon grease): 1553 cm⁻¹ (C=O), 110 cm⁻¹ lower than in free Ph₂CO.
- [4] **2**, ¹H-NMR (90 MHz, d⁶-benzene): δ 1.86 (s, 15H, Cp*); δ 1.72 (s, 3H, Me); δ 1.41 (s, 3H, Me); δ 3.01 (d, ²J 12.8 Hz, 1H, CH₂CO); δ 2.85 (d, ²J 12.8 Hz, 1H, CH₂CO); δ 1.00 (d, ²J 11.4 Hz, 1H, CH₂Hf, the other signal around 1.8 ppm partially obscured); δ 7.5-6.9 ppm (m, 10H, Ph). C₂₉H₃₅HfClO (613.53): calcd. C, 56.77; H, 5.75; Cl, 5.78; found C, 56.85; H, 5.62; Cl, 5.84.
- [5] a. Y. Kai, N. Kanehisa, K. Miki, M. Akita, H. Yasuda, and A. Nakamura, *Bull. Chem. Soc. Jpn.* **56**, 3735 (1983).
b. G. Erker, K. Engel, J.L. Atwood, and W.E. Hunter, *Ang. Chem.* **95**, 506 (1983).
- [6] **3**, ¹H-NMR (90 MHz, d⁶-benzene): δ 1.88 (s, 15H, Cp*); δ 1.27 (s, 6H, Me); δ 4.20 (d, ²J 13.8 Hz, 2H, CH₂); δ 2.84 (d, ²J 13.8 Hz, 2H, CH₂); δ 7.6-7.1 (m, 20H, Ph). ¹³C-NMR (22.6 MHz, d⁶-benzene): δ 11.7 (q, ²J 127 Hz, Cp*-Me); δ 20.10 (q, ²J 126 Hz, =C(CH₃)); δ 48.64 (t, ²J 126 Hz, CH₂); δ 90.00 (s, CO); δ 121.10 (s, Cp*-ring); δ 127.14 (d, ²J 158 Hz, Ph CH); δ 127.71 (d, ²J 157 Hz, Ph CH); δ 133.01 (s, C(Me)); δ 147.19 (s, Ph C); δ 150.65 (s, Ph C). C₄₂H₄₅HfClO₂ (795.74); calcd. C, 63.40; H, 5.70; Cl, 4.45; Hf, 22.43; found C, 63.68; H, 5.70; Cl, 4.50; Hf, 22.21.