University of Groningen

Asymmetric Strecker Synthesis of α -Amino Acids via a Crystallization-Induced Asymmetric Transformation Using (R)-Phenylglycine Amide as Chiral Auxiliary

Boesten, Wilhelmus H.J.; Seerden, Jean-Paul G.; Lange, Ben de; Dielemans, Hubertus J.A.; Elsenberg, Henk L.M.; Kaptein, Bernard; Moody, Harold M.; Kellogg, Richard M.; Broxterman, Quirinus B.

Published in:
Organic letters

DOI:
[10.1021/ol007042c](https://doi.org/10.1021/ol007042c)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2001

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Boesten, W. H. J., Seerden, J-P. G., Lange, B. D., Dielemans, H. J. A., Elsenberg, H. L. M., Kaptein, B., Moody, H. M., Kellogg, R. M., & Broxterman, Q. B. (2001). Asymmetric Strecker Synthesis of α -Amino Acids via a Crystallization-Induced Asymmetric Transformation Using (R)-Phenylglycine Amide as Chiral Auxiliary. *Organic letters*, 3(8). <https://doi.org/10.1021/ol007042c>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Asymmetric Strecker Synthesis of α -Amino Acids via a Crystallization-Induced Asymmetric Transformation Using (*R*)-Phenylglycine Amide as Chiral Auxiliary

Wilhelmus H. J. Boesten,[†] Jean-Paul G. Seerden,[†] Ben de Lange,^{*,†}
Hubertus J. A. Dielemans,[†] Henk L. M. Elsenberg,[†] Bernard Kaptein,[†]
Harold M. Moody,[†] Richard M. Kellogg,[‡] and Quirinus B. Broxterman^{*,†}

DSM Research Life Sciences-Organic Chemistry and Biocatalysis, P.O. Box 18,
6160 MD Geleen, and Syncom B.V., Kadijk 3, 9747 AT Groningen, The Netherlands

rinus.broxterman@dsm-group.com

Received December 22, 2000 (Revised Manuscript Received March 15, 2001)

ABSTRACT

Diastereoselective Strecker reactions based on (*R*)-phenylglycine amide as chiral auxiliary are reported. The Strecker reaction is accompanied by an *in situ* crystallization-induced asymmetric transformation, whereby one diastereomer selectively precipitates and can be isolated in 76–93% yield and $\text{dr} > 99/1$. The diastereomerically pure α -amino nitrile obtained from pivaldehyde ($\text{R}_1 = t\text{-Bu}$, $\text{R}_2 = \text{H}$) was converted in three steps to (*S*)-*tert*-leucine in 73% yield and $> 98\%$ ee.

The asymmetric synthesis of α -amino acids and derivatives is an important topic as a result of their extensive use in pharmaceuticals and agrochemicals and as chiral ligands. Many highly enantioselective approaches have been reported.¹ Industrial production of α -amino acids via the Strecker reaction is historically one of the most versatile methods to obtain these compounds in a cost-effective manner, making use of inexpensive and easily accessible

starting materials.² The Strecker reaction is usually followed by resolution of the racemic amino acid or amino acid amide obtained after hydrolysis of the amino nitrile.³ Either process leads to a maximum yield of 50% if the unwanted enantiomer is not racemized. In principle, asymmetric synthesis ap-

[†] DSM Research Life Sciences-Organic Chemistry & Biocatalysis. E-mail for Ben de Lange: Ben-B.Lange-de@dsm-group.com.

[‡] Syncom B.V.

(1) (a) Calmes, M.; Daunis, J. *Amino Acids* **1999**, *16*, 215. (b) Cativiela, C.; Díaz-de-Villegas, M. D. *Tetrahedron: Asymmetry* **1998**, *9*, 3517 (c) Williams, R. M. *Synthesis of Optically Active α -Amino Acids*; Pergamon: Oxford, 1989.

(2) (a) Kunz, H. In *Houben-Weyl: Stereoselective Synthesis*; Helmchen, G., Hoffmann, R. W., Mulzer, J., Schaumann, E., Eds.; Thieme Verlag: Stuttgart, 1995, 1931; Vol. E 21, D.1.4.4. (b) Shafran, Y. M.; Bakulev, V. A.; Mokrushin, V. S. *Russ. Chem. Rev.* **1989**, *58*, 148 (c) Strecker, A. *Ann. Chem. Pharm.* **1850**, *75*, 27.

(3) For reviews, see: resolution by aminoamidases. (a) Sonke, T.; Kaptein, B.; Boesten, W. H. J.; Broxterman, Q. B.; Schoemaker, H. E.; Kamphuis, J.; Formaggio, F.; Toniolo, C.; Rutjes, F. P. J. T. In *Stereoselective Biocatalysis*; Patel, R., Ed.; Dekker: New York, 2000; Chapter 2. (b) Resolution by acylases. Drauz, K.; Waldmann, H. In *Enzyme Catalysis in Organic Synthesis*; VCH: Weinheim, 1995; Vol. 1, p 393.

Table 1. Asymmetric Strecker Reactions of (*R*)-Phenylglycine Amide **1** and Pivalaldehyde **2**

entry	solvent	temp (°C)	time (h)	yield (%) ^a	dr (<i>R,S</i>)- 3 / <i>R,R</i>)- 3 ^b
1	MeOH	rt	20	80	65/35
2	MeOH/2-PrOH, 1/9 ^c	rt	22	51	99/1
3	2-PrOH	rt	22	84	88/12
4	2-PrOH/ <i>t</i> -BuOH, 4/1 ^c	rt	20	65	96/4
5	MeOH/H ₂ O, 35/1 ^c	rt	20	69	81/19
6	H ₂ O	55	24	81	85/15
7	H ₂ O	60	24	84	96/4
8	H ₂ O	65	24	84	98/2
9	H ₂ O	70	24	93	>99/1

^a Isolated yield after: evaporation of the solvent (entry 1) or filtration of precipitated amino nitrile **3** (entries 2–9). ^b The dr was determined by ¹H NMR spectroscopy. ^c Ratio in volume/volume.

proaches that lead to a maximum yield of 100% of a single enantiomer are more advantageous.

Recently several catalytic asymmetric Strecker reactions leading to N-protected amino nitriles in high ee's and high yields have been published.⁴ Alternatively, diastereoselective Strecker syntheses using a broad variety of chiral inducing agents, like α -arylethylamines,⁵ β -amino alcohols and derivatives,⁶ amino diols,⁷ sugar derivatives,⁸ and sulfinates⁹ have been reported to provide the α -amino nitriles with varying diastereoselectivities. A major drawback of these chiral auxiliaries can be cost and/or availability, because they are used in stoichiometric amounts and in principle lost during the conversion. Furthermore, in many cases the α -amino nitriles need to be purified in a separate step to obtain diastereomerically pure compounds. Purification requires, for example, crystallization or chromatography, which may lead to losses. An interesting solution to these problems would be a crystallization-induced asymmetric transformation,^{10,11} in which one diastereomer precipitates and the other epimerizes in solution via the corresponding imine. This would lead both to high yield and high diastereoselectivity in a practical one-pot procedure.

Recently, optically pure (*R*)-phenylglycine amide **1** became readily accessible as a result of application on an industrial scale as key intermediate in the enzymatic synthesis of β -lactam antibiotics.¹² Either aminopeptidase-catalyzed hydrolysis of racemic phenylglycine amide³ or asymmetric transformation of racemic phenylglycine amide with (*S*)-mandelic acid as resolving agent¹³ can be used to prepare **1**. Because of its ready availability on a large scale and its anticipated easy removal via catalytic hydrogenolysis, we decided to investigate the application of (*R*)-phenylglycine amide **1** as chiral auxiliary in asymmetric synthesis.

In this paper, the first two examples of the use of (*R*)-phenylglycine amide in asymmetric Strecker reactions are presented. Pivalaldehyde and 3,4-dimethoxyphenylacetone

have been used as starting materials, which lead, respectively, to enantiomerically enriched *tert*-leucine and α -methyl-dopa, two important nonproteogenic α -amino acids for pharmaceutical applications. In addition, *tert*-leucine has considerable utility as a chiral building block.¹⁴

The asymmetric Strecker reaction of (*R*)-phenylglycine amide **1**, pivalaldehyde **2** and HCN generated in situ from NaCN and AcOH was studied (Table 1). Amino nitriles (*R,S*)-**3** and (*R,R*)-**3** were obtained in 80% yield in a ratio of 65:35 by stirring an equimolar mixture of **1** (as AcOH salt)

(4) (a) Sigman, M. S.; Vachal, P.; Jacobsen, E. N. *Angew. Chem., Int. Ed. Engl.* **2000**, *39*, 1279. (b) Porter, J. R.; Wirsching, W. G.; Kuntz, K. W.; Snapper, M. L.; Hoveyda, A. H. *J. Am. Chem. Soc.* **2000**, *122*, 2657. (c) Ishitani, H.; Komiyama, S.; Hasegawa, Y.; Kobayashi, S. *J. Am. Chem. Soc.* **2000**, *122*, 762. (d) Vachal, P.; Jacobsen, E. N. *Org. Lett.* **2000**, *2*, 867. (e) Corey, E. J.; Grogan, M. *Org. Lett.* **1999**, *1*, 157.

(5) (a) Vincent, S. P.; Schleyer, A.; Wong, C.-H. *J. Org. Chem.* **2000**, *65*, 4440. (b) Wede, J.; Volk, F.-J.; Frahm, A. W. *Tetrahedron: Asymmetry* **2000**, *11*, 3231. (c) Juaristi, E.; Leon-Romo, J. L.; Reyes, A.; Escalante, J. *Tetrahedron: Asymmetry* **1999**, *10*, 2441. (d) Speelman, J. C.; Talma, A. G.; Kellogg, R. M.; Meetsma, A.; de Boer, A.; Beurskens, P. T.; Bosman, W. P. *J. Org. Chem.* **1989**, *54*, 1055. (e) Stout, D. M.; Black, L. A.; Matier, W. L. *J. Org. Chem.* **1983**, *48*, 5369.

(6) (a) Dave, R. H.; Hosangadi, B. D. *Tetrahedron* **1999**, *55*, 11295. (b) Ma, D.; Tian, H.; Zou, G. *J. Org. Chem.* **1999**, *64*, 120. (c) Chakraborty, T. K.; Hussain, K. A.; Reddy, G. V. *Tetrahedron* **1995**, *51*, 9179.

(7) Weinges, K.; Brachmann, H.; Stahnecker, P.; Rodewald, H.; Nixdorf, M.; Irngarter, H. *Liebigs Ann. Chem.* **1985**, 566.

(8) Kunz, H.; Sager, W.; Schanzenbach, D.; Decker, M. *Liebigs Ann. Chem.* **1991**, 649.

(9) Davis, F. A.; Fanelli, D. L. *J. Org. Chem.* **1998**, *63*, 1981.

(10) Only very few examples of crystallization-induced asymmetric transformations in Strecker reactions have been reported based on arylalkyl-methyl ketones: (a) Weinges, K.; Gries, K.; Stemmler, B.; Schrank, W. *Chem. Ber.* **1977**, *110*, 2098. (b) Weinges, K.; Klotz, K.-P.; Droste, H. *Chem. Ber.* **1980**, *113*, 710.

(11) For a broad discussion of crystallization-induced asymmetric transformation, see: Vedejs, E.; Chapman, R. W.; Lin, S.; Muller, M.; Powell, D. R. *J. Am. Chem. Soc.* **2000**, *122*, 3047 and references therein.

(12) Bruggink, A.; Roos, E. S.; de Vroom, E. *Org. Process Res. Dev.* **1998**, *2*, 128.

(13) (a) Boesten, W. H. J. European Patent Appl. EP 442584, 1991 (Chem. Abstr. **1992**, *116*, 42062r). (b) Boesten, W. H. J. European Patent Appl. EP 442585, 1991 (Chem. Abstr. **1992**, *116*: 42063s).

(14) Bommarius, A. S.; Schwarm, M.; Stingl, K.; Kottenhahn, M.; Huthmacher, K.; Drauz, K. *Tetrahedron: Asymmetry* **1995**, *6*, 2851.

with **2** and NaCN in MeOH overnight at room temperature, followed by evaporation of the solvent (entry 1). The diastereomeric ratio of (*R,S*)-**3** and (*R,R*)-**3** was determined by ^1H NMR on the basis of the relative integration between the *t*-Bu signals at 1.05 ppm for (*R,S*)-**3** and 1.15 ppm for (*R,R*)-**3**. The assignments have been made on the basis of the absolute configuration as established by X-ray analysis and conversion to (*S*)-*tert*-leucine (vide infra).

Because in methanol crystallization of amino nitrile **3** did not take place, first the solvent was varied in order to attempt to find conditions for a crystallization-induced asymmetric transformation. At a MeOH/2-PrOH ratio of 1/9 amino nitrile (*R,S*)-**3** was isolated in 51% yield and dr 99/1 (entry 2). Other combinations of alcoholic solvents failed to lead to a higher yield of precipitated (*R,S*)-**3** in high dr (entries 3 and 4). On further screening of solvents it was observed that upon addition of H_2O to the methanol solution selective precipitation of amino nitrile (*R,S*)-**3** occurred giving (*R,S*)-**3** and (*R,R*)-**3** in a ratio of 81:19 and 69% yield (entry 5). The asymmetric Strecker reaction was further studied in H_2O alone using temperature as a variable. The results of these experiments are given in Table 1 (entries 6–9). After addition of NaCN/AcOH at 23–28 °C to (*R*)-phenylglycine amide **1** and pivaldehyde **2** in H_2O , the mixture was heated to the indicated temperatures.

After approximately 24 h of stirring, the mixture was cooled to 30 °C and the precipitated amino nitrile filtered and analyzed by ^1H NMR to determine the dr. The results in Table 1 show that optimal results were achieved after 24 h of stirring in water at 70 °C. The amino nitrile (*R,S*)-**3** was obtained in 93% yield and a dr > 99/1 via a crystallization-induced asymmetric transformation (entry 6). At lower temperatures the epimerization reaction is slower.¹⁵

The crystallization-induced asymmetric transformation in water at 70 °C is verified further by the observed increase of the dr of (*R,S*)-**3** as a function of the reaction time (Figure 1). After 30 h the precipitated (*R,S*)-**3** was obtained with a dr > 99/1.

Figure 1. Crystallization-induced asymmetric transformation of amino nitrile **3** in water at 70 °C.

The observed diastereoselectivity in the asymmetric Strecker step via the crystallization-induced asymmetric transfor-

(15) At higher temperatures, lower yields of product were found, probably by degradation of amino nitrile.

Figure 2. Crystallization-induced asymmetric transformation of amino nitrile **3**.

mation can be explained as shown in Figure 2. Apparently, the *re*-face addition of CN^- to the intermediate imine **4** is preferred at room temperature in methanol and results in a dr 65/35. At elevated temperatures in water the diastereomeric outcome and yield of the process is controlled by the reversible reaction of the amino nitriles **3** to the intermediate imine and by the difference in solubilities of both diastereomers under the applied conditions.^{16,17}

The absolute configuration of amino nitrile (*R,S*)-**3** was confirmed by X-ray analysis as shown in Figure 3¹⁸ and by conversion to (*S*)-*tert*-leucine.

Figure 3. X-ray structure of amino nitrile (*R,S*)-**3**.

Conversion of the amino nitrile (*R,S*)-**3** to (*S*)-*tert*-leucine **7** was accomplished via the reaction sequence shown in Scheme 1. Hydrolysis of (*R,S*)-**3** to the diamide (*R,S*)-**5**

(16) For example, in the case of phenylacetone (not illustrated) it was found that in solution the initially formed minor isomer preferentially precipitated under crystallization conditions.

(17) For a discussion of asymmetric transformation of α -amino nitriles with mandelic acid, see: Hassan, N. A.; Bayer, E.; Jochims, J. C. *J. Chem. Soc., Perkin Trans. 1* **1998**, 3747.

(18) The crystal structure of (*R,S*)-**3** has been deposited at the Cambridge Crystallographic Data Center and allocated the deposition number CCDC 154034.

Scheme 1

proceeded smoothly in concentrated H_2SO_4 in high yield and without racemization.

Removal of the phenylacetamide group under 2 atm of H_2 with catalytic Pd/C afforded (*S*)-*tert*-leucine amide **6** in 90% yield. Finally, hydrolysis of the amide was accomplished by heating in 6 N HCl at 100 °C to give (*S*)-*tert*-leucine **7** in 86% yield and >98% ee. The absolute configuration assignment, (*S*), was made by comparison with an authentic sample.³ Obviously, other routes to convert the amino nitrile derivatives to the amino acid can be envisaged and are under investigation.

The crystallization-induced asymmetric transformation, using (*R*)-phenylglycine amide **1** as chiral auxiliary in diastereoselective Strecker reactions, was further explored with 3,4-dimethoxyphenylacetone **8** (Scheme 2).

The optimized asymmetric Strecker reaction of (*R*)-phenylglycine amide **1** (used as HCl salt) and an equimolar amount of 3,4-dimethoxyphenylacetone **8** in MeOH/ H_2O (6/1 v/v) gave, after addition of NaCN (30% aqueous solution) and stirring for 96 h at room temperature, the nearly diastereomerically pure (dr > 99/1) amino nitrile **9** as a solid in 76% isolated yield. The dr could easily be determined by

Scheme 2

^1H NMR analysis. It was found that in solution at room temperature an equilibrium of 55:45 exists between the two diastereomers (*R,S*)-**9** and (*R,R*)-**9**. Clearly, again a crystallization-induced asymmetric transformation has occurred.

In summary, (*R*)-phenylglycine amide **1** is an excellent chiral auxiliary in the asymmetric Strecker reaction with pivaldehyde or 3,4-dimethoxyphenylacetone. Nearly diastereomerically pure amino nitriles can be obtained via a crystallization-induced asymmetric transformation in water or water/methanol. This practical one-pot asymmetric Strecker synthesis of (*R,S*)-**3** in water leads to the straightforward synthesis of (*S*)-*tert*-leucine **7**. Since (*S*)-phenylglycine amide is also available, this can be used if the other enantiomer of a target molecule is required. More examples are currently under investigation to extend the scope of this procedure.¹⁹

Acknowledgment. Mr. A. Meetsma of the department of crystallography of the University of Groningen is acknowledged for the X-ray structure of (*R,S*)-**3**.

Supporting Information Available: Procedures and characterization data of all compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL007042C

(19) Several other amino nitriles could be obtained as crystalline materials from $\text{H}_2\text{O}/\text{MeOH}$ mixtures, e.g., $\text{R}_1 = \text{'Pr}$, $\text{R}_2 = \text{H}$; $\text{R}_1 = \text{Ph}$, $\text{R}_2 = \text{Me}$; $\text{R}_1 = \text{'Pr}$, $\text{R}_2 = \text{Me}$. Conditions are being sought to obtain also a crystallization-induced asymmetric transformation in these cases.