University of Groningen

Amino-substituted rylene dicarboximides and their quinoidal charge delocalization after deprotonation

Liu, Zhihong; Li, Chen; Wagner, Manfred; Avlasevich, Yuri; Herrmann, Andreas; Müllen, Klaus

Published in:
Chemical Communications

DOI:
[10.1039/b811252e](https://doi.org/10.1039/b811252e)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2008

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Liu, Z., Li, C., Wagner, M., Avlasevich, Y., Herrmann, A., & Müllen, K. (2008). Amino-substituted rylene dicarboximides and their quinoidal charge delocalization after deprotonation. *Chemical Communications*, 5028-5030. <https://doi.org/10.1039/b811252e>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Amino-substituted rylene dicarboximides and their quinoidal charge delocalization after deprotonation

Zhihong Liu, Chen Li, Manfred Wagner, Yuri Avlasevich, Andreas Herrmann and Klaus Müllen*

Max Planck Institute for Polymer Research, Ackermannweg 10, D-55128 Mainz, Germany

Supporting information

General: The solvents and chemicals used were of commercial grade. *N*-(2, 6-diisopropylphenyl)-4-bromo-naphthalene 1,8-dicarboximide (**6**), *N*-(2, 6-diisopropylphenyl)-9-bromo-perylene 3, 4-dicarboximide (**7**), *N*-(2, 6-diisopropylphenyl)-1,6,9,14-tetrakis(4-(1,1,3,3-tetramethylbutyl)phenoxy)-11-bromo-terrylene 3,4-dicarboximide (**8**) and *N*-(2, 6-diisopropylphenyl)-1,6-bis(4-(1,1,3,3-tetramethylbutyl)phenoxy)-13-bromo-quarterrylene 3, 4-dicarboximide (**9**) were synthesized as previously described¹⁻³. Column chromatography was performed on silica gel (Geduran Si₆₀, Merck). ¹H, ¹³C, H, H COSY and NOE NMR were recorded on Bruker DPX 250, DRX 500 and Avance 700 NMR spectrometers. Infrared spectra were obtained on a Nicolet FT IR320. FD mass spectra were recorded with a VG-instruments ZAB 2-SE-FPD instrument. MALDI-TOF mass spectra were recorded on a Bruker MALDI-TOF spectrometer. UV/Vis/NIR spectra were recorded in 1-cm quartz cuvette on a Perkin-Elmer Lambda 900 spectrophotometer. Elemental analyses were performed by the Department of Chemistry and Pharmacy of the University of Mainz.

General procedure for Buchwald condition.

The bromo-RI (0.18 mmol), 4-aminobenzonitrile (0.36 mmol), tris-(dibenzylideneacetone)-dipalladium(0) (17 mg, 0.018 mmol), tris-*tert*-butylphosphine (18 mg, 0.09mmol), sodium-*tert*-butoxide (67 mg, 0.69 mmol) and dry toluene (10 mL) were stirred at 80°C under argon overnight. After cooling, the mixture was evaporated in vacuo and purified by a column chromatography on silica gel using dichloromethane as an eluent.

***N*-(2,6-diisopropylphenyl)-4-(*p*-cyanophenyl)amino)naphthalene 1,8-dicarboximide (**2**)**

Using **6**, the general procedure afforded the title compound (81 mg, 95%). ¹H NMR (700MHz, acetone-*d*₆, 50°C): δ = 8.90(s, N-H, 1H), 8.76 (d, *J*=8.4 Hz, 1H), 8.63(d, *J*=7.2 Hz, 1H), 8.53(d, *J*=8.1 Hz, 1H), 7.87(t, *J*₁=7.6 Hz, *J*₂=8.1 Hz, 1H), 7.80(d, *J*=8.1 Hz, 1H), 7.74(d, *J*=8.3 Hz, 2H), 7.54 (d, *J*=8.2 Hz, 2H), 7.42(t, *J*=7.8 Hz, 1H), 7.33(d, *J*=7.8 Hz, 2H), 2.64 (sep, *J*=6.8 Hz, 2H), 1.13(d, *J*=6.8 Hz, 12H); H,H COSY NMR (700MHz, acetone-*d*₆, 50°C): coupling of δ=(8.76, 8.61, 7.87), (8.53, 7.80), (7.74, 7.54), (7.42, 7.33); ¹³C-NMR (62.5 MHz, acetone-*d*₆, 25°C): δ = 165.0, 164.3, 147.4, 146.8, 145.8, 134.6, 133.7, 132.7, 132.5, 131.3, 129.8, 129.7, 127.1, 125.4, 124.5, 124.0, 120.0, 119.7, 116.6, 114.4, 105.4, 30.6, 24.2 ppm. IR: ν = 3315, 3068, 2962, 2929, 2869, 2219, 1700, 1648, 1575, 1508, 1463, 1357, 1309, 1236, 1164, 1056, 1010, 917, 831, 777 cm⁻¹; UV-Vis (acetone) λ_{max}, nm (ε): 431 (21300); MS (FD): *m/z*, 473.4 (100%), M⁺; Found: C, 78.81; H, 5.87; N, 8.47 %. Calc. for C₃₁H₂₇N₃O₂: C, 78.62; H, 5.75; N, 8.87 %.

The NMR data of **2'** (deprotonation by NaOH). ¹H NMR (500MHz, acetone-*d*₆, 25°C): δ = 8.83(d, *J*=7.9 Hz, 1H), 8.27(d, *J*=7.4 Hz, 1H), 7.82(d, *J*=9.1 Hz, 1H), 7.47(d, *J*=8.5 Hz, 2H), 7.29(t, *J*=7.7 Hz, 1H), 7.24(t, *J*₁=7.7 Hz, *J*₂=7.6 Hz, 1H), 7.21(d, *J*=7.7 Hz, 2H), 7.08(d, *J*=8.5 Hz, 2H), 6.52 (d, *J*=9.1 Hz, 1H), 2.64 (sep, *J*=6.8 Hz, 2H), 1.13(d, *J*=6.8 Hz, 12H).

***N*-(2, 6-diisopropylphenyl)-9-(*p*-cyanophenyl)amino)perylene 3,4-dicarboximide (**3a**)**

Using **7**, the general procedure afforded the title compound (90 mg, 84%). ¹H NMR (700MHz, acetone-*d*₆, 50°C): δ = 8.78(d, *J*=7.5 Hz, 1H), 8.72(d, *J*=8.0 Hz, 1H), 8.70(d, *J*=8.3 Hz, 1H), 8.63(d, *J*=8.0 Hz, 1H), 8.60(d, *J*=8.0 Hz, 1H), 8.58(d, *J*=8.0 Hz, 1H), 8.49(s, 1N-H), 8.35(d, *J*=8.4 Hz, 1H), 7.76-7.74(m, 2H), 7.66(d, *J*=8.5 Hz, 2H), 7.44(t, *J*=7.9 Hz, 1H), 7.36-7.33(m, 4H), 2.83 (sep, *J*=6.8 Hz, 2H), 1.16(d, *J*=6.9 Hz, 12H); H,H COSY NMR (700MHz, acetone-*d*₆, 50°C): coupling of δ=(8.78, 8.35, 7.76), (8.70, 7.76), (8.72, 8.60), (8.63, 8.58), (7.66, 7.36), (7.44,7.36), (2.83, 1.16); NOE NMR (700MHz, acetone-*d*₆, 50°C): coupling of δ=(8.78, 8.72), (8.70, 8.63), (8.49, 7.36), (8.49, 8.35), (7.76, 7.36); ¹³C-NMR (62.5 MHz, CD₂Cl₂ 25°C): δ = 164.6, 164.4, 148.0, 146.5, 139.8, 137.8, 134.3, 132.3, 132.12, 132.0, 130.2, 129.7, 128.6, 127.4, 125.7, 125.2, 125.0, 124.7, 124.4, 121.5, 120.8, 120.5, 120.0, 119.7, 118.7, 117.2, 103.6, 29.5, 24.1 ppm. IR: ν = 3328, 3064, 2962, 2929, 2867, 2217, 1689, 1644, 1563, 1504, 1351, 1288,

1172, 1058, 971, 910, 804, 750 cm^{-1} ; UV-Vis (acetone) λ_{max} , nm (ϵ): 550(34800); MS (FD): m/z , 598.0 (100%), M^+ ; Found: C, 81.75; H, 5.42; N, 6.98 %. Calc. for $\text{C}_{41}\text{H}_{31}\text{N}_3\text{O}_2$: C, 82.39; H, 5.23; N, 7.03 %.

The NMR data of **3a'** (deprotonation by NaOH). ^1H NMR (700MHz, acetone- d_6 , 25°C): δ =8.81 (d, J =7.7 Hz, 1H), 8.65(d, J =7.5 Hz, 1H), 8.29(d, J =8.3 Hz, 1H), 8.18(d, J =8.4 Hz, 1H), 8.09(d, J =8.8 Hz, 1H), 8.06(d, J =9.7 Hz, 1H), 7.63(d, J =8.6 Hz, 1H), 7.57(d, J =8.4 Hz, 2H), 7.45(t, J =7.7 Hz, 1H), 7.32(t, J =7.6 Hz, 1H), 7.25(d, J =7.8 Hz, 2H), 7.10(d, J =8.4 Hz, 2H), 6.60(d, J =9.4 Hz, 1H), 2.83 (sep, J =6.8 Hz, 2H), 1.16(d, J =6.9Hz, 12H); H,H COSY NMR (700MHz, acetone- d_6 , 25°C): coupling of δ =(8.81, 8.65, 7.45), (8.29, 8.18), (8.09, 7.63), (8.06, 6.60), (7.57, 7.10), (7.32, 7.25), (2.83, 1.16); NOE NMR (700MHz, acetone- d_6 , 25°C): coupling of δ =(8.65, 8.18), (8.06, 7.63), (7.10, 6.60).

***N*-(2, 6-diisopropylphenyl)-9-(*p*-octylphenyl)amino)perylene 3,4-dicarboximide (**3b**)**

Using **7** and *p*-octylaniline instead of 4-aminobenzonitrile, the general procedure afforded the title compound (95 mg, 77%). ^1H NMR (700MHz, acetone- d_6 , 50°C): δ = 8.75 (d, J =7.3 Hz, 1H), 8.63 (d, J =8.1 Hz, 1H), 8.55 (d, J =7.9 Hz, 1H), 8.54 (d, J =8.6 Hz, 1H), 8.50 (d, J =8.1 Hz, 1H), 8.48 (d, J =8.5 Hz, 1H), 8.42 (d, J =8.2 Hz, 1H), 8.12 (s, 1N-H), 7.73 (t, J =8.2 Hz, 1H), 7.45 (t, J =7.9 Hz, 1H), 7.38 (d, J =8.5 Hz, 1H), 7.34 (d, J =7.9 Hz, 2H), 7.34 (d, J =8.4 Hz, 2H), 7.29 (d, J =8.4 Hz, 2H), 2.87-2.70 (sep, J =6.8 Hz, 2H), 2.67-2.61 (t, J =7.7 Hz, 2H), 1.68-1.62 (t, J =7.4 Hz, 2H), 1.34-1.30 (m, 10H), 1.18 (d, J =6.9 Hz, 12H), 0.92-0.87ppm (t, J =7.1Hz, 3H); H,H COSY NMR (700MHz, acetone- d_6 , 50°C): coupling of δ =(7.73, 8.75, 8.48), (7.38, 8.54), (7.29, 7.34), (7.45, 7.34), (8.50, 8.42), (8.55, 8.63), (2.87, 1.18), (2.67, 1.68), (1.62, 1.34), (1.30, 0.92); ^{13}C -NMR (62.5 MHz, CD_2Cl_2 25°C): δ =164.71, 164.55, 146.52, 144.34, 139.16, 138.96, 138.71, 138.17, 132.27, 131.88, 131.07, 129.97, 129.76, 129.16, 129.56, 129.16, 126.68, 126.47, 126.47, 125.11, 125.02, 124.35, 123.61, 121.84, 120.92, 120.83, 119.88, 118.69, 118.48, 111.08, 35.74, 32.30, 32.06, 29.89, 29.73, 29.69, 29.45, 24.14, 24.10, 23.08, 14.27 ppm. IR: ν = 3372, 2964, 2924, 2854, 1690, 1650, 1566, 1514, 1354, 1284, 804,750 cm^{-1} ; UV-Vis (acetone) λ_{max} , nm (ϵ):

596 (35233); MS (FD): m/z 683.5 (100%), M^+ ; Found: C, 83.39; H, 7.06; N, 3.96 %. Calc. for $C_{48}H_{48}N_2O_2$: C, 84.17; H, 7.06; N, 4.09 %.

The NMR data of **3b'** (deprotonation by NaOH): 1H NMR (250MHz, acetone- d_6 , 25 °C): δ = 8.83 (d, $J=7.6$ Hz, 1H), 8.67 (d, $J=8.5$ Hz, 1H), 8.25 (d, $J=8.5$ Hz, 1H), 8.11 (d, $J=8.7$ Hz, 1H), 7.99 (d, $J=9.0$ Hz, 1H), 7.94 (d, $J=10.1$ Hz, 1H), 7.48 (d, $J=9.0$ Hz, 1H), 7.46 (t, $J=7.7$ Hz, 1H), 7.32 (t, $J=7.9$ Hz, 1H), 7.25 (d, $J=7.9$ Hz, 2H), 7.16 (d, $J=8.4$ Hz, 2H), 6.89 (d, $J=8.4$ Hz, 2H), 6.52 (d, $J=10.0$ Hz, 1H), 2.77-2.66 (sep, $J=6.9$ Hz, 4H), 2.63-2.56 (t, $J=7.4$ Hz, 2H), 1.64-1.59 (t, $J=7.4$ Hz, 2H), 1.34-1.30 (m, 10H), 1.13-1.10 (d, $J=6.9$ Hz, 12H), 0.92-0.86 ppm (t, $J=7.0$ Hz, 3H); H,H COSY NMR (700MHz, acetone- d_6 , 25°C): coupling of δ =(7.50, 8.83, 8.67), (6.87, 7.12), (6.52, 6.48), (7.35, 7.25), (7.99, 7.48), (8.11, 8.25), (2.77, 1.13), (2.63, 1.64), (1.64, 1.34), (1.30, 0.92).

***N*-(2, 6-diisopropylphenyl)-1,6,9,14-tetrakis(4-(1,1,3,3-tetramethylbutyl)phenoxy)-11-(*p*-cyanophenyl)amino)terrylene 3,4-dicarboximide (4)**

Using **8**, the general procedure afforded the title compound (28 mg, 55%). 1H NMR (250MHz, acetone- d_6 , 25°C): δ =9.50-9.45 (dd, 2H), 9.22-9.13 (dd, 2H) 8.35(d, $J=8.9$ Hz, 1H), 8.17(d, $J=8.9$ Hz, 1H), 7.68-7.52(m, 10H), 7.46(d, $J=7.2$ Hz, 2H), 7.38-7.28(m, 7H), 7.39-7.11(m, 6H), 2.92(sep, $J=6.8$ Hz, 2H), 1.95-1.89 (q, 8H), 1.55-1.49 (q, 24H), 1.23-1.19 (q, 12H), 1.01(d, $J=6.9$ Hz, 12H), 0.86-0.83ppm (q, 24H); IR: ν = 3326, 3054, 2952, 2923, 2865, 2221, 1698, 1594, 1502, 1365, 1303, 1211, 1170, 1052, 1014, 958, 829, 727 cm^{-1} ; UV-Vis (acetone) λ_{max} , nm (ϵ): 668(41900); MS (FD): m/z , 1539.3 (100%), M^+ .

***N*-(2, 6-diisopropylphenyl)-1,6-bis(4-(1,1,3,3-tetramethylbutyl)phenoxy)-13-(*p*-cyanophenyl)amino)quarterrylene 3,4-dicarboximide (5)**

Using **9**, the general procedure afforded the title compound (25 mg, 53%). IR: ν =3359, 3054, 2954, 2925, 2865, 2211, 1693, 1577, 1500, 1384, 1317, 1265, 1209, 1170, 1064, 1014, 806, 744 cm^{-1} ; UV-Vis (acetone) λ_{max} , nm (ϵ): 751(51200); MS (FD): m/z , 1255.3 (100%), M^+ .

Fig. S1. ^1H NMR spectra of **2** and **2'** (after deprotonation by NaOH): in acetone- d_6 .

Table S1. Chemical shifts and H,H coupling constants of **2** before or after deprotonation in acetone- d_6 .

Proton Items	H_a	H_b	H_c	H_d	H_e	H_f	H_j	H_l	H_m	H_n
δ_1 , ppm	7.74	7.54	7.80	8.53	7.42	7.33	8.91	8.76	7.87	8.63
J_1 , Hz	8.3	8.3	8.1	8.1	7.8	7.8	X	8.4		7.2
δ_2 , ppm	7.47	7.08	6.52	7.82	7.29	7.21		8.27	7.24	8.83
J_2 , Hz	8.5	8.5	9.1	9.1	7.7	7.7		7.4		7.9
$\delta_1 - \delta_2$, ppm	0.27	0.44	1.28	0.71	0.13	0.12		0.49	0.63	-0.20

δ_1 , δ_2 , J_1 and J_2 denote the chemical shift and H,H coupling constants before and after deprotonation, respectively.

Fig. S2. ^1H NMR spectra of **3a** and **3a'** (after deprotonation by NaOH) in acetone- d_6

Table S1. Chemical shifts and H_iH_j coupling constants of **3a** before or after deprotonation in acetone- d_6 .

Proton Items	H_a	H_b	H_c	H_d	H_e	H_f	H_g	H_h	H_i	H_j	H_k	H_l	H_m	H_n
δ_1 , ppm	7.66	7.35	7.75	8.70	7.44	7.35	8.62	8.58	8.60	8.49	8.72	8.78	7.75	8.35
J_1 , Hz	8.5	8.5	8.3	8.3	7.9	7.9	8.0	8.0	8.0		8.0	7.5		8.4
δ_2 , ppm	7.57	7.10	6.60	8.06	7.32	7.25	7.63	8.09	8.29		8.18	8.65	7.45	8.81
J_2 , Hz	8.4	8.4	9.4	9.7	7.6	7.8	8.6	8.8	8.3		8.4	7.5		7.7
$\delta_1 - \delta_2$, ppm	0.09	0.25	1.15	0.64	0.12	0.10	0.99	0.49	0.11		0.54	0.13	0.30	-0.46

δ_1 , δ_2 , J_1 and J_2 denote the chemical shift and H_iH_j coupling constants before and after deprotonation, respectively.

Fig. S3. ^1H NMR spectra of **3b** and **3b'** (after deprotonation by NaOH) in acetone- d_6 .

Table S1. Chemical shifts and H,H coupling constants of **3b** and **3b'** (after deprotonation by NaOH) in acetone- d_6 .

Proton Items	H _a	H _b	H _c	H _d	H _e	H _f	H _g	H _h	H _i	H _j	H _k	H _l	H _m	H _n
δ_1 , ppm	7.29	7.34	7.38	8.54	7.44	7.34	8.50	8.42	8.55	8.12	8.63	8.75	7.71	8.48
J_1 , Hz	8.4	8.4	8.5	8.5	7.9	7.9	8.2	8.2	8.1		8.1	7.3		8.5
δ_2 , ppm	6.87	7.16	6.52	7.94	7.35	7.25	7.48	7.99	8.25		8.11	8.67	7.46	8.83
J_2 , Hz	8.2	8.2	10.0	10.0	7.5	7.5	9.0	9.0	8.5		8.5	8.5		7.6
$\delta_1 - \delta_2$, ppm	0.42	0.18	0.86	0.60	0.09	0.09	1.02	0.43	0.30		0.50	0.08	0.25	-0.35

δ_1 , δ_2 , J_1 and J_2 denote the chemical shifts and H,H coupling constants before and after deprotonation, respectively.

For comparison, the absorption spectrum of the radical cation of **3b**, generated by chemical oxidation with SbCl_5 following the previously reported procedure⁴, was measured. It shows a sharp absorption peak at 710 nm and a very broad and weak band above 1000 nm in the NIR region.

Fig. S4. The absorption spectra of the radical cation of **3b** generated by chemical oxidization upon more and more SbCl_5 addition. (SbCl_5 stock solution in dichloromethane is 0.001M)

Fig. S5. The absorption change of **3b** deprotonated by BuONa in acetone after 3 days

Figure S6. Absorption switches of **3b** in acetone for 5 cycles

References

1. F. Nolde, J.Q. Qu, C. Kohl, N. G. Pschirer, E. Reuther, K. Müllen, *Chem. Eur. J.* **2005**, *11*, 3959-3967.
2. N. G. Pschirer; J.Q. Qu; M. Koenemann; WO 2006117383
3. Y. Avlasevich, K. Müllen, *J. Org. Chem.* **2007**, *72*, 10243-10246.
4. G. Zhou, M. Baumgarten, K. Müllen, *J. Am. Chem. Soc.* **2007**, *129*, 12211-12221.

^1H NMR (250 MHz, 25 °C) spectrum of **4** in acetone- d_6

MALDI-TOF-MS spectrum of **5**

^1H , ^1H COSY NMR (700MHz, 50°C) spectrum of **2** in acetone- d_6 .

^1H , ^1H COSY NMR (700MHz, 50°C) spectrum of **3a** in acetone- d_6 .

NOE NMR (700MHz, 50°C) spectrum of **3a** in acetone- d_6

H, H COSY NMR (700MHz, 25°C) spectrum of **3a'** (deprotonated by NaOH) in acetone-d₆

NOE NMR (700MHz, 25°C) spectrum of **3a'** (deprotonated by NaOH) in acetone- d_6

H, H COSY NMR (700MHz, 50°C) spectrum of **3b** in acetone-d₆