University of Groningen

P.303 Adjusting ruminative thinking? Effects of positive fantasizing vs. stress induction on perseverative cognition in individuals with varying vulnerability for depression

Besten, M.; Van Tol, M. J.; Van Rij, J. C.; Van Vugt, M. K.

Published in:
European Neuropsychopharmacology

DOI:
[10.1016/j.euroneuro.2020.09.228](https://doi.org/10.1016/j.euroneuro.2020.09.228)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Besten, M., Van Tol, M. J., Van Rij, J. C., & Van Vugt, M. K. (2020). P.303 Adjusting ruminative thinking? Effects of positive fantasizing vs. stress induction on perseverative cognition in individuals with varying vulnerability for depression. *European Neuropsychopharmacology*, 40, S174-S175.
<https://doi.org/10.1016/j.euroneuro.2020.09.228>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

gender and education level) underwent 3T fMRI scanning during an Emotion Regulation Task. During this task, participants were instructed to passively attend to neutral, negative or positive images, or to downregulate negative, and upregulate positive images, using cognitive reappraisal techniques. Furthermore, they underwent assessment of attentional biases (using a modified attentional dot-probe task with both disengagement and engagement bias indices for negative and positive stimuli [4]) and of rumination on both negative (Rumination on Sadness Scale) and positive (Responses to Positive Affect Scale) content. Following standard preprocessing and event-related modeling of fMRI-data (using SPM12, implemented in MATLAB(R2015a)), contrast images reflecting brain activation during attend (>fixation) and regulate (>attend) conditions were entered in nonparametric permutation-based group comparisons and multiple regression analyses with attentional bias and rumination scores as predictors (using FSL Randomise). Effects were considered significant at $p < 0.05$, TFCE-corrected.

rMDD patients showed no attentional biases, but reported higher rumination on sadness and dampening of positive affect, and lower self-related savoring of positive affect than HC. During implicit processing emotional information, rMDD patients showed decreased activation in the occipital cortex, precuneus, posterior cingulate cortex (PCC), inferior frontal gyrus and dorsolateral prefrontal cortex (DLPFC) than HC. During explicit emotion regulation (vs attending), rMDD patients showed increased precuneus, PCC, frontal pole, and amygdala activity for negative emotions, and decreased right insula activity for positive emotions, though below corrected threshold (uncorrected $p < 0.001$). Within rMDD patients, during downregulating negative emotions, higher rumination was related to lower PCC activation, and lower savoring of positive affect was related to higher amygdala, hippocampus, anterior cingulate cortex, DLPFC, precentral gyrus and occipital lobe activation. During upregulating positive emotions, higher dampening of positive affect was related to higher right ventral anterior insula activation.

Our results suggest that rMDD patients tend to engage more in negative rumination styles and less in positive rumination styles, and show abnormal frontolimbic activation during processing and regulating emotional information. Inadequate rumination and regulation styles in rMDD patients might relate to aberrant self-related processing and regulatory control over emotional processing during negative emotion regulation, and interoceptive awareness during positive emotion regulation. Understanding these neurocognitive abnormalities in rMDD may add to the clinical improvement of preventive treatment.

No conflict of interest

References

- [1] Disner, S., Beevers, E., Haigh, E.A.P., Beck, A.T., 2011. Neural mechanisms of the cognitive model of depression. *Nature Reviews Neuroscience* 12, 467-477.
- [2] Marchetti, I., Koster, E.H.W., Sonuga-Barke, E.J., De Raedt, R., 2012. The Default Mode Network and recurrent depression: a neurobiological model of cognitive risk factors. *Neuropsychology Review* 22, 229-251.
- [3] Van Kleef, R.S., Bockting, C.L.H., Van Valen, E., Aleman, A., Marsman, J.B.C., Van Tol, M.J., 2019. Neurocognitive working mechanisms of the prevention of relapse in remitted recurrent depression (NEWPRIDE): protocol of a randomized controlled neuroimaging trial of preventive cognitive therapy. *BMC Psychiatry* 19 (1), 409.
- [4] Grafton, B., MacLeod, C., 2014. Enhanced probing of attentional bias: the independence of anxiety-linked selectivity in attentional engagement with and disengagement from negative information. *Cognition and Emotion* 28 (7), 1287-1302.

doi: [10.1016/j.euroneuro.2020.09.227](https://doi.org/10.1016/j.euroneuro.2020.09.227)

P.303

Adjusting ruminative thinking? Effects of positive fantasizing vs. stress induction on perseverative cognition in individuals with varying vulnerability for depression

M. Besten¹, M.J. Van Tol¹, J.C. Van Rij², M.K. Van Vugt²

¹University Medical Center Groningen, Cognitive Neuroscience Center, Groningen, The Netherlands

²University of Groningen, Artificial Intelligence and Cognitive Engineering, Groningen, The Netherlands

Background: Key characteristics of Major Depressive Disorder are repetitive negative thinking and rumination (i.e. perseverative cognition [PC]). This maladaptive type of thinking has been associated with a high risk to develop and maintain Major Depressive Disorder [1,2]. Because of the relation between PC and depressive relapse, targeting PC could be a powerful way of preventing recurrence of depression. A therapy successful in preventing depressive relapse and reducing depressive symptoms is the Preventive Cognitive Therapy [3]. One of the core techniques in the Preventive Cognitive Therapy is positive fantasising, a positive-mood inducing technique aimed at enhancing positive attitudes and/or beliefs by positive future-related thinking. Whereas positive fantasising may reduce PC, stress may increase PC. In our study, we contrasted the effects of a single session of positive fantasising and stress induction on PC, quantified in a novel task-based assay, to investigate whether positive fantasising is potent in affecting PC in contrast to PC following stress-induction in individuals varying in their vulnerability for depression.

Methods: Groups high ($n=40$) and low ($n=40$) on neuroticism and worrying, measured using the Neuroticism scale of the NEO-Five-Factor Inventory and the Penn State Worry Questionnaire indicating vulnerability for negative affect and depression, performed a Sustained Attention to Response Task after a single session of positive fantasising and a single session of stress induction in a cross-over design. The Sustained Attention to Response Task is a boring go-/no-go task with interspersed self-report questions about the content of thoughts, frequently used in mind-wandering studies. Affective states were measured before and after the interventions using the Positive and Negative Affect Schedule.

Findings: A paired t-test showed increased negative affect after stress and increased positive and reduced negative affect after fantasising compared to baseline. Additionally,

thoughts were more on-task and future-related, easier to disengage from and less negative after fantasising compared to after stress. However, these effects decreased over time and were only found when fantasising was the first intervention. No significant interaction was found in PC between intervention (after stress vs. after fantasising) and group (high vs. low vulnerability for depression).

Discussion: Results indicate that PC can be adjusted in individuals both characterised by a high and low vulnerability for depression, and that these effects on PC can be measured behaviorally. Interestingly, changes in PC were only found when fantasising was the first intervention, suggesting that fantasising makes thought content more reactive to subsequent negative affect. Fantasising furthermore had beneficial effects on both positive and negative self-reported affect. These results suggest changing PC by interventions such as positive fantasising may be potent in reducing the vulnerability for depression. Future research should investigate whether positive fantasising could serve as an intervention to treat or prevent recurrence of depression through manipulation of PC.

No conflict of interest

References

- [1] Brosschot, J.F., Verkuil, B., Thayer, J.F., 2010. Conscious and unconscious perseverative cognition: Is a large part of prolonged physiological activity due to unconscious stress? *Journal of Psychosomatic Research* 69, 407-416. doi:10.1016/j.jpsychores.2010.02.002.
- [2] Nolen-Hoeksema, S., Wisco, B.E., Lyubomirsky, S., 2008. Rethinking Rumination. *Perspectives on Psychological Science* 3, 400-424. doi:10.1111/j.1745-6924.2008.00088.x.
- [3] Bockting, C.L.H., Spinhoven, P., Wouters, L.F., Koeter, M.W.J., Schene, A.H., 2009. Long-Term Effects of Preventive Cognitive Therapy in Recurrent Depression. *The Journal of Clinical Psychiatry* 70, 1621-1628. doi:10.4088/jcp.08m04784blu.

doi: 10.1016/j.euroneuro.2020.09.228

P.304

Aberrant psychophysiological and cognitive responses to infant signals of emotion in mothers with affective disorders and implications for the infant

A. Bjertrup¹, M. Moszkowicz², I. Pedersen³,
A. Kjærbye-Thygesen⁴, R. Nielsen⁵, C. Parsons⁶,
L. Kessing⁷, A.K. Pagsberg⁸, M. Væver³, K. Miskowiak¹

¹Copenhagen Affective Disorders research Center CADIC, Psychiatric Centre Copenhagen- Faculty of Health and Medical Sciences- University of Copenhagen- Copenhagen- Denmark, Copenhagen, Denmark

²Child and Adolescent Psychiatric Center- Infant Psychiatric Unit, Faculty of Health and Medical Sciences- University of Copenhagen, Copenhagen, Denmark

³Center for Early Intervention and Family Studies, Department of Psychology- University of Copenhagen, Copenhagen, Denmark

⁴Department of Obstetrics and Gynecology, Faculty of Health and Medical Sciences- University of Copenhagen, Hvidovre, Denmark

⁵Aalborg University Hospital, Psychiatry- North Denmark Region, Aalborg, Denmark

⁶Interacting Minds Center, Department of Clinical Medicine- Aarhus University, Aarhus, Denmark

⁷Copenhagen Affective Disorders research Center CADIC, Psychiatric Centre Copenhagen- Faculty of Health and Medical Sciences- University of Copenhagen, Copenhagen, Denmark

⁸Child and Adolescent Mental Health Centre- Mental Health Services Capital Region of Denmark, Faculty of Health and Medical Sciences- University of Copenhagen, Copenhagen, Denmark

Background: Affective disorders are highly heritable, but it is unclear how subtle mother-infant interaction adversity contributes to the transmission of risk. Healthy mothers show adaptive neurocognitive changes indicating that they “tune in” to infants [1,2]. However, core features of affective disorders are aberrant neurocognitive and psychophysiological responses to emotional stimuli. Whereas unipolar disorder (UD) is associated with negative cognitive bias, which applies to processing of infant stimuli among mothers in a depressive state [3], emerging evidence indicate positively biased emotion processing in bipolar disorder (BD) [4]. Biased emotion processing may influence mother-infant interactions and have adverse effects on infant development [3,5].

Aim: This study aimed to investigate psychophysiological and cognitive responses to emotional infant stimuli and their relation to mother-infant interaction and infant development for mothers with BD or UD in full or partial remission compared with healthy control mothers (HC).

Methods: Mothers and their infants were assessed during home visits around four months after birth. First, mothers were interviewed with clinical rating scales to ensure that mothers with BD or UD were in full or partial remission at the time of testing. Then, mothers' psychophysiological and cognitive responses to emotional infant stimuli were assessed on computerised tasks during which their facial expression, galvanic skin responses (GSR) and eye-gazes and fixations were recorded. Lastly, we assessed infant development and mother-infant-interaction. Group and interaction effects were analysed using one-way and repeated measures ANOVA, and associations were investigated with correlation analyses.

We included 85 mothers: 27 with BD, 13 with UD and 36 who were healthy, and their infants.

Results: Mothers with affective disorders had fewer GSR peaks in response to infant videos ($F(1,70)=4.86$, $p=0.03$, $\eta^2=0.07$) and spent less time gazing and fixating at infant videos and faces (videos: gaze: $F(1,58)=20.44$, $p < 0.001$, $\eta^2=0.26$, fixation: $F(1,58)=4.53$, $p=0.04$, $\eta^2=0.07$; images: gaze: $F(1,72)=17.73$, $p < 0.001$, $\eta^2=0.20$). Mothers with BD showed more incongruent positive facial expressions to infant distress vs. laughter videos ($F(1,58)=4.47$, $p=0.04$, $\eta^2=0.07$) and rated infant cry less negatively than HC mothers ($F(2.0,120.5)=3.37$, $p=0.04$, $\eta^2=0.07$; $t=2.37$, $df=60$, $p=0.02$). Mothers with UD displayed more negative facial expression when listening to infant