University of Groningen

The effectiveness of rituximab in pemphigus, and the benefit of additional maintenance infusions

Rashid, Hanan; Lamberts, Aniek; van Maanen, Dian; Bolling, Maria C; Diercks, Gilles F H; Pas, Hendri H; Jonkman, Marcel F; Horváth, Barbara

Published in:
Journal of the American Academy of Dermatology

DOI:
[10.1016/j.jaad.2020.06.024](https://doi.org/10.1016/j.jaad.2020.06.024)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Rashid, H., Lamberts, A., van Maanen, D., Bolling, M. C., Diercks, G. F. H., Pas, H. H., Jonkman, M. F., & Horváth, B. (2020). The effectiveness of rituximab in pemphigus, and the benefit of additional maintenance infusions: daily practice data from a retrospective study. *Journal of the American Academy of Dermatology*, 83(5), 1503-1505. <https://doi.org/10.1016/j.jaad.2020.06.024>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Fig 1. Intra-incisional clindamycin protocol algorithm for clinician use.

subcutis, negligible amounts enter systemic circulation; therefore, there is decreased potential for antibiotic resistance, systemic drug interactions, and disruption of the intestinal microbiome.⁴

One limitation of this study was the patient retention rate; approximately 18% did not return for routine postoperative follow-up. As such, it is plausible that the true surgical-site infection rate was higher. However, this is unlikely because we

maintain close contact with our patients and referring providers for all surgical site complications.

In summary, this study demonstrates that preoperative, intra-incisional, prophylactic clindamycin is a safe and effective method to reduce postoperative surgical-site infections and may help to reduce systemic antibiotic overuse.

Teo Soleymani, MD,^a David G. Brodland, MD,^{a,b} and John A. Zitelli, MD^{a,b}

From Zitelli and Brodland Skin Cancer Center, Pittsburgh, Pennsylvania^a; and University of Pittsburgh Medical Center, Pennsylvania.^b

Funding sources: None.

Conflicts of interest: None disclosed.

Reprints not available from the authors.

Correspondence to: Teo Soleymani, MD, Zitelli and Brodland Skin Cancer Center, 5200 Centre Ave, Ste 303, Pittsburgh, PA 15232

E-mail: teo.soleymani@gmail.com

REFERENCES

1. Dohmen PM. Antibiotic resistance in common pathogens reinforces the need to minimise surgical site infections. *J Hosp Infect.* 2008;70(suppl 2):15-20.
2. Darouiche RO, Wall MJ Jr, Itani KM, et al. Chlorhexidine-alcohol versus povidone-iodine for surgical-site antisepsis. *N Engl J Med.* 2010;362(1):18-26.
3. Olds C, Spataro E, Li K, Kandathil C, Most SP. Postoperative antibiotic use among patients undergoing functional facial plastic and reconstructive surgery. *JAMA Facial Plast Surg.* 2019;21(6):491-497. <https://doi.org/10.1001/jamafacial.2019.1027>.
4. Huether MJ, Griego RD, Brodland DG, Zitelli JA. Clindamycin for intra-incisional antibiotic prophylaxis in dermatologic surgery. *Arch Dermatol.* 2002;138(9):1145-1148.
5. Barbieri JS, Etzkorn JR, Margolis DJ. Use of antibiotics for dermatologic procedures from 2008 to 2016. *JAMA Dermatol.* 2019;155(4):465-470.

<https://doi.org/10.1016/j.jaad.2020.05.146>

The effectiveness of rituximab in pemphigus and the benefit of additional maintenance infusions: Daily practice data from a retrospective study

To the Editor: Rituximab in combination with short term prednisone or other immunosuppressive therapy is recognized as the first-line therapy in pemphigus.¹ Additional maintenance infusions are often given on clinical indication, but evidence regarding these infusions is scarce. The aim of this study was to

Table I. Baseline characteristics of pemphigus patients treated with rituximab

Characteristics	Value
Male/female, n (%)	20 (38)/33 (62)
Mean age at start of rituximab, y, median (SD; range)	53.8 (14.3; 26-86)
PV/PF, n (%)	44 (83)/9 (17)
Mucosal PV, n (%)	17 (32)
Mucocutaneous PV, n (%)	27 (51)
Disease duration before rituximab, weeks, median (SD; range)	88 (192.8; 8-793)
Rituximab administered within 1 year after symptoms, n (%)	16 (30)
Rituximab naive/nonnaive,* n (%)	45 (85)/8 (15)
Time between prior and current cycle, weeks, median (SD; range)	149 (66.0; 84-248)
Received M6 and M12, [†] n (%)	25 (47)
Systematic [‡]	19 (36)
Nonsystematic [§]	6 (11)
Previous therapies, n (%)	
Prednisone	51 (96)
Azathioprine	36 (68)
Mycophenolate mofetil	12 (23)
Dapsone	6 (11)
Methotrexate	3 (6)
HIVIG	1 (2)
Number of previous therapies, median (SD; range)	2 (0.8; 0-4)
Adjuvant therapies to which rituximab was added, n (%)	
Prednisone	42 (79)
Mean cumulative dose, mg, median (SD; range)	1777 (1975.5; 0-7110)
Azathioprine	13 (25)
Mycophenolate mofetil	7 (13)
Dapsone	2 (4)
Methotrexate	2 (4)
HIVIG	2 (4)

HIVIG, Human intravenous immunoglobulin G; M6, month 6; M12, month 12; PF, pemphigus foliaceus; PV, pemphigus vulgaris; SD, standard deviation.

*Nonnaive patients previously received 2 infusions of 500 mg of rituximab.

[†]Patients who received 500 mg of rituximab at months 6 and 12.

[‡]Patients who received M6 and M12 infusions as the standard procedure.

[§]Patients who received M6 and M12 infusions on clinical indication.

analyze the effectiveness of rituximab with and without maintenance infusions. Retrospective data were obtained from patients with pemphigus vulgaris and pemphigus foliaceus treated with rituximab between 2012 and 2017. Two infusions of 1000 mg were administered with an interval of 2 weeks (month [M] 0 and M1/2). From 2014, patients received additional maintenance infusions of 500 mg

at M6 and M12 as the standard protocol (systematic infusions). Before 2014, M6 and M12 infusions were administered on clinical indication (nonsystematic infusions). The primary outcome was the relapse rate during the first 3 years after initial rituximab infusion, comparing patients with and without M6 and M12. Secondary outcomes were disease control, partial remission (PR), and complete remission on/off therapy² and adverse events according to the Common Terminology Criteria for Adverse Events grading system.

Fifty-three patients received M0 and M1/2 infusions, of whom 25 received additional M6 and M12 infusions. Table I summarizes the baseline characteristics. A total of 30 (57%) patients developed a relapse within the first 3 years after the M0 infusion, with a median time of 42 weeks (standard deviation, 41.6; range, 9-149 weeks). The relapse rate was significantly lower in the group with M6 and M12 infusions compared to the group without these infusions ($n = 10$ [40%] vs $n = 20$ [71%]; $P = .024$). Table II summarizes the clinical response, relapses, and prednisone dose per subgroup. No significant difference in relapse rate was seen between the group with systematic and nonsystematic infusions (Table II). In multivariate logistic regression, maintenance infusions at M6 and M12 were associated with a significantly lower relapse rate corrected for the following variables: pemphigus subtype, rituximab naive versus non-naive patients, disease duration before rituximab administration, and mean cumulative dose of prednisone (odds ratio, 0.126; 95% confidence interval, 0.024-0.667; $P = .015$) (Supplemental Table I; available via Mendeley at <https://doi.org/10.17632/byrr3vppdm.1>). All patients achieved disease control, of whom 12 (23%) patients achieved PR and 41 (77%) patients achieved CR. PR off therapy was achieved by 4 (8%) patients and CR off therapy by 30 (57%) patients. No significant differences in clinical response were seen between the pemphigus subtypes (Table II). Overall, 78% of the reported adverse events were mild, 15% were moderate, and 7% were severe, but none were life-threatening (Supplemental Table II; available via Mendeley at <https://doi.org/10.17632/byrr3vppdm.1>). No deaths occurred during follow-up.

In this study, additional maintenance infusions at M6 and M12 showed a beneficial effect in preventing relapses. Our study was limited by the heterogeneity of the study population and the absence of Pemphigus Disease Area Index scores. However, after correction for other variables related to disease severity, maintenance infusions were still associated with a significantly lower relapse rate.

Table II. Clinical endpoints and cumulative dose of concomitant prednisone in subgroups of patients with pemphigus treated with rituximab

Subgroups	DC, n (%)	PR, n (%)	PR off therapy, n (%)	CR, n (%)	CR off therapy, n (%)	TTR, weeks, median (SD; range)	Relapse, n (%)	Mean cumulative dose of prednisone, mg
PV	44 (100)	12 (27)	4 (9)	32 (73)	21 (48)	39 (25.6; 5-106)	27 (61)	1997
Mucocutaneous PV	27 (100)	6 (22)	2 (7)	21 (78)	14 (52)	33 (24.9; 12-106)	14 (52)	2103
Mucosal PV	17 (100)	6 (35)	2 (12)	11 (65)	7 (41)	45 (27.4; 5-92)	13 (76)	1852
PF	9 (100)	0 (0)	0 (0)	9 (100)	9 (100)	62 (48.8; 49-170)	3 (33)	970
M6 and M12 yes*	25 (100)	4 (16)	1 (4)	21 (84)	16 (64)	55 (36.8; 7-168)	10 (40) [§]	2625
M6 and M12 no*	28 (100)	8 (29)	3 (11)	20 (71)	14 (50)	39 (32.1; 5-170)	20 (71) [§]	928
Systematic M6 and M12 [†]	19 (100)	1 (5)	1 (5)	18 (95)	13 (68)	49 (27.5; 17-109)	6 (32)	2326
Nonsystematic M6 and M12 [‡]	6 (100)	3 (50)	0 (0)	3 (50)	3 (50)	69 (58.5; 7-168)	4 (67)	2045

CR, Complete remission; DC, disease control; M, month; PF, pemphigus foliaceus; PR, partial remission; PV, pemphigus vulgaris; SD, standard deviation; TTR, time to remission.

*Patients who received 500 mg of rituximab at both M6 and M12.

[†]Patients who received both M6 and M12 infusions as the standard procedure.

[‡]Patients who received both M6 and M12 infusions on clinical indication.

[§] $P = .024$. Denotes significant differences between patients who received M6 and M12 infusions versus patients without these infusions. All other variables did not show a significant difference between the subgroups.

^{||} $P = .023$. Denotes significant differences between patients who received M6 and M12 infusions versus patients without these infusions. All other variables did not show a significant difference between the subgroups.

Other studies³⁻⁵ indirectly support the importance of maintenance infusions by reporting high relapse rates varying between 50% and 100% after treatment with rituximab without additional infusions. All patients in our study achieved remission, with the majority achieving CR, proving that rituximab is an effective treatment for pemphigus. In addition, only minimal severe adverse events were reported. These findings imply that maintenance treatment should be administered in all patients with pemphigus to prevent relapses.

Hanan Rashid, MD,^a Aniek Lamberts, MD,^a Dian van Maanen, MD,^a Maria C. Bolling, MD, PhD,^a Gilles F. H. Diercks, MD, PhD,^{a,b} Hendri H. Pas, PhD,^a Marcel F. Jonkman, MD, PhD,^{a,†} and Barbara Horváth, MD, PhD^a

From the Department of Dermatology, Center for Blistering Diseases, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands^a; and Department of Pathology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands.^b

[†]Deceased.

Funding sources: None.

Conflicts of interest: None disclosed.

IRB approval status: Approved.

Reprints not available from the authors.

Correspondence to: Hanan Rashid, MD, Department of Dermatology, University Medical Center Groningen, Hanzeplein 1, 9700 RB Groningen, The Netherlands

E-mail: h.rashid@umcg.nl

REFERENCES

1. Joly P, Maho-Vaillant M, Prost-Squarcioni C, et al. First-line rituximab combined with short-term prednisone versus prednisone alone for the treatment of pemphigus (Ritux 3): a prospective, multicentre, parallel-group, open-label randomised trial. *Lancet*. 2017;389(10083):2031-2040.
2. Murrell DF, Dick S, Ahmed AR, et al. Consensus statement on definitions of disease, end points, and therapeutic response for pemphigus. *J Am Acad Dermatol*. 2008;58(6):1043-1046.
3. Cianchini G, Lupi F, Masini C, Corona R, Puddu P, De Pittà O. Therapy with rituximab for autoimmune pemphigus: results from a single-center observational study on 42 cases with long-term follow-up. *J Am Acad Dermatol*. 2012;67(4):617-622.
4. Saleh MA. A prospective study comparing patients with early and late relapsing pemphigus treated with rituximab. *J Am Acad Dermatol*. 2018;79(1):97-103.
5. Currimbhoy S, Zhu V, Dominguez AR, Pandya AG. Rituximab in the treatment of 38 patients with pemphigus with long-term follow-up. *J Eur Acad Dermatol Venereol*. 2016;30:1050-1052.