University of Groningen

Environmental change reduces body condition, but not population growth, in a high-arctic herbivore

Layton-Matthews, Kate; Grøtan, Vidar; Hansen, Brage Bremset; Loonen, Maarten J.J.E.; Fuglei, Eva; Childs, Dylan Z.

Published in:
Ecology Letters

DOI:
[10.1111/ele.13634](https://doi.org/10.1111/ele.13634)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Layton-Matthews, K., Grøtan, V., Hansen, B. B., Loonen, M. J. J. E., Fuglei, E., & Childs, D. Z. (2021). Environmental change reduces body condition, but not population growth, in a high-arctic herbivore. *Ecology Letters*, 24(2), 227-238. <https://doi.org/10.1111/ele.13634>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

LETTER

Environmental change reduces body condition, but not population growth, in a high-arctic herbivore

Kate Layton-Matthews,^{1*}
 Vidar Grøtan,¹
 Brage Bremset Hansen,¹
 Maarten J. J. E. Loonen,²
 Eva Fuglei³ and
 Dylan Z. Childs⁴

¹Centre for Biodiversity Dynamics,
 Department of Biology, Norwegian
 University of Science and Technol-
 ogy, Trondheim, Norway

²Arctic Centre, University of Gronin-
 gen, Groningen, the Netherlands

³Norwegian Polar Institute, Tromsø,
 Norway

⁴Department of Animal and Plant
 Sciences, University of Sheffield,
 Sheffield, UK

Abstract

Environmental change influences fitness-related traits and demographic rates, which in herbivores are often linked to resource-driven variation in body condition. Coupled body condition-demo- graphic responses may therefore be important for herbivore population dynamics in fluctuating environments, such as the Arctic. We applied a transient Life-Table Response Experiment ('tran- sient-LTRE') to demographic data from Svalbard barnacle geese (*Branta leucopsis*), to quantify their population-dynamic responses to changes in body mass. We partitioned contributions from direct and delayed demographic and body condition-mediated processes to variation in population growth. Declines in body condition (1980–2017), which positively affected reproduction and fledg- ling survival, had negligible consequences for population growth. Instead, population growth rates were largely reproduction-driven, in part through positive responses to rapidly advancing spring phenology. The virtual lack of body condition-mediated effects indicates that herbivore popula- tion dynamics may be more resilient to changing body condition than previously expected, with implications for their persistence under environmental change.

Keywords

Arctic, barnacle goose, climate change, integral projection models, life table response experiments, population dynamics, trait-mediated and modified effects, transient LTRE.

Ecology Letters (2021) 24: 227–238

*Correspondence: E-mail: kate.-
 matthews@nina.no

INTRODUCTION

Fluctuations in fitness-related traits and population size are jointly affected by environmental stochasticity and density dependence (Lande *et al.*, 2003; Sæther *et al.*, 2016). Body condition, often measured as proxies, for example body mass or mass/tarsus ratio (Schamber *et al.*, 2009), is a key fitness-related trait, reflecting energy reserves available to survive, grow and reproduce (Labocha and Hayes, 2012). Individual body condition is constantly responding to weather and density-dependent processes, as they influence food availability and energy expenditure (Sæther, 1997; Parker *et al.*, 2009). Particularly in herbivores, this can directly influence reproduction and survival (Festa-Bianchet *et al.*, 1997; Sæther, 1997; Post and Stenseth, 1999). Short-term effects of weather and density can also lead to cohort effects on body condition, which, in turn, can have lasting impacts on reproduction (Albon *et al.*, 1987; Choudhury *et al.*, 1996) and population size (Beckerman *et al.*, 2002). Simultaneous changes in body condition and vital rates occur when both respond to variation in weather or density (Parmesan, 2006), with potentially complex population-growth consequences (Post *et al.*, 1997; Ozgul *et al.*, 2010). However, this coupling remains largely unexplored. Linking changes in fitness-related traits to variation in population growth is particularly complex when environment-trait-demography associations vary temporally (e.g. seasonally, Paniw *et al.*, 2019b) or among life-history components (e.g. Douhard *et al.*, 2013).

Arctic environments are seasonal and stochastic. Conse- quently, Arctic herbivores are exposed to variable resource availability, causing annual fluctuations in body condition and

population size (Forchhammer *et al.*, 2002; Couturier *et al.*, 2008; Albon *et al.*, 2017). Weather patterns are being modified by climate change (Scheffer *et al.*, 2001), which is occurring most rapidly in the Arctic (Serreze and Barry, 2011). Accord- ingly, climate change effects on body condition (Albon *et al.*, 2017), reproduction (Post and Forchhammer, 2008; Layton- Matthews *et al.*, 2020), survival (Aubry *et al.*, 2013) and pop- ulation size (Forchhammer *et al.*, 2002; Albon *et al.*, 2017) have been documented in Arctic herbivores. However, climate effects on fitness-related traits and demographic rates do not act independently, and relatively little is known about the linkages among them due to the challenge of quantifying these – often complex – relationships (McLean *et al.*, 2016; Visser *et al.*, 2016; Jenouvrier *et al.*, 2018). Additionally, traits can influence environment-demography relationships via two mechanisms: trait-mediating (Ozgul *et al.*, 2010; Plard *et al.*, 2015; Albon *et al.*, 2017) or trait-modifying effects (Herfindal *et al.*, 2006; Harrison *et al.*, 2013). A mediating trait effect *explains* a relationship between environmental conditions and vital rates, for example temperature affects body condition, which in turn affects survival, whereas a modifying effect requires a body condition-temperature interaction effect on survival.

Since body condition is so influential on life-history pro- cesses in Arctic herbivores, we could expect large population- level responses to changes in this trait (Albon *et al.*, 2017). However, changes in fitness-related traits do not necessarily affect population growth (McLean *et al.*, 2020), since three conditions must be met for trait-mediated effects on popula- tion growth to arise; (1) the trait must fluctuate at the relevant life-history stage for a given, potentially age-specific,

demographic rate, (2) changes in the trait must influence the demographic rate and (3) the population growth rate must, in turn, be influenced by variation in the demographic rate. If any conditions are not met, then no pathway exists from trait variation to population growth variation (Jenouvrier *et al.*, 2018).

Populations are constantly being perturbed by short-term, temporal variation in the environment (Bierzychudek, 1999; Clutton-Brock & Coulson 2002). This can lead to 'transient population dynamics' (Hastings, 2004; Koons *et al.*, 2005; Ezard *et al.*, 2010), as fluctuating environments cause changes in underlying population structure (e.g. age structure, Koons *et al.*, 2016) or trait distributions (e.g. body mass, Ozgul *et al.*, 2010). Changes in population structure can have delayed, transient effects on future population growth, especially when there is substantial variation in the vital rates of different classes of individuals (Beckerman *et al.*, 2002; De Roos *et al.*, 2003; Hansen *et al.*, 2019). If poor conditions reduce cohort body condition, vital rates can be affected for several subsequent years through delayed effects, mediated by early-life body condition (e.g. Albon *et al.*, 1987). Recent extensions of matrix (MPM) and integral (IPM) projection models have attempted to quantify population-growth consequences of such delayed ('lagged') effects (Koons *et al.*, 2005; Kuss *et al.*, 2008). Transient life-table response experiments (transient-LTREs) partition variance in the realised population growth rate into contributions from demographic rates. Furthermore, they can be used to separate temporal variation in λ into direct demographic effects versus delayed effects from transient changes in population structure/trait distributions (e.g. cohort effects), thereby explicitly incorporating past environments (Maldonado-Chaparro *et al.* 2018).

We explored how a substantial change in body condition contributed to variation in population growth in a high-arctic herbivore, the barnacle goose (*Branta leucopsis*). We analysed female mark-recapture and body mass data (1990–2017) to quantify environmental and density effects on survival, reproduction, growth and fledgling body mass. We used the regression models to construct an environmentally driven, stochastic IPM (Rees and Ellner, 2009; Metcalf *et al.*, 2015). Using a recently developed transient-LTRE (Maldonado-Chaparro *et al.* 2018), we decomposed variation in the realised population growth rate (λ_t) into vital rate contributions through demographic and trait-mediated pathways, separated into direct and delayed effects. This methodology revealed that variation in population growth was largely reproduction-driven, through direct and delayed effects (i.e. changes in age structure), whereas body condition-mediated pathways contributed negligibly to population growth. Thus, herbivore populations appear more resilient to substantial changes in body condition than anticipated, with implications for their persistence under future environmental change.

MATERIAL AND METHODS

Study species

Arctic geese are migratory capital breeders, relying, to some extent, on accumulated body reserves for reproduction

(Jönsson, 1997; Hahn *et al.*, 2011). Therefore, an individual's body condition prior to the breeding season affects their reproductive success (Ankney and MacInnes, 1978; Ebbs and Spaans, 1995). Svalbard barnacle geese overwinter in Scotland, UK (55° N, 3.30° W). They fly to Svalbard for breeding in summer, stopping over in spring along the coast of mainland Norway. The study population breeds close to Ny-Ålesund, western Spitsbergen (78°55' N, 11°56' E). Geese arrive at the end of May and nest on islands in the fjord, Kongsfjorden. Hatching occurs from late June. Families leave the nesting islands to forage thereafter, until offspring fledge at the end of August and geese migrate back to Scotland by October.

Demographic data

All analyses were female-based. Over the main study period (mark–recapture period: 1990–2017), 1669 females were caught in July–August and ringed with unique colour and metal identification bands. Recapture data were based on daily observations of ringed individuals around Ny-Ålesund during the foraging period from late June to August ($n_{\text{obs}} = 7280$). Some years were lacking recapture or body mass data (see Appendix S1 for annual sample sizes). We assessed reproduction based on observations of sexually mature adults with fledged offspring, at the beginning of August. We attributed individuals to two age classes: individuals ringed in their first year of life as 'fledglings' (*fl*) and older individuals as 'adults' (*ad*) – a pooled age class of up to 28 years old. Body mass was measured during a catch ($n_{\text{mass}}/n_{\text{total}}$: *fl* = 691/712, *ad* = 2123/6568). We analysed body mass, rather than other body condition proxies, since body mass is a reliable measure of condition in geese (Schmutz, 1993; Lindholm *et al.*, 1994).

Analytical approach

First, we fitted (generalised) linear mixed-effects models ((G) LMMs) to describe fledgling body mass (C_0), growth (i.e. change in body mass from t to $t + 1$, G), overwinter survival (ϕ) and reproduction: the probability of reproducing (R) and fledged brood size (fec). For each model, we quantified effects of age class, body mass and covariates, using model selection. Based on the best-approximating models (see Appendix S3 for more details), we constructed an IPM to model temporal dynamics of population size and body mass distribution. We decomposed variation in the population growth rate (λ_t), using a transient-LTRE, into direct effects of demographic rates versus indirect effects through fluctuations in age class structure and body mass distribution. Contributions were further decomposed into variation from environmental and density covariates versus random effects.

Regression models

We fitted an LMM to fledgling body mass data, to model the mean and distribution of fledgling body masses (C_0), including catch date as a predictor since gosling growth exhibited a seasonal trend (Appendix S2). We also fitted an LMM to body mass data of both age classes (fledglings and adults), describing body mass-dependent growth between years, due to ontogeny

and phenotypic plasticity (G). To estimate apparent survival (ϕ), we modelled mark-recapture data with a Cormack-Jolly-Seber framework using the *RMark* interface (Laake, 2013) for program MARK (White and Burnham, 1999). We modelled detection probabilities with a fixed year effect. We fitted GLMMs to reproductive data and modelled reproduction as two response parameters. R describes the annual reproduction probability that is whether or not a female had at least one fledgling (0/1), fitted as a binomial response. Fledged brood size (fec) describes the number of fledglings per mother, fitted as a Poisson response. We included observations from 2 year-olds onwards (age of sexual maturity, Forslund and Larsson, 1992; Fjellidal *et al.*, 2020) in the reproductive models, and only successfully reproducing individuals ($R = 1$) in the model of fec . We fitted all (G)LMM's with year as a random effect using the package *lme4* in R (Bates *et al.*, 2015).

Using these regression models, we identified the most parsimonious model including effects of age class, body mass, density and environmental covariates (see below, *Covariates*) on C_0 , G , R , fec and ϕ , using Akaike's Information Criterion corrected for small sample sizes (AICc, Burnham and Anderson, 2002). For the survival model in *RMark*, we imputed missing adult body mass observations using linear interpolation between two observations of an individual. We detrended body mass for model selection of G and C_0 , to avoid spurious correlations caused by declining trends. We detrended body mass data against year by calculating the least-squares fit of a straight line and subtracting the resulting function from the data. A set of candidate models were fitted for each rate including all possible subsets of covariates and interactions between age class, body mass and covariates (global models shown in Appendix S3, Table S1). If competing models had $\Delta AICc < 2$, we considered the one with the least parameters as most parsimonious. Finally, since *RMark* does not allow for estimation of year and age-specific random effects, we fitted the most parsimonious model of survival rates including covariates (based on model selection in *RMark*), in a Bayesian framework to model age class- and year-specific random effects, and with year-specific detection probabilities. We implemented Markov Chain Monte Carlo (MCMC) simulations in JAGS via the *rjags* package (Plummer, 2013), assuming annual variation in survival originated from a random process with zero mean and age class-specific deviations (see Schaub *et al.*, 2013; Layton-Matthews *et al.*, 2019 for details). All priors were non-informative however, missing body mass observations were imputed by drawing from a normal distribution, where priors were set at the age class-specific mean body mass and variance (Gimenez *et al.*, 2006). While the Bayesian framework allows for estimation of age- and year-specific random effects, imputing missing body mass data can lead to under-estimation of individual heterogeneity (McCarthy and Masters, 2005).

Covariates

In addition to body mass, we included covariates reflecting weather and population density over the annual cycle, and predator abundance effects on barnacle goose demography (Layton-Matthews *et al.*, 2020) in the regression models of

reproduction (R , fec), survival (ϕ), growth (G) and fledgling body mass (C_0). From the overwintering grounds at Solway Firth, Scotland (*win*, i.e. winter: October_t–March_{t+1}), we included annual mean winter temperature (T_{win}) and total fly-way population counts (N_{win}). From the spring staging grounds at Helgeland, (*spr*, i.e. spring: April–May), we included spring precipitation (P_{spr}) using data from the Vega weather station (65°38' N, 11°52' E). Climate covariates from the breeding grounds in Svalbard (*sum*, i.e. summer) included temperature (mid-June–mid-July, T_{sum}) and precipitation (mid-July–mid-August, P_{sum}). Additionally, the date of spring onset (SO_{sum}) describes the onset of snowmelt and plant-growth onset at the breeding grounds, and is the (ordinal) day when the 10-day smoothed daily temperature crossed 0°C and remained above for at least 10 days (Le Moullec *et al.*, 2019). We also included estimated adult population size in Kongsfjorden (N_{sum} , Layton-Matthews *et al.*, 2019), and the proportion of occupied known dens as an index of Arctic fox (*Vulpes lagopus*) abundance (fox_{sum}), since predation by Arctic foxes affects pre-fledging survival (Fuglei *et al.*, 2003; Layton-Matthews *et al.*, 2020). More details on covariates are found in Layton-Matthews *et al.* (2020).

Stochastic integral projection model

We constructed a stochastic IPM, describing temporal dynamics of population size and body mass distribution, $n(m, t)$, of fledglings (*fl*) and adults (*ad*), following the life cycle in Fig. 1. The growth kernel, $G^{(t)}(m', m)$, describes the probability density function of body masses m' in August in year $t + 1$ of an individual of body mass m in year t . Annual age-class specific survival, $\Phi^{(t)}(m, a)$, describes the probability of an individual, of age class a (*fl* or *ad*) and body mass m , at year t , surviving to year $t + 1$. $P_{ad}^{(t)}(m', m)$ and $P_{fl}^{(t)}(m', m)$ represent survival-growth kernels for adults and fledglings describing how individuals of body mass m at time t , survive and grow to reach mass m' at $t + 1$, given by:

$$P_a^{(t)}(m', m) = \Phi^{(t)}(m, a)G^{(t)}(m', m) \text{ for } a = \text{fledglings or adults} \quad (1)$$

Annual reproduction probability, $R^{(t)}(m)$, describes the probability of a >1-year-old female of body mass m producing at least one fledgling at $t + 1$, given she survives. Fledged brood size, $fec^{(t)}$, describes the number of fledglings per mother at $t + 1$, conditional on reproduction. Fledgling body mass kernel, $C_0^{(t)}(m')$, describes the probability distribution of fledgling body masses in August at $t + 1$. This was assumed to be independent of mother body mass, since a pedigree was not available. $F_{ad}^{(t)}(m', m)$ is the reproduction kernel, describing the density of fledglings of body mass m' that adults of body mass, m , can contribute to the population at year $t + 1$;

$$F_{ad}(m', m) = \Phi^{(t)}(m, ad)R^{(t)}(m)fec^{(t)}C_0^{(t)}(m')/2 \quad (2)$$

Reproduction was divided by 2 since the model was female based. The structure of the IPM was:

$$n_{fl}(m', t+1) = \int_L^U F_{ad}^{(t)}(m', m)n_{ad}(m, t)dm \quad (3)$$

Figure 1 Life cycle of barnacle geese based on a post-breeding census (i.e. breeding occurs just before a census). Individuals in age class a must survive with a body mass(m)-dependent probability ($\phi(m, a)$) and grow to the next year ($G(m', m)$) in order to reproduce ($R(m)$, fec) and contribute fledglings of body mass ($C_0(m')$) to the population. Model predictions of body mass effects at t on: (a) fledgling (ϕ_{fl}) and adult (ϕ_{ad}) survival, (b) body mass at $t + 1$ (dashed line = 1:1 slope), reproduction probability (R) of adult females at the (c) 20th quantile, (d) mean and (e) 80th quantile of the date of spring onset (SO_{sum}). Effects of adult population size at the breeding grounds (N_{sum}) on fledgling body mass (C_0) at the (f) 20th, (g) mean and (h) 80th quantile of Arctic fox abundance (fox_{sum})

$$\begin{aligned}
 n_{ad}(m', t+1) &= \int_L^U P_{ad}^{(t)}(m', m) n_{ad}(m, t) dm \\
 &+ \int_L^U P_{fl}^{(t)}(m', m) n_{fl}(m, t) dm
 \end{aligned}
 \quad (4)$$

We simulated stochastic population dynamics, assuming density-independent population growth, and employed a two-step Monte Carlo resampling approach (Metcalf *et al.*, 2015). Details of the simulation and Monte Carlo resampling approach can be found in Appendix S4.

Transient life table response experiment

Life table response experiments (LTREs) decompose variance in a demographic response metric (typically, population growth rate) into contributions arising from spatial or temporal variation in vital rates. LTREs have become a standard approach to study population-level responses to environmental factors, with fixed, random and regression designs routinely applied to plant and animal populations (Caswell, 1989). Here, rather than studying asymptotic population growth rates, with the assumption that the population is always close to its stable structure, we decomposed variance in the realised population growth rate at time t (λ_t). Specifically, we used an extension of the Monte Carlo regression-random LTRE (Rees and Ellner, 2009), to incorporate transient fluctuations in age structure and body mass distribution (Maldonado-Chaparro *et al.* 2018). We included lagged parameter effects in the model of λ_t , to quantify delayed effects of parameters (i.e. parameter i at time t , $\theta_{i,t}$), acting through changes in age structure or body mass distribution. We compared variance decompositions of λ_t assuming either linear (linear model, LM) or nonlinear (generalised additive model, GAM) dependencies of λ_t . Following Maldonado-Chaparro *et al.* (2018), we decomposed variance in $\log(\lambda_t)$. We compared the R^2 of LMs and GAMs and incorporated delayed effects with increasing numbers of (year) lags to capture the impact of fluctuating stage-structure and body condition. We also tested whether interaction effects between parameters (pairwise smooths for GAMs) contributed to substantial variation in $\log(\lambda_t)$. We considered a model of $\log(\lambda_t)$ to be a better fit when the amount of variance explained increased by at least 1%. To summarise relative contributions of each term, we calculated scaled contributions by dividing the (co)variance contribution from a given term by the total variance in $\log(\lambda_t)$ (Maldonado-Chaparro *et al.* 2018).

Vital rate contributions were further partitioned into variation from modelled covariate effects versus random effects. In this case, (net) contributions of covariates to λ_t depended on: (1) temporal covariances among covariates, (2) their effect size on $\theta_{i,t}$ and (3) the sensitivity of λ_t to $\theta_{i,t}$. We assumed linear dependencies of λ_t on $\theta_{i,t}$ for the environmental decomposition. In the IPM, trait-mediated effects could contribute to intercept variation in each function, whereas trait-modified effects would cause variation in the slope describing the relationship between body mass and a function. Consequently, we decomposed variation in $\log(\lambda_t)$ at three hierarchical levels; (1) overall contributions from C_0 , G , R , fec and ϕ , (2) modelled

environment versus random effects and (3) slope versus intercept variation.

RESULTS

Body mass and life-history processes

The most parsimonious model of overwinter survival (ϕ) included an interaction effect between age class and body mass, with a much stronger positive effect on fledglings than adults (Fig. 1a), and additive effects of overwintering population size (N_{win} , negative effect) and winter temperature (T_{win} , positive effect) (Table 1). The best growth model (G), estimating body mass at year $t+1$, included a positive effect of body mass at t (Fig. 1b). The best model of reproduction probability (i.e. of an adult female producing fledglings, R) included spring precipitation (positive effect, P_{spr}) and date of spring onset (negative effect, SO_{sum}), and an interaction effect between mother body mass and SO_{sum} , with a strong, positive body mass effect in late springs and no effect in early springs (Fig. 1c–e). For fledged brood size (fec), the best model included negative effects of Arctic fox abundance (fox_{sum}) and summer precipitation (P_{sum}). The best model of fledgling body mass (C_0) included an interaction effect of fox_{sum} and adult population density at the breeding grounds (N_{sum}), where N_{sum} tended to have a negative effect only at high fox abundance (Fig. 1f–h). Further description of the model selection and associated tables (Tables S3, 1–3.5) can be found in Appendix S3.

Over the study period, average cohort body mass declined significantly (slope = -8.3 g per cohort, SE = 0.8 g, $p < 0.001$). Interannual changes in body mass were relatively small for adults, but larger for fledglings (Fig. 2).

Transient LTRE

The mean stochastic population growth rate λ was 1.07 (95% confidence intervals: 0.77, 1.59) (Appendix S4, Fig. S1), which was similar to the observed growth rate (1.05; 0.65, 1.53). The stable body mass-age class distribution reflected the bimodal distribution of fledgling and adult body mass (Appendix S4, Fig. S2). 94% of the variation in λ_t was explained by the main effects of the functions, including a one-year time lag, with a transient linear LTRE (LM-LTRE), which increased to 98% with a generalised additive model (GAM-LTRE). The only pairwise smooth (interaction between parameters) explaining more than 1% variance was between the intercept and slope terms of reproduction probability (R) – since SO_{sum} was a predictor in both terms. Parameter effects with more than a 1-year lag explained $<1\%$ of the variance in λ_t .

Sensitivity surfaces were estimated using the GAM-LTRE for $\log(\lambda_t)$, illustrating the influence of each function through direct (Fig. 3a) and delayed effects (Fig. 3b). Direct effects of variation operating through reproduction probability (R) were larger than survival (ϕ) or fledged brood size (fec). λ_t was sensitive to variation through survival rates, in particular adult survival (Fig. 3a). Direct effects of variation through changes in growth and fledgling body mass are not included in Fig. 3a since changes in body mass only affect λ the following year

Table 1 Linear predictors for best-approximating regression models of each function, with mean parameter estimates, used to parameterise the integral projection model (IPM)

IPM function	Model	Best-fitting regression model
Survival	$\text{logit}(\phi_{fl, ad})$	$-2.67 + 4.32a + 0.004m - 0.003a:m + 0.22T_{win} - 0.21N_{win}$
Growth	G	$648.50 + 0.60m$
Fledgling body mass	C_0	$-3357.40 + 19.46c - 2.37fox_{sum} - 46.30N_{sum} - 7.70fox_{sum}:N_{sum}$
Reproductive rate	$\text{logit}(R)$	$-3.76 + 0.002m + 0.86P_{spr} - 3.52SO_{sum} + 0.002z:SO_{sum}$
Fledged brood size	$\text{log}(fec)$	$0.77 - 0.16fox_{sum} - 0.10P_{sum}$

Functions included the covariates; winter temperature (T_{win}) and overwinter population size (N_{win}) in Scotland, Arctic fox abundance (fox_{sum}), adult population size (N_{sum}), the date of spring onset (SO_{sum}) and summer precipitation (P_{sum}) at the breeding grounds on Svalbard and spring precipitation at the spring stopover site at Helgeland (P_{spr}). a is a dummy variable equal to 0 for fledglings (fl) and 1 for adults (ad), m refers to body mass and c to catch date.

Figure 2 Temporal trends in average cohort adult (black, 1980–2016) and annual fledgling (grey, 1991–2016) body mass. Slope (bold line) with 95% confidence intervals (shading) were calculated based on a linear regression, with year as an explanatory variable

(i.e. delayed effects). Population growth was insensitive to delayed effects (Fig. 3b) through G and C_0 (i.e. trait-mediated effects). The sensitivity of λ_t to delayed effects (i.e. at $t-1$) through R and fec reflects how increased reproduction in a given year had a negative impact the following year, via a shift in the age structure towards a larger proportion of non-reproductive individuals.

Vital rate contributions, arising from variances and covariances in $\log(\lambda_t)$, were separated into direct and delayed effects. The largest vital rate contribution to variation in $\log(\lambda_t)$ stemmed from variation in reproductive probability, R , (81%, Fig. 4), through both direct effects (69%) and delayed effects (12%, i.e. changes in age structure). Remaining variation was explained by direct and delayed contributions from variation in fledged brood size (fec , 7%), adult survival (ϕ_{ad} , total contribution = 6%) and fledgling survival (ϕ_{fl} , 3%). Changes in body mass acting through fledgling body mass and growth

functions (i.e. trait-mediated effects) made negligible contributions to variance in λ_t (0.04% through G and 0.5% through C_0).

Reproductive probability and fledged brood size were negatively correlated with adult survival (ϕ_{ad}) and their covariances led to a negative contribution to the variance in $\log(\lambda_t)$ of -7% . Both reproductive parameters (R and fec) and survival parameters (ϕ_{fl} and ϕ_{ad}) positively covaried, each contributing 3% to $\log(\lambda_t)$.

Vital rate contributions were further decomposed into variance explained by environment and density covariates versus random effects. 58% of the variation in $\log(\lambda_t)$, acting through reproduction probability (R), was attributed to modelled covariates (Fig. 4). Precipitation at the spring stopover site (P_{spr}) contributed to 26% of the variation, while date of spring onset at the breeding grounds (SO_{sum}) contributed 16% through intercept and slope variance. 59% of the variation in $\log(\lambda_t)$ that was attributed to adult survival (ϕ_{ad}) was explained by temperature (T_{win}) and population size (N_{win}) at the wintering grounds, accounting for 1% and 2% of the overall variation in $\log(\lambda_t)$. In contrast, $< 1\%$ of the variation in fledgling survival was attributed to T_{win} and N_{win} although these covariates explained 95% of the positive covariation between fledgling and adult survival, leading to a positive contribution of 1% to variation in $\log(\lambda_t)$ (Fig. 4). Finally, 65% of the variation in $\log(\lambda_t)$ through fledged brood size (fec) was explained by Arctic fox abundance and summer precipitation, each contributing 3% to variation in $\log(\lambda_t)$.

DISCUSSION

Coupled trait-demography responses to environmental change may be key to understand and predict short- and long-term population dynamics, especially in Arctic herbivores. Using a transient-LTRE, we quantified population growth responses to a substantial, temporal decline in body mass (body condition proxy) in an Arctic goose population, caused by degradation of the Arctic breeding grounds. For such trait-mediated effects on population growth to arise, three conditions must be met; the trait must fluctuate, changes in the trait must influence the demographic rate and population growth must be influenced by variation in the demographic rate (Table 2). Here, trends and interannual fluctuations in body condition (Fig. 2) did not incur population growth responses because all three conditions were not met. Although body condition at

Figure 3 Sensitivity surfaces illustrating the effects of fledgling (ϕ_f) and adult (ϕ_{ad}) survival, reproductive rate (R), fledged brood size (f_{ec}), growth (G) and fledgling body mass (C_0) on the population growth rate, $\log(\lambda_t)$. Contributions from each vital rate parameter were separated into (a) direct and (b) delayed contributions, that is parameter effects at t and $t-1$ respectively, on $\log(\lambda_t)$. The x-axis and y-axis rugs show distributions of parameters (centred values) and $\log(\lambda_t)$ respectively

fledging strongly affected fledgling survival (Fig. 1a), population growth was insensitive to fledgling survival (Fig. 3a). Conversely, population growth was sensitive to changes in adult survival, which was insensitive to body condition. Variation in population growth was instead largely explained by direct and delayed effects operating through reproduction, in part shaped by annual variation in spring onset (Fig. 4) and only weakly influenced by adult body mass.

Both resident and migrant Arctic herbivores are exposed to variable weather conditions, resulting in variation in body condition and population size (Festa-Bianchet *et al.*, 1997;

Sæther, 1997). Although weather and density effects on body condition are well documented in herbivores, whether these effects extend to variation in population growth remains largely unanswered. Goose populations have expanded across the Arctic, as a result of increasing population size caused by hunting bans and agricultural change at overwintering grounds (Madsen and Cracknell, 1999; Fox and Madsen, 2017). Density-dependent processes associated with overgrazing by geese have degraded Arctic breeding grounds, leading to declining body condition in Arctic geese (Cooch *et al.*, 1991b; Loonen *et al.*, 1997; Reed and Plante, 1997; Larsson

Figure 4 Percentage contributions of the largest demographic contributions (through adult survival, ϕ_{ad} , fledgling survival, ϕ_n , reproduction probability, R and fledged brood size, fec) to variance in the population growth rate, $\log(\lambda_t)$. Contributions were decomposed into direct and delayed variances and covariances. Colours represent contributions from modelled covariates versus unexplained variation (random effects)

Table 2 Summary of which conditions for body mass-mediated effects on the population growth rate (for each demographic rate) were met or not (✓ = condition met). All conditions need to be met for body mass-mediated effects on the population growth rate to arise.

Condition	Fledgling survival (ϕ_n)	Adult survival (ϕ_{ad})	Reproduction probability (R)	Fledged brood size (fec)
Body mass fluctuates at relevant stage	✓			
Body mass change influences demographic rate	✓		✓	
Variation in demographic rate influences λ		✓	✓	✓

et al., 1998). Cohort adult body condition in the study population declined by 10% from 1980 to 2017. We documented density dependence in fledgling body condition at high Arctic fox abundance, that is the main predator of goslings and, occasionally, adults (Fuglei *et al.*, 2003). Foxes therefore influenced geese through lethal effects on fledged brood size (fec) and non-lethal effects on fledgling body condition, through density-dependent restrictions on their ability to utilise available foraging areas (Loonen *et al.*, 1998).

Despite the temporal decline in (cohort) adult body condition, interannual fluctuations in adult body condition were

small. Fledgling body condition exhibited more interannual variation, likely a result of strong resource dependence during growth (Cooch *et al.*, 1991a; Lindholm *et al.*, 1994). Body condition, in turn, positively influenced survival and reproduction. The reserves which first-year-geese accumulate at the breeding grounds affect survival during migration (Owen and Black, 1989; Menu *et al.*, 2005), reflected in the strong positive effect of fledgling body condition on first-year survival found here. Adult survival, however, was largely unaffected by body condition.

Several studies have stressed the potential importance of coupled trait-demography responses in predicting population persistence under environmental change (Ozgul *et al.*, 2009, 2010; Plard *et al.*, 2014; Jenouvrier *et al.*, 2018). Direct evidence of this is scarce, however, and several studies have indicated that trait changes may lead to more limited population-level responses than expected (Wilson and Arcese, 2003; Maldonado-Chaparro *et al.* 2018; McLean *et al.*, 2020). Here, despite the potential for coupled body condition-demography responses, changes in Arctic goose body condition did not incur population-level responses through trait-mediated effects. Over the study period, body condition declined and positively influenced demographic rates. However, these trends and interannual fluctuations in body condition did not translate into effects on population growth because all three conditions required for trait-mediated pathways to arise were not met for any age class-specific demographic rate (Table 2). Although body mass influenced fledging survival, variance in fledgling survival contributed negligibly to population growth, since fledglings represent a small fraction of the population. In contrast, population growth was sensitive to changes in adult survival, but this vital rate was insensitive to body condition. Likewise, population growth was also highly sensitive to changes in reproductive rate, which was only weakly influenced by adult body condition. Importantly, despite adult body condition showing a pronounced long-term decline, interannual fluctuations were limited. Resulting in a lack of trait-mediated effects on population growth rates through reproduction. Nevertheless, further changes in body condition could have large effects on population growth, operating through reproduction, and long-term, delayed consequences of changes in fledgling mass (i.e. ‘silver spoon effects’, Madsen and Shine, 2000) could also be important. However, in our study population no significant effect of fledgling mass was found on adult mass, adult survival or reproduction (i.e. no evidence for silver spoon effects).

According to the demographic buffering hypothesis, variability in population growth reduces fitness (Tuljapourkar, 1982), leading to selection for reduced variance (Gaillard and Yoccoz, 2003; Jongejans *et al.*, 2010). In long-lived species, individuals may, when resources are scarce, increase/maintain their survival by limiting investment in reproduction. Buffering of population growth against trait-mediated variation may occur via such mechanisms. However, in cases where body condition influences adult survival we might expect population-level responses (e.g. Pigeon *et al.*, 2017). Therefore, the magnitude of population-level responses to body condition likely depends on species’ life history and the way in which individuals utilise resources (i.e. capital versus income

breeding). In capital breeders, where body condition is important for reproductive success, we could therefore expect strong effects on population growth, given substantial annual variation in adult body mass and reproduction. However, both in barnacle geese (this study) and in other capital breeders (e.g. van Benthem *et al.*, 2017; Maldonado-Chaparro *et al.* 2018), changes in body condition did not induce population-level responses, likely as a result of low variation due to demographic buffering. A lack of population-level effects of body condition was also observed in income-breeding passerine birds (McLean *et al.*, 2020), indicating that the capital-income breeding dichotomy cannot predict species variation in trait-mediated effects on population growth. However, in several partially capital-breeding ungulate species, changes in body condition did elicit population-level responses (e.g. Soay Sheep, Pelletier *et al.*, 2007; Svalbard reindeer, Albon *et al.*, 2017; bighorn sheep, Pigeon *et al.*, 2017). In these cases, annual fluctuations in body condition were relatively large and body condition influenced adult survival as well as recruitment. This suggests that population-level responses requires body condition to induce variation in vital rates with a large impact on population growth (e.g. adult survival). Body condition may also be less constrained (and more variable) in ungulates, compared to, for example, migratory birds, where individuals carry reserves over long distances (Owen and Black, 1991).

Individual heterogeneity in body condition, which often becomes increasingly important under poor environmental conditions (Barbraud and Weimerskirch, 2005), can also help to understand the lack of trait-mediated effects on population growth. In geese, larger individuals tend to gain better access to resources due to dominance behaviour, and thus reproduce better (Stahl *et al.*, 2001). Within-year, rather than among-year, variation in body condition may thus be relevant also in understanding density-dependent effects, that is through intra-specific competition. Such discrepancies between individual- and population-level responses to trait variation have been documented previously (Reed *et al.*, 2013). Although we assumed density-independent population growth in the IPM, density dependence may explain some of the unexplained variation in population growth. Either through effects on survival, as found here and in previous studies (Layton-Matthews *et al.*, 2020), but also on reproduction since density-dependent effects have been found in previous studies (Layton-Matthews *et al.*, 2019). An additional limitation was the lack of a pedigree, that is mother body mass was not included as a predictor of fledgling body mass. If body mass is heritable, as found in barnacle geese in the Baltic (Larsson and Forslund, 1992), this would lead to a positive correlation between adult growth and fledgling mass, influencing fledgling survival. However, since variation in fledgling survival made a small contribution to population growth, the population-level impact should also be minimal.

Quantifying demographic and associated trait responses to climate change is necessary for a mechanistic and predictive understanding of population-level consequences (Jenouvrier, 2013; Paniw *et al.* 2019a). Arctic warming is advancing snow melt, with widespread effects on plant phenology, while rising summer temperatures are influencing plant productivity

(Bjorkman *et al.*, 2020). For migrating Arctic geese, spring phenology at the breeding grounds also dictates when nesting sites become snow-free as well as the onset of plant growth: both important determinants of breeding success (Reed *et al.*, 2004; Madsen *et al.*, 2007). Positive effects of earlier spring onset on fledgling production has previously been attributed to females laying bigger clutches with improved hatching success (Layton-Matthews *et al.*, 2020). Advancing spring phenology could therefore benefit reproduction and population growth, which appear unhampered by the temporal decline in body condition (since effects of body condition on reproduction diminish with earlier springs). However, neither reproductive parameter exhibited positive temporal trends (Appendix S6), likely due to contrasting direct versus indirect climate change effects, as the number of Arctic foxes – and thus gosling predation rates – are generally increasing (Layton-Matthews *et al.*, 2020; unpublished data, E. Fuglei).

In capital breeders, such as Arctic geese, the accumulation of body stores is beneficial in unpredictable environments, and this strategy is typical at higher latitudes (Varpe *et al.*, 2009; Sainmont *et al.*, 2014). Storing resources along their flyway allows geese to initiate reproduction without immediate food access (Klaassen *et al.*, 2017). Consequently, several reproductive stages depend on fat reserves (Bêty *et al.*, 2003; Guillemain *et al.*, 2008; Aubry *et al.*, 2013), reflected here in the positive relationship between body condition and reproduction. Here, spring onset operated through a trait-modified effect on reproduction, where heavier individuals were more likely to reproduce than lighter ones under poor (delayed) spring conditions, since they have more ‘capital’ to initiate reproduction. While in earlier springs (i.e. typically in more recent years), with excess nesting sites and food resources, benefits of accumulating fat reserves were reduced and the influence of body condition on reproduction was weaker. Further advanced springs due to climate change could tip the balance in favour of accumulating fewer resources for reproduction, potentially relaxing selection on body condition if individuals gain a survival advantage by requiring fewer fat reserves for migration (Larsson *et al.*, 1998).

Overall, population dynamics of Arctic migratory geese appear unaffected by the decline in body condition, thus far. Buffering population growth against changes in body condition, which is essential for survival and reproduction in Arctic herbivores, has clear implications for their resilience to future environmental change. However, as the Arctic continues to change, further declines in body condition could potentially have big effects on population growth, via trait-mediated effects through reproduction.

This work emphasises the importance of holistic approaches capturing pathways from environmental variation to individual and population-level responses. Even when environmental change substantially alters trait distributions that are correlated with vital rates, we cannot assume this will have population-level consequences.

ACKNOWLEDGEMENTS

Funding for data collection was provided by; NWO, Ministry of Foreign Affairs, BIRDHEALTH (851.40.071), Geese on

Arctic Tundra (866.12.407), EU FP7-project FRAGILE, University of Groningen and the Norwegian Polar Institute. The Research Council of Norway supported this work (FRI-MEDBIO project 276080, Centres of Excellence 223257, Arctic Field Grant 282619). We thank scientists, students and volunteers for data collection and Stefan J.G. Vriend for comments on the manuscript.

PEER REVIEW

The peer review history for this article is available at <https://publons.com/publon/10.1111/ele.13634>.

DATA AVAILABILITY STATEMENT

Data supporting the results are archived in Dryad data repository: <https://doi.org/10.5061/dryad.9p8cz8wdv>.

REFERENCES

- Albon, S., Clutton-Brock, T. & Guinness, F. (1987). Early development and population dynamics in red deer. II. Density-independent effects and cohort variation. *The Journal of Animal Ecology*, 56, 69–81.
- Albon, S.D., Irvine, R.J., Halvorsen, O., Langvatn, R., Loe, L.E., Ropstad, E. *et al.* (2017). Contrasting effects of summer and winter warming on body mass explain population dynamics in a food-limited Arctic herbivore. *Glob. Change Biol.*, 23, 1374–1389.
- Ankney, C.D. & MacInnes, C.D. (1978). Nutrient reserves and reproductive performance of female Lesser Snow Geese. *Auk*, 95, 459–471.
- Aubry, L.M., Rockwell, R.F., Cooch, E.G., Brook, R.W., Mulder, C.P. & Koons, D.N. (2013). Climate change, phenology, and habitat degradation: drivers of gosling body condition and juvenile survival in lesser snow geese. *Glob. Change Biol.*, 19, 149–160.
- Barbraud, C. & Weimerskirch, H. (2005). Environmental conditions and breeding experience affect costs of reproduction in blue petrels. *Ecology*, 86, 682–692.
- Bates, D., Machler, M., Bolker, B.M. & Walker, S.C. (2015). Fitting linear mixed-effects models using lme4. *J. Stat. Softw.*, 67, 1–48.
- Beckerman, A., Benton, T.G., Ranta, E., Kaitala, V. & Lundberg, P. (2002). Population dynamic consequences of delayed life-history effects. *Trends Ecol. Evol.*, 17, 263–269.
- Béty, J., Gauthier, G. & Giroux, J.-F. (2003). Body condition, migration, and timing of reproduction in snow geese: a test of the condition-dependent model of optimal clutch size. *Am. Nat.*, 162, 110–121.
- Bierzychudek, P. (1999). Looking backwards: assessing the projections of a transition matrix model. *Ecol. Appl.*, 9, 1278–1287.
- Bjorkman, A.D., García Criado, M., Myers-Smith, I.H., Ravolainen, V., Jónsdóttir, I.S., Westergaard, K.B. *et al.* (2020). Status and trends in Arctic vegetation: evidence from experimental warming and long-term monitoring. *Ambio*, 49, 678–692.
- Burnham, K.P. & Anderson, D.R. (2002). *Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach*, 2nd edn. Springer Science & Business Media, New York.
- Caswell, H. (1989). Analysis of life table response experiments I. Decomposition of effects on population growth rate. *Ecol. Model.*, 46, 221–237.
- Choudhury, S., Black, J.M. & Owen, M. (1996). Body size, fitness and compatibility in barnacle geese *Branta leucopsis*. *The Ibis*, 138, 700–709.
- Clutton-Brock, T. & Coulson, T. (2002). Comparative ungulate dynamics: the devil is in the detail. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 357, 1285–1298.
- Cooch, E., Lank, D., Dzubin, A., Rockwell, R. & Cooke, F. (1991a). Body size variation in lesser snow geese: environmental plasticity in gosling growth rates. *Ecology*, 72, 503–512.
- Cooch, E., Lank, D., Rockwell, R. & Cooke, F. (1991b). Long-term decline in body size in a snow goose population: evidence of environmental degradation? *The Journal of Animal Ecology*, 483–496.
- Couturier, S., Cote, S.D., Huot, J. & Otto, R.D. (2008). Body-condition dynamics in a northern ungulate gaining fat in winter. *Can. J. Zool.*, 87, 367–378.
- De Roos, A.M., Persson, L. & McCauley, E. (2003). The influence of size-dependent life-history traits on the structure and dynamics of populations and communities. *Ecol. Lett.*, 6, 473–487.
- Douhard, M., Gaillard, J.-M., Delorme, D., Capron, G., Duncan, P., Klein, F. *et al.* (2013). Variation in adult body mass of roe deer: early environmental conditions influence early and late body growth of females. *Ecology*, 94, 1805–1814.
- Ebbinge, B.S. & Spaans, B. (1995). The importance of body reserves accumulated in spring staging areas in the temperate zone for breeding in dark-bellied brent geese *Branta b. bernicla* in the high Arctic. *J. Avian Biol.*, 105–113.
- Ezard, T.H., Bullock, J.M., Dalglish, H.J., Millon, A., Pelletier, F., Ozgul, A. *et al.* (2010). Matrix models for a changeable world: the importance of transient dynamics in population management. *J. Appl. Ecol.*, 47, 515–523.
- Festa-Bianchet, M., Jorgenson, J.T., Bérubé, C.H., Portier, C. & Wishart, W.D. (1997). Body mass and survival of bighorn sheep. *Can. J. Zool.*, 75, 1372–1379.
- Fjellidal, M.A., Layton-Matthews, K., Lee, A.M., Grøtan, V., Loonen, M.J. & Hansen, B.B. (2020). High-Arctic family planning: earlier spring onset advances age at first reproduction in barnacle geese. *Biol. Lett.*, 16, 20200075.
- Forchhammer, M.C., Post, E., Stenseth, N.C. & Boertmann, D.M. (2002). Long-term responses in arctic ungulate dynamics to changes in climatic and trophic processes. *Popul. Ecol.*, 44, 113–120.
- Forslund, P. & Larsson, K. (1992). Age-related reproductive success in the barnacle goose. *J. Anim. Ecol.*, 61, 195–204.
- Fox, A.D. & Madsen, J. (2017). Threatened species to super-abundance: The unexpected international implications of successful goose conservation. *Ambio*, 46, 179–187.
- Fuglei, E., Øritsland, N.A. & Prestrud, P. (2003). Local variation in arctic fox abundance on Svalbard, Norway. *Polar Biol.*, 26, 93–98.
- Gaillard, J.-M. & Yoccoz, N.G. (2003). Temporal variation in survival of mammals: a case of environmental canalization? *Ecology*, 84, 3294–3306.
- Gimenez, O., Covas, R., Brown, C.R., Anderson, M.D., Brown, M.B. & Lenormand, T. (2006). Nonparametric estimation of natural selection on a quantitative trait using mark-recapture data. *Evolution*, 60, 460–466.
- Guillemain, M., Elmberg, J., Arzel, C., Johnson, A.R. & Simon, G. (2008). The income-capital breeding dichotomy revisited: late winter body condition is related to breeding success in an income breeder. *The Ibis*, 150, 172–176.
- Hahn, S., Loonen, M.J.J.E. & Klaassen, M. (2011). The reliance on distant resources for egg formation in high Arctic breeding barnacle geese *Branta leucopsis*. *J. Avian Biol.*, 42, 159–168.
- Hansen, B.B., Gamelon, M., Albon, S.D., Lee, A.M., Stien, A., Irvine, R.J. *et al.* (2019). More frequent extreme climate events stabilize reindeer population dynamics. *Nat. Commun.*, 10, 1616.
- Harrison, X.A., Hodgson, D.J., Inger, R., Colhoun, K., Gudmundsson, G.A., McElwaine, G. *et al.* (2013). Environmental conditions during breeding modify the strength of mass-dependent carry-over effects in a migratory bird. *PLoS One*, 8, e77783.
- Hastings, A. (2004). Transients: the key to long-term ecological understanding? *Trends Ecol. Evol.*, 19, 39–45.
- Herfindal, I., Sæther, B.E., Solberg, E.J., Andersen, R. & Høgda, K.A. (2006). Population characteristics predict responses in moose body mass to temporal variation in the environment. *J. Anim. Ecol.*, 75, 1110–1118.
- Jenouvrier, S. (2013). Impacts of climate change on avian populations. *Glob. Change Biol.*, 19, 2036–2057.

- Jenouvrier, S., Desprez, M., Fay, R., Barbraud, C., Weimerskirch, H., Delord, K. *et al.* (2018). Climate change and functional traits affect population dynamics of a long-lived seabird. *J. Anim. Ecol.*, 87, 906–920.
- Jongejans, E., De Kroon, H., Tuljapurkar, S. & Shea, K. (2010). Plant populations track rather than buffer climate fluctuations. *Ecol. Lett.*, 13, 736–743.
- Jönsson, K.I. (1997). Capital and income breeding as alternative tactics of resource use in reproduction. *Oikos*, 78, 57–66.
- Klaassen, M., Hahn, S., Korthals, H. & Madsen, J. (2017). Eggs brought in from afar: Svalbard-breeding pink-footed geese can fly their eggs across the Barents Sea. *J. Avian Biol.*, 48, 173–179.
- Koons, D.N., Grand, J.B., Zinner, B. & Rockwell, R.F. (2005). Transient population dynamics: relations to life history and initial population state. *Ecol. Model.*, 185, 283–297.
- Koons, D.N., Iles, D.T., Schaub, M. & Caswell, H. (2016). A life-history perspective on the demographic drivers of structured population dynamics in changing environments. *Ecol. Lett.*, 19, 1023–1031.
- Kuss, P., Rees, M., ġisdóttir, H.H., Ellner, S.P. & Stöcklin, J. (2008). Evolutionary demography of long-lived monocarpic perennials: a time-lagged integral projection model. *J. Ecol.*, 96, 821–832.
- Labocha, M.K. & Hayes, J.P. (2012). Morphometric indices of body condition in birds: a review. *J. Ornithol.*, 153, 1–22.
- Lande, R., Engen, S. & Saether, B.-E. (2003). *Stochastic population dynamics in ecology and conservation*. Oxford University Press, Oxford, UK.
- Larsson, K. & Forslund, P. (1992). Genetic and social inheritance of body and egg size in the barnacle goose (*Branta leucopsis*). *Evolution*, 46, 235–244.
- Larsson, K., Van der Jeugd, H.P., Van der Veen, I.T. & Forslund, P. (1998). Body size declines despite positive directional selection on heritable size traits in a barnacle goose population. *Evolution*, 52, 1169–1184.
- Layton-Matthews, K., Hansen, B.B., Grøtan, V., Fuglei, E. & Loonen, M.J.J.E. (2020). Contrasting consequences of climate change for migratory geese: predation, density dependence and carryover effects offset benefits of high-arctic warming. *Glob. Change Biol.*, 26, 642–657.
- Layton-Matthews, K., Loonen, M.J.J.E., Hansen, B.B., Saether, B.-E., Coste, C.F.D. & Grøtan, V. (2019). Density-dependent population dynamics of a high Arctic capital breeder, the barnacle goose. *J. Anim. Ecol.*, 88, 1191–1201.
- Le Moullec, M., Buchwal, A., van der Wal, R., Sandal, L. & Hansen, B.B. (2019). Annual ring growth of a widespread high arctic shrub reflects past fluctuations in community-level plant biomass. *J. Ecol.*, 107, 436–451.
- Lindholm, A., Gauthier, G. & Desrochers, A. (1994). Effects of hatch date and food supply on gosling growth in arctic-nesting Greater Snow Geese. *Condor*, 96, 898–908.
- Loonen, M.J.J.E., Oosterbeek, K. & Drent, R. (1997). Variation in growth of young and adult size in barnacle geese *Branta leucopsis*: evidence for density dependence. *Ardea*, 85, 177–192.
- Loonen, M.J.J.E., Tombre, I.M. & Mehlum, F. (1998). Development of an arctic barnacle goose colony: interactions between density and predation. *Nor. Polarinst. Skr.*, 200, 67–80.
- Laake, J.L. (2013). RMark: an R interface for analysis of capture-recapture data with MARK. AFSC Processed Report. Alaska Fisheries Science Center, National Marine Fisheries Service, NOAA Seattle, Washington, USA, pp. 2013–2101.
- Madsen, J. & Cracknell, G. (1999). Goose populations of the Western Palearctic: a review of status and distribution. National Environmental Research Institute, Denmark and Wetlands International, Wageningen, The Netherlands.
- Madsen, J., Tamstorf, M., Klaassen, M., Eide, N., Glahder, C., Rigét, F. *et al.* (2007). Effects of snow cover on the timing and success of reproduction in high-Arctic pink-footed geese *Anser brachyrhynchus*. *Polar Biol.*, 30, 1363–1372.
- Madsen, T. & Shine, R. (2000). Silver spoons and snake body sizes: prey availability early in life influences long-term growth rates of free-ranging pythons. *J. Anim. Ecol.*, 69, 952–958.
- Maldonado-Chaparro, A.A., Blumstein, D.T., Armitage, K.B. & Childs, D.Z. (2018). Transient LTRE analysis reveals the demographic and trait-mediated processes that buffer population growth. *Ecol. Lett.*, 21, 1693–1703.
- McCarthy, M.A. & Masters, P. (2005). Profiting from prior information in Bayesian analyses of ecological data. *J. Appl. Ecol.*, 42, 1012–1019.
- McLean, N., Lawson, C.R., Leech, D.I. & van de Pol, M. (2016). Predicting when climate-driven phenotypic change affects population dynamics. *Ecol. Lett.*, 19, 595–608.
- McLean, N.M., van der Jeugd, H.P., Van Turnhout, C.A., Lefcheck, J.S. & Van de Pol, M. (2020). Reduced avian body condition due to global warming has little reproductive or population consequences. *Oikos*, 129 (5), 714–730.
- Menu, S., Gauthier, G. & Reed, A. (2005). Survival of young greater snow geese (*Chen caerulescens atlantica*) during fall migration. *Auk*, 122, 479–496.
- Metcalfe, C.J.E., Ellner, S.P., Childs, D.Z., Salguero-Gómez, R., Merow, C., McMahon, S.M. *et al.* (2015). Statistical modelling of annual variation for inference on stochastic population dynamics using Integral Projection Models. *Methods Ecol. Evol.*, 6, 1007–1017.
- Owen, M. & Black, J. (1991). The importance of migration mortality in non-passerine birds. In *Bird Population Studies: Relevance to Conservation and Management* (eds Perrins, C., Lebreton, J.-D., Horins, G.). Oxford University Press, New York, NY, pp. 360–372.
- Owen, M. & Black, J.M. (1989). Factors affecting the survival of barnacle Geese on migration from the breeding grounds. *J. Anim. Ecol.*, 58, 603–617.
- Ozgul, A., Childs, D.Z., Oli, M.K., Armitage, K.B., Blumstein, D.T., Olson, L.E. *et al.* (2010). Coupled dynamics of body mass and population growth in response to environmental change. *Nature*, 466, 482.
- Ozgul, A., Tuljapurkar, S., Benton, T.G., Pemberton, J.M., Clutton-Brock, T.H. & Coulson, T. (2009). The Dynamics of Phenotypic Change and the Shrinking Sheep of St. Kilda. *Science*, 325, 464–467.
- Paniw, M., James, T., Archer, C.R., Roemer, G., Levin, S., Compagnoni, A., *et al.* (2019a). Global analysis reveals complex demographic responses of mammals to climate change. *bioRxiv*.
- Paniw, M., Maag, N., Cozzi, G., Clutton-Brock, T. & Ozgul, A. (2019b). Life history responses of meerkats to seasonal changes in extreme environments. *Science*, 363, 631–635.
- Parker, K.L., Barboza, P.S. & Gillingham, M.P. (2009). Nutrition integrates environmental responses of ungulates. *Funct. Ecol.*, 23, 57–69.
- Parnesan, C. (2006). Ecological and evolutionary responses to recent climate change. *Annu. Rev. Ecol. Evol. Syst.*, 37, 637–669.
- Pelletier, F., Clutton-Brock, T., Pemberton, J., Tuljapurkar, S. & Coulson, T. (2007). The evolutionary demography of ecological change: linking trait variation and population growth. *Science*, 315, 1571–1574.
- Pigeon, G., Ezard, T.H., Festa-Bianchet, M., Coltman, D.W. & Pelletier, F. (2017). Fluctuating effects of genetic and plastic changes in body mass on population dynamics in a large herbivore. *Ecology*, 98, 2456–2467.
- Plard, F., Gaillard, J.-M., Coulson, T., Hewison, A.M., Delorme, D., Warnant, C. *et al.* (2014). Mismatch between birth date and vegetation phenology slows the demography of roe deer. *PLoS Biol.*, 12, e1001828.
- Plard, F., Gaillard, J.-M., Coulson, T., Hewison, A.M., Douhard, M., Klein, F. *et al.* (2015). The influence of birth date via body mass on individual fitness in a long-lived mammal. *Ecology*, 96, 1516–1528.
- Plummer, M. (2013). rjags: Bayesian graphical models using MCMC. R package version, 3.
- Post, E. & Forchhammer, M.C. (2008). Climate change reduces reproductive success of an Arctic herbivore through trophic mismatch. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, 363, 2367–2373.
- Post, E. & Stenseth, N.C. (1999). Climatic variability, plant phenology, and northern ungulates. *Ecology*, 80, 1322–1339.

- Post, E., Stenseth, N.C., Langvatn, R. & Fromentin, J.-M. (1997). Global climate change and phenotypic variation among red deer cohorts. *Proc. R. Soc. Lond. B Biol. Sci.*, 264, 1317–1324.
- Reed, A. & Plante, N. (1997). Decline in body mass, size, and condition of greater snow geese, 1975–94. *J. Wildl. Manag.*, 61(2), 413–419.
- Reed, E.T., Gauthier, G. & Giroux, J.-F. (2004). Effects of spring conditions on breeding propensity of greater snow goose females. *Anim. Biodivers. Conserv.*, 27, 35–46.
- Reed, T.E., Jenouvrier, S. & Visser, M.E. (2013). Phenological mismatch strongly affects individual fitness but not population demography in a woodland passerine. *J. Anim. Ecol.*, 82, 131–144.
- Rees, M. & Ellner, S.P. (2009). Integral projection models for populations in temporally varying environments. *Ecol. Monogr.*, 79, 575–594.
- Sainmont, J., Andersen, K.H., Varpe, Ø. & Visser, A.W. (2014). Capital versus income breeding in a seasonal environment. *Am. Nat.*, 184, 466–476.
- Schamber, J.L., Esler, D. & Flint, P.L. (2009). Evaluating the validity of using unverified indices of body condition. *J. Avian Biol.*, 40, 49–56.
- Schaub, M., Jakober, H. & Stauber, W. (2013). Strong contribution of immigration to local population regulation: evidence from a migratory passerine. *Ecology*, 94, 1828–1838.
- Scheffer, M., Straile, D., van Nes, E.H. & Hosper, H. (2001). Climatic warming causes regime shifts in lake food webs. *Limnol. Oceanogr.*, 46, 1780–1783.
- Schmutz, J.A. (1993). Survival and pre-fledging body mass in juvenile Emperor Geese. *The Condor*, 95, 222–225.
- Serreze, M.C. & Barry, R.G. (2011). Processes and impacts of Arctic amplification: A research synthesis. *Global Planetary Change*, 77, 85–96.
- Stahl, J., Tolsma, P.H., Loonen, M.J.J.E. & Drent, R.H. (2001). Subordinates explore but dominants profit: resource competition in high Arctic barnacle goose flocks. *Anim. Behav.*, 61, 257–264.
- Sæther, B.-E. (1997). Environmental stochasticity and population dynamics of large herbivores: a search for mechanisms. *Trends Ecol. Evol.*, 12, 143–149.
- Sæther, B.-E., Visser, M.E., Grøtan, V. & Engen, S. (2016). Evidence for r- and K-selection in a wild bird population: a reciprocal link between ecology and evolution. *Proceedings of the Royal Society B: Biological Sciences*, 283, 20152411.
- Tuljapurkar, S.D. (1982). Population dynamics in variable environments. III. Evolutionary dynamics of r-selection. *Theor. Popul. Biol.*, 21, 141–165.
- van Benthem, K.J., Froy, H., Coulson, T., Getz, L.L., Oli, M.K. & Ozgul, A. (2017). Trait–demography relationships underlying small mammal population fluctuations. *J. Anim. Ecol.*, 86, 348–358.
- Varpe, Ø., Jørgensen, C., Tarling, G.A. & Fiksen, Ø. (2009). The adaptive value of energy storage and capital breeding in seasonal environments. *Oikos*, 118, 363–370.
- Visser, M.D., Bruijning, M., Wright, S.J., Muller-Landau, H.C., Jongejans, E., Comita, L.S. *et al.* (2016). Functional traits as predictors of vital rates across the life cycle of tropical trees. *Funct. Ecol.*, 30, 168–180.
- White, G.C. & Burnham, K.P. (1999). Program MARK: survival estimation from populations of marked animals. *Bird Study*, 46, S120–S139.
- Wilson, S. & Arcese, P. (2003). El Niño drives timing of breeding but not population growth in the song sparrow (*Melospiza melodia*). *Proc. Natl Acad. Sci.*, 100, 11139–11142.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

Editor, Dave Hodgson

Manuscript received 22 April 2020

First decision made 10 June 2020

Second decision made 16 September 2020

Manuscript accepted 6 October 2020