University of Groningen

Prediction of Psychological Distress Among Persons With Spinal Cord Injury or Acquired Brain Injury and Their Significant Others

POWER Grp; Scholten, Eline W. M.; Ketelaar, Marjolijn; Visser-Meily, Johanna M. A.; Roels, Ellen H.; Kouwenhoven, Mirjam; Post, Marcel W. M.

Published in:
Archives of Physical Medicine and Rehabilitation

DOI:
[10.1016/j.apmr.2020.05.023](https://doi.org/10.1016/j.apmr.2020.05.023)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

POWER Grp, Scholten, E. W. M., Ketelaar, M., Visser-Meily, J. M. A., Roels, E. H., Kouwenhoven, M., & Post, M. W. M. (2020). Prediction of Psychological Distress Among Persons With Spinal Cord Injury or Acquired Brain Injury and Their Significant Others. *Archives of Physical Medicine and Rehabilitation*, 101(12), 2093-2102. <https://doi.org/10.1016/j.apmr.2020.05.023>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

ORIGINAL RESEARCH

Prediction of Psychological Distress Among Persons With Spinal Cord Injury or Acquired Brain Injury and Their Significant Others

Eline W.M. Scholten, MSc,^a Marjolijn Ketelaar, PhD,^a
Johanna M.A. Visser-Meily, MD, PhD,^{a,b} Ellen H. Roels, MD,^c Mirjam Kouwenhoven, MD,^d
POWER Group, Marcel W.M. Post, PhD^{a,c}

From the ^aCenter of Excellence for Rehabilitation Medicine, UMC Utrecht Brain Center, University Medical Center Utrecht, and De Hoogstraat Rehabilitation, Utrecht, The Netherlands; ^bDepartment of Rehabilitation, Physical Therapy Science & Sports, UMC Utrecht Brain Center, University Medical Center Utrecht, The Netherlands; ^cUniversity of Groningen, University Medical Center Groningen, Center for Rehabilitation, Department of Rehabilitation Medicine, Groningen, The Netherlands; and ^dDepartment of Neurological Rehabilitation, De Hoogstraat Rehabilitation, Utrecht, The Netherlands.

Abstract

Objectives: To identify intra- and interpersonal sociodemographic, injury-related, and psychological variables measured at admission of inpatient rehabilitation that predict psychological distress among dyads of individuals with spinal cord injury (SCI) or acquired brain injury (ABI) and their significant others (ie, individuals close to the individual with a disability, mostly family members) 6 months after discharge. Differences in predictors were investigated for persons with SCI or ABI and their significant others and were compared between diagnoses.

Design: Prospective longitudinal study.

Setting: Twelve Dutch rehabilitation centers.

Participants: Dyads (N=157) consisting of adults with SCI or ABI who were admitted to inpatient rehabilitation and their adult significant others.

Interventions: Not applicable.

Main Outcome Measures: Psychological distress (Hospital Anxiety and Depression Scale).

Results: Sociodemographic and injury-related variables were not or were only weakly associated with psychological distress among individuals with SCI or ABI and their significant others 6 months after discharge. Bivariately, higher baseline psychological distress, lower scores on adaptive psychological characteristics (combination of self-efficacy, proactive coping, purpose in life, resilience), and higher scores on maladaptive psychological characteristics (combination of passive coping, neuroticism, appraisals of threat and loss) were related to higher psychological distress, as well as crosswise between individuals with SCI or ABI and their significant others, although less strongly. Combined prediction models showed that psychological distress among persons with SCI or ABI was predicted by education level of their significant other, their own baseline psychological distress, and their own maladaptive psychological characteristics (explained variance, 41.9%). Among significant others, only their own baseline psychological distress predicted psychological distress (explained variance, 40.4%). Results were comparable across diagnoses.

Conclusions: Although a dyadic connection was shown, primarily one's own baseline psychological distress and psychological characteristics were important in the prediction of later psychological distress among both individuals with SCI or ABI and their significant others. Screening based on these variables could help to identify persons at risk for psychological distress.

Archives of Physical Medicine and Rehabilitation 2020;101:2093-102

© 2020 by the American Congress of Rehabilitation Medicine

Chronic conditions such as spinal cord injury (SCI) or acquired brain injury (ABI) are important causes of chronic injury. In 2016, there were 0.9 million new cases of SCI worldwide and 43.6 million cases of ABI (ie, traumatic brain injury, stroke, meningitis).^{1,2} In the Netherlands, nearly two-thirds of all adult patients in inpatient rehabilitation have SCI (11.0%) or ABI (53.2%).³

Having SCI or ABI may have consequences on the well-being of the individuals involved and their significant others (ie, persons close to the individual with a disability, mostly family members). Compared with the general population, persons with SCI (pwSCI) or persons with ABI (pwABI) and their significant others more often experience psychological distress,⁴⁻⁷ defined as “a state of emotional suffering characterized by symptoms of depression or anxiety.”^{8(p107)} It is important to pay attention to psychological distress not only among pwSCI and pwABI, but also among their significant others, even more so because they play an important role in supporting the pwSCI or pwABI.⁹⁻¹¹ To be able to support the pwSCI, pwABI, and their significant others properly, it is important to identify those individuals with a higher risk of psychological distress.

The literature has shown that injury-related (eg, pain severity, physical impairment, and motor function) and sociodemographic variables (eg, age, sex, educational level) are poor and inconsistent predictors of psychological distress among pwSCI, pwABI, and their significant others.¹⁰⁻¹⁴ Early postinjury symptoms of anxiety and depression were found to be important predictors of later psychological distress.^{10,12,13,15} Furthermore, psychological factors including neuroticism, appraisals, coping, optimism, resilience, and self-efficacy were found to be related to psychological distress among pwSCI, pwABI, and their significant others in previous studies.^{10-14,16-19}

Most research on psychological distress after the onset of SCI or ABI focuses on either pwSCI or pwABI or their significant others. However, it is also important to focus on the interdependence of individuals within a dyad.²⁰ For example, previous research has shown that anxiety and depression of pwABI were related to later anxiety and depression among caregivers.^{10,21} A dyadic relation was also found between stroke survivors' self-esteem and partner depression.²² These findings underline the importance of investigating distress among pwSCI or pwABI and their significant others in relation to each other. Furthermore, more insight is needed regarding the variables that are most important in the prediction of psychological distress. SCI and ABI are both conditions characterized by a sudden onset, but with diverging commonly reported consequences. For example, pain is an often reported consequence among pwSCI, and pwABI often experience problems regarding cognitive functioning. Therefore, it is also important to investigate diagnosis-based differences in predictors of psychological distress.

The present study aimed to identify intra- and interpersonal sociodemographic, injury-related, and psychological factors measured at admission of inpatient rehabilitation that predict psychological distress among dyads of pwSCI or pwABI and their significant others 6 months after discharge. The hypotheses were as follows: (1) psychological variables relate more strongly to psychological distress among pwSCI or pwABI and their significant others than sociodemographic and injury-related variables, and (2) intra- and interpersonal psychological variables are both significant predictors of psychological distress.

As an explorative investigation, differences in predictors between diagnoses (ie, SCI, ABI) were studied. Insight in predictors and differences between diagnoses can contribute to the early identification of individuals who are vulnerable for psychological distress after discharge from inpatient rehabilitation and, therefore, provides knowledge to help optimize rehabilitation care.

Methods

Design

The current analyses are part of the POWER study.²³ The overall aim of the cohort part of this study was to identify predictors of empowerment and adjustment among dyads of pwSCI or pwABI and their significant others. The study was conducted in 12 Dutch rehabilitation centers. Dyads were included between April 2016 and July 2018. The Medical Ethics Committee of the University Medical Center Utrecht declared that this study did not require approval according to the Dutch Law on Medical Research (protocol 15-617/C). Boards of all participating rehabilitation centers provided approval to perform the study.

Participants

PwSCI and pwABI were eligible to participate if they met the inclusion criteria: first inpatient rehabilitation after the onset of injury (no inpatient readmission), expected inpatient stay of 4 weeks or longer, age of 18 years or older, and ability to name a significant other (partner, other family member, or close friend). Exclusion criteria were expectation of a full or nearly full recovery, discharge to a long-term care facility, or limited life expectation based on clinical judgment by rehabilitation physicians, or when they were not able to respond to questionnaires because of severe cognitive disabilities, as assessed by nurses based on their clinical view and the Dutch aphasia scale.²⁴ Significant others had to be at least 18 years of age. All participants signed informed consent. Data of dyads in which both individuals completed the baseline and follow-up assessments were used in the current study.

Procedure

Soon after admission, dyads were informed about the study and a few days later their willingness to participate was investigated. Participants completed self-report questionnaires (print or online, according to personal preference). The baseline assessment was completed on average 2 weeks after inpatient admission. Baseline injury-related information was obtained from the medical file. The follow-up assessment was conducted 6 months after discharge (print or online).

List of abbreviations:

ABI	acquired brain injury
A-PC	adaptive psychological characteristics scale
HADS	Hospital Anxiety and Depression Scale
MANCOVA	multivariate analysis of covariance
M-PC	maladaptive psychological characteristics scale
pwABI	persons with ABI
pwSCI	persons with SCI
SCI	spinal cord injury
USER	Utrecht Scale for Evaluation of Rehabilitation

Measures

At both baseline and follow-up, psychological distress was measured with the Hospital Anxiety and Depression Scale (HADS),²⁵ which is an effective measure of general psychological distress.^{26,27} Fourteen questions answered on a 4-point scale assess symptoms of anxiety and depression. A total HADS sum score was computed in which higher scores reflected greater psychological distress (range, 0-42). The HADS has been used in various populations and has shown good clinimetric properties.²⁸

Baseline sociodemographic variables included sex (male, 0; female, 1), age (y), and education (low, 0 [ie, <bachelor degree]; high, 1 [ie, ≥bachelor degree]), and type of relationship with the pwSCI or pwABI (ie, partner, parent, child, other family, friend, or other).

Baseline injury-related variables included diagnosis (SCI, 0; ABI, 1), level of SCI (paraplegia, 0; tetraplegia, 1), completeness of SCI (A-D, 0-3),²⁹ and location of ABI. Physical independence was measured with the sum score of the mobility and self-care scales of the Utrecht Scale for Evaluation of Clinical Rehabilitation (USER).³⁰ Higher scores indicated greater physical independence (range, 0-70). Causes of SCI and ABI were categorized as traumatic (0) or non-traumatic (1).

The psychological measures assessed at baseline are shown in table 1.³¹⁻⁴³ All measures were found to be valid and reliable,^{28,32,37,42-45} and the internal consistency figures of all measures were satisfactory (Cronbach's alpha, ≥0.7) in the current study.⁴⁶

Statistical analysis

Data were analyzed with IBM SPSS Statistics 25.^a A significance level of *P* less than .05 (2-tailed) was used. To reduce the number of psychological variables, we clustered these in 2 scales based on exploratory factor analysis (principal components extraction, oblim rotation) using data of all participants who had completed the baseline assessment (n=223), similar to the method used in a previous study.⁴⁷ The adaptive psychological characteristics scale (A-PC) included self-efficacy, proactive coping, purpose in life, and resilience (pwSCI/pwABI: Eigenvalue, 3.58; 51.14% explained variance; factor loadings, .64-.87; significant others: Eigenvalue, 3.66; 52.26% explained variance; factor loadings, .59-.85). The maladaptive psychological characteristics scale (M-PC) included passive coping,

neuroticism, and appraisals of threat and loss (pwSCI/pwABI: Eigenvalue, .95; 13.55% explained variance; factor loadings, .72-.90; significant others: Eigenvalue, .94; 13.48% explained variance; factor loadings, .72-.96). Both psychological variables were calculated as the mean of the scores of the underlying scales, which were first standardized into z scores to obtain a common metric. A maximum of 1 missing score on an underlying scale was allowed.

The HADS scores were positively skewed and were transformed (square root). Descriptive statistics report raw data, and statistical analyses were carried out on transformed data.

Missing sociodemographic data (total, 8 data points) were imputed with the score of the other person in the dyad. Other missing baseline scores (6 USER, 1 HADS, and 1 A-PC and M-PC) were imputed with the mean score within the corresponding scale, adjusted for diagnosis. Missing follow-up HADS-scores were not imputed.

Independent samples *t* tests and Pearson chi-square tests were conducted to investigate sociodemographic, injury-related, psychological differences between dropped-out and analyzed dyads, as well as between SCI and ABI.

To investigate our first hypothesis, we calculated Pearson correlations between potential predictors and follow-up psychological distress among pwSCI or pwABI and their significant others. Multivariate analyses of covariance (MANCOVA) were used to investigate our second hypothesis. MANCOVA takes into account nonindependence between individuals within a dyad with the possibility to analyze 2 interdependent outcome variables simultaneously, in our case psychological distress of pwSCI or pwABI and their significant others.²⁰ The results provided insight into the independent predictors of psychological distress at dyad level (multivariate effects) and at the level of pwSCI or pwABI and their significant others separately (between-subjects effects). Only variables that were bivariately significantly related with at least 1 dependent variable (follow-up HADS of the pwSCI, pwABI, or significant other) were added in the MANCOVA. In the first model, only sociodemographic and injury-related variables were entered as predictors. In model 2a through 2c, respectively, early post-injury psychological distress, A-PC, and M-PC were additionally entered. The final model (model 3) contained all variables together in a single model. All analyses were repeated for SCI and ABI separately to explore diagnosis-based differences.

Table 1 Psychological variables and the measures used

Independent Variable	Measure	No. of Items	Range Score	Higher Score Indicates
Self-efficacy	ALCOS-12 ^{31,32,*}	12	12-60	Higher self-efficacy
Proactive coping competencies	Shortened Utrecht Proactive Coping Competence scale ³⁴	7	1-4	Better proactive coping competency
Purpose in life	Purpose in Life Scale-Short Form ^{35,36}	4	4-28	Higher purpose in life
Resilience	Connor-Davidson Resilience Scale ^{37,38}	10	0-40	Higher resilience capacity
Passive coping	Passive reaction pattern subscale of the Utrecht Coping List ^{39,40}	7	7-28	Greater tendency to adopt a passive coping style
Neuroticism	Neuroticism subscale of the Eysenck Personality Questionnaire-Revised Short Form ^{41,42}	12	0-12	Higher levels of neuroticism
Appraisals	Threat and loss subscales of the Appraisals of Life Events scale ^{43,†}	10	0-5	More appraisals of threat and loss

* The ALCOS-12 is the Dutch version of the General Self Efficacy originally developed by Sherer.³³

† The threat subscale contains 6 items, and the loss subscale contains 4 items. Subscale scores were computed as the mean of the respective item scores. A total score was computed as the mean of the 2 subscale scores so that both subscales contributed equally to the total score.

Fig 1 Flowchart showing the inclusion of dyads of pwSCI or pwABI and their significant others.

Results

Background: participants and main outcomes

Data of 157 dyads were analyzed (fig 1). The main reasons for excluding dyads were expected stay in inpatient rehabilitation less than 4 weeks (26.0%), limited life expectation (16.3%), no significant other (15.2%), or severe cognitive disabilities (13.3%). The main reasons for declined participation were “no interest” (45.2%) or “too burdensome” (34.0%). Table 2 shows sociodemographic and injury-related information of the analyzed dyads. The cause of SCI was traumatic in half the cases (fall, 21.5%; sport/leisure accident, 15.2%; traffic accident, 12.7%; occupational accident, 1.3%). Non-traumatic causes included spinal degeneration (11.4%), tumor (11.4%), inflammation (10.1%), medical complication (8.9%), and vascular diseases (7.6%). Causes of ABI were mostly nontraumatic (infarction, 48.7%; hemorrhage, 26.9%; or “other” such as meningitis, 11.5%). Table 2 shows the differences between sociodemographic and injury-related variables between the analyzed dyads and those who dropped out during follow-up. Score distributions of the psychological variables are shown in table 3. No differences were found regarding psychological variables between the analyzed dyads and those that dropped out.

Hypothesis 1: correlations

Of all sociodemographic and injury-related variables, only higher education level of the significant other was related to lower psychological distress among pwSCI or pwABI. A diagnosis of ABI was related to lower psychological distress among the significant others (table 4). With respect to psychological factors, higher early postinjury psychological distress, higher M-PC, and lower A-PC were related to higher psychological distress at follow-up, both intrapersonal and crosswise, although less strongly, between pwSCI or pwABI and their significant others.

Hypothesis 2: prediction at dyad level (multivariate effect)

The first MANCOVA model (table 5) included only educational level (significant other) and diagnosis. Both variables were

significant predictors of psychological distress at the dyad level. Model 2a through 2c showed that early post-injury psychological distress (of pwSCI/pwABI and significant others), A-PC (of both), and M-PC (of both) were significant predictors of psychological distress at the dyad level, adjusted for significant others’ education level and diagnosis. The final model, which contained all variables together, showed that only diagnosis, early postinjury psychological distress (of both), and M-PC (of the pwSCI or pwABI) remained significant predictors of psychological distress at the dyad level.

Hypothesis 2: predictors for pwSCI/ABI (between-subjects effects)

Adjusted for educational level of the significant other and diagnosis, early postinjury psychological distress, A-PC, and M-PC among pwSCI and pwABI were significant predictors of psychological distress among pwSCI and pwABI when tested in separated models (see table 5, model 2a-c). The final model showed that, when adjusted for other variables, educational level of the significant other, early postinjury psychological distress (pwSCI/pwABI), and M-PC (pwSCI/pwABI) were significant predictors of psychological distress at follow-up (explained variance, 41.9%).

Hypothesis 2: predictors for significant others (between-subjects effects)

Adjusted for significant others’ educational level and diagnosis, significant others’ scores on early postinjury psychological distress, A-PC, and M-PC were significant predictors of follow-up psychological distress when tested in separated models (see table 5, model 2a-c). The final model showed that, when adjusted for other variables, early postinjury psychological distress of the significant others themselves was the only significant predictor of psychological distress at follow-up (explained variance, 40.4%).

Exploratory: SCI vs ABI

Differences between SCI and ABI were found in the level of physical independence (SCI was lower), cause of injury (SCI was more often traumatic), and sex of the significant other (SCI was more often female) (see table 2). Significant others of pwSCI reported higher baseline and follow-up psychological distress than significant others of pwABI (see table 3). PwSCI and their significant others reported higher levels of appraisals of threat and loss (resulting in higher M-PC scores) than individuals in the ABI subgroup.

Correlation and MANCOVA analyses were repeated for SCI and ABI separately (supplemental tables S1-S3, available online only at <http://www.archives-pmr.org/>). In the final MANCOVA model for SCI, early postinjury psychological distress of the pwSCI and that of the significant other were found to be significant predictors of psychological distress at the dyad level. Among pwSCI, only the individual’s own early postinjury psychological distress and M-PC were found to be significant predictors. Among significant others, their own early postinjury psychological distress was found to be the only significant predictor.

In ABI, the final MANCOVA model showed that early post-injury psychological distress of the significant other was the only predictor of psychological distress at the dyad level. Among

Table 2 Sociodemographic and injury-related characteristics of pwSCI or pwABI and their significant others

Variable	Total (N=157)		Dropped Out (n=66)	SCI (n=79)	ABI (n=78)
	n	n (%) or Mean $\pm$ SD, Range	n (%) or Mean $\pm$ SD, Range	n (%) or Mean $\pm$ SD, Range	n (%) or Mean $\pm$ SD, Range
pwSCI/pwABI					
Sex (female)	157	67 (42.7)	18 (27.3)	28 (35.4)	39 (50.0)
Age, y	157	56.3 $\pm$ 14.9, 18-87	57.8 $\pm$ 14.4, 22-84	55.3 $\pm$ 16.3, 18-81	57.3 $\pm$ 13.3, 26-87
Education (high)*	154	58 (37.7)	15 (23.4)	25 (32.9)	33 (42.3)
Physical independence (USER, 0-70) [†]	151	36.4 $\pm$ 19.0, 1-70	29.3 $\pm$ 15.8, 1-70 [‡]	28.8 $\pm$ 17.9, 1-70 [§]	44.4 $\pm$ 16.9, 5-70 [§]
Cause of injury (non-traumatic)	—	—	—	39 (49.4) [§]	68 (87.2) [§]
AIS (SCI only)					
A	—	—	—	10 (12.7)	—
B	—	—	—	12 (15.2)	—
C	—	—	—	17 (21.5)	—
D	—	—	—	40 (50.6)	—
Tetraplegia (SCI only)	—	—	—	44 (55.7)	—
Location (ABI only)					
Left	—	—	—	—	31 (39.7)
Right	—	—	—	—	27 (34.6)
Both sides	—	—	—	—	13 (16.7)
Brainstem	—	—	—	—	3 (3.9)
Unknown	—	—	—	—	4 (5.2)
Weeks between admission and completing baseline assessment	157	2.0 (2.0) [§]	2.0 (2.0) [¶]	2.0 (2.0) [¶]	2.0 (2.0) [¶]
Significant others					
Sex (female)	157	97 (61.8)	52 (78.8)	55 (69.6) [§]	42 (53.8) [§]
Age, y	155	55.5 $\pm$ 12.4, 23-82	51.0 $\pm$ 13.1, 25-79 [‡]	56.6 $\pm$ 12.9, 25-82	54.4 $\pm$ 11.9, 23-75
Education (high)*	155	64 (41.3)	19 (30.6)	31 (39.7)	33 (42.9)
Relationship with pwSCI/pwABI					
Partner	157	123 (78.3)	40 (62.5) [‡]	60 (75.9)	63 (80.8)
Parent	—	14 (8.9)	5 (7.8)	9 (11.4)	5 (6.4)
Child	—	11 (7.0)	16 (25.0)	5 (6.3)	6 (7.7)
Other family	—	5 (3.2)	2 (3.1)	2 (2.5)	3 (3.8)
Friend	—	3 (1.9)	—	2 (2.5)	1 (1.3)
Other	—	1 (0.6)	1 (1.6)	1 (1.3)	—

Abbreviation: AIS, American Spinal Injury Association Impairment Scale.

* High education indicates a bachelor degree or higher.

[†] Physical independence was measured with the Utrecht Scale for Evaluation of Clinical Rehabilitation. Higher scores indicated greater physical independence (range, 0-70).

[‡] Independent samples *t* tests or Pearson χ^2 test showed a significant difference between analyzed and dropped out dyads.

[§] Independent samples *t* tests or Pearson χ^2 test showed a significant difference between SCI and ABI.

^{||} American Spinal Injury Association Impairment Scale. A, complete SCI; B, sensory incomplete; C, motor incomplete with less than half of key muscle functions below the single neurological level of injury having a muscle grade ≥ 3 ; D, motor incomplete with at least half of key muscle functions below the single neurological level of injury having a muscle grade ≥ 3 .²⁹

[¶] Median and interquartile range is given rather than mean and SD.

pwABI, none of the variables was a significant predictor of psychological distress when adjusted for other variables. Among significant others, their own early postinjury psychological distress was found to be the only significant predictor.

Discussion

By using a dyadic approach and comparing 2 diagnostic groups, this study extended upon earlier research aimed to identify sociodemographic, injury-related, and psychological predictors of later psychological distress among pwSCI or pwABI and significant others.

The hypothesis that psychological variables related more strongly to psychological distress among pwSCI or pwABI and significant others than sociodemographic and injury-related variables was supported and in line with previous research findings.^{10-14,16-19} It was additionally hypothesized that intra- and interpersonal psychological variables were both important in the prediction of psychological distress. As an indication of interdependence between pwSCI or pwABI and significant others,²⁰ correlations were found between psychological variables of 1 person in a dyad and outcome psychological distress of the other person. At the dyad level, psychological variables of both individuals within a dyad did indeed add in the prediction of psychological distress. On the other hand, the crosswise correlations

Table 3 Psychological scores of pwSCI or pwABI and their significant others

	Total (N = 157)		Dropped Out (n = 66)*	SCI (n = 79)	ABI (n = 78)
Variable	N	Mean $\pm$ SD	Mean $\pm$ SD	Mean $\pm$ SD	Mean $\pm$ SD
pwSCI/pwABI					
Psychological distress (HADS, 0-42)					
Baseline [†]	156	12.0 (6.0-17.0) [‡]	12.0 (7.0-19.0) [‡]	13.0 (6.0-17.0) [‡]	12.0 (6.7-17.0) [‡]
Follow-up [§]	157	11.0 (5.0-16.0) [‡]	NA	11.0 (5.0-12.0) [‡]	11.0 (5.8-16.0) [‡]
Adaptive psychological characteristics [§]					
Total	157	-0.0 $\pm$ 0.8	-0.02 $\pm$ 0.8	0.0 $\pm$ 0.8	-0.0 $\pm$ 0.7
Self-efficacy (blinded for peer review, 12-60)	155	48.1 $\pm$ 8.1	48.9 $\pm$ 8.0	49.1 $\pm$ 7.9	47.1 $\pm$ 8.3
Proactive coping (UPCC-SF, 1-4)	157	2.9 $\pm$ 0.5	2.9 $\pm$ 0.5	2.9 $\pm$ 0.5	3.0 $\pm$ 0.4
Purpose in life (PIL-SF, 4-28)	156	22.8 $\pm$ 3.8	22.9 $\pm$ 3.5	23.0 $\pm$ 3.9	22.5 $\pm$ 3.7
Resilience (CD-RISC-10, 0-40)	157	27.7 $\pm$ 6.9	27.2 $\pm$ 6.8	27.6 $\pm$ 7.0	27.8 $\pm$ 6.8
Maladaptive psychological characteristics [§]					
Total	157	-0.0 $\pm$ 0.8	0.1 $\pm$ 0.9	0.1 $\pm$ 0.9	-0.1 $\pm$ 0.8
Passive coping (UCL, 7-28)	156	10.5 $\pm$ 2.9	10.9 $\pm$ 3.3	10.8 $\pm$ 3.1	10.1 $\pm$ 2.7
Neuroticism (EPQ-RSS-N, 0-12)	155	3.1 $\pm$ 3.1	3.1 $\pm$ 3.4	3.3 $\pm$ 3.2	2.9 $\pm$ 3.0
Appraisals of threat and loss (ALE, 0-5)	157	1.8 $\pm$ 1.2	2.0 $\pm$ 1.4	2.0 $\pm$ 1.3	1.5 $\pm$ 1.1
Significant others					
Psychological distress (HADS, 0-42)					
Baseline [†]	157	12.0 (7.0-18.0) [‡]	12.0 (8.0-19.0) [‡]	13.0 (8.0-21.0) ^{‡,}	10.7 (6.0-17.0) ^{‡,}
Follow-up [†]	157	9.0 (4.0-15.0) [‡]	NA	11.0 (5.0-16.0) ^{‡,}	8.0 (3.0-12.0) ^{‡,}
Adaptive psychological characteristics [§]					
Total	156	-0.0 $\pm$ 0.8	0.1 $\pm$ 0.8	-0.1 $\pm$ 0.7	0.1 $\pm$ 0.8
Self-efficacy (blinded for peer review, 12-60)	156	49.7 $\pm$ 6.9	48.8 $\pm$ 8.0	48.8 $\pm$ 6.7	50.5 $\pm$ 7.1
Proactive coping (UPCC-SF, 1-4)	156	3.0 $\pm$ 0.4	3.1 $\pm$ 0.4	3.0 $\pm$ 0.4	3.0 $\pm$ 0.4
Purpose in life (PIL-SF, 4-28)	154	22.7 $\pm$ 3.6	23.0 $\pm$ 3.8	22.4 $\pm$ 3.4	23.0 $\pm$ 3.7
Resilience (CD-RISC-10, 0-40)	155	28.2 $\pm$ 6.1	28.2 $\pm$ 6.0	27.4 $\pm$ 6.2	29.1 $\pm$ 5.8
Maladaptive psychological characteristics [§]					
Total	156	0.0 $\pm$ 0.8	0.1 $\pm$ 1.0	0.2 $\pm$ 0.9	-0.2 $\pm$ 0.7
Passive coping (UCL, 7-28)	156	10.4 $\pm$ 2.7	10.6 $\pm$ 3.0	10.7 $\pm$ 2.9	10.0 $\pm$ 2.4
Neuroticism (EPQ-RSS-N, 0-12)	153	2.8 $\pm$ 2.6	3.6 $\pm$ 3.4	3.0 $\pm$ 2.7	2.7 $\pm$ 2.5
Appraisals of threat and loss (ALE, 0-5)	155	1.3 $\pm$ 1.1	1.4 $\pm$ 2.3	1.6 $\pm$ 1.1	0.9 $\pm$ 0.9

Abbreviations: ALE, Appraisals of Life Events scale; CD-RISC, Connor-Davidson Resilience; EPQ-RSS-N, Neuroticism subscale of the Eysenck Personality Questionnaire-Revised Short Form; NA, not applicable; PIL-SF, Purpose in Life Scale-Short Form; UCL, Utrecht Coping List; UPCC-SF, Shortened Utrecht Proactive Coping Competence scale.

* Independent samples *t* tests showed no significant difference between analyzed and dropped out dyads for any of the variables.

† Assessed at the start of inpatient rehabilitation (baseline).

‡ Median and interquartile range are presented rather than mean and SD.

§ Assessed at 6 months after inpatient discharge (6mo follow-up).

|| Independent samples *t* tests or Pearson χ^2 test showed a significant difference between SCI and ABI.

were relatively weak ($r < .3$) and in combination with the individual's own psychological variables, they were not of added value in the prediction of psychological distress at individual level, which contradicts our hypothesis. This could possibly be explained by the strong correlation between one's own baseline and follow-up psychological distress in combination with a relatively small change in psychological distress scores over time, which might limit the impact of other variables in the explanation.

Separately, early postinjury psychological distress, A-PC, and M-PC were all significant predictors of psychological distress, both at the dyad and individual levels. However, altogether, early postinjury psychological distress and M-PC were the most important predictors of psychological distress among pwSCI and pwABI, and early post-injury psychological distress was the most important among significant others. Possibly, the concept of "maladaptive psychological characteristics" measured with the

M-PC is more similar to the HADS (measuring anxiety and depression) than the concept measured with the A-PC, which may explain why A-PC was found to be a less important predictor of psychological distress. This idea is supported by the relatively strong correlation between the baseline HADS and M-PC (pwSCI/pwABI: $r = .76$; significant others: $r = .80$).

Exploratory differences between the diagnostic groups were investigated. We did find some diagnosis-based differences in levels of variables and correlations, but not in the predictors of psychological distress. In both diagnosis subgroups, the individual's own early postinjury psychological distress and M-PC appeared to be the most important predictors. Results were less consistent in the ABI subgroup owing to an absence of significant predictors to predict psychological distress among pwABI. However, *F* values found among pwABI were comparable to those found among pwSCI and *P* values were close to .05. Future

Table 4 Bivariate correlation coefficients between independent variables and psychological distress of the pwSCI or pwABI and significant others 6 months after discharge from inpatient rehabilitation (N=157)

Variable	Psychological Distress pwSCI/pwABI*	Psychological Distress Significant Others*
	Correlation Coefficient (P Value)	Correlation Coefficient (P Value)
Sex (female)		
pwSCI/pwABI	.10 (.20)	-.08 (.33)
Significant others	-.04 (.61)	.05 (.56)
Age		
pwSCI/pwABI	-.06 (.46)	.07 (.41)
Significant others	.02 (.83)	.10 (.24)
Education (high)		
pwSCI/pwABI	-.13 (.12)	.03 (.72)
Significant others	-.24 (<.01) [†]	-.07 (.40)
Significant other is partner	.02 (.85)	.04 (.60)
Diagnosis (ABI)	.01 (.94)	-.22 (<.01) [†]
Physical independence	.02 (.78)	-.12 (.14)
Psychological distress [‡]		
pwSCI/pwABI	.57 (<.001) [§]	.12 (.12)
Significant others	.21 (<.01) [†]	.61 (<.001) [§]
Adaptive psychological characteristics [‡]		
pwSCI/pwABI	-.46 (<.001) [§]	-.19 (.02)
Significant others	-.25 (.001) [†]	-.37 (<.001) [§]
Maladaptive psychological characteristics [‡]		
pwSCI/pwABI	.57 (<.001) [§]	.21 (<.01) [†]
Significant others	.17 (.03)	.49 (<.001) [§]

* Assessed at 6 months after inpatient discharge (6mo follow-up).

† $P < .01$.

‡ Assessed at the start of inpatient rehabilitation (baseline).

§ $P < .001$.

|| $P < .05$.

research should be conducted to explore whether the differences reflect diagnosis-based differences in the prediction of psychological distress, or if the differences could be explained by the limited power in our sample.

Study limitations

First, to increase the power, we combined different psychological scales, with the result that no statements can be made about the predictive value of individual psychological variables. Exploratory factor analysis and a comparable application in previous research supported our decision to cluster the scales.⁴⁷ Second, a relatively large number of the dyads did not meet the inclusion criteria or declined to participate, which may have resulted in selection or nonresponse bias. Unfortunately, we have no information on the excluded and declined dyads. However, regarding age, sex, and injury-specific information (completeness of SCI, level of SCI, physical independence pwABI), our sample appears to be highly comparable to the general Dutch SCI and stroke population in an inpatient setting.^{48,49} Third, selective loss to follow-up may have influenced the results. However, this seems to be unlikely as scores on the significant predictors of psychological distress did not differ between the analyzed and dropped out dyads. Fourth, half of the pwSCI had an AIS D score. They have a better prognosis than pwSCI with a score of A through C, possibly resulting in lower psychological distress levels at baseline and follow-up. However, because we found that injury-related factors did not

add to the prediction of psychological distress, we do not expect major differences in the results when a smaller proportion of the pwSCI would have had an AIS score of D. Fifth, different kinds of significant others participated in the study. Most significant others (78.3%) were partners and, therefore, the results apply mainly to situations in which the significant other is the partner. Lastly, previous research has shown that a history of preinjury psychological problems is an important predictor of postinjury psychological distress.¹¹⁻¹⁴ We have not assessed preinjury psychological problems. Alternatively, we included the HADS at baseline, which was considered an indicator for vulnerability at admission.

Implications

Our results highlight the importance of one's own early postinjury psychological distress and adaptive and maladaptive psychological characteristics in the prediction of later psychological distress. A screening based on these variables conducted at the start of inpatient rehabilitation could help to identify individuals more at risk for psychological distress earlier. Screening only the pwSCI or pwABI does not appear to be sufficient to accurately predict psychological distress among both individuals within the dyad. It is also important to obtain insight into distress among significant others because significant others play an important, and often necessary, role in providing practical and emotional support to pwSCI and pwABI.^{50,51} If significant others become overloaded, this has adverse consequences for themselves, but also for the

Table 5 MANCOVA to predict psychological distress 6 months after discharge from inpatient rehabilitation (N=157)

Independent Variables	Model 1: Basic Model*			Model 2a: Psychological Distress†			Model 2b: Adaptive Psychological Characteristics‡		
	Between-Subjects Effects			Between-Subjects Effects			Between-Subjects Effects		
	Multivariate	Distress pwSCI/pwABI	Distress SO	Multivariate	Distress pwSCI/pwABI	Distress SO	Multivariate	Distress pwSCI/pwABI	Distress SO
	V [‡] (P Value)	F (P Value)	F (P Value)	V [‡] (P Value)	F (P Value)	F (P Value)	V [‡] (P Value)	F (P Value)	F (P Value)
Education (high)									
pwSCI/pwABI	—	—	—	—	—	—	—	—	—
SO	.06 (<.01) [§]	9.78 (<.01) [§]	.54 (.47)	.03 (.08)	4.90 (.03)	.14 (.71)	.04 (.04)	6.33 (.01)	.00 (.99)
Diagnosis (ABI)	.06 (.01)	.05 (.83)	7.63 (<.01) [§]	.04 (.05)	.74 (.39)	3.68 (.06)	.04 (.04)	.02 (.89)	5.89 (.02)
Psychological distress†									
pwSCI/pwABI	—	—	—	.32 (<.001) [¶]	61.95 (<.001) [¶]	.00 (.96)	—	—	—
SO	—	—	—	.35 (<.001) [¶]	2.96 (.09)	80.73 (<.001) [¶]	—	—	—
Adaptive psychological characteristics‡									
pwSCI/pwABI	—	—	—	—	—	—	.16 (<.001) [¶]	29.72 (<.001) [¶]	1.54 (.22)
SO	—	—	—	—	—	—	.10 (<.001) [¶]	1.73 (.19)	16.24 (<.001) [¶]
Maladaptive psychological characteristics‡									
pwSCI/pwABI	—	—	—	—	—	—	—	—	—
SO	—	—	—	—	—	—	—	—	—
Explained variance (%)	NA	6.0	5.2	NA	36.0	38.6	NA	25.5	17.7
	Model 2c: Maladaptive Psychological Characteristics‡						Model 3: Final Model		
	Between-Subjects Effects			Between-Subjects Effects			Between-Subjects Effects		
	Multivariate	Distress pwSCI/pwABI	Distress SO	Multivariate	Distress pwSCI/pwABI	Distress SO	Multivariate	Distress pwSCI/pwABI	Distress SO
	V [‡] (P Value)	F (P Value)	F (P Value)	V [‡] (P Value)	F (P Value)	F (P Value)	F (P Value)	F (P Value)	F (P Value)
Education (high)									
pwSCI/pwABI	—	—	—	—	—	—	—	—	—
SO	.05 (.02)	8.21 (<.01) [§]	.41 (.52)	.04 (.07)	5.23 (.02)	.05 (.82)	.05 (.03)	1.75 (.19)	3.27 (.07)
Diagnosis (ABI)	.05 (.03)	3.23 (.07)	2.09 (.15)	.05 (.03)	1.75 (.19)	3.27 (.07)	.05 (.03)	1.75 (.19)	3.27 (.07)
Psychological distress†									
pwSCI/pwABI	—	—	—	—	.07 (<.01) [§]	6.07 (.02)	6.07 (.02)	1.43 (.23)	1.43 (.23)
SO	—	—	—	—	.18 (<.001) [¶]	.29 (.59)	.29 (.59)	31.46 (<.001) [¶]	31.46 (<.001) [¶]
Adaptive psychological characteristics‡									
pwSCI/pwABI	—	—	—	—	.01 (.45)	.84 (.36)	.84 (.36)	1.27 (.26)	1.27 (.26)
SO	—	—	—	—	.01 (.55)	.31 (.58)	.31 (.58)	1.15 (.29)	1.15 (.29)
Maladaptive psychological characteristics‡									
pwSCI/pwABI	.32 (<.001) [¶]	70.20 (<.001) [¶]	1.65 (.20)	.07 (<.01) [§]	10.90 (<.01) [§]	.41 (.52)	10.90 (<.01) [§]	.41 (.52)	.41 (.52)
SO	.20 (<.001) [¶]	1.09 (.30)	36.53 (<.001) [¶]	.00 (1.00)	.00 (1.00)	.00 (.98)	.00 (1.00)	.00 (.98)	.00 (.98)
Explained variance (%)	NA	37.8	26.0	NA	41.9	40.4	41.9		40.4

Abbreviations: NA, not applicable; SO, significant other.

* Only bivariately significant variables ($P < .05$) were added in the MANCOVA models.[†] Assessed at the start of inpatient rehabilitation (baseline).[‡] Pillai's trace value.[§] $P < .01$.^{||} $P < .05$.[¶] $P < .001$.

pwSCI or pwABI, possibly even resulting in institutionalization.^{52,53} Therefore, we recommend screening both pwSCI or pwABI and their significant others.

Risk screening also helps to personalize the support provided to pwSCI, pwABI, and their significant others, thereby making the most efficient use of available resources.⁵⁴ Previous research has shown that tailored interventions provide more positive outcomes than the application of standard interventions.⁵⁵ Interventions that apply psychological techniques (eg, cognitive behavioral therapy, coping skill-training, problem-solving therapy) appear to be most effective in the reduction of psychological distress.⁵⁵

Finally, identifying pwSCI or pwABI and significant others at greater risk for psychological distress earlier and providing them with appropriate support could be crucial in reducing healthcare system burden and costs.⁵⁶⁻⁵⁸

Conclusions

There is a dyadic connection between early postinjury psychological distress, psychological variables, and follow-up psychological distress among pwSCI or pwABI and their significant others. However, primarily one's own early postinjury psychological distress and psychological variables were important in the prediction of follow-up psychological distress. The results were comparable for SCI and ABI. A psychological screening of patients and significant others could help with early identification of individuals more at risk for later psychological distress.

Acknowledgments

The members of the POWER Group are: Rehabilitation enter Adelante: C.C.M. van Laake-Geelen; Rehabilitation Center De Hoogstraat: J. Stolwijk, C.A. Dijkstra, E. Agterhof; Rehabilitation Center Helimare: D. Gobets; Rehabilitation Center Het Roesingh: E.M. Maas; Rehabilitation Center Merem/De Trappenberg: H. van der Werf, C.E. de Boer; Rehabilitation Center Reade: M. Beurskens; Rehabilitation Center Sint Maartenskliniek: I. van Nes, T. van Diemen; Rehabilitation Friesland: K.H. Woldendorp, J. Hurkmans; Revant Rehabilitation Center: M. Luijkx; Rijndam Rehabilitation: D.C.M. Spijkerman, R. Osterthun; UMCG Center for Rehabilitation: J. Sprik-Bakker, E.H. Roels; and Vogellanden Center for Rehabilitation: M. Hoonhorst.

Supplier

a. SPSS; IBM Corp.

Keywords

Brain injuries; Caregivers; Mental health; Rehabilitation; Spinal cord injuries

Corresponding author

Marcel W.M. Post, PhD, De Hoogstraat Rehabilitation, Rembrandtkade 10, 3583 TM, Utrecht, The Netherlands. *E-mail address:* m.post@dehoogstraat.nl.

References

1. GBD 2016 Traumatic Brain Injury and Spinal Cord Injury Collaborators. Global, regional, and national burden of traumatic brain injury and spinal cord injury, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet Neurol* 2019;18:56–87.
2. GBD 2016 Stroke Collaborators. Global, regional, and national burden of stroke, 1990–2016: A systematic analysis for the Global Burden of Disease Study 2016. *Lancet Neurol* 2019;18:439–58.
3. Revalidatie Nederland, Branchgegevens 2017 Revalidatie Nederland [Branch report 2017 Rehabilitation The Netherlands], Utrecht. 2018. Available at: https://www.revalidatie.nl/userfiles/File/publicaties/Brancher_apport_RN_2017_DIGI.pdf. Accessed July 16, 2020.
4. Robinson RG, Jorge RE. Post-stroke depression: a review. *Am J Psychiatry* 2016;173:221–31.
5. Loh AZ, Tan JS, Zhang MW, Ho RC. The global prevalence of anxiety and depressive symptoms among caregivers of stroke survivors. *J Am Med Dir Assoc* 2017;18:111–6.
6. Post MWM, Van Leeuwen CMC. Psychosocial issues in spinal cord injury: a review. *Spinal Cord* 2012;50:382–9.
7. Lynch J, Cahalan R. The impact of spinal cord injury on the quality of life of primary family caregivers: a literature review. *Spinal Cord* 2017;55:964–78.
8. Drapeau A, Marchand A, Beaulieu-Prévost D. Epidemiology of psychological distress. In: L'Abate L, editor. *Mental illnesses - understanding, prediction and control*. Rijeka, Croatia: IntechOpen; 2012.
9. Müller R, Peter C, Cieza A, Geyh S. The role of social support and social skills in people with spinal cord injury: a systematic review of the literature. *Spinal Cord* 2012;50:94–106.
10. Kruijthof WJ, Post MWM, Van Mierlo ML, Van den Bos GAM, De Man-van Ginkel JM, Visser-Meily JMA. Caregiver burden and emotional problems in partners of stroke patients at two months and one year post-stroke: determinants and prediction. *Patient Educ Couns* 2016;99:1632–40.
11. Davis LC, Sander AM, Struchen MA, Sherer M, Nakase-Richardson R, Malec JF. Medical and psychosocial predictors of caregiver distress and perceived burden following traumatic brain injury. *J Head Trauma Rehabil* 2009;24:145–54.
12. Menlove L, Crayton E, Kneebone I, Allen-Crooks R, Otto E, Harder H. Predictors of anxiety after stroke: a systematic review of observational studies. *J Stroke Cerebrovasc Dis* 2015;24:1107–17.
13. Pollock K, Dorstyn D, Butt L, Prentice S. Posttraumatic stress following spinal cord injury: a systematic review of risk and vulnerability factors. *Spinal Cord* 2017;55:800–11.
14. Shi Y, Yang D, Zeng Y, Wu W. Risk factors for post-stroke depression: a meta-analysis. *Front Aging Neurosci* 2017;9:218.
15. Kouwenhoven SE, Kirkevold M, Engedal K, Kim HS. Depression in acute stroke: prevalence, dominant symptoms and associated factors. A systematic literature review. *Disabil Rehabil* 2011;33:539–56.
16. Van Mierlo ML, Van Heugten CM, Post MW, De Kort PL, Visser-Meily JM. Psychological factors determine depressive symptomatology after stroke. *Arch Phys Med Rehabil* 2015;96:1064–70.
17. Greenop KR, Almeida OP, Hankey GJ, Van Bockxmeer F, Lautenschlager NT. Premorbid personality traits are associated with post-stroke behavioral and psychological symptoms: a three-month follow-up study in Perth, Western Australia. *Int Psychogeriatrics* 2009;21:1063–71.
18. Noh S-M, Chung SJ, Kim K-K, et al. Emotional disturbance in CADASIL: its impact on quality of life and caregiver burden. *Cerebrovasc Dis* 2014;37:188–94.
19. Elliott TR, Berry JW, Richards JS, Shewchuk RM. Resilience in the initial year of caregiving for a family member with a traumatic spinal cord injury. *J Consult Clin Psychol* 2014;82:1072–86.
20. McCarthy MJ, Lyons KS, Powers LE. Expanding poststroke depression research: movement toward a dyadic perspective. *Top Stroke Rehabil* 2011;18:450–60.

21. Hultman MT, Everson-Rose SA, Tracy MF, Lindquist R, Niakosari Hadidi N, Hadidi NN. Associations between characteristics of stroke survivors and caregiver depressive symptoms: a critical review. *Top Stroke Rehabil* 2019;26:528-37.
22. Chung ML, Bakas T, Plue LD, Williams LS. Effects of self-esteem, optimism, and perceived control on depressive symptoms in stroke survivor-spouse dyads. *J Cardiovasc Nurs* 2016;31:E8-16.
23. Hillebregt CF, Scholten EWM, Ketelaar M, Post MWM, Visser-Meily JMA. Effects of family group conferences among high-risk patients of chronic disability and their significant others: study protocol for a multicentre controlled trial. *BMJ Open* 2018;8:e026768.
24. Deelman BG, Koning-Haanstra M, Liebrand WBG, Van den Burg W. SAN Test, een afasie test voor auditief en mondeling taalgebruik [SAN Test, an aphasia test for auditory and oral language] [in Dutch]. Lisse, the Netherlands: Swets & Zeitlinger; 1981.
25. Zigmond AS, Snaith RP. The Hospital Anxiety and Depression Scale. *Acta Psychiatr Scand* 1983;67:361-70.
26. Cosco TD, Doyle F, Ward M, McGee H. Latent structure of the Hospital Anxiety and Depression Scale: a 10-year systematic review. *J Psychosom Res* 2012;72:180-4.
27. Norton S, Cosco T, Doyle F, Done J, Sacker A. The Hospital Anxiety and Depression Scale: a meta confirmatory factor analysis. *J Psychosom Res* 2013;74:74-81.
28. Spinhoven P, Ormel J, Sloekers PPA, Kempen GIJM, Speckens AEM, Van Hemert AM. A validation study of the Hospital Anxiety and Depression Scale (HADS) in different groups of Dutch subjects. *Psychol Med* 1997;27:363-70.
29. Kirshblum SC, Burns SP, Biering-Sorensen F, et al. International standards for neurological classification of spinal cord injury. *Spinal Cord* 2011;34:535-46.
30. Post MWM, Van de Port IGL, Kap B, Van Berlekom BSH. Development and validation of the Utrecht Scale for Evaluation of Clinical Rehabilitation (USER). *Clin Rehabil* 2009;23:909-17.
31. Bosscher RJ, Smit JH, Kempen G. Algemene competentieverwachtingen bij ouderen [General competence expectations in the elderly] [in Dutch]. *Ned Tijdschr voor Psychol* 1997;52:239-48.
32. Bosscher RJ, Smit JH. Confirmatory factor analysis of the General Self-Efficacy Scale. *Behav Res Ther* 1998;36:339-43.
33. Sherer M, Maddux JE, Mercandante B, Prentice-Dunn S, Jacobs B, Rogers RW. The Self-Efficacy Scale: construction and validation. *Psychol Rep* 1982;51:663-71.
34. Bode C, Thoolen B, De Ridder D. Measuring proactive coping. Psychometric characteristics of the Utrecht Proactive Coping Competence scale (UPCC). *Psychol Gezondh* 2008;36:81-91.
35. Crumbaugh JC, Maholick LT. An experimental study in existentialism: the psychometric approach to Frankl's concept of noogenic neurosis. *J Clin Psychol* 1964;20:200-7.
36. Schulenberg SE, Schnetzer LW, Buchanan EM. The Purpose in Life Test-Short Form: development and psychometric support. *J Happiness Stud* 2011;12:861-76.
37. Campbell-Sills L, Stein MB. Psychometric analysis and refinement of the Connor-Davidson Resilience Scale (CD-RISC): validation of a 10-item measure of resilience. *J Trauma Stress* 2007;20:1019-28.
38. Connor KM, Davidson JRT. Development of a new resilience scale: the Connor-Davidson Resilience Scale (CD-RISC). *Depress Anxiety* 2003;18:76-82.
39. Schreurs PJG, Tellegen B, Van de Willige G. Gezondheid, stress en coping: De ontwikkeling van de Utrechtse Coping Lijst [Health, stress, and coping: The development of the Dutch Coping List] [in Dutch]. *Tijdschr voor Psychol* 1984;12:101-11.
40. Schreurs PJG, Van de Willige G, Brosschot JF, Tellegen B, Graus GMH, editors. Handleiding Utrechtse Coping Lijst UCL (herziene versie) [Manual Utrecht Coping List UCL (revised version)] [in Dutch]. Lisse, the Netherlands: Swets & Zeitlinger; 1993.
41. Eysenck HJ, Eysenck SBG. Manual of the Eysenck Personality Scales (EPS Adult). London: Hodder & Stoughton; 1991.
42. Sanderman R, Arrindell W, Ranchor A, Eysenck H, Eysenck S. Het meten van persoonlijkheidskenmerken met de Eysenck Personality Questionnaire (EPQ), een handleiding [Measuring personality characteristics with the Eysenck Personality Questionnaire (EPQ), a guide] [in Dutch]. 2nd ed. Groningen, the Netherlands: UMC-G/Rijksuniversiteit Groningen, Research Institute SHARE; 2012.
43. Ferguson E, Matthews G, Cox T. The Appraisal of Life Events (ALE) scale: reliability and validity. *Br J Health Psychol* 1999;4:97-116.
44. Schaufeli W, Van Dierendonck D. De betrouwbaarheid en validiteit van de Utrechtse Coping Lijst: Een longitudinaal onderzoek bij schoolverlaters [Reliability and validity of the Utrecht Coping List. A longitudinal study in high school alumni] [in Dutch]. *Gedrag en Gezondh* 1992;20:38-45.
45. Tielemans NS, Visser-Meily JM, Schepers VP, Post MW, Van Heugten CM. Proactive coping poststroke: psychometric properties of the Utrecht proactive coping competence scale. *Arch Phys Med Rehabil* 2014;95:670-5.
46. Tavakol M, Dennick R. Making sense of Cronbach's alpha. *Int J Med Educ* 2011;2:53-5.
47. Wijenberg MLM, Van Heugten CM, Van Mierlo ML, Visser-Meily JMA, Post MWM. Psychological factors after stroke: are they stable over time? *J Rehabil Med* 2019;51:18-25.
48. Post MWM, Nachtegaal J, Van Langeveld SA, et al. Progress of the Dutch spinal cord injury database: completeness of database and profile of patients admitted for inpatient rehabilitation in 2015. *Top Spinal Cord Inj Rehabil* 2018;24:141-50.
49. Ten Brink AF, Hajos TRS, Van Bennekom C, et al. Predictors of physical independence at discharge after stroke rehabilitation in a Dutch population. *Int J Rehabil Res* 2017;40:37-45.
50. Smith EM, Boucher E, Miller WC, SCIRE, Boucher N. Caregiving services in spinal cord injury: a systematic review of the literature. *Spinal Cord* 2016;54:562-569.
51. Van Exel NJA, Koopmanschap MA, Van Den Berg B, Brouwer WBF, Van Den Bos GAM. Burden of informal caregiving for stroke patients: identification of caregivers at risk of adverse health effects. *Cerebrovasc Dis* 2005;19:11-7.
52. Kokorelias KM, Gignac MAM, Naglie G, Cameron JI. Towards a universal model of family centered care: a scoping review. *BMC Health Serv Res* 2019;564.
53. Camak DJ. Addressing the burden of stroke caregivers: a literature review. *J Clin Nurs* 2015;24:2376-82.
54. Kneebone II. Stepped psychological care after stroke. *Disabil Rehabil* 2016;38:1836-43.
55. Panzeri A, Rossi Ferrario S, Vidotto G. Interventions for psychological health of stroke caregivers: a systematic review. *Front Psychol* 2019;10:2045.
56. Schofield D, Shrestha RN, Zeppel MJB, et al. Economic costs of informal care for people with chronic diseases in the community: lost income, extra welfare payments, and reduced taxes in Australia in 2015–2030. *Health Soc Care Community* 2019;27:493-501.
57. Dunbar SB, Khavjou OA, Bakas T, et al. Projected costs of informal caregiving for cardiovascular Disease: 2015 to 2035: a policy statement from the American Heart Association. *Circulation* 2018;137:558-77.
58. Joo H, Zhang P, Wang G. Cost of informal care for patients with cardiovascular disease or diabetes: current evidence and research challenges. *Qual Life Res* 2017;26:1379-86.