University of Groningen

Comparison of predicted and real propofol and remifentanyl concentrations in plasma and brain tissue during target-controlled infusion

Van Hese, L; Cuypers, E; Theys, T; Absalom, A; Rex, S

Published in:
Anaesthesia

DOI:
[10.1111/anae.15452](https://doi.org/10.1111/anae.15452)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Van Hese, L., Cuypers, E., Theys, T., Absalom, A., & Rex, S. (2021). Comparison of predicted and real propofol and remifentanyl concentrations in plasma and brain tissue during target-controlled infusion: a reply. *Anaesthesia*. <https://doi.org/10.1111/anae.15452>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Correspondence

Comparison of predicted and real propofol and remifentanyl concentrations in plasma and brain tissue during target-controlled infusion: a reply

We thank White et al. [1] for their interest in our study [2] and their thoughts on the implications it may have for the clinical use of total intravenous anaesthesia.

Online Supporting information Table S1 provides details of the exact type of surgery and medications of the patients in our study.

When discussing the development of pharmacokinetic models from data, it should be acknowledged that a balance must be sought between model complexity and parsimony. Some very simple models function very well clinically. An example is the Gepts model for sufentanil, which contains no covariates, meaning that for the same target concentration all patients receive the same dose. There is probably a very large number of factors that influence the pharmacokinetics of any one drug. The authors have suggested adding comorbidities and the effect of interacting drugs as covariates. While this may superficially seem simple, it is far from it, as the complexity can become almost infinite. If, for example, one considers adding interacting drugs as a covariate, there are several drugs that can cause PK and PD interactions, and for each there are additional variable factors (such as class of drug, exact drug used, dose of drug, duration of use, route of administration and duration since last dose). Similar complexity applies to comorbidities. Even when a drug has been used and studied in a very large population, and includes large numbers of patients in whom information about the covariables is available, there will always remain pitfalls associated with under-fitting (how would one then accurately model the influence of an interacting drug or comorbidity on PK parameters), but also with over-fitting (adding a covariate to a model structure, having wrongly attributed some of the residual variation to that covariate).

L. Van Hese

University Hospitals Leuven,
Belgium

E. Cuypers

Maastricht University,
Maastricht, The Netherlands
Email: e.cuypers@maastrichtuniversity.nl

T. Theys

University Hospitals Leuven,
Belgium

A. Absalom

University of Groningen,
The Netherlands

S. Rex

University Hospitals Leuven,
Belgium

No competing interests declared.

References

1. White S, Chirvasuta R, Eusuf D. Brain concentrations of anaesthetic agents: the implications of epilepsy surgery. *Anaesthesia* 2020. Epub 10 December. <https://doi.org/10.1111/anae.15344>.
2. Van Hese L, Theys T, Absalom A, Rex S, Cuypers E. Comparison of predicted and real propofol and remifentanyl concentrations in plasma and brain tissue during target-controlled infusion: a prospective observational study. *Anaesthesia* 2020; **75**: 1626–34.

doi:10.1111/anae.15452

Supporting Information

Additional supporting information may be found online via the journal website.

Table S1. Type of surgery and medications of the patients.