University of Groningen

Letter to the editor concerning Valstar et al., [Radiother Oncol 2020 Sep 23;S0167-8140(20)30809-4.doi

Bikker, Floris J; Vissink, Arjan

Published in:
Radiotherapy and Oncology

DOI:
[10.1016/j.radonc.2020.12.005](https://doi.org/10.1016/j.radonc.2020.12.005)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Bikker, F. J., & Vissink, A. (2021). Letter to the editor concerning Valstar et al., [Radiother Oncol 2020 Sep 23;S0167-8140(20)30809-4.doi: 10.1016/j.radonc.2020.09.034. *Radiotherapy and Oncology*, 154, 318-318. <https://doi.org/10.1016/j.radonc.2020.12.005>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Letter to the Editor

Letter to the editor concerning Valstar et al.
[Radiother Oncol 2020 Sep 23;S0167-8140(20)30809-4. doi: 10.1016/j.radonc.2020.09.034]

Dear editor,

We read the paper of Valstar et al. [1] on the discovery of the tubarial salivary glands with great interest. The authors discovered a pair of previously overlooked, clinically relevant macroscopic glands. We agree with the authors that these glands potentially may play an important role with regard to the efficient working and protection of the nasopharyngeal and oropharyngeal mucosa, comparable to the role of saliva for the oral cavity [2]. It is known that treatment of head and neck cancer with radiotherapy might result in swallowing problems [3]. This points towards damage of structures lubricating the nasopharyngeal and oropharyngeal mucosa. The so-called tubarial salivary glands may play an important in this lubricating process. This can be deducted from the observation that a lower cumulative radiation dose to these glands is accompanied by less swallowing problems [1].

We do not support, however, to yet call these tubarial glands, salivary glands. First, saliva is a slightly alkaline secretion of water, salts and proteins such as mucins, agglutinin, lactoferrin, lysozyme and amylase that is secreted into the mouth by salivary glands [4]. Dawes [5] showed that while saliva enters the mouth at several locations, the different secretions are hardly mixed. In this respect, tubarial glands should not be considered salivary glands as they do not contribute to the oral fluid, but at best ectopic salivary glands when they indeed secrete a fluid with a composition comparable to saliva [6]. Second, the histopathology points towards a conglomerate of minor glands rather than a major gland. Major glands have a main excretory duct and not multiple macroscopically visible draining duct openings, although the sublingual gland has some macroscopically visible draining ducts in addition. The duct openings macroscopically shown in their paper more resemble the duct endings of e.g., the labial glands. Third, salivary gland glands are known to secrete amylase. The discovered glands, however, were negative for amylase. Therefore, we would have expected that other salivary markers would have been tested in addition. As the histopathology of the discovered glands has a high resemblance with sublingual and minor salivary gland tissue, agglutinin, lactoferrin and lysozyme could have been used to better proof whether the glands indeed should be considered to be salivary glands [7–9]. Fourth, it is not yet known on what stimuli these glands react to start secretion. Are this, e.g., chewing, gustatory or olfactory stimuli? These are the common stimuli for salivary glands. Fifth, it would be important to collect secretions of the tubarial glands to unravel its composition to be able to gain a better insight in the potential functions of the secreted fluid. Collection of such secretion presumably is possible when

endoscopically inspecting the nasopharynx with methods used to study minor salivary gland secretions.

Resuming, the tubarial glands presumably exert an important function in the nasopharynx and oropharynx, but it is too early to name these glands, ectopic salivary glands.

References

- [1] Valstar MH, de Bakker BS, Steenbakkers RJHM, de Jong KH, Smit LA, Klein Nulst TJW, et al. The tubarial salivary glands: A potential new organ at risk for radiotherapy. *Radiother Oncol* 2020;S0167-8140(20)30809-4. <https://doi.org/10.1016/j.radonc.2020.09.034>. Online ahead of print.
- [2] Dawes C, Pedersen AM, Villa A, Ekström J, Proctor GB, Vissink A, et al. The functions of human saliva: A review sponsored by the World Workshop on Oral Medicine VI. *Arch Oral Biol* 2015;60:863–74. <https://doi.org/10.1016/j.archoralbio.2015.03.004>.
- [3] Christianen ME, van der Schaaf A, van der Laan HP, Verdonck-de Leeuw IM, Doornaert P, Chouvalova O, et al. Swallowing sparing intensity modulated radiotherapy (SW-IMRT) in head and neck cancer: Clinical validation according to the model-based approach. *Radiother Oncol* 2016;118:298–303. <https://doi.org/10.1016/j.radonc.2015.11.009>.
- [4] <https://www.merriam-webster.com/dictionary/saliva>.
- [5] Dawes C. Salivary flow patterns and the health of hard and soft oral tissues. *J Am Dent Assoc* 2008(139 Suppl):185–245. <https://doi.org/10.14219/jada.archive.2008.0351>.
- [6] Veerman ECI, van den Keybus PAM, Vissink A, Nieuw Amerongen AV. Human glandular saliva: their separate collection and analysis. *Eur J Oral Sci* 1996;104:346–52. <https://doi.org/10.1111/j.1600-0722.1996.tb00090.x>.
- [7] Moro I, Umemura S, Crago SS, Mestecky J. Immunohistochemical distribution of immunoglobulins, lactoferrin, and lysozyme in human minor salivary glands. *J Oral Pathol* 1984;13:97–104. <https://doi.org/10.1111/j.1600-0714.1984.tb01405.x>.
- [8] Chomette G, Aurioi M, Vaillant JM, Kasai T, Niwa M, Mori M. An immunohistochemical study of the distribution of lysozyme, lactoferrin, alpha 1-antitrypsin and alpha 1-antichymotrypsin in salivary adenoid cystic carcinoma. *Pathol Res Pract* 1991;187:1001–8. [https://doi.org/10.1016/s0344-0338\(11\)81072-1](https://doi.org/10.1016/s0344-0338(11)81072-1).
- [9] Bikker FJ, Ligtnerberg AJM, van der Wal JE, van den Keijbus PAM, Holmskov U, Veerman ECI, et al. Immunohistochemical detection of salivary agglutinin/gp-340 in human parotid, submandibular, and labial salivary glands. *J Dent Res* 2002;81:134–9.

Floris J. Bikker

Department of Oral Biochemistry, ACTA, Amsterdam, Free University
and University of Amsterdam, The Netherlands

Arjan Vissink

Department of Oral and Maxillofacial Surgery, University of Groningen
and University Medical Center Groningen, Groningen, The Netherlands

Received 24 October 2020

Accepted 2 December 2020

Available online 10 December 2020