University of Groningen

The predictive value of quantitative sensory testing

Petersen, Kristian; Vaegter, Henrik B; Stubhaug, A; Wolff, André; Scammell, Brigitte; Arendt-Nielsen, Lars; Larsen, Dennis B

Published in:
Pain

DOI:
[10.1097/j.pain.0000000000002019](https://doi.org/10.1097/j.pain.0000000000002019)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Petersen, K., Vaegter, H. B., Stubhaug, A., Wolff, A., Scammell, B., Arendt-Nielsen, L., & Larsen, D. B. (2021). The predictive value of quantitative sensory testing: a systematic review on chronic postoperative pain and the analgesic effect of pharmacological therapies in patients with chronic pain. *Pain*, 162(1), 31-44. <https://doi.org/10.1097/j.pain.0000000000002019>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

The predictive value of quantitative sensory testing: a systematic review on chronic postoperative pain and the analgesic effect of pharmacological therapies in patients with chronic pain

Kristian Kjær Petersen^{a,b,*}, Henrik B. Vaegter^{c,d}, Audun Stubhaug^{e,f}, André Wolff^g, Brigitte E. Scammell^{h,i}, Lars Arendt-Nielsen^{a,b}, Dennis B. Larsen^{a,b}

Abstract

Studies have suggested that quantitative sensory testing (QST) might hold a predictive value for the development of chronic postoperative pain and the response to pharmacological interventions. This review systematically summarizes the current evidence on the predictive value of QST for chronic postoperative pain and the effect of pharmacological interventions. The main outcome measures were posttreatment pain intensity, pain relief, presence of moderate-to-severe postoperative pain, responders of 30% and 50% pain relief, or validated questionnaires on pain and disability. A systematic search of MEDLINE and EMBASE yielded 25 studies on surgical interventions and 11 on pharmacological interventions. Seventeen surgical and 11 pharmacological studies reported an association between preoperative or pretreatment QST and chronic postoperative pain or analgesic effect. The most commonly assessed QST modalities were pressure stimuli (17 studies), temporal summation of pain (TSP, 14 studies), and conditioned pain modulation (CPM, 16 studies). Of those, the dynamic QST parameters TSP (50%) and CPM (44%) were most frequently associated with chronic postoperative pain and analgesic effects. A large heterogeneity in methods for assessing TSP ($n = 4$) and CPM ($n = 7$) was found. Overall, most studies demonstrated low-to-moderate levels of risk of bias in study design, attrition, prognostic factors, outcome, and statistical analyses. This systematic review demonstrates that TSP and CPM show the most consistent predictive values for chronic postoperative pain and analgesic effect, but the heterogeneous methodologies reduce the generalizability and hence call for methodological guidelines.

Keywords: Quantitative sensory testing, Chronic pain, Chronic postoperative pain, Analgesic effect

1. Introduction

Chronic pain is a major problem in the adult population,¹⁸ and treatment is difficult because of the limited amount of available efficient drugs and the undesired side effects. It is evident that chronic postoperative pain is present in 10% to 50% of patients

after different surgical treatments,^{49,87} and the effect of available pharmacological treatments remains low.^{29,86} Identifying patients responding to standard pain treatments is warranted as a start to implement individualized pain treatment.

Quantitative sensory testing (QST) can be combined using different sensory stimuli, and multiple QST protocols have been developed to probe the activity in specific nerve fibre populations and as proxies for spinal and pain modulatory functions.^{11,83} Most QST protocols allow the assessment of thermal, electrical, tactile, or pressure pain modalities.^{12,23,35,83,102} Reduced pain thresholds or increased sensory intensity ratings assessed at a local painful site mainly reflect modality-specific peripheral hyperalgesia, whereas assessments distant to a painful site may reflect widespread hyperalgesia as a surrogate measure of central hypersensitivity.^{10,34} Studies have found reduced pressure pain thresholds at distal sites to the knee in patients with knee osteoarthritis,^{8,37} increased sensitivity to thermal and mechanical stimuli in a subgroup of patients with neuropathies,^{82,99} and reduced electrical pain thresholds at the dorsal pancreatic referred dermatomes and at distant dermatomes in patients with chronic pancreatitis⁵⁵ compared with pain-free subjects. These studies indicate that patients with chronic pain of different aetiologies show generalised widespread hyperalgesia.⁷

Dynamic QST protocols have been developed to explore the central wind-up process using the proxy temporal summation of pain (TSP).³⁴ Facilitated TSP has been reported across multiple

Sponsorships or competing interests that may be relevant to content are disclosed at the end of this article.

^a SMI, Department of Health Science and Technology, Faculty of Medicine, Aalborg University, Aalborg, Denmark, ^b Department of Health Science and Technology, Center for Neuroplasticity and Pain, Faculty of Medicine, Aalborg University, Aalborg, Denmark, ^c Pain Research Group, Pain Center, Department of Anesthesiology and Intensive Care Medicine, University Hospital Odense, Odense, Denmark, ^d Department of Clinical Research, Faculty of Health Sciences, University of Southern Denmark, Denmark, ^e Department of Pain Management and Research, Oslo University Hospital, Oslo, Norway, ^f Institute of Clinical Medicine, Faculty of Medicine, University of Oslo, Oslo, Norway, ^g Department of Anaesthesiology, University of Groningen, University Medical Center Groningen, Pain Center, Groningen, The Netherlands, ^h Nottingham Elective Orthopaedic Service, Nottingham University Hospitals NHS Trust, Nottingham, United Kingdom, ⁱ Pain Centre Versus Arthritis, School of Medicine, The University of Nottingham, United Kingdom

*Corresponding author. Address: SMI, Department of Health Science and Technology, Faculty of Medicine, Aalborg University, Fredrik Bajers Vej 7 D3, DK-9220 Aalborg, Denmark. Tel.: +45 9940 7529; fax: +45 9815 4008. E-mail address: kkp@hst.aau.dk (K.K. Petersen).

PAIN 162 (2021) 31–44

© 2020 International Association for the Study of Pain
<http://dx.doi.org/10.1097/j.pain.0000000000002019>

chronic pain disorders such as osteoarthritis,¹⁰ fibromyalgia,³⁶ irritable bowel syndrome,⁷ and in subgroups of patients with neuropathic pain.⁶² Aspects of descending pain inhibitory control can most likely be mechanistically evaluated using the human proxies of conditioned pain modulation (CPM),¹⁰⁷ exercise-induced hypoalgesia,⁷⁰ or offset analgesia⁴⁴ protocols. Impaired CPM has been reported in several chronic painful conditions when compared with pain-free subjects.^{7,107} Exercise-induced hypoalgesia seems functional in asymptomatic subjects^{93–96} and impaired in different pain populations,^{28,57,67,92} although the current literature is inconclusive.^{19,28,92} Similarly, offset analgesia seems functional in asymptomatic subjects⁴⁴ and impaired in patients with chronic pain.⁹⁰

Studies have suggested a possible association between preoperative QST parameters and chronic postoperative pain^{47,56,74} and that pretreatment QST may predict the analgesic effect of pharmacological interventions.^{75,110} Previous systematic reviews^{2,38,46,84,101} have investigated the predictive value of QST on postoperative pain, the most recent in 2017,⁸⁴ and the predictive value of QST on the analgesic effect, the most recent in 2013.³⁸ The most recent review from 2017 on surgical studies⁸⁴ indicated that preoperative QST mainly predicts chronic postoperative pain but not acute postoperative pain. As multiple studies have been published since 2017, this review systematically summarizes the current literature on the possible predictive role of QST on (1) chronic postoperative pain and (2) the analgesic effect of pharmacological interventions in patients with chronic pain.

2. Methods

In accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement, this systematic review investigated the predictive role of QST on chronic postoperative pain and the analgesic effect of pharmacological interventions in patients with chronic pain. The systematic review has been registered on the Open Science Framework web site (OSF.IO, registration citation: Ref. 58, link to protocol: DOI: 10.17605/OSF.IO/HSVYK).

2.1. Outcomes

The primary outcomes for chronic postoperative pain were postoperative pain intensity, postoperative pain relief, presence of moderate-to-severe postoperative pain, or validated questionnaires on pain and disability. The primary outcomes of

pharmacological studies were pretreatment and posttreatment changes in pain scores, classification of responders to 30% or 50% pain relief, end of treatment pain intensity, or validated questionnaires on pain and disability.

2.2. Search strategy and selection of studies

A literature search was conducted in April 2020 in the databases PubMed and EMBASE. The search was limited to the literature published in the past 20 years (April 2000–April 2020). Only peer-reviewed studies published in English and with available full-text articles were considered eligible for the systematic review. Two searches were conducted to identify (1) the predictive value of QST for chronic postoperative pain outcomes and (2) the predictive value of QST on analgesic effect outcomes of pharmacological interventions. The MeSH terms and/or text word combinations are shown in **Table 1**.

All citations were exported to EndNote X4 (Thomson Reuters, Philadelphia, PA), and duplicates were removed. Owing to the large number of potential studies, the initial screening was conducted on title and abstract to remove citations that did not meet the scope of the systematic review. The screening process was independently performed by 2 reviewers (K.K.P. and D.B.L.) after the initial systematic database search. Disagreements in relevancy were solved by consensus. In case consensus was not reached, a third reviewer (H.B.V.) was consulted who made the final decision. After the screening, all full-text articles were obtained. Cross-referencing the included studies and the authors' own article collection was conducted for additional relevant literature. The inclusion and exclusion of the relevant literature are shown in **Figure 1** (PRISMA flowchart).

2.3. Eligibility criteria

The studies included had to report the predictive model of a postoperative or postpharmacological outcome of at least 1 QST modality, including electrical, thermal, mechanical or pressure, pain, tolerance and suprathreshold stimuli, TSP, allodynia, exercise-induced hypoalgesia, offset analgesia, CPM, or the full German Research Network on Neuropathic Pain (DFNS) protocol. Studies were included if they reported associations between QST and pain-related outcome after surgery or pharmacological intervention using correlations (Spearman or Pearson correlations), regression models, or other predictive models such as support vector machine or reduction in numbers needed to treat. Studies on animals and healthy subjects were

Table 1
Search strategy.

	Focus	Example key words (PubMed search)
1	Analgesic effect	("Analgesia"[tw] OR "Drugs"[tw] OR "Drug therapy"[tw])
2	Postoperative pain	("Postoperative pain"[MeSH] OR "Postsurgical pain"[tw])
3	Quantitative sensory testing	{{("Quantitative sensory testing"[tw] OR "QST"[tw]) OR ("Conditioned pain modulation"[tw] OR "CPM"[tw]) OR ("Temporal summation"[tw] OR "TSP"[tw])}}
4	Surgery	("Surgery"[tw])
5	Limits	"2000/01/01"[PDat], "English"[lang]

The MeSH and text word (tw) strings were permuted dependent on database.

Figure 1. Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram. QST, quantitative sensory testing.

excluded. Surgical studies were included if they assessed follow-up pain at least 3 months after surgery. Acute postoperative pain and studies in the subacute phase were not considered. Pharmacological studies were included if they studied the long-term effect (weeks of treatment) of pharmacological interventions. Therefore, studies investigating the acute effect (minutes or hours after administration) were excluded. All studies were required to have a detailed description of the used QST paradigm.

2.4. Data extraction, synthesis, and assessment

The data extraction was conducted by K.K. Peterson. For studies on chronic postoperative pain, the data included type of surgery, patient cohort, QST parameters tested, follow-up time after surgery, the dependent outcomes of the prediction model, and preoperative predictors and their associated predictive value for the dependent variable. For the pharmacological studies, the

data involved type of pharmacological intervention, patient cohort, QST parameters tested, treatment period, dependent variable in the prediction model, and predictive value of the QST parameters for the dependent variable. All outcomes were narratively synthesized to provide an overview of each QST paradigm assessed and presented within each domain of QST in its prediction of chronic postoperative pain or analgesic effect of pharmacological interventions. This was conducted because of the large heterogeneity in both the QST paradigms used and the clinical cohorts included in the studies, which meant that a meta-analysis would not be appropriate. The methodological and the overall quality of the included studies were assessed by 2 reviewers (K.K.P. and D.B.L.) using the Quality In Prognostic Studies (QUIPS) tool.⁴³ Following the guidelines, each study was assessed on methodological quality in the major categories of "Study participation," "Study attrition," "Prognostic factor measurement," "Outcome measurement," "Study confounding," and

“Statistical analysis and reporting.” Disagreements in the risk of bias analysis were solved by consensus, and if no consensus was reached, a third reviewer (H.B.V.) was consulted.

3. Results

The literature searches identified 1811 surgical and 2689 pharmacological studies, of which 25 surgical and 11 pharmacological studies were included (**Fig. 1**). The outcome parameters of the surgical studies were postoperative pain intensity (12 studies^{1,4,39,48,56,61,66,74,76,98,103,109}), postoperative pain relief (3 studies^{9,78,92}), presence of moderate-to-severe postoperative pain (6 studies^{16,41,69,71,81,89}), or validated questionnaires on pain and disability (3 studies^{40,105,106}) (**Table 2**). The outcome parameters in the pharmacological studies were pretreatment and posttreatment changes in pain scores (4 studies^{25,75,77,110}), migraine scores (1 study⁵¹), and classification of responders to 30% (3 studies^{5,63,72}) or 50% (2 studies^{20,21}) pain relief, and end of treatment pain intensity²⁴ (**Table 3**).

Most studies (27 studies) reported using multivariate statistical models, and univariate analyses were reported in 9 studies.

3.1. Quality assessment

The quality assessment is presented in **Table 4** (surgical) and **Table 5** (pharmaceutical). The reviewers (K.K.P. and D.B.L.) initially agreed on 89% of the ratings. Consensus was reached on all ratings after discussion.

3.2. Quantitative sensory testing variables

3.2.1. Electrical stimuli

Electrical stimuli were reported as electrical detection threshold, electrical pain threshold, or electrical pain tolerance threshold. Electrical stimuli were reported in 4 studies and found predictive in 2 studies (2/4 = 50%).^{61,72}

3.2.1.1. Electrical detection threshold

Electrical detection threshold was reported in 1 surgical study and no pharmacological studies. Preoperative electrical pain threshold was not found significantly associated with the chronic postoperative pain intensity after total knee arthroplasty.⁶¹

3.2.1.2. Electrical pain threshold

Electrical pain threshold was reported in 2 surgical studies and 1 pharmacological study. Low preoperative electrical pain threshold and high pain at rest predicted the chronic postoperative pain intensity after total knee arthroplasty.⁶¹ Electrical pain threshold was not significantly associated with persistent chronic postoperative pain or disability after segment spinal surgery of patients with chronic low back pain.⁶⁹

In the pharmacological studies, using a support vector machine analysis, the pretreatment ratio between electrical pain threshold at a pancreatic referred dermatome vs a nonaffected dermatome predicted the response to pregabalin (reduction in the clinical pain score of 30% or more after 3 weeks of treatment compared with placebo) with a sensitivity of 87.5% and a specificity of 80.0%.⁷²

3.2.1.3. Electrical pain tolerance threshold

Electrical pain tolerance threshold was reported in 1 surgical study in which the preoperative electrical pain tolerance threshold was not associated with the chronic postoperative pain intensity

after major abdominal surgery.¹⁰³ No pharmacological studies reported electrical pain tolerance threshold.

3.2.2. Thermal stimuli

Thermal stimuli were reported as cold and warm detection threshold, cold and heat pain threshold, and suprathreshold heat and cold stimuli. Thermal stimuli were reported in 11 studies and found predictive in 5 studies (5/11 ≈ 45%).^{1,25,51,63,76}

3.2.2.1. Warm detection threshold

Warm detection thresholds were reported in 3 surgical studies and 1 pharmacological study. One surgical study reported a linear regression model demonstrating that low preoperative warm detection threshold, low heat pain threshold, low degree of radiologically assessed osteoarthritis, and high TSP predicted a high chronic postoperative pain intensity after total knee arthroplasty in patients with knee osteoarthritis.⁷⁶ Two surgical studies did not find an association between preoperative warm detection thresholds and chronic postoperative pain intensity (groin hernia repair surgery¹ and breast cancer surgery⁴). One pharmacological (duloxetine for the treatment of diabetic peripheral neuropathy¹¹⁰) study did not find associations between the preoperative warm detection threshold and analgesic effect.

3.2.2.2. Heat pain threshold

Heat pain thresholds were reported in 9 surgical and 3 pharmacological studies. One surgical study reported a linear regression model including low preoperative heat pain threshold, low warm detection threshold, low degree of radiologically assessed osteoarthritis, and high TSP, which predicted a high chronic postoperative pain intensity after total knee arthroplasty in patients with knee osteoarthritis.⁷⁶ Eight surgical studies did not find associations between preoperative heat pain thresholds and chronic postoperative pain intensity (thoracic surgery,¹⁰⁹ groin hernia repair surgery,¹ arthroscopic surgery of the shoulder,⁹⁸ and breast cancer surgery⁴), postoperative Western Ontario and McMaster Universities Osteoarthritis Index scores (total knee arthroplasty¹⁰⁵), and the presence of moderate-to-severe chronic postoperative pain (total knee arthroplasty⁷¹ and segmental spinal surgery⁶⁹).

In the pharmacological studies, a hierarchical regression model demonstrated that low heat pain threshold was associated with a small analgesic effect of opioids in patients with postherpetic neuralgia.²⁵ Furthermore, using multivariate regression models, a great analgesic response to duloxetine in patients with migraine⁵¹ and painful diabetic neuropathy¹¹⁰ was not significantly associated with pretreatment heat pain thresholds.

3.2.2.3. Cold detection threshold

Cold detection thresholds were reported in 2 surgical studies and 1 pharmacological study. Preoperative or pretreatment cold detection thresholds were not statistically significantly associated with the chronic postoperative pain intensity (total knee arthroplasty⁷⁶ and breast cancer surgery⁴) or the analgesic effect (duloxetine for diabetic peripheral neuropathy¹¹⁰), respectively.

3.2.2.4. Cold pain threshold

Cold pain thresholds were reported in 4 surgical and no pharmacological studies. Preoperative cold pain thresholds were not significantly associated with the chronic postoperative pain intensity (total knee arthroplasty^{66,76}) and the presence of moderate-to-severe chronic postoperative pain (thoracic surgery¹⁶ and segmental spinal surgery⁶⁹).

Table 2**Surgical studies assessing quantitative sensory testing (QST) as predictors of chronic postoperative pain.**

Reference	Year	Type of surgery	Patients	QST parameters	Follow-up time	Postoperative dependent parameter	Models and preoperative predictors
Joint-related surgeries							
Martinez et al. ⁶⁶	2007	Total knee arthroplasty	Knee osteoarthritis: N = 20	ALL, MPT, HPT, CPT	4 mo	Pain intensity	Correlations (U): No prediction
Lundblad et al. ⁶¹	2008	Total knee arthroplasty	Knee osteoarthritis: N = 69	EDT, EPT	18 mo	Pain intensity	Regression (M): Pre-op pain (OR: 6.48) and EPT (OR: 9.19)
Gwilym et al. ⁴⁰	2011	Arthroscopic subacromial decompression	Shoulder impingement syndrome: N = 17	MPT	3 mo	Oxford shoulder score	Correlation (U): No prediction
Wylde et al. ¹⁰⁵	2013	Total knee arthroplasty	Knee osteoarthritis: N = 51	PPTs, HPT	13 mo	WOMAC	Correlations (U): PPT: R = 0.37
Valencia et al. ⁹⁸	2014	Arthroscopic shoulder surgery	Rotator cuff tendinopathy: N = 73	HPT, hTSP, hCPM*	3 and 6 mo	Pain intensity	Regressions (M): No prediction
Noiseux et al. ⁷¹	2014	Total knee arthroplasty	Knee osteoarthritis: N = 193	MPT, HPT, PPT	6 mo	Presence of moderate-to-severe postoperative pain	Regression (M): No prediction
Petersen et al. ⁷⁴	2015	Total knee arthroplasty	Knee osteoarthritis: N = 78	PPTs, mTSP, pCPM†	12 mo	Pain intensity	Regression (M): mTSP and Pre-op VAS: R ² = 0.13
Wylde et al. ¹⁰⁶	2015	Total knee and hip arthroplasty	Knee osteoarthritis: N = 239, hip osteoarthritis: N = 254	PPT	12 mo	WOMAC	Regression (M): THA: PPT: β = 0.091 (WOMAC) – 0.114 (movement pain) TKA: No prediction
Petersen et al. ⁷⁸	2016	Total knee arthroplasty	Knee osteoarthritis: N = 103	cPPT, cPTT, cTSP, cCPM, PPTs	12 mo	Pain relief	Regression (M): R ² = 0.379, using the cPPT and VAS
Bossmann et al. ¹⁷	2017	Total knee arthroplasty	Knee osteoarthritis: N = 47	mTSP, pCPM†	6 mo	WOMAC pain subscale	Regression (M): pCPM, heart rate variability, and gender: R ² = 0.09
Vaegter et al. ⁹²	2017	Total knee arthroplasty	Knee osteoarthritis: N = 14	cPPT, cPTT, PPT, pCPM†, EIH	6 mo	Pain relief	Correlations (U): pCPM: R = 0.57 EIH: R = 0.53
Arendt-Nielsen et al. ⁹	2018	Total knee arthroplasty	Knee osteoarthritis: N = 70	PPT	12 mo	Pain relief	Regression (M): PPT: R ² = 0.09–0.110
Petersen et al. ⁷⁶	2018	Total knee arthroplasty	Knee osteoarthritis: N = 130	CDT, CPT, WDT, HPT, mTSP	12 mo	Pain intensity	Regression (M): Pre-op mTSP, WDT, HPT, and KL: R ² = 0.119
Rice et al. ⁸¹	2018	Total knee arthroplasty	Knee osteoarthritis: N = 288	PPT, mTSP, pCPM†	6 and 12 mo	WOMAC pain >30/100	Regression (M): WOMAC pain, mTSP, Trait anxiety, and expected pain: AUC: 0.70 Sensitivity: 0.72 Specificity: 0.64 Correctly classified: 65.67%
Kurien et al. ⁵⁶	2018	Total knee arthroplasty	Knee osteoarthritis: N = 50	PPTs, cPPT, cPTT, cTSP, cCPM, mTSP,		Pain intensity	Correlations (U): mTSP: R = 0.343 PPT: R = –0.262
Müller et al. ⁶⁹	2019	Segment spinal surgery	Chronic low back pain: N = 141	PPT, PTT, HPT, CPT, CPM, EPT	12 mo	Persistent pain or persistent disability	Regression (M): No prediction
Thoracic-related surgeries							
Yarnitsky et al. ¹⁰⁹	2008	Thoracic surgery	Posterolateral and muscle-sparing lateral thoracotomy: N = 62	hCPM*, HPT, STHS	29 wk	Pain intensity	Regression (M): hCPM* (OR: 0.55)
Grosen et al. ³⁹	2014	Surgical correction of funnel chest	Patients scheduled for surgical correction of funnel chest: N = 41	ALL, MDT, PPT, pCPM†	6 mo	Pain intensity	Regressions (M): No prediction
Bayman et al. ¹⁶	2017	Thoracic surgery	Patients scheduled for thoracic surgery: N = 99	CPT, STHS	6 mo	Presence of postoperative pain	Regression (M): No prediction

(continued on next page)

Table 2 (continued)

Reference	Year	Type of surgery	Patients	QST parameters	Follow-up time	Postoperative dependent parameter	Models and preoperative predictors
Abdominal and gynecology-related surgery							
Aasvang et al. ¹	2010	Groin hernia repair (open and laparoscopic)	Primary unilateral hernia: N = 442	WDT, HPT, STHS	6 mo	Pain intensity	Regression (M): Activity (OR: 1.16–7.37), STHS (OR: 1.34) Surgical type (OR: 0.45)
Wilder-Smith et al. ¹⁰³	2010	Major abdominal surgery	Lower and upper gastrointestinal or genitourinary tract issues: N = 20	EPTT, PTT, eCPM†, pCPM†	6 mo	Pain intensity	Regression (M): Pre-op eCPM: R ² = 0.46
Jarrell et al. ⁴⁸	2014	Gynecologic laparoscopy	Patients with gynecological pain: N = 77	MDT, ALL	6 mo	Pain intensity	Regressions (M): The presence of pre-op ALL (R ² = 0.590)
Sng et al. ⁸⁹	2018	Elective abdominal or laparoscopic hysterectomy for benign conditions	Patients with benign conditions scheduled for surgery: N = 159	mTSP	6 mo	VAS >3/10	Regression (M): mTSP: OR = 1.078 Pain during sexual intercourse: OR = 5.312 Morphine consumption (24–48 hours post-op): OR: 1.172
Breast cancer surgeries							
Andersen et al. ⁴	2017	Breast cancer surgery	Patients with breast cancer: N = 290	MDT, MPT, WDT, CDT, HPT, PPT, size of the hypoesthesia area using: ALL and cold (25°C) and warm (40°C) rolls.	12 mo	Pain intensity	Regression (M): Size of the hypoesthesia area: OR: 1.003–1.006
Habib et al. ⁴¹	2019	Breast cancer surgery	Patients scheduled for breast cancer surgery: N = 124	mTSP, hCPM*	6 and 12 mo	VAS >3/10 and impact of pain on daily living	Regression (M): No prediction

QST modalities: ALL, dynamic mechanical allodynia; ANOVA, analysis of variance; CDT, cold detection threshold; CPT, cold pain threshold; cPPT, cuff-induced pressure pain threshold; cPTT, cuff-induced pressure tolerance threshold; CPM, conditioning pain modulation (c, cuff pressure test and condition stimuli; e, electrical test stimulus; heat, heat test stimulus; p, pressure test stimulus; *, hot water condition stimulus; †, cold-pressor tests as condition stimulus; ‡, pinching conditioning stimulus); EDT, electrical detection threshold; EIH, exercise-induced hypoalgesia; EPT, electrical pain threshold; EPTT, electrical pain tolerance threshold; HPT, heat pain threshold; KOOS, Knee Injury and Osteoarthritis Outcome Score (assesses pain, stiffness and function, and daily living of patients with knee osteoarthritis); (M), indicates multivariate analysis; MDT, mechanical detection threshold; MPT, mechanical pain threshold; N, number of patients participating in the study; OR, odds ratio; PPT, pressure pain threshold; PTT, pressure tolerance threshold; R, coefficient of determination; STCS, suprathreshold cold stimulus; STHS, suprathreshold heat stimulus; TSP, temporal summation of pain (c, using cuff stimuli; e, using electrical stimuli; h, using heat stimuli; m, using mechanical pinprick stimuli); (U), indicates univariate analysis; VAS, visual analog scale; WDT, warm detection threshold; WOMAC, Western Ontario and McMaster Universities Osteoarthritis Index (assesses pain, as well as stiffness and function of the knee and hip).

§ Calculated for this review alone and not presented in the original article.

3.2.2.5. Suprathreshold heat and cold stimuli

Suprathreshold heat and cold stimuli were reported in 3 surgical and no pharmacological studies. In a logistic regression model, preoperative high pain intensities to suprathreshold heat stimuli along with lowered warm detection thresholds and pain-related impairment of activity were predictive of the presence of moderate-to-severe chronic postoperative pain after hernia repair.¹ Preoperative suprathreshold heat stimuli and suprathreshold cold stimuli were not significantly associated with postoperative pain intensity (thoracic surgery¹⁰⁹) and the presence of moderate-to-severe chronic postoperative pain (thoracic surgery¹⁶).

3.2.3. Cutaneous mechanical stimuli

Cutaneous mechanical stimuli were reported as mechanical detection and pain threshold. Cutaneous mechanical stimuli were reported in 7 and predictive in no studies (0/7 = 0%).

3.2.3.1. Mechanical detection threshold

Mechanical detection thresholds were reported in 2 surgical studies and 1 pharmacological study. Preoperative and

pretreatment mechanical detection thresholds were not statistically significantly associated with the chronic postoperative pain intensity (surgical correction of funnel chest³⁹ and breast cancer surgery⁴) or analgesic effect (duloxetine for the treatment of diabetic peripheral neuropathy¹¹⁰).

3.2.3.2. Mechanical pain threshold

Mechanical pain thresholds were reported in 4 surgical studies and 1 pharmacological study. Preoperative and pretreatment mechanical pain thresholds were not associated with the chronic postoperative pain intensity (total knee arthroplasty⁶⁶ and breast cancer surgery⁴), presence of moderate-to-severe postoperative pain (total knee arthroplasty⁷¹), postoperative Oxford shoulder score (arthroscopic subacromial decompression⁴⁰), or analgesic effect (duloxetine for the treatment of diabetic peripheral neuropathy¹¹⁰).

3.2.4. Pressure stimuli

Pressure stimuli were studied as pressure pain and tolerance threshold as well as cuff-induced pressure pain and tolerance thresholds. Deep pressure stimuli were reported in 17 studies and were predictive in 5 studies (5/17 ≈ 29%).^{9,56,78,105,106}

Table 3**Pharmacological studies using quantitative sensory testing (QST) to predict analgesic effects.**

Reference	Year	Type of intervention(s)	Patients	QST parameters	Treatment period	Dependent parameter	Model and pretreatment predictor
Edwards et al. ²⁵	2006	Opioids: N = 16 TCAs: N = 14 Placebo: N = 14	Postherpetic neuralgia	HPT	8 wk	Change in pain intensity	Regression (M): Opioids: $R^2 = 0.35$ using HPT TCA: no prediction Placebo: no prediction
Yarnitsky et al. ¹¹⁰	2012	Duloxetine: N = 30	Painful diabetic neuropathy	CDT, WDT, HPT, MDT, MPT, mTSP, hCPM*	5 wk	Change in pain intensity	Regression (M): CPM: $R^2 = 0.673$
Olesen et al. ⁷²	2013	Pregabalin: N = 31 Placebo: N = 29	Painful chronic pancreatitis	EPT, PPT, differences in EPT and PPT in the affected and the unaffected area (EPT/PPT ratio)	3 wk	Classify responders (>30% pain reduction) and nonresponders	Support vector machine (M): EPT ratio: Sensitivity: 87.5% Specificity: 80.0%
Demant et al. ²¹	2014	Oxcarbazepine: N = 48 Placebo: N = 35	Peripheral neuropathic pain	ALL, WDT, CDT, HPT, CPT, PHS MDT, MPT, MPS, mTSP, VDT, PPT (DFNS protocol)	6 wk	Patients classified into irritable nociceptor (IN) or nonirritable nociceptor (NIN). Numbers needed to treat (NNTs) to obtain 50% pain relief for the IN and NIN.	NNT for the IN: 3.9 NNT for the NIN: 13 (U)
Demant et al. ²⁰	2015	Lidocaine 5% patch: N = 40 Placebo: N = 40	Peripheral neuropathic pain	ALL, WDT, CDT, HPT, CPT, PHS MDT, MPT, MPS, mTSP, VDT, PPT (DFNS protocol)	4 wk	Patients classified into irritable nociceptor (IN) or nonirritable nociceptor (NIN). Numbers needed to treat (NNTs) to obtain 50% pain relief for the IN and NIN.	NNT for the IN: 7.5 NNT for the NIN: not possible to determine due to recruitment issues (U)
Arendt-Nielsen et al. ⁵	2016	COX-2 inhibitor: N = 37 Placebo: N = 37	Knee osteoarthritis	PPT, pTSP, pCPM	4 wk	Change in pain intensity for nonresponders (<30% and <50% reduction)	Correlation (U): pTSP predicting a nonresponse: $R = 0.421-0.639$
Edwards et al. ²⁴	2016	NSAID (topical gel): N = 35	Knee osteoarthritis	PPT, mTSP, pCPM†, pCPM	4 wk	Change in pain average daily pain intensity (ADP) and KOOS pain	Regressions (M): ADP: CPM: $R = -0.38$ CPM
Mainka et al. ⁶³	2016	Capsaicin (topical, 8%): N = 20	Peripheral neuropathic pain	ALL, WDT, CDT, HPT, CPT, PHS MDT, MPT, MPS, mTSP, VDT, PPT (DFNS protocol)	8 wk	Classifying responders or nonresponders; responders were defined as +30% reduction in pain or 2 points on a 0-10 NRS	Regression (M): Sensitivity: 70%, specificity: 100% for patients with CPT and MPT >0.8 compared with the z value from DFNS.
Petersen et al. ⁷⁵	2019	NSAID and paracetamol (oral): N = 132	Knee osteoarthritis	cPPT, cPTT, cTSP	3 wk	Change in pain intensity	Regression (M): $R^2 = 0.269$ using the VAS and cTSP
Petersen et al. ⁷⁷	2019	NSAID and paracetamol (oral): N = 42	Knee osteoarthritis	OA, cCPM	3 wk	Change in pain intensity	Correlation (U): $R^2 = 0.186$ using the cCPM and VAS
Kisler et al. ⁵¹	2019	Duloxetine: N = 27 Placebo: N = 28	Migraine	Tonic heat pain (47°C), mTSP, OA, TS_{alone} , $TS_{conditioned}$, hCPM†	8 wk	Migraine improvement (pain reduction)	Regression (M): TS_{alone} ($R = 0.47$), $TS_{conditioned}$ ($R = 0.49$)

QST modalities: ALL, dynamic mechanical allodynia; CDT, cold detection threshold; CPT, cold pain threshold; cPPT, cuff-induced pressure pain threshold; cPTT, cuff-induced pressure tolerance threshold; CPM, conditioning pain modulation (c, cuff pressure test and condition stimuli; e, electrical test stimulus; heat, heat test stimulus; p, pressure test stimulus; *, hot water condition stimulus; †, cold-pressor tests as condition stimulus); EDT, electrical detection threshold; EIH, exercise-induced hypoalgesia; EPT, electrical pain threshold; EPTT, electrical pain tolerance threshold; HPT, heat pain threshold; (M), indicates multivariate analysis; MDT, mechanical detection threshold; MPT, mechanical pain threshold; N, number of patients participating in the study; s, offset analgesia; OR, odds ratio; PPT, pressure pain threshold; PTT, pressure tolerance threshold; R, coefficient of determination; STCS, suprathreshold cold stimulus; STHS, suprathreshold heat stimulus; TSP, temporal summation of pain (c, using cuff stimuli; e, using electrical stimuli; h, using heat stimuli; m, using mechanical pinprick stimuli; p, using pressure stimuli); (U), indicates univariate analysis; VAS, visual analog scale; WDT, warm detection threshold.

3.2.4.1. Pressure pain threshold

Pressure pain thresholds were reported in 11 surgical and 3 pharmacological studies. Low pressure pain thresholds assessed at the osteoarthritic affected knee⁵⁶ were associated with postoperative pain, low pressure pain thresholds at the nonaffected knee⁹ were associated with postoperative pain relief, and low pressure pain thresholds at the forearm¹⁰⁵ were associated with postoperative high Western Ontario and

McMaster Universities Osteoarthritis Index scores after total knee arthroplasty. A study found that low pressure pain thresholds assessed at the forearm in patients with hip osteoarthritis were associated with high postoperative Western Ontario and McMaster Universities Osteoarthritis Index scores after total hip arthroplasty but not in patients with knee osteoarthritis after total knee arthroplasty.¹⁰⁶ Eight studies did not find a statistically significant association between

Table 4**Risk of bias (RoB) for studies investigating the prognostic value of QST parameters on chronic postoperative pain.**

	Study participation	Study attrition	Prognostic factor measurement	Outcome measurement	Study confounding	Statistical analysis and reporting
Martinez et al. ⁶⁶	M	L	M	L	M	L
Lundblad et al. ⁶¹	M	H	M	M	H	L
Yarnitsky et al. ¹⁰⁹	M	L	L	M	L	M
Aasvang et al. ¹	L	L	L	M	H	L
Wilder-Smith et al. ¹⁰³	L	L	L	M	M	M
Gwilym et al. ⁴⁰	M	L	L	L	H	M
Wylde et al. ¹⁰⁵	M	L	L	L	M	M
Grosen et al. ³⁹	L	M	L	L	L	L
Jarrell et al. ⁴⁸	L	H	M	M	H	L
Valencia et al. ⁹⁸	M	M	L	L	L	L
Noiseux et al. ⁷¹	M	L	L	L	M	L
Petersen et al. ⁷⁴	M	L	L	M	M	L
Wylde et al. ¹⁰⁶	L	M	L	M	L	L
Petersen et al. ⁷⁸	L	M	L	M	M	L
Bayman et al. ¹⁶	M	M	L	L	L	L
Bossmann et al. ¹⁷	M	M	L	L	L	L
Andersen et al. ⁴	L	L	M	M	L	L
Vaegter et al. ⁹²	M	L	L	L	M	L
Arendt-Nielsen et al. ⁹	L	M	L	M	L	M
Petersen et al. ⁷⁶	L	M	M	L	M	L
Rice et al. ⁸¹	L	L	L	L	M	L
Sng et al. ⁸⁹	L	M	L	M	M	L
Kurien et al. ⁵⁶	M	L	L	M	M	L
Habib et al. ⁴¹	L	L	L	L	M	H
Müller et al. ⁶⁹	L	M	L	L	L	L

Using the Quality in Prognostic Studies (QUIPS) tool, the RoB assessment was based on study participation, study attrition, prognostic factor measurement, outcome measurement, study confounding, and statistical analysis. In general, low-to-moderate risk of bias was observed in most studies distributed across all factors. Risk of bias was evaluated as low (L), medium (M) or high (H). QST, quantitative sensory testing.

preoperative pressure pain thresholds and chronic postoperative pain intensity (total knee arthroplasty,⁵⁶ surgical correction of funnel chest,³⁹ and breast cancer surgery⁴), postoperative pain relief (total knee arthroplasty^{78,92}), or the presence of moderate-to-severe postoperative pain (total knee arthroplasty^{71,81} and segmental spinal surgery⁶⁹).

For the pharmacological studies, 3 studies found no association between pretreatment pressure pain thresholds and analgesic effect (COX-2⁵ inhibitors and NSAID gels²⁴ for the treatment of painful knee osteoarthritis and pregabalin for the treatment of painful chronic pancreatitis⁷²).

3.2.4.2. Pressure tolerance threshold

Pressure tolerance thresholds were reported in 2 surgical and no pharmacological studies. Preoperative pressure tolerance threshold was not associated with the chronic postoperative pain intensity (major abdominal surgery¹⁰³) and presence of persistent pain and disability (segmental spinal surgery⁶⁹).

3.2.4.3. Cuff-induced pain detection threshold

Cuff-induced pressure pain thresholds were reported in 3 surgical studies and 1 pharmacological study. Low cuff-induced pressure pain threshold assessed at the lower leg in patients with knee osteoarthritis was associated with chronic postoperative pain relief after total knee arthroplasty.⁷⁸ Two studies did not find a significant association between preoperative cuff-induced pressure pain thresholds and the chronic postoperative pain intensity (total knee arthroplasty⁵⁶) or postoperative pain relief (total knee arthroplasty⁹²). One pharmacological study demonstrated no association between the pretreatment cuff-induced pressure pain threshold and analgesic effect (oral NSAIDs and paracetamol for the treatment of pain in knee osteoarthritis⁷⁵).

3.2.4.4. Cuff-induced pressure tolerance threshold

Cuff-induced pain tolerance thresholds were reported in 3 surgical studies and 1 pharmacological study. Three surgical studies did not find a significant association between the preoperative cuff-induced pain tolerance threshold and the

Table 5

Risk of bias (RoB) for studies investigating the prognostic value of QST parameters on the analgesic effect after nonsurgical treatments.

	Study participation	Study attrition	Prognostic factor measurement	Outcome measurement	Study confounding	Statistical analysis and reporting
Edwards et al. ²⁵	M	M	M	L	M	L
Yarnitsky et al. ¹¹⁰	M	M	L	L	M	L
Olesen et al. ⁷²	L	L	L	L	M	M
Demant et al. ²¹	L	L	L	M	M	L
Demant et al. ²⁰	L	L	M	L	M	L
Arendt-Nielsen et al. ⁵	L	M	L	L	M	L
Edwards et al. ²⁴	M	M	L	L	M	L
Mainka et al. ⁶³	L	L	L	L	M	M
Petersen et al. ⁷⁵	L	M	L	L	M	L
Petersen et al. ⁷⁷	L	L	L	L	M	L
Kisler et al. ⁵¹	M	M	L	M	M	L

Using the Quality in Prognostic Studies (QUIPS) tool, the RoB assessment was based on study participation, study attrition, prognostic factor measurement, outcome measurement, study confounding, and statistical analysis. In general, low-to-moderate risk of bias was observed in most studies distributed across all factors.

QST, quantitative sensory testing.

chronic postoperative pain intensity (total knee arthroplasty⁵⁶) or postoperative pain relief (total knee arthroplasty^{78,92}). The pharmacological study did not find a significant association between the pretreatment cuff-induced pain tolerance threshold and analgesic effect (oral NSAIDs and paracetamol for the treatment of pain in knee osteoarthritis⁷⁵).

3.2.5. Dynamic mechanical allodynia

Dynamic mechanical allodynia was reported in 3 surgical and no pharmacological studies and was reported predictive in 1 study (1/3 $\approx$ 33%). The presence of preoperative dynamic mechanical allodynia (yes/no) was associated with the chronic postoperative pain intensity in females undergoing gynecologic laparoscopy.⁴⁸ Two studies did not find associations between preoperative dynamic mechanical allodynia and the postoperative pain intensity (surgical correction of funnel chest³⁹ and total knee arthroplasty⁶⁶).

3.2.6. Temporal summation of pain

Temporal summation of pain was reported in 14 studies (9 surgical and 5 pharmacological studies) and predictive in 7 studies (7/14 = 50%).^{5,56,74–76,81,89}

In the surgical studies, TSP was assessed using mechanical stimuli,^{56,74,76,81,89} heat stimuli,⁹⁸ and cuff stimuli.^{56,78} In 5 surgical studies, high preoperative TSP was associated with the chronic postoperative pain intensity after total knee arthroplasty^{56,74,76} and the presence of moderate-to-severe chronic postoperative pain (total knee arthroplasty⁸¹ and abdominal or laparoscopic hysterectomy⁸⁹). Four studies reported that preoperative TSP was not associated with the chronic postoperative pain intensity (total knee arthroplasty⁷⁸ and arthroscopic shoulder surgery⁹⁸), postoperative WOMAC (total knee arthroplasty¹⁷), and the presence of moderate-to-severe chronic postoperative pain (breast cancer surgery⁴¹).

In the pharmacological studies, TSP was assessed using mechanical stimuli,^{24,51,110} computer-controlled pressure stimuli,⁵ and manual cuff stimuli.⁷⁵ In 2 studies, high TSP was

reported to be associated with poor analgesic effect after 4 weeks of treatment with COX-2 inhibitors⁵ and 3 weeks of treatment with NSAIDs and paracetamol⁷⁵ in patients with knee osteoarthritis. Three studies did not find an association between pretreatment TSP and the analgesic effect of duloxetine for diabetic peripheral neuropathy,¹¹⁰ migraine,⁵¹ or topical NSAIDs for painful knee osteoarthritis.²⁴

3.2.7. Conditioned pain modulation

Conditioned pain modulation was reported in 17 studies (12 surgical and 5 pharmacological studies) and predictive in 7 studies (7/17 $\approx$ 41%).^{24,77,92,103,109,110}

In the surgical studies, the test stimulus was assessed using pressure,^{17,39,69,74,81,92,103} cuff,^{56,78} heat,^{41,69,98,109} and electrical^{69,103} stimuli, with the conditioning stimuli being hot water,^{41,98,109} cold water,^{39,69,74,81,92,103} tonic cuff pressure,^{56,78} or pinching.¹⁷ In 4 studies, preoperative impaired CPM was associated with chronic postoperative pain intensity (thoracic surgery¹⁰⁹ and major abdominal surgery¹⁰³), high postoperative WOMAC scores (total knee arthroplasty¹⁷), and a reduction in postoperative pain relief (total knee arthroplasty⁹²). Eight studies did not find an association between preoperative CPM and the chronic postoperative pain intensity (total knee arthroplasty,^{56,74,78,81} surgical correction of funnel chest,³⁹ and arthroscopic shoulder surgery⁹⁸), postoperative pain relief (total knee arthroplasty⁷⁸), and the presence of moderate-to-severe chronic postoperative pain (total knee arthroplasty,⁸¹ breast cancer surgery,⁴¹ and segmental spinal surgery⁶⁹).

In the pharmacological studies, the test stimulus was assessed using pressure,^{5,24} cuff,⁷⁷ and heat^{51,110} stimuli, with the conditioning stimuli being hot water,¹¹⁰ cold water,⁵¹ or tonic cuff pressure.^{5,24,77} Impaired CPM was associated with a great analgesic effect of duloxetine in patients with diabetic neuropathy.¹¹⁰ This was also the case with topical NSAIDs²⁴ or NSAIDs and paracetamol⁷⁷ in patients with knee osteoarthritis. Pretreatment CPM was not associated with the analgesic effect of COX-2 inhibitors in patients with knee osteoarthritis⁵ or duloxetine in patients with migraine.⁵¹ Of note, Kisler et al.⁵¹ found that the

pretreatment test stimulus and the conditioned test stimulus in the CPM paradigm predicted the analgesic effect of duloxetine in patients with migraine but not the calculated CPM effect itself.

3.2.7.1. Offset analgesia

Offset analgesia was reported in 1 pharmacological study in which the pretreatment offset analgesia showed no association with the analgesic effect of NSAID and paracetamol for patients with knee osteoarthritis.⁷⁷

3.2.7.2. Exercise-induced hypoalgesia

Exercise-induced hypoalgesia was reported in 1 surgical study in which low preoperative exercise-induced hypoalgesia was associated with low postoperative pain relief after total knee arthroplasty.⁹²

3.2.8. Hypoesthesia area

One study assessed the preoperative size of the hypoesthesia area using warm (40°C) and cold (25°C) rolls and found an association between the size of the hypoesthesia and the chronic postoperative pain intensity after breast cancer surgery.⁴

3.2.9. The German Research Network on Neuropathic Pain (DFNS) protocol

The German Research Network on Neuropathic Pain (DFNS) protocol consists of a wide range of QST modalities including allodynia, thermal detection and pain thresholds, paradoxical heat sensations, mechanical detection, pain thresholds, mechanical suprathreshold, TSP (wind-up), vibration detection threshold, and pressure pain threshold. The DFNS protocol was assessed in 3 pharmacological studies. In 2 pharmacological studies, the DFNS protocol was used to define the irritable nociceptor (IN) or non-IN (NIN) in patients with peripheral neuropathic pain. One study found that the number needed to treat for 50% pain relief of oxcarbazepine was 3.9 for the IN and 13 for the NIN.²¹ A study found that the number needed to treat for 50% pain relief of lidocaine 5% patch was 7.5 for the IN and not definable for the NIN due to recruitment issues. Finally, the DFNS protocol was used to predict the responders (+30% pain alleviation) and nonresponders to capsaicin patch treatment in patients with peripheral neuropathy and found a sensitivity of 70% and a specificity of 100% for patients with cold pain thresholds and mechanical pain thresholds >0.8 compared with z values from DFNS.⁶³

3.2.10. Prediction of specific surgical procedures

In this review, 16 studies addressed joint-related surgeries, 3 studies addressed thoracic-related surgeries, 4 studies addressed abdominal and gynecology-related surgeries, and 2 studies addressed breast cancer surgeries. A significant preoperative prediction was demonstrated for 11 studies (69%) in the joint-related surgeries, 1 study (33%) in the thoracic-related surgeries, 4 studies (100%) in the abdominal and gynecology-related surgeries, and 1 study (50%) related to breast cancer surgery.

4. Discussion

The current systematic review describes the predictive role of QST on pain after surgical and pharmacological interventions. Twenty-five surgical (10 new studies since the latest review⁸⁴) and

11 pharmacological (8 studies since the latest review³⁸) studies published since 2000 were identified. Seventeen studies demonstrated an association between preoperative QST and chronic postoperative pain, and 11 studies demonstrated an association between pretreatment QST and the analgesic effect of pharmacological interventions but with a large heterogeneity in the QST paradigms used. Significant preoperative predictions were most often presented for joint-related surgeries and abdominal and gynecology-related surgeries. Temporal summation of pain, CPM, and different variations of pressure thresholds were the most frequently reported methods, and TSP and CPM were most frequently found as predictors of the chronic postoperative pain intensity, the presence of moderate-to-severe chronic postoperative pain, postoperative pain relief, and the analgesic effect to pharmacological interventions.

4.1. The predictive value of quantitative sensory testing

This review suggests a possible association between the selected QST parameters and chronic postoperative pain and the analgesic response to pharmacological interventions, but the results are not consistent. Overall, the most used QST paradigms were mechanical and pressure stimuli, TSP, and CPM, which were also most frequently associated with chronic postoperative pain or analgesic effect. In addition, the strength of the predictive value of QST varied with R^2 values ranging from 0.13 to 0.673, which further underlines that variance explanation remains suboptimal at best. Finally, it is important to acknowledge that the literature on the predictive role of QST for chronic postoperative pain and analgesic response to pharmacological interventions is conflicting, and therefore, QST might not be appropriate as clinical guiding tool yet.

Decades of research has focused on the difference in QST parameters comparing pain-free subjects and patients with chronic pain; yet, the differences have not been established to be specific for the pain diagnosis.^{7,34,82,99,104} It is evident that some patients with chronic pain are generally more pain sensitive than others,^{6,91,99} but the underlying factors driving the increased pain sensitivity are still largely unclear. Studies have suggested that the pain sensitivity can be increased in patients with chronic pain because of, for example, sleep impairment,⁸⁸ increased pain catastrophizing,⁶⁸ or comorbidities such as diabetes.²⁷ These factors are often observed in patients in chronic pain populations and warrant consideration when addressing the predictive value of QST in future studies.

A previous review⁸⁴ suggested a link between different QST modalities and certain pain disorders. To exemplify this, cutaneous stimuli have been found to activate cutaneous fibres,³ and pressure stimuli using algometers have been found to target muscles³⁰ or fascia.⁶⁴ This could indicate that, for example, cutaneous activation would be suited for dermatological disorders, and pressure stimuli would be suited for patients with muscle or joint pain. The most studied population of the current review was patients with osteoarthritis, and the hypothesis that pressure stimuli should adhere better to these patients is not supported by the current review because pressure stimuli were rarely (approx. 29%) predictive of chronic postoperative pain or analgesic effect. In addition, joint-related surgeries and abdominal and gynaecology-related surgeries were most often studied, and preoperative QSTs were most often associated with chronic postoperative pain, which could indicate that preoperative QSTs are more frequently predictive of certain surgical procedures. Future studies should pursue this hypothesis to clarify whether there is an interaction between certain QST modalities and certain pain disorders.

This review investigated the predictive role of QST for chronic postoperative pain and analgesic response to pharmacological interventions, although these 2 outcomes are different. In a pain mechanistic context, surgical procedures aim to remove the peripheral pain driver, and it has been demonstrated that a pain-free recovery after removal of such peripheral drives by, for example, total knee³⁷ or hip arthroplasty⁵⁴ does normalize the responses to CPM and TSP when comparing preoperative and 6-month postoperative assessments. The advantage of certain pharmacological interventions is the possibility to target central pain mechanisms. As an example, preclinical trials have established that serotonin and noradrenalin are important for descending pain inhibition,^{13,14,60} and human administration of duloxetine (a serotonin and noradrenaline reuptake inhibitor) does improve CPM in patients with painful diabetic neuropathy.¹¹⁰ Likewise, the *N*-methyl-D-aspartate receptors are important for dorsal horn neuron excitability, and human administration of ketamine (an *N*-methyl-D-aspartate antagonist) reduces TSP in patients with fibromyalgia.³⁶ Therefore, specific QST modalities might be better predictors for a certain treatment if the pain mechanistic profile is matched with the intervention.

The most consistent predictive QST paradigms were TSP and CPM, but these studies used multiple different protocols (assessment parameters and modalities). The reliability of CPM has been questioned,⁵⁰ and studies comparing CPM protocols have highlighted that different test and conditioning stimuli combinations will yield different reliability results.^{45,97} Some CPM protocols operate under the premise that the conditioning stimulus should be painful (visual analog scale > 3) and that the CPM effect increases with increased conditioning stimulus intensity.³⁵ Conversely, other protocols are based on the notion that the intensity of the conditioning stimulus is independent of the CPM effect.³³ The 2015 recommendation for CPM¹⁰⁸ called for standardization of CPM testing across laboratories, which would greatly increase the generalizability of CPM in future studies. However, the current review highlights that the variability in different QST modalities has the potential to be used as a predictor of pain after surgery and pharmacological interventions. It is well known that dynamic QST modalities such as TSP and CPM are influenced by multiple factors such as sleep deprivation,^{26,59} use of opioids,⁶⁵ or high clinical pain intensities.¹⁰ Therefore, standardized protocols based on pain-free subjects might not represent the clinical use of QST.

4.2. Quality assessment

In this systematic review, the included studies overall demonstrated a low-to-moderate bias distributed among the 4 categories for bias assessment. For study participation, missing information on sampling frame and recruitment and place for assessment were the main causes for bias. For study attrition, lack of description of loss to follow-up and differences in participants completing and not completing the study were the main reasons for bias. In the category of prognostic factor measurement, validity and reliability of the used methods together with the uncertainty with respect to the cutoffs chosen were the main reasons for bias. Confounding was another category that revealed substantial bias, in that confounders in either study design or analysis were rarely accounted for and that multivariate analyses were not performed consistently.

4.3. Future directions

The current review highlights inconsistencies in QST as predictors of outcomes after surgical treatments and pharmacological

interventions. The most consistent findings include studies assessing central pain mechanisms, and therefore, future studies are encouraged to include parameters such as TSP or CPM and to understand the variability in the assessment of TSP and CPM. Fjeld et al., 2019,³¹ studied patients with acute admission due to sciatica and found that the assessment of CPM 6 weeks after discharge did not predict pain at 12 months of follow-up, which does highlight that the timing of assessment might be crucial. In addition, other non-QST parameters such as pain catastrophizing or epigenetic markers have been associated with chronic postoperative pain^{15,22,32,79,80} and analgesic effects.³⁹ Including these parameters will likely increase the strength of the predictive models in future studies. Finally, because of the large heterogeneity in choice of QST parameters, standardization is needed and assessment of both painful and nonpainful anatomic regions should be considered because this could influence the predictive value of QST.^{105,106}

In 2016, a task force appointed by the International Association for the Study of Pain⁵³ suggested “nociplastic” as a new pain descriptor to describe a state in which the nociceptive functions are changed. Several studies have now identified that specific subgroups of patients with chronic pain exist^{6,91} and that these patients respond poorly to standard care treatments.^{42,73–75} Understanding whether the modulation of specific central pain pathways before standard care treatments does increase the likelihood of positive analgesic responses would move this field forward. In this regard, acknowledging that a large variety of sensitization manifestations may represent a large variety of natural or acquired sensorimotor function modalities might be important.

Studies have found that, for example, duloxetine can improve CPM¹¹⁰ and ketamine can reduce TSP.³⁶ A recent study⁵² on patients scheduled for total knee arthroplasty recruited “pain sensitive” patients using the Central Sensitization Inventory Questionnaire and found that preoperative and 6-week postoperative administration of duloxetine reduced the postoperative pain intensity at 2 to 12 weeks of follow-up compared with placebo. Studies combining preoperative predictors with pharmacological interventions targeting these predictions might pave the way for the future of personalized pain medicine.

4.4. Limitations

The current review is limited to studies assessing chronic postoperative pain defined as pain reported at least 3 months after surgery. Several studies on the predictive role of QST for acute postoperative pain have been conducted (see review from 2017⁸⁴). Of note, studies such as Lunn et al., 2013, and Izumi et al., 2017, studied the postoperative period from 30 days to 6 weeks after surgery, and these studies are not included in the current review.

Furthermore, the current review is limited by not addressing studies using QST to predict the acute effect of pharmacological interventions such as Wasner et al.¹⁰⁰ and Schliessbach et al.⁸⁵

5. Conclusion

This systematic review identified 17 surgical and 11 pharmacological studies reporting a predictive role of QST modalities for chronic postoperative pain or the analgesic effect of pharmacological interventions. The strengths of the predictive models vary, and no consistency was found for a single QST parameter. Pressure stimuli and dynamic QST parameters such as TSP and CPM were the most frequently assessed, and thermal stimuli,

TSP, and CPM were most frequently associated with chronic postoperative pain or analgesic effects.

Conflict of interest statement

The authors have no conflicts of interest to declare.

Acknowledgements

The authors thank the Aalborg University Talent Management Programme (j.no. 771126) for providing the funding to initiate this work. The Center for Neuroplasticity and Pain (CNAP) is supported by the Danish National Research Foundation (DNRF121).

Article history:

Received 4 June 2020

Received in revised form 5 July 2020

Accepted 14 July 2020

Available online 21 July 2020

References

- [1] Aasvang EK, Gmaehle E, Hansen JB, Gmaehle B, Forman JL, Schwarz J, Bittner R, Kehlet H. Predictive risk factors for persistent postherniotomy pain. *Anesthesiology* 2010;112:957–69.
- [2] Abrishami A, Chan J, Chung F, Wong J. Preoperative pain sensitivity and its correlation with postoperative pain and analgesic consumption. *Anesthesiology* 2011;114:445–57.
- [3] Andersen HH, Elberling J, Sølvsten H, Yosipovitch G, Arendt-Nielsen L. Nonhistaminergic and mechanical itch sensitization in atopic dermatitis. *PAIN* 2017;158:1780–91.
- [4] Andersen KG, Duriaud HM, Kehlet H, Aasvang EK. The relationship between sensory loss and persistent pain 1 year after breast cancer surgery. *J Pain* 2017;18:1129–38.
- [5] Arendt-Nielsen L, Egsgaard LL, Petersen KK. Evidence for a central mode of action for etoricoxib (COX-2 inhibitor) in patients with painful knee osteoarthritis. *PAIN* 2016;157:1634–44.
- [6] Arendt-Nielsen L, Egsgaard LL, Petersen KK, Eskehave TN, Graven-Nielsen T, Hoeck HC, Simonsen O. A mechanism-based pain sensitivity index to characterize knee osteoarthritis patients with different disease stages and pain levels. *Eur J Pain* 2015;19:1406–17.
- [7] Arendt-Nielsen L, Morlion B, Perrot S, Dahan A, Dickenson A, Kress HG, Wells C, Bouhassira D, Mohr Drewes A. Assessment and manifestation of central sensitisation across different chronic pain conditions. *Eur J Pain* 2018;22:216–41.
- [8] Arendt-Nielsen L, Nie H, Laursen MB, Laursen BS, Madeleine P, Simonsen OH, Graven-Nielsen T. Sensitization in patients with painful knee osteoarthritis. *PAIN* 2010;149:573–81.
- [9] Arendt-Nielsen L, Simonsen O, Laursen MB, Roos EM, Rathleff MS, Rasmussen S, Skou ST. Pain and sensitization after total knee replacement or non-surgical treatment in patients with knee osteoarthritis: identifying potential predictors of outcome at 12 months. *Eur J Pain* 2018;22:1088–102.
- [10] Arendt-Nielsen L, Skou STST, Nielsen TA, Petersen KK. Altered central sensitization and pain modulation in the CNS in chronic joint pain. *Curr Osteoporos Rep* 2015;13:225–34.
- [11] Arendt-Nielsen L, Yarnitsky D. Experimental and clinical applications of quantitative sensory testing applied to skin, muscles and viscera. *J Pain* 2009;10:556–72.
- [12] Arezzo JC, Bolton CF, Boulton A, Zimmerman I, Casey KL, Dyck PJ, Comblath D, Field SH, Hanson N, Greenspan J. Quantitative sensory testing—a consensus report from the peripheral neuropathy association. *Neurology* 1993;43:1050–2.
- [13] Bannister K, Lockwood S, Goncalves L, Patel R, Dickenson AH. An investigation into the inhibitory function of serotonin in diffuse noxious inhibitory controls in the neuropathic rat. *Eur J Pain* 2017;21:750–60.
- [14] Bannister K, Patel R, Goncalves L, Townson L, Dickenson AH. Diffuse noxious inhibitory controls and nerve injury: restoring an imbalance between descending monoamine inhibitions and facilitations. *PAIN* 2015;156:1803–11.
- [15] Baranoff J, Hanrahan SJ, Connor JP. The roles of acceptance and catastrophizing in rehabilitation following anterior cruciate ligament reconstruction. *J Sci Med Sport* 2015;18:250–4.
- [16] Bayman EO, Parekh KR, Keech J, Selte A, Brennan TJ. A prospective study of chronic pain after thoracic surgery. *Anesthesiology* 2017;126:938–51.
- [17] Bossmann T, Brauner T, Wearing S, Horstmann T. Predictors of chronic pain following total knee replacement in females and males: an exploratory study. *Pain Manag* 2017;7:391–403.
- [18] Breivik H, Collett B, Ventafridda V, Cohen R, Gallacher D. Survey of chronic pain in Europe: prevalence, impact on daily life, and treatment. *Eur J Pain* 2006;10:287–333.
- [19] Burrows NJ, Booth J, Sturmeiks DL, Barry BK. Acute resistance exercise and pressure pain sensitivity in knee osteoarthritis: a randomised crossover trial. *Osteoarthritis Cartil* 2014;22:407–14.
- [20] Demant DT, Lund K, Finnerup NB, Vollert J, Maier C, Segerdahl MS, Jensen TS, Sindrup SH. Pain relief with lidocaine 5% patch in localized peripheral neuropathic pain in relation to pain phenotype: a randomised, double-blind, and placebo-controlled, phenotype panel study. *PAIN* 2015;156:2234–44.
- [21] Demant DT, Lund K, Vollert J, Maier C, Segerdahl M, Finnerup NB, Jensen TS, Sindrup SH. The effect of oxcarbazepine in peripheral neuropathic pain depends on pain phenotype: a randomised, double-blind, placebo-controlled phenotype-stratified study. *PAIN* 2014;155:2263–73.
- [22] Dworkin RH, Peirce-Sandner S, Turk DC, McDermott MP, Gibofsky A, Simon LS, Farrar JT, Katz NP. Outcome measures in placebo-controlled trials of osteoarthritis: responsiveness to treatment effects in the REPORT database. *Osteoarthritis Cartil* 2011;19:483–92.
- [23] Dyck PJ, Zimmerman I, Gillen DA, Johnson D, Karnes JL, O'Brien PC. Cool, warm, and heat-pain detection thresholds: testing methods and inferences about anatomic distribution of receptors. *Neurology* 1993;43:1500–8.
- [24] Edwards RR, Dolman AJ, Martel MO, Finan PH, Lazaridou A, Cornelius M, Wasan AD. Variability in conditioned pain modulation predicts response to NSAID treatment in patients with knee osteoarthritis. *BMC Musculoskelet Disord* 2016;17:284.
- [25] Edwards RR, Haythornthwaite JA, Tella P, Max MB, Raja S. Basal heat pain thresholds predict opioid analgesia in patients with postherpetic neuralgia. *Anesthesiology* 2006;104:1243–8.
- [26] Eichhorn N, Treede R-D, Schuh-Hofer S. The role of sex in sleep deprivation related changes of nociception and conditioned pain modulation. *Neuroscience* 2018;1:191–200.
- [27] Eitner A, Pester J, Vogel F, Marintschev I, Lehmann T, Hofmann GO, Schaible H-G. Pain sensation in human osteoarthritic knee joints is strongly enhanced by diabetes mellitus. *PAIN* 2017;158:1743–53.
- [28] Fingleton C, Smart KM, Doody CM. Exercise-induced hypoalgesia in people with knee osteoarthritis with normal and abnormal conditioned pain modulation. *Clin J Pain* 2017;33:395–404.
- [29] Finnerup NB, Attal N, Haroutounian S, McNicol E, Baron R, Dworkin RH, Gilron I, Haanpää M, Hansson P, Jensen TS, Kamerman PR, Lund K, Moore A, Raja SN, Rice ASC, Rowbotham M, Sena E, Siddall P, Smith BH, Wallace M. Pharmacotherapy for neuropathic pain in adults: a systematic review and meta-analysis. *Lancet Neurol* 2015;14:162–73.
- [30] Finocchietti S, Takahashi K, Okada K, Watanabe Y, Graven-Nielsen T, Mizumura K. Deformation and pressure propagation in deep tissue during mechanical painful pressure stimulation. *Med Biol Eng Comput* 2013;51:113–22.
- [31] Fjeld OR, Grotle M, Matre D, Pedersen LM, Lie MU, Småstuen MC, Storheim K, Heuch I, Stubhaug A, Zwart J-A, Nilsen KB. Predicting the outcome of persistent sciatica using conditioned pain modulation: 1-year results from a prospective cohort study. *Scand J Pain* 2019;20:69–75.
- [32] Giordano R, Petersen KK, Andersen HH, Lichota J, Simonsen OH, Arendt-Nielsen L. MicroRNAs expression as circulating genomic biomarkers in patients with chronic postoperative pain after total knee replacement. In: MicroRNAs expression as circulating genomic biomarkers in patients with chronic postoperative pain after total knee replacement. Boston: 17th World Congress on Pain, IASP 2018, 2018.
- [33] Granot M, Weissman-Fogel I, Crispel Y, Pud D, Granovsky Y, Sprecher E, Yarnitsky D. Determinants of endogenous analgesia magnitude in a diffuse noxious inhibitory control (DNIC) paradigm: do conditioning stimulus painfulness, gender and personality variables matter? *PAIN* 2008;136:142–9.
- [34] Graven-Nielsen T, Arendt-Nielsen L. Assessment of mechanisms in localized and widespread musculoskeletal pain. *Nat Rev Rheumatol* 2010;6:599–606.
- [35] Graven-Nielsen T, Izumi M, Petersen KK, Arendt-Nielsen L. User-independent assessment of conditioning pain modulation by cuff pressure algometry. *Eur J Pain* 2017;21:552–61.

- [36] Graven-Nielsen T, Kendall SA, Henriksson KG, Bengtsson M, Sörensen J, Johnson A, Gerdle B, Arendt-Nielsen L. Ketamine reduces muscle pain, temporal summation, and referred pain in fibromyalgia patients. *PAIN* 2000;85:483–91.
- [37] Graven-Nielsen T, Wodehouse T, Langford RM, Arendt-Nielsen L, Kidd BL. Normalization of widespread hyperesthesia and facilitated spatial summation of deep-tissue pain in knee osteoarthritis patients after knee replacement. *Arthritis Rheum* 2012;64:2907–16.
- [38] Grosen K, Fischer IWD, Olesen AE, Drewes AM. Can quantitative sensory testing predict responses to analgesic treatment? *Eur J Pain* 2013;17:1267–80.
- [39] Grosen K, Vase L, Pilegaard HK, Pfeiffer-Jensen M, Drewes AM. Conditioned pain modulation and situational pain catastrophizing as preoperative predictors of pain following chest wall surgery: a prospective observational cohort study. *PLoS One* 2014;9:e90185.
- [40] Gwilym SE, Oag HCL, Tracey I, Carr AJ. Evidence that central sensitisation is present in patients with shoulder impingement syndrome and influences the outcome after surgery. *J Bone Joint Surg Br* 2011;93-B:498–502.
- [41] Habib AS, Kertai MD, Cooter M, Greenup RA, Hwang S. Risk factors for severe acute pain and persistent pain after surgery for breast cancer: a prospective observational study. *Reg Anesth Pain Med* 2019;44:192–9.
- [42] Hansen S, Vaegter HB, Petersen KK. Pretreatment exercise-induced hypoalgesia is associated with change in pain and function after standardized exercise therapy in painful knee osteoarthritis. *Clin J Pain* 2020;36:16–24.
- [43] Hayden JA, van der Windt DA, Cartwright JL, Côté P, Bombardier C. Assessing bias in studies of prognostic factors. *Ann Intern Med* 2013;158:280–6.
- [44] Hermans L, Calders P, Van Oosterwijck J, Verschelde E, Bertel E, Meeus M. An overview of offset analgesia and the comparison with conditioned pain modulation: a systematic literature review. *Pain Physician* 2016;19:307–26.
- [45] Imai Y, Petersen KK, Mørch CD, Arendt Nielsen L. Comparing test–retest reliability and magnitude of conditioned pain modulation using different combinations of test and conditioning stimuli. *Somatosens Mot Res* 2016;33:169–77.
- [46] Ip HYV, Abrishami A, Peng PWH, Wong J, Chung F. Predictors of postoperative pain and analgesic consumption. *Anesthesiology* 2009;111:657–77.
- [47] Izumi M, Petersen KK, Laursen MB, Arendt-Nielsen L, Graven-Nielsen T. Facilitated temporal summation of pain correlates with clinical pain intensity after hip arthroplasty. *PAIN* 2017;158:323–32.
- [48] Jarrell J, Ross S, Robert M, Wood S, Tang S, Stephanson K, Giamberardino MA. Prediction of postoperative pain after gynecologic laparoscopy for nonacute pelvic pain. *Am J Obstet Gynecol* 2014;211:360.e1–8.
- [49] Kehlet H, Jensen TS, Woolf CJ, Centre M. Persistent postsurgical pain: risk factors and prevention. *Lancet* 2006;367:1618–25.
- [50] Kennedy DL, Kemp HI, Ridout D, Yarnitsky D, Rice AS. Reliability of conditioned pain modulation. *PAIN* 2016;157:2410–19.
- [51] Kisler LB, Weissman-Fogel I, Coghill RC, Sprecher E, Yarnitsky D, Granovsky Y. Individualization of migraine prevention. *Clin J Pain* 2019;35:753–65.
- [52] Koh JJ, Kim MS, Sohn S, Song KY, Choi NY, In Y. Duloxetine reduces pain and improves quality of recovery following total knee arthroplasty in centrally sensitized patients. *J Bone Joint Surg* 2019;101:64–73.
- [53] Kosek E, Cohen M, Baron R, Gebhart GF, Mico JA, Rice ASC, Rief W, Sluka AK. Do we need a third mechanistic descriptor for chronic pain states? *PAIN* 2016;157:1382–6.
- [54] Kosek E, Ordeberg G. Lack of pressure pain modulation by heterotopic noxious conditioning stimulation in patients with painful osteoarthritis before, but not following, surgical pain relief. *PAIN* 2000;88:69–78.
- [55] Kuhlmann L, Olesen SS, Grönlund D, Olesen AE, Phillips AE, Faghni M, Drewes AM. Patient and disease characteristics associate with sensory testing results in chronic pancreatitis. *Clin J Pain* 2019;35:786–93.
- [56] Kurien T, Arendt-Nielsen L, Petersen KK, Graven-Nielsen T, Scammell BE. Preoperative neuropathic pain-like symptoms and central pain mechanisms in knee osteoarthritis predicts poor outcome 6 months after total knee replacement surgery. *J Pain* 2018;19:1329–41.
- [57] Lannersten L, Kosek E. Dysfunction of endogenous pain inhibition during exercise with painful muscles in patients with shoulder myalgia and fibromyalgia. *PAIN* 2010;151:77–86.
- [58] Larsen DB. The Predictive Value of Quantitative Sensory Testing—A Systematic Review on Chronic Postoperative Pain and the Analgesic Effect of Pharmaceutical Therapies for Chronic Pain Patients. 2020:4. doi: 10.17605/OSF.IO/HSVYK.
- [59] Lee YC, Lu B, Edwards RR, Wasan AD, Nassikas NJ, Clauw DJ, Solomon DH, Karlson EW. The role of sleep problems in central pain processing in rheumatoid arthritis. *Arthritis Rheum* 2013;65:59–68.
- [60] Lockwood SM, Bannister K, Dickenson AH. An investigation into the noradrenergic and serotonergic contributions of diffuse noxious inhibitory controls in a monoiodoacetate model of osteoarthritis. *J Neurophysiol* 2019;121:96–104.
- [61] Lundblad H, Kreicbergs A, Jansson K-Å. Prediction of persistent pain after total knee replacement for osteoarthritis. *J Bone Joint Surg* 2008;90-B:166–71.
- [62] Maier C, Baron R, Tölle TR, Binder A, Birbaumer N, Birklein F, Gierthmühlen J, Flor H, Geber C, Hüge V, Krumova EK, Landwehrmeyer GB, Magerl W, Maihöfner C, Richter H, Rolke R, Scherens A, Schwarz A, Sommer C, Tronnier V, Üeyler N, Valet M, Wasner G, Treede RD. Quantitative sensory testing in the German Research Network on Neuropathic Pain (DFNS): somatosensory abnormalities in 1236 patients with different neuropathic pain syndromes. *PAIN* 2010;150:439–50.
- [63] Mainka T, Malewicz NM, Baron R, Enax-Krumova EK, Treede R-D, Maier C. Presence of hyperalgesia predicts analgesic efficacy of topically applied capsaicin 8% in patients with peripheral neuropathic pain. *Eur J Pain* 2016;20:116–29.
- [64] Manafí Khanian B, Arendt-Nielsen L, Kjær Petersen K, Samani A, Graven-Nielsen T. Interface pressure behavior during painful cuff algometry. *Pain Med* 2016;17:915–23.
- [65] Martel MO, Petersen K, Cornelius M, Arendt-Nielsen L, Edwards R. Endogenous pain modulation profiles among individuals with chronic pain: relation to opioid use. *J Pain* 2019;20:462–71.
- [66] Martinez V, Fletcher D, Bouhassira D, Sessler DI, Chauvin M. The evolution of primary hyperalgesia in orthopedic surgery: quantitative sensory testing and clinical evaluation before and after total knee arthroplasty. *Anesth Analg* 2007;105:815–21.
- [67] Meeus M, Roussel NA, Truijien S, Nijs J. Reduced pressure pain thresholds in response to exercise in chronic fatigue syndrome but not in chronic low back pain: an experimental study. *J Rehabil Med* 2010;42:884–90.
- [68] Meints SM, Mawla I, Napadow V, Kong J, Gerber J, Chan S-T, Wasan AD, Kaptchuk TJ, McDonnell C, Carriere J, Rosen B, Gollub RL, Edwards RR. The relationship between catastrophizing and altered pain sensitivity in patients with chronic low-back pain. *PAIN* 2019;160:833–43.
- [69] Müller M, Limacher A, Agten CA, Treichel F, Heini P, Seidel U, Andersen OK, Arendt-Nielsen L, Jüni P, Curatolo M. Can quantitative sensory tests predict failed back surgery? *Eur J Anaesthesiol* 2019;36:695–704.
- [70] Naugle KM, Fillingim RB, Riley JL III. A meta-analytic review of the hypoalgesic effects of exercise. *J Pain* 2012;13:1139–50.
- [71] Noiseux NO, Callaghan JJ, Clark CR, Zimmerman MB, Sluka KA, Rakel BA. Preoperative predictors of pain following total knee arthroplasty. *J Arthroplasty* 2014;29:1383–7.
- [72] Olesen SS, Graversen C, Bouwense SAW, van Goor H, Wilder-Smith OHG, Drewes AM. Quantitative sensory testing predicts pregabalin efficacy in painful chronic pancreatitis. *PLoS One* 2013;8:e57963.
- [73] O’Leary H, Smart KM, Moloney NA, Blake C, Doody CM. Pain sensitization associated with nonresponse after physiotherapy in people with knee osteoarthritis. *PAIN* 2018;159:1877–86.
- [74] Petersen KK, Arendt-Nielsen L, Simonsen O, Wilder-Smith O, Laursen MB. Presurgical assessment of temporal summation of pain predicts the development of chronic postoperative pain 12 months after total knee replacement. *PAIN* 2015;156:55–61.
- [75] Petersen KK, Olesen AE, Simonsen O, Arendt-Nielsen L. Mechanistic pain profiling as a tool to predict the efficacy of 3-week nonsteroidal anti-inflammatory drugs plus paracetamol in patients with painful knee osteoarthritis. *PAIN* 2019;160:486–92.
- [76] Petersen KK, Simonsen O, Laursen MB, Arendt-Nielsen L. The role of preoperative radiologic severity, sensory testing, and temporal summation on chronic postoperative pain following total knee arthroplasty. *Clin J Pain* 2018;34:193–7.
- [77] Petersen KK, Simonsen O, Olesen AE, Mørch CD, Arendt-Nielsen L. Pain inhibitory mechanisms and response to weak analgesics in patients with knee osteoarthritis. *Eur J Pain* 2019;23:1904–12.
- [78] Petersen KKKK, Graven-Nielsen T, Simonsen O, Laursen MBMB, Arendt-Nielsen L. Preoperative pain mechanisms assessed by cuff algometry are associated with chronic postoperative pain relief after total knee replacement. *PAIN* 2016;157:1400–6.
- [79] Pinto PR, McIntyre T, Araújo-Soares V, Almeida A, Costa P. Psychological factors predict an unfavorable pain trajectory after hysterectomy. *PAIN* 2018;159:956–67.

- [80] Pinto PR, McIntyre T, Araújo-Soares V, Costa P, Almeida A. Differential predictors of acute post-surgical pain intensity after abdominal hysterectomy and major joint arthroplasty. *Ann Behav Med* 2015;49:384–97.
- [81] Rice DA, Kluger MT, McNair PJ, Lewis GN, Somogyi AA, Borotkanics R, Barratt DT, Walker M. Persistent postoperative pain after total knee arthroplasty: a prospective cohort study of potential risk factors. *Br J Anaesth* 2018;121:804–12.
- [82] Rolke R, Baron R, Maier C, Tolle TR, Treede RD, Beyer A, Binder A, Birbaumer N, Birklein F, Botefur IC, Braune S, Flor H, Hüge V, Klug R, Landwehrmeyer GB, Magerl W, Maihofner C, Rolko C, Schaub C, Scherens A, Sprenger T, Valet M, Wasserka B. Quantitative sensory testing in the German Research Network on Neuropathic Pain (DFNS): standardized protocol and reference values. [Erratum appears in *Pain*. 2006 Nov;125(1-2):197]. *PAIN* 2006;123:231–43.
- [83] Rolke R, Magerl W, Campbell KA, Schalber C, Caspari S, Birklein F, Treede RD. Quantitative sensory testing: a comprehensive protocol for clinical trials. *Eur J Pain* 2006;10:77–88.
- [84] Sangesland A, Støren C, Vaegter HB. Are preoperative experimental pain assessments correlated with clinical pain outcomes after surgery? A systematic review. *Scand J Pain* 2017;15:44–52.
- [85] Schliessbach J, Siegenthaler A, Bütikofer L, Vuilleumier P, Jüni P, Stamer U, Arendt-Nielsen L, Curatolo M. Predicting drug efficacy in chronic low back pain by quantitative sensory tests. *Eur J Pain* 2018;22:973–88.
- [86] Schnitzer TJ, Marks JA. A systematic review of the efficacy and general safety of antibodies to NGF in the treatment of OA of the hip or knee. *Osteoarthritis Cartil* 2015;23:S8–S17.
- [87] Schug SA, Pogatzki-Zahn EM. Chronic pain after surgery or injury. *Pain Clin Updat* 2011;19:19.
- [88] Sivertsen B, Lallukka T, Petrie KJ, Steingrimsdottir OA, Stubhaug A, Nielsen CS. Sleep and pain sensitivity in adults. *PAIN* 2015;156:1433–9.
- [89] Sng BL, Ching YY, Han NLR, Ithnin FB, Sultana R, Assam PN, Sia ATH. Incidence and association factors for the development of chronic post-hysterectomy pain at 4- and 6-month follow-up: a prospective cohort study. *J Pain Res* 2018;11:629–36.
- [90] Szikszay TM, Adamczyk WM, Luedtke K. The magnitude of offset analgesia as a measure of endogenous pain modulation in healthy participants and patients with chronic pain. *Clin J Pain* 2019;35:189–204.
- [91] Vaegter HB, Graven-Nielsen T. Pain modulatory phenotypes differentiate subgroups with different clinical and experimental pain sensitivity. *PAIN* 2016;157:1480–8.
- [92] Vaegter HB, Handberg G, Emmeluth C, Graven-Nielsen T. Preoperative hypoalgesia after cold pressor test and aerobic exercise is associated with pain relief 6 months after total knee replacement. *Clin J Pain* 2017;33:475–84.
- [93] Vaegter HB, Handberg G, Graven-Nielsen T. Isometric exercises reduce temporal summation of pressure pain in humans. *Eur J Pain* 2015;19:973–83.
- [94] Vaegter HB, Handberg G, Graven-Nielsen T. Similarities between exercise-induced hypoalgesia and conditioned pain modulation in humans. *PAIN* 2014;155:158–67.
- [95] Vaegter HB, Handberg G, Jørgensen MN, Kinly A, Graven-Nielsen T. Aerobic exercise and cold pressor test induce hypoalgesia in active and inactive men and women. *Pain Med* 2015;16:923–33.
- [96] Vaegter HB, Lyng KD, Yttereng FW, Christensen MH, Sørensen MB, Graven-Nielsen T. Exercise-induced hypoalgesia after isometric wall squat exercise: a test-retest reliability study. *Pain Med* 2019;20:129–37.
- [97] Vaegter HB, Petersen KK, Mørch CD, Imai Y, Arendt-Nielsen L. Assessment of CPM reliability: quantification of the within-subject reliability of 10 different protocols. *Scand J Pain* 2018;18:729–37.
- [98] Valencia C, Fillingim RB, Bishop M, Wu SS, Wright TW, Moser M, Farmer K, George SZ. Investigation of central pain processing in postoperative shoulder pain and disability. *Clin J Pain* 2014;30:775–86.
- [99] Vollert J, Attal N, Baron R, Freynhagen R, Haanpää M, Hansson P, Jensen TS, Rice ASC, Segerdahl M, Serra J, Sindrup SH, Tölle TR, Treede R-D, Maier C. Quantitative sensory testing using DFNS protocol in Europe: an evaluation of heterogeneity across multiple centers in patients with peripheral neuropathic pain and healthy subjects. *PAIN* 2016;157:750–8.
- [100] Wasner G, Naleschinski D, Binder A, Schattschneider J, McLachlan EM, Baron R. The effect of menthol on cold allodynia in patients with neuropathic pain. *Pain Med* 2008;9:354–8.
- [101] Werner MU, Mjöbo HN, Nielsen PR, Rudin Å. Prediction of postoperative pain. *Anesthesiology* 2010;112:1494–502.
- [102] Wilder-Smith OH. A paradigm-shift in pain medicine. Aalborg: Center for Sensory-Motor Interaction (SMI), Department of Health Science and Technology, Aalborg University, 2013.
- [103] Wilder-Smith OH, Schreyer T, Scheffer GJ, Arendt-Nielsen L. Patients with chronic pain after abdominal surgery show less preoperative endogenous pain inhibition and more postoperative hyperalgesia: a pilot study. *J Pain Palliat Care Pharmacother* 2010;24:119–28.
- [104] Woolf CJ, Salter MW. Neuronal plasticity: increasing the gain in pain. *Science* 2000;288:1765–9.
- [105] Wylde V, Palmer S, Learmonth ID, Dieppe P. The association between pre-operative pain sensitisation and chronic pain after knee replacement: an exploratory study. *Osteoarthritis Cartil* 2013;21:1253–6.
- [106] Wylde V, Sayers A, Lenguerrand E, Gooberman-Hill R, Pyke M, Beswick AD, Dieppe P, Blom AW. Preoperative widespread pain sensitization and chronic pain after hip and knee replacement. *PAIN* 2015;156:47–54.
- [107] Yarnitsky D. Conditioned pain modulation (the diffuse noxious inhibitory control-like effect): its relevance for acute and chronic pain states. *Curr Opin Anaesthesiol* 2010;23:611–15.
- [108] Yarnitsky D, Bouhassira D, Drewes AM, Fillingim RB, Granot M, Hansson P, Landau R, Marchand S, Matre D, Nilsen KB, Stubhaug A, Treede RD, Wilder-Smith OHG. Recommendations on practice of conditioned pain modulation (CPM) testing. *Eur J Pain* 2015;19:805–6.
- [109] Yarnitsky D, Crispel Y, Eisenberg E, Granovsky Y, Ben-Nun A, Sprecher E, Best LA, Granot M. Prediction of chronic post-operative pain: pre-operative DNIC testing identifies patients at risk. *PAIN* 2008;138:22–8.
- [110] Yarnitsky D, Granot M, Nahman-Averbuch H, Khamaisi M, Granovsky Y. Conditioned pain modulation predicts duloxetine efficacy in painful diabetic neuropathy. *PAIN* 2012;153:1193–8.