University of Groningen

Probing Structural Information of Gas-Phase Peptides by Near-Edge X-ray Absorption Mass Spectrometry

Dörner, Simon; Schwob, Lucas; Atak, Kaan; Schubert, Kaja; Boll, Rebecca; Schlathölter, Thomas; Timm, Martin; Bülow, Christine; Zamudio-Bayer, Vicente; Von Issendorf, Bernd

Published in:
Journal of the American Society for Mass Spectrometry

DOI:
[10.1021/jasms.0c00390](https://doi.org/10.1021/jasms.0c00390)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Dörner, S., Schwob, L., Atak, K., Schubert, K., Boll, R., Schlathölter, T., Timm, M., Bülow, C., Zamudio-Bayer, V., Von Issendorf, B., Lau, J. T., Techert, S., & Bari, S. (2021). Probing Structural Information of Gas-Phase Peptides by Near-Edge X-ray Absorption Mass Spectrometry. *Journal of the American Society for Mass Spectrometry*, 32(3), 670-684. <https://doi.org/10.1021/jasms.0c00390>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Probing Structural Information of Gas-Phase Peptides by Near-Edge X-ray Absorption Mass Spectrometry

Simon Dörner,* Lucas Schwob,* Kaan Atak, Kaja Schubert, Rebecca Boll, Thomas Schlathöler, Martin Timm, Christine Bülow, Vicente Zamudio-Bayer, Bernd von Issendorff, J. Tobias Lau, Simone Techert, and Sadia Bari*

Cite This: *J. Am. Soc. Mass Spectrom.* 2021, 32, 670–684

Read Online

ACCESS |

Metrics & More

Article Recommendations

Supporting Information

ABSTRACT: Near-edge X-ray absorption mass spectrometry (NEXAMS) is an action-spectroscopy technique of growing interest for investigations into the spatial and electronic structure of biomolecules. It has been used successfully to give insights into different aspects of the photodissociation of peptides and to probe the conformation of proteins. It is a current question whether the fragmentation pathways are sensitive toward effects of conformational isomerism, tautomerism, and intramolecular interactions in gas-phase peptides. To address this issue, we studied the cationic fragments of cryogenically cooled gas-phase leucine enkephalin ([LeuEnk+H]⁺) and methionine enkephalin ([MetEnk+H]⁺) produced upon soft X-ray photon absorption at the carbon, nitrogen, and oxygen K-edges. The interpretation of the experimental ion yield spectra was supported by density-functional theory and restricted-open-shell configuration interaction with singles (DFT/ROCIS) calculations. The analysis revealed several effects that could not be rationalized based on the peptide's amino acid sequences alone. Clear differences between the partial ion yields measured for both peptides upon C 1s → π*(C=C) excitations in the aromatic amino acid side chains give evidence for a sulfur–aromatic interaction between the methionine and phenylalanine side chain of [MetEnk+H]⁺. Furthermore, a peak associated with N 1s → π*(C=N) transitions, linked to a tautomeric keto-to-enol conversion of peptide bonds, was only present in the photon energy resolved ion yield spectra of [MetEnk+H]⁺.

INTRODUCTION

Proteins and peptides are chains of amino acid residues linked by peptide bonds. They have a variety of biological functions in living organisms, including the catalysis of biochemical reactions, signal transduction, and providing structure to cells. Their functions depend on both the amino acid sequence and the three-dimensional conformation resulting from intramolecular interactions like hydrogen bonds. Investigations of protein and peptide spatial structures are of great importance for research in cell biology, biocatalysis, and drug development. Since the spatial and electronic structures of a molecule are inextricably linked, it is crucial to obtain detailed information about the electronic structure of proteins and peptides as well. This, however, remains a major experimental challenge.

The past two decades saw major advances in the use of near-edge X-ray absorption fine structure (NEXAFS) spectroscopy for studies on both the spatial and electronic structure of biomolecular systems. In NEXAFS spectroscopy, the photoabsorption cross sections for photoionization and resonant photoexcitation of electrons from the atomic core level to unoccupied molecular orbitals (MOs) are determined by measuring, for example, the Auger-electron yield or the X-ray fluorescence while scanning the photon energy across an

atomic absorption edge. The positions and intensities of spectral features in a NEXAFS spectrum allow for the identification of chemical bonds and groups and the characterization of their chemical environments.

NEXAFS spectroscopy in the condensed phase has been used to investigate the electronic and spatial structure of amino acids, peptides, and proteins^{1–11} as well as nucleobases, nucleotides, and strands of deoxyribonucleic acid (DNA).^{12–20} NEXAFS spectra for thin films of all standard amino acids are available for the carbon, nitrogen and oxygen K-edges.^{3,21} The spectral changes caused by the formation of a peptide bond have been identified and characterized.^{1,22} It has been proposed that, in a building block approach, NEXAFS spectra of polypeptides can be approximated from the spectra of their constituents.^{2,5,7} Furthermore, calculations for neutral diglycine suggest that NEXAFS is sensitive toward the conformation of the peptide bond.¹

Received: October 22, 2020

Revised: February 1, 2021

Accepted: February 1, 2021

Published: February 12, 2021

There is a growing interest in NEXAFS studies on biomolecular systems in the gas phase, in order to investigate their intrinsic properties. Electrospray ionization (ESI) has proven to be a reliable technique to produce charged gas-phase biomolecules. Working with charged sample molecules enables the use of advanced ion manipulation tools, including quadrupole mass filters and ion traps as well as separation techniques based on ion mobility, thus allowing the study of fragile molecules in well-defined states and controlled molecular environment. By combining gas-phase NEXAFS spectroscopy with ESI tandem mass spectrometry, the mass-to-charge ratios (m/z) and the yields of the photoionization and photodissociation products of biomolecular ions can be measured as a function of the photon energy. To highlight the position of this technique at the overlap between NEXAFS spectroscopy and (biomolecular) mass spectrometry (MS), it is also referred to as near-edge X-ray absorption mass spectrometry (NEXAMS^{8,23–25}). The photon energy dependent ion yields can reveal correlations between electronic transitions and structurally informative fragmentation pathways. In this direction, NEXAMS could help identify unknown compounds, similar to the 2D UV-MS fingerprinting technique,^{26,27} and could, furthermore, be complementary to IR spectroscopy by determining certain aspects of a molecular structure^{28,29} and, therefore, constraining the search of the calculated conformational space. Additionally, when the dissociation becomes site-selective,²⁵ NEXAMS may then be employed to follow ultrafast charge migration and energy transfer in pump–probe schemes at XUV/X-ray Free-Electron Laser facilities.

In the first NEXAMS study on a protein, Milosavljević et al.⁹ demonstrated that absorption of a single soft X-ray photon by gas-phase protonated cytochrome *c* predominantly leads to nondissociative photoionization. In contrast, absorption of soft X-ray photons by gas-phase protonated leucine enkephalin (LeuEnk; Tyr-Gly-Gly-Phe-Leu) at the carbon K-edge induces fragmentation into immonium ions, short amino acid sequence ions, and side-chain fragments.⁸ By measuring the photo-fragmentation mass spectra of different peptides and proteins in the C 1s excitation and C 1s ionization regime, Egorov et al.³⁰ showed that the correlation between the size of a polypeptide and the degree of fragmentation is based on intramolecular vibrational redistribution (IVR) over the molecule's degrees-of-freedom. The authors also proposed that, in peptides, dissociation driven by IVR is in competition with fast local fragmentation processes involving Auger decay. Furthermore, a study on gas-phase protonated collagen peptides highlighted the involvement of radical-induced dissociation with a low energetic barrier (1–2 eV) in the relaxation processes following photoionization.³¹ Recently, we presented a NEXAMS study on gas-phase protonated methionine enkephalin (MetEnk; Tyr-Gly-Gly-Phe-Met), demonstrating site-selective dissociation of a peptide upon site-specific photoexcitation of the methionine sulfur atom at the sulfur L-edge.²⁵

By measuring the ionization products of multiply protonated ubiquitin (76 residues) at the carbon K-edge, Milosavljević et al.³² demonstrated an interplay between the electronic and spatial structure of a protein. While, overall, the ionization threshold increased with the protonation state, over the region of charge states +4 to +8 the unfolding of the protein compensated for the introduced charge, leading to the effective Coulomb field being virtually constant. In contrast, for the

smaller, largely α -helical polypeptide melittin (26 residues), even though the conformation changed drastically over the charge states of +2 to +4, the ionization energy still increased.¹¹ Additionally, the authors observed an increased rate of secondary ionization by inelastic scattering of Auger electrons in the compact conformer.

The current challenge is to find evidence for the sensitivity of NEXAMS toward the conformation of and intramolecular interactions within small peptides, i.e., investigate whereas the related absorption bands are accompanied by specific cleavage in peptides. In order to address this question, we carried out a comprehensive investigation into the yields of cationic dissociation products of gas-phase protonated LeuEnk and MetEnk ($[\text{LeuEnk}+\text{H}]^+$ and $[\text{MetEnk}+\text{H}]^+$) measured upon soft X-ray photon absorption at the carbon, nitrogen and oxygen K-edges. The analysis of the experimental data is supported by X-ray absorption spectra calculated using density-functional theory (DFT) and restricted-open-shell configuration interaction with singles (ROCIS).

The only difference between the amino acid sequences of LeuEnk and MetEnk is the C-terminal residue, which is leucine or methionine, respectively (see Figure 1a,b). LeuEnk is a well-

Figure 1. Structural formulas of (a) $[\text{LeuEnk}+\text{H}]^+$, (b) $[\text{MetEnk}+\text{H}]^+$ and (c) $[\text{MetEnk}+\text{H}]^+$ with the Gly-Gly peptide bond in enol form. All peptides are N-terminal-protonated. The red lines and blue lines indicate the cleavage sites of the $[\text{c}_2 - \text{Y}_{\text{sc}}]^+$ and y_2b_4^+ fragment, respectively. The green boxes indicate the enol and keto form of the Gly-Gly peptide bond of $[\text{MetEnk}+\text{H}]^+$.

established standard in ESI-based biomolecular mass spectrometry and has been investigated extensively using non-ionizing activation methods, including collision- (CID^{33–35}), surface- (SID³⁶), and ultraviolet photon-induced dissociation,^{37–39} IR–UV double resonance photofragment spectroscopy⁴⁰ and infrared multiphoton dissociation (IRMPD⁴¹) as well as ionizing activation methods, involving vacuum-ultraviolet (VUV) photon-,^{42,43} soft X-ray photon-,⁸ keV ion-⁴⁴ and high-intensity femtosecond IR pulse-induced

dissociation.⁴⁵ The ion yields measured upon soft X-ray absorption by gas-phase $[\text{LeuEnk}+\text{H}]^+$ at the carbon K-edge⁸ are qualitatively in agreement with data reported for gas-phase multiply protonated proteins.^{9,10,30} The characteristic spectral features of the peptide backbone and the aromatic residues tyrosine and phenylalanine were observed in the total ion yield spectrum as well as in the partial ion yield spectra of all reported fragments. From the peak intensity ratios, the conclusion was drawn that $\text{C } 1s \rightarrow \pi^*(\text{CONH})$ excitations in the backbone promote the formation of small immonium ions and side-chain fragments and $\text{C } 1s \rightarrow \pi^*(\text{C}=\text{C})$ excitations in the aromatic side chains contribute more strongly to the formation of larger backbone fragments.

METHODS

Near-Edge X-ray Absorption Mass Spectrometry. The experiments were performed at the NanoclusterTrap end-station^{46,47} at the UES2_PGM beamline at the BESSY II synchrotron (HZB Berlin, Germany). The sample molecules were introduced into the gas phase by electrospray ionization in positive ion mode. The ions were focused into a beam using a radio frequency ion funnel and guided through a radio frequency hexapole ion guide to a quadrupole mass filter. For the measurements on $[\text{MetEnk}+\text{H}]^+$ we combined the NanoclusterTrap with our self-made electrospray ionization source and high-fluence ion funnel.⁴⁸ The data sets of $[\text{LeuEnk}+\text{H}]^+$ were gathered during a different beamtime, where the electrospray ionization source and ion funnel of the NanoclusterTrap were used.

After being filtered according to their mass-to-charge ratio (m/z), the ions passed a 90° bender to prevent neutral molecules from entering the interaction region. Subsequently, the ions were accumulated and stored in a cryogenic linear ion trap while being cooled to ~ 20 K with helium buffer gas. After accumulation, the ions were irradiated with the soft X-ray photons for 100 ms. Afterward, trapped cationic products were analyzed using a reflectron time-of-flight mass spectrometer. The photon flux was measured by a photodiode behind the interaction region and was in the order of 10^{12} s^{-1} . The design of the NanoclusterTrap instrument constraints the detection of the photoproducts only on a small m/z window. In order to detect, for both peptides, most of the immonium ions and side-chain fragments, which have reported to be the main photoproducts observed for $[\text{LeuEnk}+\text{H}]^+$ at the carbon K-edge,⁸ the trapping parameters were adjusted for a mass range of about m/z 50–180. Consequently, expected backbone fragments with m/z -ratios outside of the detection window (b_3 , b_2 , GF) were not detected.

Photon energy scans on $[\text{LeuEnk}+\text{H}]^+$ and $[\text{MetEnk}+\text{H}]^+$ were performed, scanning the carbon (280–320 eV), nitrogen (395–420 eV), and oxygen K-edges (530–550 eV) in steps of 125 meV. The monochromator bandwidth was set to 150 meV at the carbon K-edge and to 250 meV at the nitrogen and oxygen K-edges. To remove contributions by fragments arising from collisions with the buffer gas, for each scan a background spectrum without photons was recorded and subtracted from each individual mass spectrum. The mass resolution $m/\Delta m$ was 590 at m/z 77, 214 at m/z 107, 317 at m/z 120, and 800 at m/z 147. Each mass spectrum was normalized to the photon flux. The peptide samples consisted of 30 μM of peptide in 1:1 water/methanol with 1 vol % of formic acid. Leucine-enkephalin acetate salt hydrate (purity $\geq 95\%$), $[\text{Met}^5]$ -enkephalin acetate salt hydrate (purity $\geq 95\%$) as well as

methanol and formic acid were purchased from Merck/Sigma-Aldrich and were used without additional purification.

Data Treatment and Fitting. Photon-energy-dependent ion-yield spectra, in the following referred to as partial NEXAMS spectra, were calculated from the photon-flux-normalized mass spectra measured at the different photon energies by summarizing the data points within a window of $\pm 0.4 m/z$ around the center of a mass peak. The total NEXAMS spectrum of a scan, calculated by summing up the partial NEXAMS spectra of all detected fragments (m/z 57–147), was used as an approximation for the total ion yield.

The experimental broadening of the spectroscopic peak can be attributed to several factors. Contributions to the broadening from the $1s$ -core-hole lifetime have values for carbon, nitrogen and oxygen which are expected to be in the order of 0.1,⁴⁹ 0.1,⁵⁰ and 0.15 eV,⁵¹ respectively. The photon energy bandwidth, defined by the opening of the exit slit of the beamline, accounts for additional broadening of 0.15 eV at the carbon K-edge and 0.25 eV at the nitrogen and oxygen K-edges. Other main sources of broadening are intrinsic to the size of the molecule and to X-ray spectroscopy. The former leads to the overlap of electronic excitations due to multiple virtually equivalent photon absorbing sites and the latter to vibrational excitations in the core-hole excited state (Franck–Condon principle). Finally, the contribution of the vibrational excitations in the initial state of the peptides ions is limited by the cooling of the ions in the cryogenic trap.

In order to identify and separate overlapping spectral features, the total NEXAMS spectra were fitted using Gaussian functions. The number of functions as well as the starting points and expected boundaries for the fit parameters were chosen based on reference data from literature and on the DFT/ROCIS-calculated X-ray absorption spectra (see the “Total NEXAMS spectra” section). In addition, a sigmoid function was applied to fit the peptide’s respective core-ionization edge. The core-ionization edge of an element within a molecule is the superposition of the ionization edges of all corresponding atoms, influenced by their chemical environments. Therefore, instead of the sigmoid function’s point of inflection E_{infl} the x -intercept of the tangent of E_{infl} was considered the peptide’s ionization potential IP .

The area A_{peak} under a Gaussian function was calculated directly from the fitting parameters. Thus, A_{peak} is the ion yield attributed to an ensemble of electronic transitions integrated over the whole spectrum. In contrast, the area of an ionization edge A_{edge} was calculated by integrating the respective sigmoid fit function between zero and $IP + 2\Delta E$ with $\Delta E = E_{\text{infl}} - IP$. The sum of the areas of all Gaussian functions plus the area of the ionization edge was considered as the total area under a spectrum A_{spec} . The partial NEXAMS spectra of individual fragments were fitted in the same way, with the exception that the peak positions and peak widths were taken from the fits of the total NEXAMS spectra.

The trapping efficiency of the NanoclusterTrap is not constant across the trapping window. Therefore, all NEXAMS spectra were normalized. The total NEXAMS spectra were normalized to the maximum intensity of the ionization edge and the partial NEXAMS spectra of individual fragments were normalized to A_{spec} .

DFT/ROCIS Calculations. All calculations were carried out with the ORCA program package.⁵² The molecular geometries were optimized by DFT using the Ahlrichs TZVP basis set⁵³ in combination with the B3LYP exchange-correlation func-

Figure 2. Photodissociation mass spectra of gas-phase $[\text{LeuEnk}+\text{H}]^+$ (blue line) and $[\text{MetEnk}+\text{H}]^+$ (red line; inverted) measured at 288 eV. sc: side-chain fragment, vin: vinyl cation, im: immonium ion, acr: acryloyl cation.

tional.^{54,55} The optimized geometries are shown in the Supporting Information (Figures S1 and S7). DFT/ROCIS calculations using parameters $c_1 = 0.18$, $c_2 = 0.20$, and $c_3 = 0.40$,⁵⁶ the TZVP basis set, and the B3LYP functional were performed to obtain the transition energies and dipole moments for the carbon, nitrogen, and oxygen K-edges. The resolution-of-identity approximation^{57–61} in combination with an auxiliary basis set generated by the AutoAux generation scheme⁶² was used during all calculations. Furthermore, all DFT calculations were performed on a dense integration grid (ORCA Grid4). In all calculations relativistic effects were taken into account using zeroth-order regular approximation (ZORA).⁶³ Vibronic effects were not taken into account.

In order to determine the energy shifts of the calculated transition energies, inherent to the DFT/ROCIS calculation, theoretical X-ray absorption spectra were calculated and matched with the experimental total NEXAMS spectra. The theoretical absorption spectra were produced by broadening the calculated transition lines using a pseudo-Voigt profile, approximated by linear combination of a Gaussian (fwhm = photon energy bandwidth + 0.4 eV) and Lorentzian term (fwhm = 1s-core-hole-lifetime broadening).^{64–66} The coefficient for the linear combination and the fwhm of the Voigt profile f_V were calculated according to the work of Thompson et al.⁶⁷ The resulting f_V values are ~ 0.6 , ~ 0.7 , and ~ 0.75 eV for the carbon, nitrogen, and oxygen K-edge measurement, respectively.

RESULTS AND DISCUSSION

Fragment Identification. At all absorption edges, qualitatively, the same fragments were detected. Thus, for the identification of the detected fragments the mass spectra measured at 288 eV, shown in Figure 2, are representative for all scans. Compared to $[\text{MetEnk}+\text{H}]^+$ the trapping window for $[\text{LeuEnk}+\text{H}]^+$ was shifted about 10 m/z to lower masses. The mass spectra are in line with the carbon K-edge data reported by González-Magaña et al.⁸ and largely in agreement with spectra produced by keV ion-induced dissociation⁴⁴ and VUV photon-induced dissociation at photon energies higher than the ionization threshold.⁴³

The most intense peaks at m/z 91, 107, 120, and 136 are common for molecules containing phenylalanine (Phe) and tyrosine (Tyr) residues. They are attributed to the Phe and Tyr immonium ions (F_{im} : 120 u, Y_{im} : 136 u) and the respective side-chain fragments (F_{sc} : 91 u, Y_{sc} : 107 u).^{68,69} The less intense peaks at m/z 57 and 86 measured for $[\text{LeuEnk}+\text{H}]^+$ stem from the leucine (Leu) side chain (L_{sc} : 57 u) and the

leucine immonium ion (L_{im} : 86 u). For $[\text{MetEnk}+\text{H}]^+$, instead, peaks matching the methionine (Met) side chain (M_{sc} : 75 u) and methionine immonium ion (M_{im} : 104 u) are present.

The peaks at m/z 131 and 147 have previously been assigned to the $[c_2 - Y_{\text{sc}}]^+_{8,45}$ and $y_2b_4^+$ fragment^{38,39} of $[\text{LeuEnk}+\text{H}]^+$ (see Figure 1a). The fragments are labeled according to the nomenclature established by Roepstorff and Fohlmann.⁷⁰ Alternatively, these peaks might be attributed to acryloyl cations of Phe and Tyr (F_{acr} : 131 u, Y_{acr} : 147 u). The peaks at m/z 103 and 119 indicate vinyl cations of Phe and Tyr (F_{vin} : 103 u, Y_{vin} : 119 u). The vinyl and acryloyl cations of Phe have been reported for low energy CID of Ag^+ -phenylalanine complexes.⁷¹ Peaks at m/z 119 and 147 have been observed upon UV-induced dissociation of cold protonated Tyr.⁷² The low energy CID spectrum of protonated 3-iodotyrosine⁷³ shows corresponding peaks at m/z 245 (119 + 126) and m/z 273 (147 + 126).

The peaks at m/z 65, 77, and 79 can be assigned to different cyclic fragments of the aromatic side chains. From organic mass spectrometry it is known that a toluene cation (F_{sc}) can rearrange into a tropylium cation^{74,75} which can then undergo a loss of C_2H_2 , leading to the formation of the cyclopentadienyl cation (C_5H_5^+ : 65 u).^{76,77} This cation also appears in the electron ionization dissociation (EID) spectra of Phe and toluene.⁷⁸ The Y_{sc} fragment can rearrange into a hydroxy-tropylium cation and a subsequent loss of CO can lead to the arenium ion (C_6H_7^+ : 79 u).^{79,80} Fragments at m/z 79 have been observed upon UV/VUV-induced dissociation of protonated Tyr⁸¹ and EID of p-cresol.⁷⁸ A peak at m/z 77 is a strong indication for the phenyl cation (C_6H_5^+ : 77 u), which has been measured as a fragment of both Tyr and Phe upon EID⁷⁸ and photon-induced dissociation in the UV/VUV range.⁸¹ By EID of 1-phenylethanol it has been shown that a loss of H_2 from the arenium ion can lead to the phenyl cation.⁸² However, direct bond cleavage and decomposition of larger Phe and Tyr related fragments are plausible origins for this cation as well. The remaining weak peaks across the mass spectra are due to small internal and terminal fragments. A list of the discussed fragments as well as their chemical structures can be found in the Supporting Information (Table S9 and Figure S16).

Total NEXAMS Spectra. Carbon K-Edge. The total NEXAMS spectra of $[\text{LeuEnk}+\text{H}]^+$ (Figure 3a) and $[\text{MetEnk}+\text{H}]^+$ (Figure 3c) measured at the carbon K-edge appear very similar at first. The spectra show two intense narrow main features at ~ 285 and ~ 288 eV, a small feature at ~ 287 eV, creating a shoulder on the left flank of the second main peak, and two broad features centered at ~ 291.5 and ~ 294 eV. The

Figure 3. Experimental carbon K-edge ion yield spectra of (a) [LeuEnk+H]⁺ and (c) [MetEnk+H]⁺ as well as DFT/ROCIS-calculated carbon K-edge X-ray absorption lines of (b) [LeuEnk+H]⁺ (shifted +11.3 eV) and (d) [MetEnk+H]⁺ (shifted +11.7 eV). The experimental spectra are normalized to the maximum intensity of the C 1s ionization edge. The calculated discrete absorption lines were broadened using a pseudo-Voigt profile with 0.6 eV fwhm. IP: ionization potential.

spectrum of [LeuEnk+H]⁺ was fitted with three intense narrow peaks CK1a, CK1b, and CK3 located at 285.0, 285.4, and 288.1 eV, two weak narrow peaks CK2 and CK4 at 287.2 and 288.6 eV, and two broad features CK5 and CK6 centered at 290.1 and 291.8 eV. The respective peak positions determined for [MetEnk+H]⁺ are CK1a: 285.1, CK1b: 285.4, CK2: 287.3, CK3: 288.2, CK4: 288.7, CK5: 290.1, and CK6: 291.8 eV. The assignments of these peaks, listed in Table 1, are discussed in the following.

Peaks CK1a and CK1b are both attributed to C 1s $\rightarrow \pi^*(C=C)$ transitions in the aromatic side chains of the Tyr and Phe residues.^{21,22,86} In the DFT/ROCIS-calculated data of both peptides the energies for transitions in the Tyr side chain were slightly higher than for the respective transitions in the Phe side chain. This suggests an assignment of CK1a to transitions in the Phe side chain and of CK1b to transitions in

the Tyr side chain. However, across all fragments no significant differences between the ratios of these peaks to each other were observed. Therefore, in the following both peaks are combined into one feature CK1 with its maximum at 285.1 eV. Features at similar positions as CK1 have also been reported for gas-phase benzene⁸³ as well as thin films of benzene and phenol.⁸⁴ Feature CK2 is attributed to C 1s $\rightarrow \pi^*(C=C)OH$ transitions in the Tyr side chain only.^{21,22,85,86} The second main feature CK3 originates from C 1s $\rightarrow \pi^*(CONH)$ transitions in the peptide backbone.^{1,4,22}

The energy position of peak CK4 is in agreement with C 1s $\rightarrow \sigma^*(C-NH_3^+)$ transitions reported for thin film data of Tyr (288.8 eV)²¹ as well as C 1s $\rightarrow \pi^*(C=O)OH$ transitions in the carboxylic acid groups of Leu,²¹ Met²¹ and isoleucine⁴ (all 288.6 eV). The DFT/ROCIS-calculated data show transitions into MOs with high $\pi^*(C=O)OH$ contribution at ~288.7 eV for [LeuEnk+H]⁺ (LUMO+11 Figure S4) and at ~288.4 eV for [MetEnk+H]⁺ (LUMO+8 Figure S10, LUMO+9 Figure S11). First significant transitions into σ^* orbitals of the C-NH₃⁺ group appear at 288.9 eV for [LeuEnk+H]⁺ (LUMO Figure S2, LUMO+1 Figure S3) and at 289.4 eV for [MetEnk+H]⁺ (LUMO Figure S8, LUMO+1 Figure S9) in the calculated data. Since the N-terminal regions of the DFT-optimized geometries of both peptides are fairly similar, the shift between these energies is a theoretical evidence for a high sensitivity of the energy of the $\sigma^*(C-NH_3^+)$ orbital toward the peptide conformation. Noteworthy electronic transitions into MOs of the Met side chain (mainly LUMO+20, LUMO+23, LUMO+27 and LUMO+28 Figures S12–S15) and MOs of the Leu side chain (mainly LUMO+23 and LUMO+26 Figures S5 and S6) also appear in the energy region of peak CK4 at 288.0–289.2 eV and 287.4–288.6 eV, respectively. However, for both peptides the intensities calculated for C 1s $\rightarrow \pi^*(C=O)OH$ transitions are 1 order of magnitude higher than for the transitions into MOs of the Met or Leu side chains. In conclusion, because of a better agreement with our calculations and peak positions reported in literature we consider the assignment of peak CK4 to C 1s $\rightarrow \pi^*(C=O)OH$ transitions as more plausible. Features CK5 and CK6 cannot be assigned to single transitions and represent the accumulation of various transitions into σ^* orbitals and potentially Rydberg states and shape resonances.

The area of peak CK1 normalized to the maximum intensity of the C 1s ionization edge is ~0.71 for [LeuEnk+H]⁺ and ~0.80 for [MetEnk+H]⁺. This means, overall, the ion yield upon excitations in the aromatic side chains is ~14.3% higher for [MetEnk+H]⁺ than for [LeuEnk+H]⁺. Experimental data reported for Leu and Met do not show transitions in the energy range of CK1.^{3,21} The DFT/ROCIS-calculated carbon K-edge absorption spectra of both peptides show only transitions related to the aromatic side chains in this energy regime. Thus, we propose that there is an intramolecular interaction in play, promoting fragmentation upon photoexcitation in the aromatic side chains. This idea is discussed in more detail in the “Evidence for Sulfur-Aromatic Interactions” section based on the partial ion yields of different fragments.

Nitrogen K-Edge. The total NEXAMS spectra of [LeuEnk+H]⁺ and [MetEnk+H]⁺ measured at the nitrogen K-edge are shown in Figure 4a,c. For both peptides there are two narrow peaks at ~401.4 and ~402.8 eV and a broad but distinct peak at ~406 eV. [MetEnk+H]⁺ additionally exhibits one weak, isolated feature at 399.2 eV. From about 408 to 411 eV the intensities in both spectra increase only slightly because the

Table 1. Experimental Energy Positions (Expt) and Proposed Assignments of the Spectral Features Observed in the Total Carbon K-Edge NEXAMS Spectra of [LeuEnk+H]⁺ and [MetEnk+H]⁺ as Well as Energies of the Corresponding Prominent DFT/ROCIS-Calculated Absorption Lines (Calcd)

peak	position (eV)				assignment	ref
	[LeuEnk+H] ⁺		[MetEnk+H] ⁺			
	expt	calcd	expt	calcd		
CK1a	285.0	285.0–285.6	285.1	285.1–285.6	→ $\pi^*(C=C)^a$	83, 84
CK1b	285.4	285.3–286.0	285.4	285.9–286.1, 286.6	→ $\pi^*(C=C)^b$	84
CK2	287.2	286.9	287.3	287.5	→ $\pi^*(C=C)OH$	21, 22, 85, 86
CK3	288.1	288.0–288.6	288.2	287.9–288.0, 289.1	→ $\pi^*(CONH)$	1, 4, 22
CK4	288.6	288.7	288.7	288.4	→ $\pi^*(C=O)OH$	4, 21
		288.9, 289.0		289.4, 289.5	→ $\sigma^*(C-NH_3^+)$	21
		287.4–288.6			→ $\sigma^*(C-C)$ Leu	
				287.9, 288.6, 288.9–289.2	→ $\sigma^*(C-S)$ Met	
CK5	290.1		290.1		(→ σ^*)	
CK6	291.8		291.8		(→ σ^*)	
IP ^c	292.0		292.1		C 1s IP	

^aMainly Phe side chain. ^bMainly Tyr side chain. ^cIP: ionization potential.

Figure 4. Experimental nitrogen K-edge ion yield spectra of (a) [LeuEnk+H]⁺ and (c) [MetEnk+H]⁺ as well as DFT/ROCIS-calculated carbon K-edge X-ray absorption lines of (b) [LeuEnk+H]⁺ (shifted +12.2 eV) and (d) [MetEnk+H]⁺ (shifted +12.6 eV). The experimental spectra are normalized to the maximum intensity of the N 1s ionization edge. The calculated discrete absorption lines were broadened using a pseudo-Voigt profile with 0.7 eV fwhm. IP: ionization potential.

falling slope of the 406 eV feature is compensated by the rising slope of the N 1s ionization edge. The total NEXAMS

spectrum of [LeuEnk+H]⁺ was fitted with three peaks NK1, NK2, and NK3 positioned at 401.4, 402.6, and 406.0 eV, respectively. The positions of peaks NK1 and NK2 were identical for [MetEnk+H]⁺ and [LeuEnk+H]⁺. The peak NK3 of [MetEnk+H]⁺ was shifted by 0.4 eV to an energy position of 406.4 eV. The isolated feature at 399.2 eV in the spectrum of [MetEnk+H]⁺ was fitted with one Gaussian peak NK4 centered at the same energy position. During the fitting procedure it became apparent that in the case of [MetEnk+H]⁺ there is a shoulder on the low-energy slope of the ~406 eV feature. Therefore, another peak NK5 at 404.4 eV was added. In the following the assignments of all peaks are discussed. The assignments are listed in Table 2.

Peak NK1 is attributed to N 1s → $\pi^*(CONH)$ transitions in peptide bonds.^{1,3,4} It is important to note that because of the delocalization of the π -electrons in a peptide bond ($O=C-NH \rightleftharpoons O^--C=N^+H$), both the C=O and C-N bond have partial π -bond character. The assignment of peak NK2 is controversial. Features at this position have been assigned to N 1s → $\pi^*(N-C\alpha)$ transitions in gas-phase alanine (402.4 eV⁸⁷) and diglycine (402.44 eV⁶) as well as N 1s → $\pi^*(-NH=)$ transitions in thin films of leucoemeraldine ([C₆H₄NH]_n; 402.1 eV).⁸⁸ For different homopolypeptides, Zubavichus et al.⁴ suggested N 1s → $\pi^*(CONH)$ transitions (402.7 eV) that are energetically shifted because of an α -helical peptide conformation. Another common assignment for peaks around 402.5 eV are transitions into 3p-Rydberg orbitals.¹ The DFT/ROCIS calculations for [MetEnk+H]⁺ (Figure 4d) show a small but distinct feature at 402.7 eV. The probed MOs are, however, highly delocalized. Ultimately, we rated the assignment of NK2 to N 1s → $\pi^*(N-C\alpha)$ transitions as most plausible.

The spectral feature NK3 involves mainly N 1s → $\sigma^*(N-C\alpha)$ and N 1s → $\sigma^*(N-H)$ transitions in the peptide backbone.^{1,4} In the energy regime of NK3 the DFT/ROCIS calculations for both peptides (Figure 4b,d) show prominent absorption lines attributed to transitions from the N-terminal nitrogen atom into MOs delocalized over the NH₃⁺ group and the tyrosine residue. Most noteworthy are intense absorption lines at 405.8 eV for [LeuEnk+H]⁺ and 406.4 eV for [MetEnk+H]⁺, leading to distinct peaks at about the same photon energies. In the experimental spectra of both peptides such

Table 2. Experimental Energy Positions (Expt) and Proposed Assignments of the Spectral Features Observed in the Total Nitrogen K-Edge NEXAMS Spectra of [LeuEnk+H]⁺ and [MetEnk+H]⁺ as Well as Energies of the Corresponding Prominent DFT/ROCIS-Calculated Absorption Lines (Calcd)

peak	position (eV)				assignment	ref
	[LeuEnk+H] ⁺		[MetEnk+H] ⁺			
	expt	calcd	expt	calcd		
NK1	401.4	400.6–401.7	401.4	401.2–401.6, 402.7	→ $\pi^*(\text{CONH})$	1, 3, 4
NK2	402.6		402.6		→ $\pi^*(\text{N}-\text{Ca})$	6, 87, 88
NK3	406.0		406.4		(→ σ^*)	1, 4
NK4			399.2		→ $\pi^*(\text{C}=\text{N})$	89
NK5		403.1	404.4	403.7, 403.9	NH ₂ → $\sigma^*(\text{N}-\text{H})$	90
IP ^a	407.4		407.6		NH ₃ ⁺ → $\sigma^*(\text{N}-\text{H})$	1, 4
					N 1s IP	

^aIP: ionization potential.

peaks were not observed. Furthermore, the calculated absorption spectrum of [LeuEnk+H]⁺ features a peak at 403.1 eV stemming mainly from a single absorption line attributed to N 1s → $\sigma^*(\text{N}-\text{H})$ transitions in the NH₃⁺ group. For [MetEnk+H]⁺ this peak is shifted to 403.8 eV. Corresponding peaks were not observed in the experimental spectra. However, peak NK5 in the ion yield spectrum of [MetEnk+H]⁺ might be connected to these transitions.

For peak NK4 we consider N 1s → $\pi^*(\text{C}=\text{N})$ transitions in peptide bonds in enol form and N 1s → $\sigma^*(\text{N}-\text{H})$ transitions in unprotonated amino groups as plausible reasons. The assignment is discussed in detail in the “Discussion of the 399.2 eV feature of [MetEnk+H]⁺” section.

Oxygen K-Edge. At the oxygen K-edge, qualitatively, the total NEXAMS spectra of both peptides (Figure 5a,c) exhibit the same features: a dominant feature at ~532 eV and three weaker features centered at ~535, ~538, and ~543 eV. For [LeuEnk+H]⁺ the fitting procedure led to one intense peak OK1 at 532.1 eV and two less intense peaks OK2 and OK3 located at 534.4 and 539.0 eV. The respective peak positions determined for [MetEnk+H]⁺ are OK1: 532.2, OK2: 535.2, and OK3: 538.9 eV. The assignments of these peaks are discussed in the following and are listed in Table 3.

Peak OK1 can be assigned to transitions of carbonyl oxygen 1s electrons into $\pi^*(\text{CONH})$ and $\pi^*(\text{C}=\text{O})\text{OH}$ orbitals. Transitions of these types have been reported for gas-phase diglycine ($\pi^*(\text{CONH})$: 532.2 eV;^{1,6} $\pi^*(\text{C}=\text{O})\text{OH}$: 531.7 eV,¹ 531.85 eV⁶). Furthermore, O 1s → $\pi^*(\text{CONH})$ transitions have been measured for N-methylacetamide (~532 eV^{95,96}) and O 1s → $\pi^*(\text{C}=\text{O})\text{OH}$ transitions for formic acid (532.17 eV⁹²). A peak in the energy region of OK1 has also been observed for gas-phase substance P (532.15 eV⁹¹), a peptide composed of 11 amino acid residues. It should be noted that the calculated absorption line attributed to O 1s → $\pi^*(\text{CONH})$ transitions in the Tyr-Gly peptide bond of [MetEnk+H]⁺ is shifted by 0.5 eV to higher energies compared to the energy of the respective transition in [LeuEnk+H]⁺, which leads to a peak at 533.3 eV in the calculated absorption spectrum of [MetEnk+H]⁺. This effect was, however, not observed in the experimental spectra.

The position of peak OK2 in the total NEXAMS spectrum of [LeuEnk+H]⁺ (534.4 eV) is shifted 0.8 eV toward lower photon energies in comparison to [MetEnk+H]⁺ (535.2 eV). The DFT/ROCIS-calculated oxygen K-edge absorption spectrum of [LeuEnk+H]⁺ shows a feature at ~534.7 eV which is caused mainly by excitations of hydroxy oxygen 1s

Figure 5. Experimental oxygen K-edge ion yield spectra of (a) [LeuEnk+H]⁺ and (c) [MetEnk+H]⁺ as well as DFT/ROCIS-calculated carbon K-edge X-ray absorption lines of (b) [LeuEnk+H]⁺ (shifted +14.1 eV) and (d) [MetEnk+H]⁺ (shifted +14.6 eV). The experimental spectra are normalized to the maximum intensity of the O 1s ionization edge. The calculated discrete absorption lines were broadened using a pseudo-Voigt profile with 0.75 eV fwhm. IP: ionization potential

electrons into adjacent π^* -orbitals of the Tyr side chain and the carboxylic acid group. For [MetEnk+H]⁺ this feature appears at ~535 eV. An assignment to O 1s → $\pi^*(\text{C}=\text{O})\text{OH}$ transitions is in line with experimental data of gas-phase formic acid (535.37 eV⁹²), gas-phase diglycine (535.3 eV,¹ 535.4 eV⁶) and gas-phase Met (535.4 eV).⁹⁷ O 1s → $\pi^*(\text{C}=\text{C})\text{OH}$

Table 3. Experimental Energy Positions (Expt) and Proposed Assignments of the Spectral Features Observed in the Total Oxygen K-Edge NEXAMS Spectra of [LeuEnk+H]⁺ and [MetEnk+H]⁺ as Well as Energies of the Corresponding Prominent DFT/ROCIS-Calculated Absorption Lines (Calcd)

peak	position (eV)				assignment	ref
	[LeuEnk+H] ⁺		[MetEnk+H] ⁺			
	expt	calcd	expt	calcd		
OK1	532.1	531.9–532.1, 532.8 532.5	532.2	531.9–532.1, 533.3 532.4	CO → π^* (CONH) CO → π^* (C=O)OH	1, 6, 91 1, 6
OK2	534.4	534.7–534.9 534.3	535.2	534.8, 535.0 535.0–535.2	OH → π^* (C=O)OH OH → π^* (C=C)OH	1, 6, 92 93, 94
OK3	539.0		538.9		(→ σ^*)	
IP ^a	539.9		539.9		O 1s IP	

^aIP: ionization potential.

Figure 6. Normalized partial ion yields of the phenylalanine (F), tyrosine (Y), and methionine (M) immonium ions (im) and complete side-chain fragments (sc) as well as the phenyl cation (C_6H_5^+) and arenium ion (C_6H_7^+) measured upon photofragmentation of protonated methionine enkephalin at the (a–c) carbon, (d–f) nitrogen, and (g–i) oxygen K-edges. Each spectrum is normalized to its total area.

transitions have been reported for gas-phase phenol (534.9 eV⁹³) and hydroquinone (534.6 eV⁹³) as well as tyrosine-terminated propanethiol assembled on a gold surface (535.5 eV⁹⁴).

The strong shift of peak OK2 in the experimental data might be due to hydrogen bonding. For water^{98–100} and different alcohols¹⁰¹ it has been demonstrated that there can be significant differences, especially in the 535 eV region, between the oxygen K-edge absorption spectra of an H-donating and H-accepting molecule. The 535 eV region of the ion yield spectrum of [MetEnk+H]⁺ could furthermore be influenced by O 1s → $\sigma^*(\text{C–OH})$ transitions in peptide bonds in enol form. The broad feature OK3 represents the entirety of excitations into σ^* orbitals and, potentially, excitations leading to Rydberg states and shape resonances.

The areas of peaks OK1 and OK2 normalized to the maximum intensity of the O 1s ionization edge are 2.45 and 0.64 for [LeuEnk+H]⁺ and 2.00 and 0.53 for [MetEnk+H]⁺. This means the total measured ion yield upon resonant O 1s → $\pi^*(\text{CONH})$ and O 1s → $\pi^*(\text{C=O})\text{OH}$ excitations is ~18% lower for [MetEnk+H]⁺ than for [LeuEnk+H]⁺. With ~17% this relative difference is very similar for peak OK2. For the immonium ions and side-chain fragments of Tyr and Phe, which govern the biggest part of the total measured ion yield, the area of peak OK1 is 12 to 38% smaller for [MetEnk+H]⁺ than for [LeuEnk+H]⁺ (see Tables S4 and S8). This shows that the difference between the total NEXAMS spectra is not

caused by the decreased yield of a single fragment. These findings are surprising because similar effects were not observed for the transitions into $\pi^*(\text{CONH})$ orbitals at the carbon and nitrogen absorption edges. Also, the DTF/ROCIS-calculated oxygen K-edge absorption spectra (Figure 5b,d) do not reflect this difference. The reduced ion yield for [MetEnk+H]⁺ upon O 1s → $\pi^*(\text{C=O})$ excitation either means that overall the rate of these excitations is reduced or that the rates in both peptides are on a similar level but in [MetEnk+H]⁺ there are competing dissociation channels leading to fragments with m/z ratios that are not within the trapping window. The reason for this effect could be connected to C=O...H–N hydrogen bonds between different peptide bonds. More information about the gas-phase structure of protonated MetEnk is needed for a conclusive interpretation of these data.

Partial NEXAMS Spectra Measured for [MetEnk+H]⁺

At each absorption edge, qualitatively, all partial NEXAMS spectra exhibit the same spectral features as the respective total NEXAMS spectrum (see the “Total NEXAMS spectra” section). The intensities of the features, however, show clear differences. The normalized yields of all detected immonium ions and complete side-chain fragments as well as the arenium ion (C_6H_7^+) and phenyl cation (C_6H_5^+) of [MetEnk+H]⁺ are shown in Figure 6. The non-normalized peak areas and intensities determined for these fragments by fitting are listed in the Supporting Information (Tables S2–S4 and S6–S8).

At all absorption edges the ratio between the maximum intensities of the $\pi^*(\text{CONH})$ feature and the ionization edge is smaller for the complete side-chain fragments than for the immonium ions. For the side-chain daughter fragments this ratio is even smaller. This effect is in general agreement with observations made at the carbon K-edge of $[\text{LeuEnk}+\text{H}]^+$.⁸ Overall, the normalized yields of F_{im} and M_{im} only show small differences (Figures 6a,d,g). In comparison, at all absorption edges, the normalized yields of Y_{im} measured upon excitations into $\pi^*(\text{CONH})$ orbitals are increased. Immonium ions are protonated fragments. Thus, migration of a proton to the fragmentation site is necessary in order to produce these ions. The Tyr residue is the N-terminal residue of $[\text{MetEnk}+\text{H}]^+$. The N-terminal amino group is, commonly, the most favorable protonation site in a peptide without basic amino acid side chains. Therefore, the increased yields of Y_{im} observed after excitations into the peptide bond is very likely connected to the close proximity to the N-terminal proton.

The normalized yields of the arenium ion and phenyl cation are almost identical at all absorption edges (Figures 6c,f,i). This suggests that both fragments stem from the same fragmentation pathway. The fragmentation of Y_{sc} into the arenium ion and the subsequent formation of the phenyl cation has been demonstrated in a CID study by Bauerle and Brodbelt⁸⁰ on cationic methylene-substituted phenol (equivalent to Y_{sc}). In contrast, there is no evidence for the arenium ion as a fragment of the Phe residue or toluene. From this we conclude that upon soft X-ray photon absorption in $[\text{MetEnk}+\text{H}]^+$ the phenyl cation is produced mainly as a daughter fragment of the Tyr side chain with the arenium ion as an intermediate.

All amino acid side chains in $[\text{MetEnk}+\text{H}]^+$ contain at least three carbon atoms. Thus, at the carbon K-edge more prominent effects caused by excitations in the amino acid side chains can be expected than at the other absorption edges. The normalized yields of the complete side-chain fragments (F_{sc} , Y_{sc} and M_{sc}) indeed show the biggest differences at the carbon K-edge (Figure 6b). Upon direct ionization the normalized yield of Y_{sc} is higher and the normalized yield of F_{sc} is lower than for M_{sc} . In the σ^* and ionization edge regime the yield of F_{sc} appears to rise in the form of one continuous spectral feature, while the yields of Y_{sc} and M_{sc} are clearly rising in two steps. Visually, this creates the effects that the ionization potentials in the partial NEXAMS spectra of these fragments are at different energy positions. However, the fitting procedure gave more consistent results when keeping the energy positions of all spectral features and the ionization edge at similar positions.

The numbers of carbon atoms in the Phe and Tyr side chain are identical. The rate of direct ionizations should therefore not be the reason for the different normalized yields of these fragments at the ionization edge. One explanation might be that the additional degree of freedom in Y_{sc} compared to F_{sc} makes this fragment more robust toward dissociation into daughter fragments. Also the localized positive Coulomb field of the N-terminal proton or an involvement of the proton itself might have an influence on the yield of Y_{sc} upon direct ionization. A positive charge created by a direct ionization of the $\text{C}\alpha$ atom of the Tyr residue is in immediate proximity to the N-terminus. At this distance Coulomb repulsion might come into play.

At the oxygen K-edge the normalized yields of F_{sc} and M_{sc} are very similar (Figure 6h). In comparison, the normalized

yield of Y_{sc} is higher upon $\text{O } 1s \rightarrow \pi^*(\text{CONH})$ excitation and direct ionization but lower in the σ^* regime. When normalizing the spectra to the maximum of the $\pi^*(\text{CONH})$ feature the only strong difference between the ion yields is a significantly lower yield for Y_{sc} in the σ^* region. Since in this energy region the yields of the side-chain daughter fragments (C_6H_7^+ and C_6H_5^+) are the highest, it could be possible that the yield of Y_{sc} is decreased because excitations from the tyrosine hydroxy oxygen atom to σ^* orbitals lead to fragmentation within the Tyr side chain.

The normalized ion yield spectra of F_{sc} , Y_{sc} , and M_{sc} measured at the nitrogen K-edge are again very similar (Figure 6e). Only the normalized yield of Y_{sc} is slightly higher at the ionization edge. None of the amino acid side chains in MetEnk contain nitrogen. Thus, fragmentation at the nitrogen K-edge takes place exclusively after excitations in the peptide backbone and the production of side-chain fragments should be driven mainly by IVR.

Evidence for Sulfur–Aromatic Interactions. In order to identify the cause for the increased ion yield upon $\text{C } 1s \rightarrow \pi^*(\text{C}=\text{C})$ excitation of $[\text{MetEnk}+\text{H}]^+$ compared to $[\text{LeuEnk}+\text{H}]^+$ the ratios between the normalized areas of peak CK1 determined for both peptides were calculated for different fragments and compared to each other (Figure 7a). The peak areas of fragments M_{sc} and M_{im} were compared to those of L_{sc} and L_{im} .

Figure 7. (a) Ratios between normalized partial ion yields of different fragments of $[\text{MetEnk}+\text{H}]^+$ and $[\text{LeuEnk}+\text{H}]^+$ measured upon $\text{C } 1s \rightarrow \pi^*(\text{C}=\text{C})$ excitation. (b) Normalized partial ion yields of different fragments of $[\text{MetEnk}+\text{H}]^+$ measured upon $\text{N } 1s \rightarrow \pi^*(\text{C}=\text{N})$ excitation.

The only area ratio smaller than one can be observed between M_{im} and L_{im} . Most fragments show area ratios between 1.02 and 1.15. With values of (1.26 ± 0.03) and (1.58 ± 0.11) the area ratios of F_{sc} and between M_{sc} and L_{sc} are significantly higher. The area ratios for peak CK3 ($\text{C } 1s \rightarrow \pi^*(\text{CONH})$) of these fragments are (1.03 ± 0.03) and (1.03 ± 0.07) . This shows that the high area ratios for CK1 are not caused by a biased normalization.

The increased relative yield of M_{sc} (compared to I_{sc}) upon $C\ 1s \rightarrow \pi^*(C=C)$ excitation is surprising because the Met residue does not contain an aromatic carbon ring. Since the Met and Phe residue are direct neighbors, the reason for the increased yields of M_{sc} and F_{sc} might be a direct intramolecular charge or energy transfer between the sulfur atom and the aromatic π -system of the Phe side chain. Another indication for a direct interaction between these side chains instead of a charge migration or energy transfer through the peptide backbone is the fact that no increased yields were observed for F_{im} and M_{im} . Sulfur–aromatic interactions play a unique role in structure stabilizing binding motifs in proteins and were focus of numerous computational and crystallographic studies.^{102–107} To the best of our knowledge, effects of sulfur–aromatic interactions in peptides have so far never been observed in X-ray absorption spectroscopy experiments.

Discussion of the 399.2 eV Feature of [MetEnk+H]⁺

Features around the energy position of NK4 (see Figure 4c) are characteristic for $N\ 1s \rightarrow \pi^*(C=N)$ transitions, as reported for example for plasma-polymerized propylamine (399.1 eV¹⁰⁸) and allylamine (399.1 eV,¹⁰⁸ 398.7 eV¹⁰⁹) as well as plasma-nitrogenated polyolefin surfaces (398.9 eV).¹¹⁰ The common chemical structure of protonated MetEnk (Figure 1b), where all peptide bonds are in keto form ($-C(=O)NH-$; also known as amide form), does not contain any $C=N$ groups. However, another tautomer of the peptide bond is the enol form (also known as imidic acid form), where the hydrogen atom is bound to the oxygen atom ($-C(OH)=N-$). The energetically more favorable tautomer is the keto form.^{111,112} An X-ray absorption spectroscopic signature of the keto-to-enol conversion of peptide bonds has been previously observed in the N K-edge XPS spectrum of the tripeptide glypromate isolated in an argon matrix (399.2 eV⁸⁹) and in the N K-edge XAS spectra of S-Layer protein substrates interacting with metal atoms (398.4 eV¹¹³). In both cases, the enol form was not the natural tautomer and was found to be produced either by UV irradiation or by protein–metal interactions, respectively.

To evaluate the effects of a tautomeric keto-to-enol conversion of a peptide bond on the theoretical nitrogen K-edge absorption spectrum of [MetEnk+H]⁺ we performed DFT/ROCIS calculations for N-terminal protonated MetEnk with its Gly-Gly bond in enol-tautomer form (Figure 1c). The Gly-Gly bond was chosen for this structural change because a variety of computational studies have been performed on the keto–enol-tautomerism of peptide bonds in polyglycine peptides.^{114–116} The calculated spectrum shows a very intense peak at 398.8 eV (Figure 8c) originating exclusively from $N\ 1s \rightarrow \pi^*(C=N)$ transitions in the altered peptide bond. The ratio between the maximum intensities of this peak and the $\pi^*(CONH)$ feature is about 0.8:1. This is in harsh contrast to the experimental data where this ratio is about 0.17:1. Consequently, assuming that peak NK4 is in fact due to transitions into MOs of a peptide bond in enol form, the equilibrium between the two tautomers still lies heavily to the keto form.

It could be possible that enol tautomers have already been present in the MetEnk sample solution. The solvent used was a 1:1 mixture of water and methanol and the pH value was 2.7. Calculations have shown that both water and methanol drastically reduce the potential barrier between the peptide bond's enol and keto form.¹¹⁴ Furthermore, the enol form is a known intermediate in the acid-catalyzed proton exchange in

Figure 8. DFT/ROCIS-calculated X-ray absorption spectra for (a) N-terminal protonated MetEnk with all peptide bonds in keto form (shifted +12.6 eV), (b) neutral MetEnk with all peptide bonds in keto form (shifted +12.3 eV), and (c) N-terminal protonated MetEnk with the Gly-Gly peptide bond in enol form (shifted +12.2 eV). The calculated discrete absorption lines were broadened using a pseudo-Voigt profile with 0.7 eV fwhm.

peptide bonds.^{117–119} However, based on this explanation the enol tautomer of [LeuEnk+H]⁺ should be expected as well, because the compositions of all sample solutions were identical. It remains unclear why the structural change leading to peak NK4 did not take place in [LeuEnk+H]⁺ but since the peptides were measured using different ion funnels it might have been induced by collisions of precursor ions with residual gas molecules in the ion-funnel region where the pressure is still in the order of millibars.

Another possible reason for peak NK4 are $N\ 1s \rightarrow \sigma^*(N-H)$ transitions in an unprotonated amino group. The peak positions of the protonated and unprotonated amino groups in the XPS spectra of different zwitterionic amino acids reported by Zubavichus et al.⁹⁰ are (401.4 ± 0.2) eV and (399.7 ± 0.2) eV. When the N-terminus is unprotonated the proton must, consequently, sit at another site of the peptide. While it has been shown that the most favorable protonation site of [LeuEnk+H]⁺ is the N-terminus,⁴¹ the protonation site of [MetEnk+H]⁺ might be influenced by the proton affinity of the Met residue, which is slightly higher than for the Leu residue.¹²⁰ Experimental evidence has been reported for backbone protonation in a doubly protonated synthetic peptide¹²¹ and in singly protonated triglycine for which it was demonstrated that protonation occurs at an amide oxygen atom.^{122,123}

We performed DFT/ROCIS calculations for unprotonated MetEnk in all-keto form (Figure 1b) in order to evaluate the effect of the unprotonated amino group on the theoretical nitrogen K-edge absorption spectrum. The calculated spectrum features a peak at 399.7 eV (Figure 8b) attributed to $N\ 1s \rightarrow \sigma^*(N-H)$ transitions in the terminal NH_2 group. It should be noted that Zubavichus et al.¹²⁴ observed a peak at ~400 eV appearing in the X-ray photoelectron spectra and peaks at 398.5 and 399.7 eV emerging in the X-ray fluorescence spectra of solid zwitterionic Tyr and Phe after partial decomposition of the samples using soft X-ray radiation. While they considered the formation of $C=N$ bonds as well, ultimately, the authors assigned these features to transitions in unprotonated amino groups.

In order to locate the site of the electronic transition behind peak NK4 the areas of this peak in the normalized partial

NEXAMS spectra of different fragments were compared (Figure 7b). With (0.032 ± 0.002) , Y_{im} exhibits the largest peak area for this feature. The second highest area (0.021 ± 0.002) was observed for Y_{sc} . The peak areas of F_{sc} and the fragments at m/z 131 and 147 are at a comparable level at around 0.019. Since F_{sc} can also occur as a fragment of the Tyr side chain and the fragments at m/z 131 and 147 can be assigned to fragments $[c_2 - Y_{sc}]^+$ and Y_{acr} , all fragments with an increased area of peak NK4 can be associated with the Tyr residue. This indicates that the transitions leading to peak NK4 take place near the N-terminus. Assuming that there is one peptide bond in enol form it would, therefore, most likely be the Tyr-Gly bond. However, since Tyr is the most N-terminal amino acid residue in MetEnk N 1s $\rightarrow \sigma^*(N-H)$ transitions in an unprotonated amino group remain a plausible reason for peak NK4 as well.

Based on our DFT/ROCIS calculations alone, both presented types of transitions are possible reasons for peak NK4. Since the experimental energy position of this spectral feature is in slightly better agreement with the values for N 1s $\rightarrow \pi^*(C=N)$ transitions reported in literature we consider this assignment as more likely. Nevertheless, further experiments are needed to undoubtedly assess the origin of this feature.

CONCLUSIONS

We have conducted a NEXAMS study on cryogenically cooled gas-phase $[LeuEnk+H]^+$ and $[MetEnk+H]^+$ by measuring their cationic fragments produced upon soft X-ray photon absorption at the carbon, nitrogen and oxygen K-edges. Our results show differences between the total and partial NEXAMS spectra of both peptides which could be interpreted in relation to aspects of their structure, including amino acid side-chain interactions and the tautomerism of peptide bonds.

A spectral feature at 399.2 eV was observed only in the nitrogen K-edge NEXAMS spectra of $[MetEnk+H]^+$. Increased areas of this peak in the partial NEXAMS spectra of fragments related to tyrosine suggest that the electronic transitions behind this peak takes place near the N-terminal Tyr residue. We discussed both N 1s $\rightarrow \pi^*(C=N)$ transitions in the enol form of the Tyr-Gly peptide bond and N 1s $\rightarrow \sigma^*(N-H)$ transitions in unprotonated amino groups as plausible reasons for this feature. The DFT/ROCIS-calculated nitrogen K-edge absorption spectra for different MetEnk isomers gave evidence for both electronic transitions but the better agreement with peaks reported for this energy region in literature suggests that N 1s $\rightarrow \pi^*(C=N)$ transitions are the more probable reason. For a final assignment measurements on individual isomers of gas-phase $[MetEnk+H]^+$ are necessary. Especially C=N bonds could serve as dedicated excitation sites in studies on local effects of site-specific photoexcitation. In this context, also an investigation on the dynamics of the UV photon-induced tautomeric keto-to-enol conversion of peptide bonds, demonstrated by Mateo-Marti and Pradier,⁸⁹ in an UV pump-soft X-ray probe experiment is of high interest.

Furthermore, the total measured ion yield for $[MetEnk+H]^+$ upon C 1s $\rightarrow \pi^*(C=C)$ excitation was 13.7% higher than for $[LeuEnk+H]^+$. Among all fragments this effect was the most prominent for the complete Phe and Met side-chain fragments. Therefore, we propose an involvement of a direct sulfur-aromatic interaction between the Phe and Met side chains, promoting the formation of the respective side-chain fragments. Investigations on custom peptides with different pairs

of neighboring sulfur-containing and aromatic amino acid residues will be conducted to gain more information about the effects of sulfur-aromatic interactions on the soft X-ray photon-induced dissociation of peptides.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jasms.0c00390>.

DFT-optimized geometries, isosurfaces of DFT-calculated LUMOs, intensities and areas of fitted spectral features, and assignments and structural formulas of detected fragments (PDF)

AUTHOR INFORMATION

Corresponding Authors

Simon Dörner – Deutsches Elektronen-Synchrotron DESY, 22607 Hamburg, Germany; orcid.org/0000-0001-9628-0147; Email: simon.doerner@desy.de

Lucas Schwob – Deutsches Elektronen-Synchrotron DESY, 22607 Hamburg, Germany; orcid.org/0000-0002-4274-365X; Email: lucas.schwob@desy.de

Sadia Bari – Deutsches Elektronen-Synchrotron DESY, 22607 Hamburg, Germany; Email: sadia.bari@desy.de

Authors

Kaan Atak – Deutsches Elektronen-Synchrotron DESY, 22607 Hamburg, Germany

Kaja Schubert – Deutsches Elektronen-Synchrotron DESY, 22607 Hamburg, Germany

Rebecca Boll – Deutsches Elektronen-Synchrotron DESY, 22607 Hamburg, Germany; European XFEL, 22869 Schenefeld, Germany

Thomas Schlathöler – Zernike Institute for Advanced Materials, University of Groningen, 9747AG Groningen, The Netherlands; orcid.org/0000-0001-9923-172X

Martin Timm – Abteilung Hochempfindliche Röntgenspektroskopie, Helmholtz-Zentrum Berlin für Materialien und Energie, 12489 Berlin, Germany

Christine Bülow – Abteilung Hochempfindliche Röntgenspektroskopie, Helmholtz-Zentrum Berlin für Materialien und Energie, 12489 Berlin, Germany

Vicente Zamudio-Bayer – Abteilung Hochempfindliche Röntgenspektroskopie, Helmholtz-Zentrum Berlin für Materialien und Energie, 12489 Berlin, Germany

Bernd von Issendorff – Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, 79104 Freiburg, Germany

J. Tobias Lau – Abteilung Hochempfindliche Röntgenspektroskopie, Helmholtz-Zentrum Berlin für Materialien und Energie, 12489 Berlin, Germany; Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, 79104 Freiburg, Germany

Simone Techert – Deutsches Elektronen-Synchrotron DESY, 22607 Hamburg, Germany; Institut für Röntgenphysik, Georg-August-Universität Göttingen, 37077 Göttingen, Germany; orcid.org/0000-0002-7549-7651

Complete contact information is available at: <https://pubs.acs.org/doi/10.1021/jasms.0c00390>

Author Contributions

S.D. and L.S. contributed equally to this work.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank the Helmholtz-Zentrum Berlin for the provision of synchrotron radiation at the UE52-PGM beamline. S.D., L.S., K.S., and S.B. acknowledge funding from the Helmholtz Initiative and Networking Fund through the Young Investigators Group Program (VH-NG-1104). Furthermore, K.S., S.T., and S.B. were supported by the Deutsche Forschungsgemeinschaft, project B03 in the SFB 755-Nanoscale Photonic Imaging. The NanoclusterTrap endstation at UE52_PGM is partially funded by the BMBF grant 05K16VF1.

■ REFERENCES

- (1) Gordon, M. L.; Cooper, G.; Morin, C.; Araki, T.; Turci, C. C.; Kaznatcheev, K.; Hitchcock, A. P. Inner-shell excitation spectroscopy of the peptide bond: Comparison of the C 1s, N 1s, and O 1s spectra of glycine, glycyl-glycine, and glycyl-glycyl-glycine. *J. Phys. Chem. A* **2003**, *107*, 6144–6159.
- (2) Stewart-Ornstein, J.; Hitchcock, A. P.; Hernández Cruz, D.; Henklein, P.; Overhage, J.; Hilpert, K.; Hale, J. D.; Hancock, R. E. W. Using Intrinsic X-ray Absorption Spectral Differences To Identify and Map Peptides and Proteins. *J. Phys. Chem. B* **2007**, *111*, 7691–7699.
- (3) Zubavichus, Y.; Shaporenko, A.; Grunze, M.; Zharnikov, M. Inner-shell absorption spectroscopy of amino acids at all relevant absorption edges. *J. Phys. Chem. A* **2005**, *109*, 6998–7000.
- (4) Zubavichus, Y.; Shaporenko, A.; Grunze, M.; Zharnikov, M. NEXAFS spectroscopy of homopolypeptides at all relevant absorption edges: Polyisoleucine, polytyrosine, and polyhistidine. *J. Phys. Chem. B* **2007**, *111*, 9803–9807.
- (5) Zubavichus, Y.; Shaporenko, A.; Grunze, M.; Zharnikov, M. Is X-ray absorption spectroscopy sensitive to the amino acid composition of functional proteins? *J. Phys. Chem. B* **2008**, *112*, 4478–4480.
- (6) Feyer, V.; Plekan, O.; Richter, R.; Coreno, M.; Prince, K. C.; Carravetta, V. Photoemission and Photoabsorption Spectroscopy of Glycyl-Glycine in the Gas Phase. *J. Phys. Chem. A* **2009**, *113*, 10726–10733.
- (7) Zubavichus, Y.; Shaporenko, A.; Grunze, M.; Zharnikov, M. NEXAFS spectroscopy of biological molecules: From amino acids to functional proteins. *Nucl. Instrum. Methods Phys. Res., Sect. A* **2009**, *603*, 111–114.
- (8) González-Magaña, O.; Reitsma, G.; Tiemens, M.; Boschman, L.; Hoekstra, R.; Schlathöller, T. Near-edge X-ray absorption mass spectrometry of a gas-phase peptide. *J. Phys. Chem. A* **2012**, *116*, 10745–10751.
- (9) Milosavljević, A.; Canon, F.; Nicolas, C.; Miron, C.; Nahon, L.; Giuliani, A. Gas-phase protein inner-shell spectroscopy by coupling an ion trap with a soft X-ray beamline. *J. Phys. Chem. Lett.* **2012**, *3*, 1191–1196.
- (10) Egorov, D.; Bari, S.; Boll, R.; Dörner, S.; Deinert, S.; Techert, S.; Hoekstra, R.; Zamudio-Bayer, V.; Lindblad, R.; Bülow, C.; Timm, M.; von Issendorff, B.; Lau, J. T.; Schlathöller, T. Near-Edge Soft X-ray Absorption Mass Spectrometry of Protonated Melittin. *J. Am. Soc. Mass Spectrom.* **2018**, *29*, 2138–2151.
- (11) Bari, S.; Egorov, D.; Jansen, T. L.; Boll, R.; Hoekstra, R.; Techert, S.; Zamudio-Bayer, V.; Bülow, C.; Lindblad, R.; Leistner, G.; Ławicki, A.; Hirsch, K.; Miedema, P. S.; von Issendorff, B.; Lau, J. T.; Schlathöller, T. Soft X-ray Spectroscopy as a Probe for Gas-Phase Protein Structure: Electron Impact Ionization from Within. *Chem. - Eur. J.* **2018**, *24*, 7631–7636.
- (12) Fujii, K.; Akamatsu, K.; Muramatsu, Y.; Yokoya, A. X-ray absorption near edge structure of DNA bases around oxygen and nitrogen K-edge. *Nucl. Instrum. Methods Phys. Res., Sect. B* **2003**, *199*, 249–254.
- (13) Samuel, N. T.; Lee, C.-Y.; Gamble, L. J.; Fischer, D. A.; Castner, D. G. NEXAFS characterization of DNA components and molecular-orientation of surface-bound DNA oligomers. *J. Electron Spectrosc. Relat. Phenom.* **2006**, *152*, 134–142.
- (14) Lee, C.-Y.; Gong, P.; Harbers, G. M.; Grainger, D. W.; Castner, D. G.; Gamble, L. J. Surface Coverage and Structure of Mixed DNA/Alkylthiol Monolayers on Gold: Characterization by XPS, NEXAFS, and Fluorescence Intensity Measurements. *Anal. Chem.* **2006**, *78*, 3316–3325.
- (15) Petrovykh, D. Y.; Pérez-Dieste, V.; Opdahl, A.; Kimura-Suda, H.; Sullivan, J. M.; Tarlov, M. J.; Himpsel, F. J.; Whitman, L. J. Nucleobase Orientation and Ordering in Films of Single-Stranded DNA on Gold. *J. Am. Chem. Soc.* **2006**, *128*, 2–3.
- (16) Plekan, O.; Feyer, V.; Richter, R.; Coreno, M.; de Simone, M.; Prince, K.; Trofimov, A.; Gromov, E.; Zaytseva, I.; Schirmer, J. A theoretical and experimental study of the near edge X-ray absorption fine structure (NEXAFS) and X-ray photoelectron spectra (XPS) of nucleobases: Thymine and adenine. *Chem. Phys.* **2008**, *347*, 360–375.
- (17) Hua, W.; Gao, B.; Li, S.; Ågren, H.; Luo, Y. Refinement of DNA Structures through Near-Edge X-ray Absorption Fine Structure Analysis: Applications on Guanine and Cytosine Nucleobases, Nucleosides, and Nucleotides. *J. Phys. Chem. B* **2010**, *114*, 13214–13222.
- (18) Howell, C.; Hamoudi, H.; Zharnikov, M. Thymine/adenine diblock-oligonucleotide monolayers and hybrid brushes on gold: a spectroscopic study. *Biointerphases* **2013**, *8*, 6.
- (19) Lin, Y.-S.; Lin, H.-R.; Liu, W.-L.; Lee, Y. T.; Tseng, C.-M.; Ni, C.-K.; Liu, C.-L.; Tsai, C.-C.; Chen, J.-L.; Hu, W.-P. Measurement and prediction of the NEXAFS spectra of pyrimidine and purine and the dissociation following the core excitation. *Chem. Phys. Lett.* **2015**, *636*, 146–153.
- (20) Tsud, N.; Bercha, S.; Ševčíková, K.; Acres, R. G.; Prince, K. C.; Matolin, V. Adenine adlayers on Cu(111): XPS and NEXAFS study. *J. Chem. Phys.* **2015**, *143*, 174704.
- (21) Kaznatcheyev, K.; Osanna, A.; Jacobsen, C.; Plashkevych, O.; Vahtras, O.; Ågren, H.; Carravetta, V.; Hitchcock, A. P. Inner-shell absorption spectroscopy of amino acids. *J. Phys. Chem. A* **2002**, *106*, 3153–3168.
- (22) Boese, J.; Osanna, A.; Jacobsen, C.; Kirz, J. Carbon edge XANES spectroscopy of amino acids and peptides. *J. Electron Spectrosc. Relat. Phenom.* **1997**, *85*, 9–15.
- (23) Schlathöller, T.; Hoekstra, R. In *Photophysics of Ionic Biochromophores*; Brøndsted Nielsen, S., Wyer, J. A., Eds.; Physical Chemistry in Action; Springer Berlin Heidelberg: Berlin, Heidelberg, 2013; Chapter 11, pp 209–226.
- (24) Reitsma, G.; Boschman, L.; Deuzeman, M. J.; Hoekstra, S.; Hoekstra, R.; Schlathöller, T. Near edge X-ray absorption mass spectrometry on coronene. *J. Chem. Phys.* **2015**, *142*, 024308.
- (25) Schwob, L.; Dörner, S.; Atak, K.; Schubert, K.; Timm, M.; Bülow, C.; Zamudio-Bayer, V.; von Issendorff, B.; Lau, J. T.; Techert, S.; Bari, S. Site-Selective Dissociation upon Sulfur L-Edge X-ray Absorption in a Gas-Phase Protonated Peptide. *J. Phys. Chem. Lett.* **2020**, *11*, 1215–1221.
- (26) Kopysov, V.; Makarov, A.; Boyarkin, O. V. Colors for Molecular Masses: Fusion of Spectroscopy and Mass Spectrometry for Identification of Biomolecules. *Anal. Chem.* **2015**, *87*, 4607–4611.
- (27) Saparbaev, E.; Kopysov, V.; Yamaletdinov, R.; Pereverzev, A. Y.; Boyarkin, O. V. Interplay of H-Bonds with Aromatics in Isolated Complexes Identifies Isomeric Carbohydrates. *Angew. Chem., Int. Ed.* **2019**, *58*, 7346–7350.
- (28) Giuliani, A.; Milosavljević, A.; Canon, F.; Nahon, L. Contribution of Synchrotron Radiation to Photoactivation Studies of Biomolecular Ions in the Gas Phase. *Mass Spectrom. Rev.* **2014**, *33*, 424–441.
- (29) Abdelmouleh, M.; Lalande, M.; El Feghaly, J.; Vizcaino, V.; Rebelo, A.; Eden, S.; Schlathöller, T.; Pouilly, J.-C. Mass Spectral Signatures of Complex Post-Translational Modifications in Proteins: A Proof-of-Principle Based on X-ray Irradiated Vancomycin. *J. Am. Soc. Mass Spectrom.* **2020**, *31*, 1738–1743.
- (30) Egorov, D.; Schwob, L.; Lalande, M.; Hoekstra, R.; Schlathöller, T. Near edge X-ray absorption mass spectrometry of

gas phase proteins: The influence of protein size. *Phys. Chem. Chem. Phys.* **2016**, *18*, 26213–26223.

(31) Schwob, L.; Lalande, M.; Egorov, D.; Rangama, J.; Hoekstra, R.; Vizcaino, V.; Schlathölter, T.; Pouilly, J.-C. Radical-driven processes within a peptidic sequence of type I collagen upon single-photon ionisation in the gas phase. *Phys. Chem. Chem. Phys.* **2017**, *19*, 22895–22904.

(32) Milosavljević, A. R.; Nicolas, C.; Ranković, M. L.; Canon, F.; Miron, C.; Giuliani, A. K-Shell Excitation and Ionization of a Gas-Phase Protein: Interplay between Electronic Structure and Protein Folding. *J. Phys. Chem. Lett.* **2015**, *6*, 3132–3138.

(33) Alexander, A.; Boyd, R. Experimental investigations of factors controlling the collision induced dissociation spectra of peptide ions in a tandem hybrid mass spectrometer. I. Leucine enkephalin. *Int. J. Mass Spectrom. Ion Processes* **1989**, *90*, 211–240.

(34) Vachet, R. W.; Bishop, B. M.; Erickson, B. W.; Glish, G. L. Novel peptide dissociation: Gas-phase intramolecular rearrangement of internal amino acid residues. *J. Am. Chem. Soc.* **1997**, *119*, 5481–5488.

(35) Rakov, V. S.; Borisov, O. V.; Whitehouse, C. M. Establishing low-energy sequential decomposition pathways of leucine enkephalin and its N- and C-terminus fragments using multiple-resonance CID in quadrupolar ion guide. *J. Am. Soc. Mass Spectrom.* **2004**, *15*, 1794–1809.

(36) Laskin, J. Energetics and dynamics of fragmentation of protonated leucine enkephalin from time- and energy-resolved surface-induced dissociation studies. *J. Phys. Chem. A* **2006**, *110*, 8554–8562.

(37) Li, L.; Lubman, D. M. Resonant Two-Photon Ionization of Enkephalins and Related Peptides Volatilized by Using Pulsed Laser Desorption in Supersonic Beam Mass Spectrometry. *Anal. Chem.* **1988**, *60*, 1409–1415.

(38) Tabarin, T.; Antoine, R.; Broyer, M.; Dugourd, P. Specific photodissociation of peptides with multi-stage mass spectrometry. *Rapid Commun. Mass Spectrom.* **2005**, *19*, 2883–2892.

(39) Herburger, A.; van der Linde, C.; Beyer, M. K. Photodissociation spectroscopy of protonated leucine enkephalin. *Phys. Chem. Chem. Phys.* **2017**, *19*, 10786–10795.

(40) Burke, N. L.; Redwine, J. G.; Dean, J. C.; Mcluckey, S. A.; Zwier, T. S. UV and IR spectroscopy of cold protonated leucine enkephalin. *Int. J. Mass Spectrom.* **2015**, *378*, 196–205.

(41) Polfer, N. C.; Oomens, J.; Suhai, S.; Paizs, B. Infrared spectroscopy and theoretical studies on gas-phase protonated Leu-enkephalin and its fragments: Direct experimental evidence for the mobile proton. *J. Am. Chem. Soc.* **2007**, *129*, 5887–5897.

(42) Ranković, M. L.; Canon, F.; Nahon, L.; Giuliani, A.; Milosavljević, A. R. VUV action spectroscopy of protonated leucine-enkephalin peptide in the 6–14 eV range. *J. Chem. Phys.* **2015**, *143*, 244311.

(43) Bari, S.; González-Magaña, O.; Reitsma, G.; Werner, J.; Schippers, S.; Hoekstra, R.; Schlathölter, T. Photodissociation of protonated leucine-enkephalin in the VUV range of 8–40 eV. *J. Chem. Phys.* **2011**, *134*, 1–9.

(44) Bari, S.; Hoekstra, R.; Schlathölter, T. Peptide fragmentation by keV ion-induced dissociation. *Phys. Chem. Chem. Phys.* **2010**, *12*, 3376.

(45) Reitsma, G.; González-Magaña, O.; Versolato, O.; Door, M.; Hoekstra, R.; Suraud, E.; Fischer, B.; Camus, N.; Kremer, M.; Moshhammer, R.; Schlathölter, T. Femtosecond laser induced ionization and dissociation of gas-phase protonated leucine enkephalin. *Int. J. Mass Spectrom.* **2014**, *365–366*, 365–371.

(46) Niemeyer, M.; Hirsch, K.; Zamudio-Bayer, V.; Langenberg, A.; Vogel, M.; Kossick, M.; Ebrecht, C.; Egashira, K.; Terasaki, A.; Möller, T.; von Issendorff, B.; Lau, J. T. Spin Coupling and Orbital Angular Momentum Quenching in Free Iron Clusters. *Phys. Rev. Lett.* **2012**, *108*, 057201.

(47) Hirsch, K.; Lau, J. T.; Klar, P.; Langenberg, A.; Probst, J.; Rittmann, J.; Vogel, M.; Zamudio-Bayer, V.; Möller, T.; von Issendorff, B. X-ray spectroscopy on size-selected clusters in an ion

trap: from the molecular limit to bulk properties. *J. Phys. B: At., Mol. Opt. Phys.* **2009**, *42*, 154029.

(48) Egorov, D.; Bari, S.; Hoekstra, R.; Lawicki, A.; Hirsch, K.; Zamudio-Bayer, V.; Lau, J. T.; von Issendorff, B.; Schlathölter, T. An intense electrospray ionization source for soft X-ray photoionization of gas phase protein ions. *Journal of Physics: Conference Series* **2015**, *635*, 9–10.

(49) Carroll, T. X.; Hahne, J.; Thomas, T. D.; Sæthre, L. J.; Berrah, N.; Bozek, J.; Kukk, E. Carbon 1s core-hole lifetime in CO₂. *Phys. Rev. A: At., Mol., Opt. Phys.* **2000**, *61*, 042503.

(50) Kempgens, B.; Kivimäki, A.; Neeb, M.; Köppe, H. M.; Bradshaw, A. M.; Feldhaus, J. A high-resolution N 1s photoionization study of the N₂ molecule in the near-threshold region. *J. Phys. B: At., Mol. Opt. Phys.* **1996**, *29*, 5389–5402.

(51) Coreno, M.; de Simone, M.; Prince, K.; Richter, R.; Vondráček, M.; Avaldi, L.; Camilloni, R. Vibrationally resolved oxygen K → Π^* spectra of O₂ and CO. *Chem. Phys. Lett.* **1999**, *306*, 269–274.

(52) Neese, F. Software update: the ORCA program system, version 4.0. *WIREs Computational Molecular Science* **2018**, *8*, 4–9.

(53) Weigend, F.; Ahlrichs, R. Balanced basis sets of split valence, triple zeta valence and quadruple zeta valence quality for H to Rn: Design and assessment of accuracy. *Phys. Chem. Chem. Phys.* **2005**, *7*, 3297.

(54) Becke, A. D. Density-functional exchange-energy approximation with correct asymptotic behavior. *Phys. Rev. A: At., Mol., Opt. Phys.* **1988**, *38*, 3098–3100.

(55) Becke, A. D. Density-functional thermochemistry. III. The role of exact exchange. *J. Chem. Phys.* **1993**, *98*, 5648–5652.

(56) Roemelt, M.; Maganas, D.; DeBeer, S.; Neese, F. A combined DFT and restricted open-shell configuration interaction method including spin-orbit coupling: Application to transition metal L-edge X-ray absorption spectroscopy. *J. Chem. Phys.* **2013**, *138*, 204101.

(57) Baerends, E.; Ellis, D.; Ros, P. Self-consistent molecular Hartree-Fock-Slater calculations I. The computational procedure. *Chem. Phys.* **1973**, *2*, 41–51.

(58) Dunlap, B. I.; Connolly, J. W. D.; Sabin, J. R. On some approximations in applications of X α theory. *J. Chem. Phys.* **1979**, *71*, 3396–3402.

(59) Vahtras, O.; Almlöf, J.; Feyereisen, M. Integral approximations for LCAO-SCF calculations. *Chem. Phys. Lett.* **1993**, *213*, 514–518.

(60) Eichkorn, K.; Treutler, O.; Öhm, H.; Häser, M.; Ahlrichs, R. Auxiliary basis sets to approximate Coulomb potentials. *Chem. Phys. Lett.* **1995**, *240*, 283–290.

(61) Eichkorn, K.; Weigend, F.; Treutler, O.; Ahlrichs, R. Auxiliary basis sets for main row atoms and transition metals and their use to approximate Coulomb potentials. *Theor. Chem. Acc.* **1997**, *97*, 119–124.

(62) Stoychev, G. L.; Auer, A. A.; Neese, F. Automatic Generation of Auxiliary Basis Sets. *J. Chem. Theory Comput.* **2017**, *13*, 554–562.

(63) van Lenthe, E.; Snijders, J. G.; Baerends, E. J. The zero-order regular approximation for relativistic effects: The effect of spin-orbit coupling in closed shell molecules. *J. Chem. Phys.* **1996**, *105*, 6505–6516.

(64) Wertheim, G. K.; Butler, M. A.; West, K. W.; Buchanan, D. N. E. Determination of the Gaussian and Lorentzian content of experimental line shapes. *Rev. Sci. Instrum.* **1974**, *45*, 1369–1371.

(65) Young, R. A.; Wiles, D. B. Profile shape functions in Rietveld refinements. *J. Appl. Crystallogr.* **1982**, *15*, 430–438.

(66) Ida, T.; Ando, M.; Toraya, H. Extended pseudo-Voigt function for approximating the Voigt profile. *J. Appl. Crystallogr.* **2000**, *33*, 1311–1316.

(67) Thompson, P.; Cox, D. E.; Hastings, J. B. Rietveld Refinement of Debye-Scherrer Synchrotron X-ray Data from Al₂O₃. *J. Appl. Crystallogr.* **1987**, *20*, 79–83.

(68) Falick, A. M.; Hines, W. M.; Medzihradsky, K. F.; Baldwin, M. A.; Gibson, B. W. Low-mass ions produced from peptides by high-energy collision-induced dissociation in tandem mass spectrometry. *J. Am. Soc. Mass Spectrom.* **1993**, *4*, 882–893.

- (69) Papayannopoulos, I. A. The interpretation of collision-induced dissociation tandem mass spectra of peptides. *Mass Spectrom. Rev.* **1995**, *14*, 49–73.
- (70) Roepstorff, P.; Fohlman, J. Letter to the editors. *Biological Mass Spectrometry* **1984**, *11*, 600–601.
- (71) Shoeib, T.; Cunje, A.; Hopkinson, A. C.; Siu, K. W. M. Gas-phase fragmentation of the Ag⁺-phenylalanine complex: Cation- π interactions and radical cation formation. *J. Am. Soc. Mass Spectrom.* **2002**, *13*, 408–416.
- (72) Soorkia, S.; Broquier, M.; Grégoire, G. Conformer- and mode-specific excited state lifetimes of cold protonated tyrosine ions. *J. Phys. Chem. Lett.* **2014**, *5*, 4349–4355.
- (73) Zhao, J.; Shoeib, T.; Siu, K. M.; Hopkinson, A. C. The fragmentation of protonated tyrosine and iodotyrosines: The effect of substituents on the losses of NH₃ and of H₂O and CO. *Int. J. Mass Spectrom.* **2006**, *255–256*, 265–278.
- (74) Von E. Doering, W.; Knox, L. H. The Cycloheptatrienylum (Tropylium) Ion. *J. Am. Chem. Soc.* **1954**, *76*, 3203–3206.
- (75) Rylander, P. N.; Meyerson, S.; Grubb, H. M. Organic Ions in the Gas Phase. II. The Tropylium Ion. *J. Am. Chem. Soc.* **1957**, *79*, 842–846.
- (76) Ausloos, P. Structure and isomerization of C₇H₇⁺ ions formed in the charge-transfer-induced fragmentation of ethylbenzene, toluene, and norbornadiene. *J. Am. Chem. Soc.* **1982**, *104*, 5259–5265.
- (77) Zhao, G.; Gäumann, T. Isomeric [C₇H₈]⁺ and [C₇H₇]⁺ ions. *Org. Mass Spectrom.* **1992**, *27*, 428–434.
- (78) NIST Standard Reference Database Number 69, 2018; <https://webbook.nist.gov/chemistry/>.
- (79) Russell, D. H.; Freiser, B. S.; McBay, E. H.; Canada, D. C. The structure of decomposing [C₇H₇O]⁺ ions: Benzyl versus tropylium ion structures. *Org. Mass Spectrom.* **1983**, *18*, 474–485.
- (80) Bauerle, G. F.; Brodbelt, J. S. Evaluation of steric and substituent effects in phenols by competitive reactions of dimethyl ether ions in a quadrupole ion trap. *J. Am. Soc. Mass Spectrom.* **1995**, *6*, 627–633.
- (81) Plekan, O.; Feyer, V.; Richter, R.; Coreno, M.; Prince, K. C. Valence photoionization and photofragmentation of aromatic amino acids. *Mol. Phys.* **2008**, *106*, 1143–1153.
- (82) Budzikiewicz, H.; Drabner, G.; Hammes, C. Rearrangements accompanying the fragmentation of ionized 1-phenylalkan-1-ols. *Org. Mass Spectrom.* **1993**, *28*, 1326–1328.
- (83) Horsley, J. A.; Stöhr, J.; Hitchcock, A. P.; Newbury, D. C.; Johnson, A. L.; Sette, F. Resonances in the K shell excitation spectra of benzene and pyridine: Gas phase, solid, and chemisorbed states. *J. Chem. Phys.* **1985**, *83*, 6099–6107.
- (84) Solomon, J.; Madix, R.; Stöhr, J. Orientation and absolute coverage of benzene, aniline, and phenol on Ag(110) determined by NEXAFS and XPS. *Surf. Sci.* **1991**, *255*, 12–30.
- (85) Petoral, R. M.; Uvdal, K. XPS and NEXAFS study of tyrosine-terminated propanethiol assembled on gold. *J. Electron Spectrosc. Relat. Phenom.* **2003**, *128*, 159–164.
- (86) Zhang, W.; Carravetta, V.; Plekan, O.; Feyer, V.; Richter, R.; Coreno, M.; Prince, K. C. Electronic structure of aromatic amino acids studied by soft x-ray spectroscopy. *J. Chem. Phys.* **2009**, *131*, 035103.
- (87) Feyer, V.; Plekan, O.; Richter, R.; Coreno, M.; Prince, K. C.; Carravetta, V. Core Level Study of Alanine and Threonine. *J. Phys. Chem. A* **2008**, *112*, 7806–7815.
- (88) Hennig, C.; Hallmeier, K.; Szargan, R. XANES investigation of chemical states of nitrogen in polyaniline. *Synth. Met.* **1998**, *92*, 161–166.
- (89) Mateo-Marti, E.; Pradier, C. UV irradiation study of a tripeptide isolated in an argon matrix: A tautomerism process evidenced by infrared and X-ray photoemission spectroscopies. *Spectrochim. Acta, Part A* **2013**, *109*, 247–252.
- (90) Zubavichus, Y.; Fuchs, O.; Weinhardt, L.; Heske, C.; Umbach, E.; Denlinger, J. D.; Grunze, M. Soft X-Ray-Induced Decomposition of Amino Acids: An XPS, Mass Spectrometry, and NEXAFS Study. *Radiat. Res.* **2004**, *161*, 346–358.
- (91) Milosavljević, A. R.; Jänkälä, K.; Ranković, M. L.; Canon, F.; Bozek, J.; Nicolas, C.; Giuliani, A. Oxygen K-shell spectroscopy of isolated progressively solvated peptide. *Phys. Chem. Chem. Phys.* **2020**, *22*, 12909–12917.
- (92) Prince, K. C.; Richter, R.; de Simone, M.; Alagia, M.; Coreno, M. Near Edge X-ray Absorption Spectra of Some Small Polyatomic Molecules. *J. Phys. Chem. A* **2003**, *107*, 1955–1963.
- (93) Francis, J. T.; Hitchcock, A. P. Inner-shell spectroscopy of p-benzoquinone, hydroquinone, and phenol: distinguishing quinoid and benzenoid structures. *J. Phys. Chem.* **1992**, *96*, 6598–6610.
- (94) Petoral, R. M., Jr.; Uvdal, K. NEXAFS Study of Amino Acid Analogues Assembled on Gold. *Phys. Scr.* **2005**, *T115*, 851.
- (95) Salén, P.; Kamińska, M.; Squibb, R. J.; Richter, R.; Alagia, M.; Stranges, S.; Van Der Meulen, P.; Eland, J. H.; Feifel, R.; Zhaunerchyk, V. Selectivity in fragmentation of N-methylacetamide after resonant K-shell excitation. *Phys. Chem. Chem. Phys.* **2014**, *16*, 15231–15240.
- (96) Li, C.; Salén, P.; Yatsyna, V.; Schio, L.; Feifel, R.; Squibb, R.; Kamińska, M.; Larsson, M.; Richter, R.; Alagia, M.; Stranges, S.; Monti, S.; Carravetta, V.; Zhaunerchyk, V. Experimental and theoretical XPS and NEXAFS studies of N-methylacetamide and N-methyltrifluoroacetamide. *Phys. Chem. Chem. Phys.* **2016**, *18*, 2210–2218.
- (97) Plekan, O.; Feyer, V.; Richter, R.; Coreno, M.; de Simone, M.; Prince, K. C.; Carravetta, V. An X-ray absorption study of glycine, methionine and proline. *J. Electron Spectrosc. Relat. Phenom.* **2007**, *155*, 47–53.
- (98) Cavalleri, M.; Ogasawara, H.; Pettersson, L. G.; Nilsson, A. The interpretation of X-ray absorption spectra of water and ice. *Chem. Phys. Lett.* **2002**, *364*, 363–370.
- (99) Myneni, S.; Luo, Y.; Näslund, L. Å.; Cavalleri, M.; Ojamäe, L.; Ogasawara, H.; Pelmenchikov, A.; Wernet, P.; Väterlein, P.; Heske, C.; Hussain, Z.; Pettersson, L. G. M.; Nilsson, A. Spectroscopic probing of local hydrogen-bonding structures in liquid water. *J. Phys.: Condens. Matter* **2002**, *14*, L213–L219.
- (100) Hetényi, B.; De Angelis, F.; Giannozzi, P.; Car, R. Calculation of near-edge x-ray-absorption fine structure at finite temperatures: Spectral signatures of hydrogen bond breaking in liquid water. *J. Chem. Phys.* **2004**, *120*, 8632–8637.
- (101) Pykkänen, T.; Lehtola, J.; Hakala, M.; Sakko, A.; Monaco, G.; Huotari, S.; Hämäläinen, K. Universal Signature of Hydrogen Bonding in the Oxygen K-Edge Spectrum of Alcohols. *J. Phys. Chem. B* **2010**, *114*, 13076–13083.
- (102) Viguera, A. R.; Serrano, L. Side-chain interactions between sulfur-containing amino acids and phenylalanine in α -helices. *Biochemistry* **1995**, *34*, 8771–8779.
- (103) Zauhar, R. J.; Colbert, C. L.; Morgan, R. S.; Welsh, W. J. Evidence for a strong sulfur-aromatic interaction derived from crystallographic data. *Biopolymers* **2000**, *53*, 233–248.
- (104) Morgan, R. S.; Tatsch, C. E.; Gushard, R. H.; McAdon, J. M.; Warme, P. K. Chains of Alternating sulfur and π -bonded atoms in eight small proteins. *Int. J. Pept. Protein Res.* **1978**, *11*, 209–217.
- (105) Morgan, R. S.; McAdon, J. M. Predictor for sulfur-aromatic interactions in globular proteins. *Int. J. Pept. Protein Res.* **1980**, *15*, 177–180.
- (106) Valley, C. C.; Cembran, A.; Perlmutter, J. D.; Lewis, A. K.; Labello, N. P.; Gao, J.; Sachs, J. N. The Methionine-aromatic Motif Plays a Unique Role in Stabilizing Protein Structure. *J. Biol. Chem.* **2012**, *287*, 34979–34991.
- (107) Forbes, C. R.; Sinha, S. K.; Ganguly, H. K.; Bai, S.; Yap, G. P.; Patel, S.; Zondlo, N. J. Insights into thiol-aromatic interactions: A stereoelectronic basis for S-H/ π interactions. *J. Am. Chem. Soc.* **2017**, *139*, 1842–1855.
- (108) Shard, A. G.; Whittle, J. D.; Beck, A. J.; Brookes, P. N.; Bullett, N. A.; Talib, R. A.; Mistry, A.; Barton, D.; McArthur, S. L. A NEXAFS Examination of Unsaturation in Plasma Polymers of Allylamine and Propylamine. *J. Phys. Chem. B* **2004**, *108*, 12472–12480.
- (109) Beck, A. J.; Whittle, J. D.; Bullett, N. A.; Eves, P.; Mac Neil, S.; McArthur, S. L.; Shard, A. G. Plasma Co-Polymerisation of Two

Strongly Interacting Monomers: Acrylic Acid and Allylamine. *Plasma Processes Polym.* **2005**, *2*, 641–649.

(110) Khosravi, Z.; Kotula, S.; Lippitz, A.; Unger, W. E. S.; Klages, C.-P. IR- and NEXAFS-spectroscopic characterization of plasma-nitrogenated polyolefin surfaces. *Plasma Processes Polym.* **2018**, *15*, 1700066.

(111) Zabicky, J. In *The Chemistry of Functional Groups*; Zabicky, J., Ed.; John Wiley & Sons, Ltd.: Chichester, UK, 1970; Vol. 2.

(112) Patai, S. In *The Chemistry of Amidines and Imidates*; Patai, S., Ed.; John Wiley & Sons, Ltd.: Chichester, UK, 1975.

(113) Makarova, A. A.; Grachova, E. V.; Neudachina, V. S.; Yashina, L. V.; Blüher, A.; Molodtsov, S. L.; Mertig, M.; Ehrlich, H.; Adamchuk, V. K.; Laubschat, C.; Vyalikh, D. V. Insight into Bio-metal Interface Formation in vacuo: Interplay of S-layer Protein with Copper and Iron. *Sci. Rep.* **2015**, *5*, 8710.

(114) Rodriguez, C. F.; Cunje, A.; Shoeib, T.; Chu, I. K.; Hopkinson, A. C.; Siu, K. W. Solvent-assisted rearrangements between tautomers of protonated peptides. *J. Phys. Chem. A* **2000**, *104*, 5023–5028.

(115) Kamiya, K.; Boero, M.; Shiraishi, K.; Oshiyama, A. Enol-to-keto tautomerism of peptide groups. *J. Phys. Chem. B* **2006**, *110*, 4443–4450.

(116) Isaev, A. N. Keto-enol tautomerization of a peptide group with proton transfer through a water bridge. *Russian Journal of Physical Chemistry A* **2015**, *89*, 1360–1367.

(117) Hvidt, A.; Nielsen, S. O. Hydrogen Exchange in Proteins. *Adv. Protein Chem.* **1966**, *21*, 287–386.

(118) Perrin, C. L. Proton Exchange in Amides: Surprises from Simple Systems. *Acc. Chem. Res.* **1989**, *22*, 268–275.

(119) Eriksson, M. A.; Härd, T.; Nilsson, L. On the pH dependence of amide proton exchange rates in proteins. *Biophys. J.* **1995**, *69*, 329–339.

(120) Gorman, G. S.; Speir, J. P.; Turner, C. A.; Amster, I. J. Proton affinities of the 20 common α -amino acids. *J. Am. Chem. Soc.* **1992**, *114*, 3986–3988.

(121) Cox, K. A.; Gaskell, S. J.; Morris, M.; Whiting, A. Role of the site of protonation in the low-energy decompositions of gas-phase peptide ions. *J. Am. Soc. Mass Spectrom.* **1996**, *7*, 522–531.

(122) Wu, R.; McMahon, T. B. Protonation Sites and Conformations of Peptides of Glycine ($\text{Gly}_{1-5}\text{H}^+$) by IRMPD Spectroscopy. *J. Phys. Chem. B* **2009**, *113*, 8767–8775.

(123) Voss, J. M.; Fischer, K. C.; Garand, E. Revealing the structure of isolated peptides: IR-IR predissociation spectroscopy of protonated triglycine isomers. *J. Mol. Spectrosc.* **2018**, *347*, 28.

(124) Zubavichus, Y.; Zharnikov, M.; Shaporenko, A.; Fuchs, O.; Weinhardt, L.; Heske, C.; Umbach, E.; Denlinger, J. D.; Grunze, M. Soft X-ray induced decomposition of phenylalanine and tyrosine: A comparative study. *J. Phys. Chem. A* **2004**, *108*, 4557–4565.