University of Groningen

17 β -Estradiol Treatment Protects Lungs Against Brain Death Effects in Female Rat Donor

Yamamoto Ricardo da Silva, Fernanda; Armstrong Junior, Roberto; Vidal-dos-Santos, Marina; Correia, Cristiano de Jesus; Coutinho e Silva, Raphael Santos; da Anunciacao, Lucas Ferreira; Moreira, Luiz Felipe Pinho; Leuvenink, Henri G.D.; Breithaupt-Faloppa, Ana Cristina

Published in:
Transplantation

DOI:
[10.1097/TP.0000000000003467](https://doi.org/10.1097/TP.0000000000003467)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Yamamoto Ricardo da Silva, F., Armstrong Junior, R., Vidal-dos-Santos, M., Correia, C. D. J., Coutinho e Silva, R. S., da Anunciacao, L. F., Moreira, L. F. P., Leuvenink, H. G. D., & Breithaupt-Faloppa, A. C. (2021). 17 β -Estradiol Treatment Protects Lungs Against Brain Death Effects in Female Rat Donor. *Transplantation*, 105(4), 775-784. <https://doi.org/10.1097/TP.0000000000003467>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

17 β -Estradiol Treatment Protects Lungs Against Brain Death Effects in Female Rat Donor

Fernanda Yamamoto Ricardo-da-Silva, MsC,¹ Roberto Armstrong Jr,¹ Marina Vidal-dos-Santos,¹ Cristiano de Jesus Correia, PhD,¹ Raphael dos Santos Coutinho e Silva,¹ Lucas Ferreira da Anunciação,¹ Luiz Felipe Pinho Moreira, MD, PhD,¹ Hendrik Gerrit Derk Leuvenink, PhD,² and Ana Cristina Breithaupt-Faloppa, PhD¹

Background. Brain death (BD) affects the viability of lungs for transplantation. A correlation exists between high-lung inflammation after BD and the decrease in female sex hormones, especially estradiol. Therefore, we investigated the effects of 17 β -estradiol (E2) treatment on the lungs of female brain dead rats. **Methods.** Female Wistar rats were divided into 4 groups: BD (submitted to BD for 6 h), sham (false operated), E2-T0 (treated with E2 immediately after BD; 50 μ g/mL, 2 mL/h), and E2-T3 (treated with E2 after 3 h of BD; 50 μ g/mL, 2 mL/h). Lung edema, hemorrhage, and leukocyte infiltration were analyzed. Adhesion molecules were evaluated, and analysis of NO synthase gene and protein expression was performed using real-time PCR and immunohistochemistry, respectively. Release of chemokines and matrix degradation in the lungs was analyzed. **Results.** BD increased leukocyte infiltration, as shown by intravital microscopy ($P=0.017$), bronchoalveolar lavage cell count ($P=0.016$), the release of inflammatory mediators ($P=0.02$), and expression of adhesion molecules. BD also increased microvascular permeability and the expression and activity of matrix metalloproteinase-9 in the lungs. E2 treatment reduced leukocyte infiltration, especially in the E2-T3 group, release of inflammatory mediators, adhesion molecules, and matrix metalloproteinase activity in the lungs. **Conclusions.** E2 treatment was successful in controlling the lung inflammatory response in females submitted to BD. Our results suggest that E2 directly decreases the release of chemokines, restraining cell traffic into the lungs. Thus, E2 has a therapeutic potential, and its role in improving donor lung quality should be explored further.

(*Transplantation* 2021;105: 775–784).

INTRODUCTION

Lung transplantation is often the only therapy option for many patients with end-stage failure. There is a low

number of multiorgan brain death (BD) donors, among which only about 15%–20% of potential lungs are considered suitable for transplantation.¹ BD initiates a systemic inflammatory process with innate immune system activation,² cytokine release, and expression of adhesion molecules, which mostly affects the lungs.³ Important hormonal and metabolic changes occur after hypothalamic/pituitary failure in experimental models of BD associated with previous events such as trauma and hemorrhage, thereby adding to the final inflammatory result of the organ donor.⁴

Although lungs from female donors are considered to have a greater risk for transplant in male recipients,^{5,6} lung transplantation between women shows potentially better results.⁵ An experimental study⁷ showed that female rats presented greater lung inflammation, associated with an acute reduction of female sex hormones. The reduction of female sex hormones is known to be detrimental to the cardiovascular system,⁸ and clinical and experimental studies reinforce the E2 protector effect on the immune system, pulmonary inflammation, and pro-inflammatory mediator release.^{9,10} In this context, this study evaluated E2 treatment as a therapeutic alternative to mitigate the deleterious effects of BD, by investigating the pulmonary inflammatory response to its application in female rats subjected to BD.

Received 25 March 2020. Revision received 25 July 2020.

Accepted 7 August 2020.

¹ Laboratório de Cirurgia Cardiovascular e Fisiopatologia da Circulação (LIM-11), Instituto do Coração (InCor), Faculdade de Medicina da Universidade de São Paulo, São Paulo, Brazil.

² Department of Surgery, University Medical Centre Groningen, University of Groningen, Groningen, The Netherlands.

This work was supported by the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP 2016/0351-0). The author F.Y.R.S. was a scholarship holder by FAPESP 2016/03692-9 and 2018/07289-0. M.V.S. was a scholarship holder by FAPESP 2017/11798-4.

The authors declare no conflicts of interest.

F.Y.R.-d.-S. and R.A. Jr. performed the study, analyzed data, and wrote the article. M.V.-d.-S., C.d.J.C., R.d.S.C.e.S., and L.F.d.A. contributed obtaining and analyzing data. A.C.B.-F. designed the study, analyzed data, and wrote the article. H.G.D.L. and L.F.P.M. contributed analyzing data and wrote the article.

Correspondence: Ana Cristina Breithaupt-Faloppa, PhD, Laboratório de Cirurgia Cardiovascular e Fisiopatologia da Circulação (LIM/11)—HC-FMUSP, Universidade de São Paulo, Av Dr Arnaldo, 455 2º andar—sala 2146—01246-903 São Paulo, Brazil. (ana.breithaupt@hc.fm.usp.br).

Copyright © 2020 Wolters Kluwer Health, Inc. All rights reserved.

ISSN: 0041-1337/21/1054-775

DOI: 10.1097/TP.00000000000003467

MATERIALS AND METHODS

Animals

Female Wistar rats (60 d, $n = 52$) were kept in a $23 \pm 2^\circ\text{C}$, 12 hour light-dark cycle, with free access to food and water. All rats received humane care in compliance with the “Principles of Laboratory Animal Care” formulated by the National Society for Medical Research and the Guide for the Care, and the “Use of Laboratory Animals” prepared by the Institute of Laboratory Animal Resources, published by the National Institute of Health (NIH Publication No. 86-23, revised 1996). The experiments were approved by the University of São Paulo Medical School Ethics Committee for Research Projects Analysis (SDC No. 4350/16/016). The animals were divided into 4 groups: *sham*—animals submitted to cranial perforation, BD—animals submitted to BD, E2-T0—animals treated with E2 immediately after BD induction, and E2-T3—animals treated with E2, 3 hours after BD induction.

Anesthesia and BD Model

Female rats in estrus or proestrus phases of the estral cycle (identified by vaginal smears) were incubated under isoflurane anesthesia. A catheter was placed in the carotid artery for measurement of mean arterial pressure (MAP) and the jugular vein for treatment and fluid administration (saline solution, 2 mL/h with cyclodextrin [E2 vehicle—Sigma-Aldrich, St. Louis, MO]).

A Fogarty-4F catheter (Baxter Healthcare Co., Marion, NC) was inserted intracranially, and BD was induced by rapid inflation of the balloon with 400 μL of saline solution. BD was confirmed by maximally dilated and fixed pupils, apnea, absence of reflexes, and a drop in MAP. After BD, anesthesia was interrupted, and fluid administration was started. All animals were maintained for 6 hours. At the end of experimental period, arterial blood samples were obtained via catheter to analyze oxygen and carbon dioxide (CO_2) partial pressures using gas analyzer (ABL555, Radiometer Medical ApS, Copenhagen, Denmark).

E2 Treatment and Serum Quantification of Hormones

The rats received 50 $\mu\text{g/mL}$ (Sigma-Aldrich) intravenous E2, IV in saline solution (2 mL/h). The E2-T0 group started fluid replacement with E2 right after BD confirmation. The E2-T3 group started fluid replacement immediately after BD confirmation but E2 treatment commenced after 3 hours. The sham and BD groups received fluid replacement only. Serum E2 and corticosterone concentrations were quantified using an ELISA kit (Cayman Chemical Company, Ann Arbor, MI) following the manufacturer's protocols.

Cell Counts on Bronchoalveolar Lavage

After euthanasia by exsanguination, 5 mL of culture medium (Roswell Park Memorial Institute, Sigma-Aldrich) was repeatedly injected in the airways. The bronchoalveolar lavage (BAL) fluid recovered was then centrifuged ($239 \times g$, 15 min), the pellet resuspended in 1 mL of Roswell Park Memorial Institute and the total cell number determined using a Neubauer chamber.

In Vivo Analysis of Leukocyte Infiltrate on Lung Microcirculation

The intravital microscopy technique was based on Schmidt et al.¹¹ Accordingly, an incision was made on the left lateral decubitus to expose the intercostal muscles, and a small incision was made between the ribs to expose the inferior right lung lobule. Leukocytes were stained based on Baatz et al.¹² with Rhodamine 6G (9 $\mu\text{mol/kg}$, IV, Sigma-Aldrich) injected 10 minutes before the end of BD. Leukocyte infiltrate was accessed in images extracted from 1-minute videos, counted in 3 distinct frames of the video, obtained with an epifluorescence microscope coupled camera. The results are expressed as cell/ mm^2 of lung tissue.

Histopathological Lung Evaluation

Lung sections were stained with hematoxylin/eosin then digitized. Tissue areas ($n = 5$ per group, 2 sections per animal, 5 areas per section) were randomly analyzed for leukocyte infiltrate, edema, and hemorrhage with NIS-Elements Software Basic Research (Nikon, Tokyo, Japan) by 2 examiners. The air space/lung tissue ratio was measured, the air space area was disconsidered and the leukocyte number reported in cells/ μm^2 of lung tissue. A reticle with 130 intersecting lines (total area 27648 μm^2) was placed covering the images, and each intersection when over edema or hemorrhage was considered as a unit, the result was expressed as units/ μm^2 .

Ex Vivo Lung Tissue Culture

Lungs were perfused via the pulmonary artery and tissue culture performed according to Breithaupt-Faloppa et al.¹³ Fragments of lung parenchyma (4 pieces per well) were incubated in 1 mL of DMEM (DMEM, Vitrocell, Sao Paulo, Brazil) for 24 hours at 37°C in a humidified atmosphere with 5% CO_2 . After incubation, the medium was stored at -80°C and lung pieces were dried and weighed.

Quantification of Inflammatory Mediators in Lung Cultures and Lung Homogenate Samples

Cytokine-induced neutrophil chemoattractant (CINC)-1, macrophage inflammatory protein (MIP)-1 α , and -2 were quantified in lung culture medium aliquots using a Milliplex MAP kit (EMD Millipore Corporation, Burlington, MA) and expressed as pg/mL. Measurements were performed using Luminex 200-Software xPonent/Analyst 4.2 version (EMD Millipore Corporation). Interleukin (IL)-1 β and tumor necrosis factor (TNF)- α were quantified in lung homogenates by ELISA following the manufacturer's instructions (R&D Systems, Inc., Minneapolis, MN).

Immunohistochemistry

The left lung was expanded with 1:3 optimal cutting temperature media diluted in PBS and snap-frozen in nitrogen-hexane solution. Lung cryosections (10 μm) were fixed in acetone for 10 minutes. Endogenous peroxidase blocking (H_2O_2 , 2%) was followed by unspecific site blockage with bovine serum albumin (1%) in Tris-buffered saline-tween. The sections were incubated at 37°C for 2 hours with primary antibodies in Tris-buffered saline-tween/bovine serum albumin: antiintercellular adhesion molecule (ICAM)-1 (1:100; BOOSTER), antivascular cell adhesion molecule (VCAM) (1:200, Abcam, Cambridge, United

Kingdom), and antiendothelial nitric oxide synthase (eNOS) (1:250; Novus Biologicals, Littleton, CO). Next, a 1:200 conjugated horseradish peroxidase (HRP) secondary antibody was applied to the sections at 37°C for 1 hour (EMD Millipore Corporation). Finally, staining was performed using HRP as substrate (3-amino-9-ethylcarbazole; Vector Laboratories, Burlingame, CA) for 5–10 minutes. Stained areas were identified in vessel walls (VCAM and eNOS) or whole lung tissues (ICAM-1) after determining a threshold and analyzing the staining area fractions with an image analyzer (NIS—Elements; Nikon). Background was determined by sections incubated in the absence of the primary antibody (negative control).

To investigate matrix metalloproteinase (MMP)-9 and inducible nitric oxide synthase (iNOS) expression in lung tissue, paraffin sections (4 µm) were rehydrated and treated for antigen retrieval (citrate buffer [pH 6.0], 20 min, 95°C). Endogenous peroxidase was blocked (H₂O₂, 2%), then goat serum (5%) was used for blocking nonspecific sites. MMP-9 immunohistochemistry was performed with 1:100 primary antibody (anti-MMP-9, Abcam) and secondary antibody linked to alkaline phosphatase (MULTIVIEW IHC kit, Enzo Life Sciences, Farmingdale, NY). iNOS immunohistochemistry was performed using a 1:100 primary antibody (anti-iNOS, Abcam) and secondary antibody conjugated to HRP (EMD Millipore Corporation). Negative controls were performed by incubating the sections in the absence of the primary antibody. Stained areas were quantified by analysis of tissue with NIS-Elements software (Nikon).

Gene Expression Analysis

Lung samples were stored in stabilization reagent at –80°C. Using commercial kit mirVana Kit (Ambion, Austin, TX) RNA was extracted. Then, cDNA was transcribed with reverse transcriptase (High-Capacity Reverse Transcriptase Kit; Applied Biosystems, Foster City, CA). Real-time PCR was performed using TaqMan Universal PCR Master Mix (Applied Biosystems) and TaqMan primers (Applied Biosystems) against glyceraldehyde-3-phosphate dehydrogenase, β -actin, ICAM-1, VCAM, and MMP-9 (Table 1). Targets were amplified over 1 cycle for

2 minutes at 50°C, 1 cycle for 10 minutes at 95°C, and 40 cycles for 15 seconds at 95°C and for 1 minute at 60°C.

In parallel, β -actin, IL-1 β , and TNF- α target genes (Table 1) were amplified with SYBR Green primers (Applied Biosystems) and SYBR Green PCR Master Mix (Applied Biosystems) over 1 cycle for 2 minutes at 50°C, 1 cycle for 10 minutes at 95°C, and 40 cycles for 15 seconds at 95°C and for 1 minute at 60°C.

MMP Activity Assay

The MMP activity assay was carried out with lung homogenate samples using a commercial fluorescent and specific substrate kit (Gelatinase Assay kit; BioVision, Milpitas, CA) according to the manufacturer's instructions. Analysis was performed using a fluorescence microplate reader (Multi-Mode Reader; Biotek, Winooski, VT) in kinetic mode.

Lung Microvascular Leakage

Evans Blue (EB) dye extravasation was used to evaluate lung microvascular permeability.⁷ The EB dye was injected (20 mg/kg, intravenously; Sigma-Aldrich) 20 minutes before the end of the BD period. Dye was extracted from tissue then quantified by spectrophotometry (620 nm) with a standard curve (0.3–100 µg/mL). Data were expressed as µg/mg of dry weight.

Homogenization of the Lung Tissue

Lung fragments were weighed and dissociated in PBS containing 0.5% hexadecyltrimethylammonium bromide and 5 mmol/L EDTA (pH 6.0, 3 mL/kg; gentleMACS Dissociator [Miltenyi Biotec, Bergisch Gladbach, Germany]). The supernatants were stored at –80°C for analysis.

Determination of Total Nitric Oxide

Samples of lung homogenate were used to determine the total nitric oxide (NO_x[–]), nitrite (NO₂[–]), and nitrate (NO₃[–]). We followed the method described by Breithaupt-Faloppa et al.¹³ Accordingly, nitrates were reduced to nitrites then measured with Griess reagent reaction (540 nm). The level of nitrites was determined with a standard curve of NaNO₂ (5–60 µmol/L).

Statistical Analysis

All data are presented as mean $\pm$ SEM. The data were analyzed by ANOVA followed by Tukey, Dunnett's, or Sidak multiple comparison tests. Abnormally distributed data were analyzed using the Kruskal-Wallis test followed by a post hoc Dunn multiple comparison test. Statistical analysis was performed using GraphPad Prism Software v.8.3.1.

RESULTS

Arterial Blood Gases and Serum Concentrations of Corticosterone and Estradiol

The BD model applied here involves the sudden increase in intracranial pressure, leading to a transient hypertensive crisis followed by sustained hypotension, as previously shown.⁷ This MAP pattern was observed in all studied groups (data not shown). Additionally, measurements of

TABLE 1.
Primer sequence used for real-time PCR analysis

Real-time PCR TaqMan		
GAPDH	Rn01775763_g1	
β -Actin	Rn00667869_m1	
ICAM-1	Rn005642227_m1	
VCAM	Rn00563627_m1	
MMP-9	Rn00579162_m1	
Real-time PCR SYBR Green		
β -Actin	RN b-act fw	5'-GGAAATCGTGCCTGACATTA-3'
	RN b-act rv	5'-GCGGCAGGGCCATCTC-3'
IL-1 β	RN IL-1B fw	5'-CAGCAATGGTCGGGACATAGTT-3'
	RN IL-1B rv	5'-GCATTAGGAATAGTCGACCCATCT-3'
TNF- α	TB TNF-a fw	5'-AGGCTGTGCTACATCACTGAA-3'
	RN TNF-a rv	5'-TGACCCGTAGGGCGATTACA-3'

GAPDH, glyceraldehyde-3-phosphate dehydrogenase; ICAM, intercellular adhesion molecule; IL, interleukin; MMP, matrix metalloproteinase; PCR, polymerase chain reaction; TNF, tumor necrosis factor; VCAM, vascular cell adhesion molecule.

TABLE 2.**Quantification of estradiol and corticosterone in serum samples (pg/mL), and oxygen and carbon dioxide partial pressure (mm Hg)**

	Sham	BD	E2-T0	E2-T3	P ANOVA
Estradiol	482.2 ± 54.2 ^a	68.2 ± 8.0	1713.5 ± 44.7 ^a	1717.9 ± 29.2 ^a	<0.0001
Corticosterone	3458.4 ± 929.9 ^a	22.6 ± 10.4	16.8 ± 8.1	40.7 ± 31.1	0.0001
Pao ₂	255.29 ± 34.48	195.15 ± 18.08	185.57 ± 22.16	186.68 ± 39.92	0.3176
PACo ₂	26.7 ± 3.75	33.56 ± 4.33	32.07 ± 3.84	38.84 ± 5.29	0.3630

The female rats were subjected to brain death (BD). In the treated groups, animals were submitted to BD and continuously treated with 17 β -estradiol (E2) immediately after BD confirmation (E2-T0) or after 3 h of BD (E2-T3). Controls were false operated animals (Sham).

Data expressed as mean $\pm$ SEM from 8 animals.

^a $P \leq 0.05$ in relation to BD.

Po₂ (O₂) and CO₂ were taken after the end of the 6 hours experiment in all groups (Table 2). Partial pressure of O₂ or CO₂ were not modified by the BD, compared with Sham and remained similar in animals treated with E2.

Data on estradiol and corticosterone serum concentrations are shown in Table 2. BD rats presented reduction of both hormones in comparison to Sham. The E2-treated rats presented similarly elevated estradiol serum concentration compared to BD group.

Estradiol Effects on Lung Leukocyte Infiltrate, Edema, and Hemorrhage

Figure 1A illustrates the BAL cell count. After BD, the BAL cell counts increased and the E2 treatments (E2-T0 and E2-T3) reduced the cell numbers in BAL. The lung in vivo quantification of leukocyte infiltration (intravital microscopy) after 6 hours of BD showed increased leukocyte infiltration into the BD group and that the 3 hours treatment (E2-T3) reduced leukocyte infiltration (Figure 1B).

Figure 2 presents the histopathologic analysis of leukocyte infiltrate, edema, and hemorrhage in lung sections. Both E2 treatments (E2-T0 and E2-T3) significantly decreased lung leukocyte infiltration. Furthermore, the lungs of E2-T3 animals presented significantly less edema and hemorrhage.

Analysis of Long-lasting Chemokine Release

Table 3 shows the concentrations of MIP-1 α , MIP-2, and CINC-1 in lung culture medium after 24 hours. MIP-1 α and CINC-1 concentrations were increased in media from BD lung culture, and MIP-2 concentration was higher in the BD group than in the sham group ($P = 0.063$). Conversely, both E2 treated groups (E2-T0 and E2-T3) significantly reduced the release of all studied chemokines.

Gene and Protein Expression of Adhesion Molecules

Figure 3 shows that gene expression of ICAM (A) and VCAM (B) was significantly increased by BD when compared with the sham group. E2 treatment decreased ICAM gene expression in both treated groups. Regarding protein expression, VCAM was increased in brain dead animals, but there was no difference in ICAM-1 levels. E2 treatment was able to reduce protein expression of both ICAM-1 and VCAM in group E2-T3 (Figures 3C,D and 4).

Analysis of Systemic Inflammatory Mediator Release

IL-1 β and TNF- α were also analyzed for gene expression and protein concentration in lung tissue homogenates

(Figure 5). In IL-1 β , both gene expression and protein concentrations in lung tissue were increased after BD. E2 treatment reduced IL-1 β gene expression in both groups (E2-T0 and E2-T3) and reduced IL-1 β levels in the lung, showing a significant decrease in the E2-T0 group.

In relation to TNF- α , lung gene expression was not altered after 6 hours of BD. Nevertheless, E2 treatment downregulated TNF- α gene expression in E2-T0 and E2-T3 animals, which was reduced in group E2-T3 compared with the BD group. The lung TNF- α protein concentration after 6 hours was not increased by BD, but it was reduced in the E2-T3 group.

MMP Expression and Activity

MMP-9 protein and gene expression levels were analyzed in the lungs (Figure 6A,B). After BD, the lungs showed increased expression of both the MMP-9 enzyme and gene. E2 treatment did not alter the enzyme expression, but the E2-T0 treatment reduced the gene expression of MMP-9. An activity assay for MMPs was performed (Figure 6C), and no differences were observed between sham and BD group animals. On the other hand, both E2 treatments reduced MMP (MMP-9 and -2) activities.

Analysis of Lung Microvascular Permeability and NO Production

BD increased the microvascular permeability in the lungs (Figure 7A), as observed by the increase in EB dye extravasation. Concurrently, there was an increase in nitrate and nitrites in lung homogenates (Figure 7B). Nitric oxide synthase expression was analyzed, as well as eNOS protein expression in lung vessels (Figure 7C) and iNOS in lung tissue (Figure 7D). eNOS was found to be reduced after BD, whereas iNOS was increased. E2 treatment had no effect on the permeability, NO synthases, or metabolites.

DISCUSSION

BD produces a systemic inflammatory response, of which the lung is one of the most vulnerable organs³ through parenchyma degradation, activation of lung inflammatory cells, and recruitment of other leukocytes to the tissue.¹⁴ The BD experimental model used by our study generated leukocyte infiltration as well as edema and hemorrhage in the lungs, resulting in lung injury. A study from our research showed that BD systemic injury affects men and women differently, whereby females have a greater inflammatory response characterized by a pronounced infiltrate in the lungs and increased permeability, associated with the

FIGURE 1. Cell counts in the BAL (A) and lung in vivo evaluation of leukocyte infiltrate using intravital macroscopy. The female rats were subjected to BD. In the treated groups, animals were submitted to BD and continuously treated with E2 immediately after BD confirmation (E2-T0) or after 3 h of BD (E2-T3). Controls were false operated animals (Sham). Data are expressed as mean $\pm$ SEM from 8 animals (A) and 5 animals (B). $^{\gamma}P \leq 0.05$ in relation to BD. BAL, bronchoalveolar lavage; BD, brain death; E2, 17 β -estradiol.

Lung histopathology					
	Sham	BD	E2-T0	E2-T3	P VALUE
Leukocytes infiltrate (leukocytes/mm ²)	2127 $\pm$ 149.5	2343 $\pm$ 96.27	1847 $\pm$ 102.4 ^{γ}	1851 $\pm$ 164.3 ^{α}	0.0025
Edema (Units/mm ²)	3213 $\pm$ 277.3	3048 $\pm$ 140.8	2628 $\pm$ 160.4	2379 $\pm$ 97.8 ^{α}	0.0087
Hemorrhage (Units/mm ²)	596.8 $\pm$ 50.4	681.3 $\pm$ 57.8	501.9 $\pm$ 58.4	488.7 $\pm$ 43.8 ^{α}	0.0499

FIGURE 2. Lung histopathology analysis of leukocyte infiltrate, edema, and hemorrhage. Photomicrographs of lung sections stained with standard hematoxylin and eosin staining. The female rats were subjected to BD. In the treated groups, animals were submitted to BD and continuously treated with E2 immediately after BD confirmation (E2-T0) or after 3 h of BD (E2-T3). Controls were false operated animals (Sham). Data are expressed as mean $\pm$ SEM from 5 photos/sample, 3 samples/animal, and 8 animals/group animals on lung histopathology. Leukocyte infiltrate: ^{γ} $P=0.012$ in relation to BD, ^{α} $P=0.005$ in relation to BD. Edema: ^{α} $P=0.0022$ in relation to BD. Hemorrhage: ^{α} $P=0.045$ in relation to BD. BD, brain death; E2, 17 β -estradiol.

acute reduction of E2, a known protector of lung inflammation.⁷ The aim of this study was to understand the effect of E2 treatment on females submitted to BD, since it is known that E2 modulates lung inflammation by regulating infiltration of inflammatory cells into the lung.¹⁵ We used 2

different time points of E2 treatment to analyze its effects on BD-induced lung injury in females (E2-T0) and its possible therapeutic application (E2-T3). To maintain high E2 concentration during the studied period, a low therapeutic dose was used, as previously reported.¹⁶

TABLE 3.**Chemokines measured in culture (explant) medium of lung fragments for 24 h at 37°**

(pg/mg)	Sham	BD	E2-T0	E2-T3	P ANOVA
MIP-1 α	5198 $\pm$ 1415 ^a	21 226 $\pm$ 4626	5329 $\pm$ 106.1 ^a	6484 $\pm$ 1037 ^a	0.0009
MIP-2	73 756 $\pm$ 39 945	417 038 $\pm$ 206 036	23 967 $\pm$ 8556 ^a	20 731 $\pm$ 7115 ^a	0.0296
CINC-1	6754 $\pm$ 264.5 ^a	29 657 $\pm$ 9097	10 062 $\pm$ 1422 ^a	9988 $\pm$ 2572 ^a	0.0181

Quantified with multiplex assay. The female rats were subjected to brain death (BD). In the treated groups, animals were submitted to BD and continuously treated with 17 β -estradiol (E2) immediately after BD confirmation (E2-T0) or after 3 h of BD (E2-T3). Controls were false operated animals (Sham). Data expressed as mean $\pm$ SEM from 4–6 animals.

^a $P \leq 0.05$ in relation to BD.

BD, brain death; CINC-1, cytokine-induced neutrophil chemoattractant 1; E2, 17 β -estradiol; MIP, macrophage inflammatory protein.

FIGURE 3. Gene and protein expression of ICAM-1 (A, C) and VCAM (B, D). The female rats were subjected to BD. In the treated groups, animals were submitted to BD and continuously treated with E2 immediately after BD confirmation (E2-T0) or after 3 h of BD (E2-T3). Controls were false operated animals (Sham). Data are expressed as mean $\pm$ SEM from 8 (A, B) and 5 photos/sample, 2 samples/animal, and 5 animals/group in (C, D). ^a $P \leq 0.05$ in relation to BD. BD, brain death; E2, 17 β -estradiol; ICAM, intercellular adhesion molecule; VCAM, vascular cell adhesion molecule.

Our first analysis was BAL cellular content, which showed that BD is responsible for a higher cell count. It seems that not only were many leukocytes recruited to the lung, but they also passed into the alveolus. Leukocyte infiltration into the alveolar tree was confirmed by the increased number of cells observed *in vivo* on the lung parenchyma. It is important to point out that E2 treatment decreased leukocyte infiltration to the airways and lung parenchyma. Our study further investigated the production and release of cytokines, chemokines, and expression of adhesion molecules responsible for leukocyte mobilization during the inflammatory process. First, we analyzed the local release of MIP-1 α , MIP-2, and CINC-1, which are relevant macrophage and neutrophil chemokines.^{17,18} BD indeed caused a long-lasting production of MIP-1 α and CINC-1, as observed in the lung in culture media. Similarly, a study from our laboratory

observed an increased release of systemic CINC-1, influencing the blood white cell numbers (data not published). BD recruited leukocytes into the blood, predominantly neutrophils as a response. In turn, E2 treatment had a role in reducing recruited white blood cells, mainly by reducing the neutrophil population in the lung tissue in both treatments through reduction of chemokines in the lung.

In the event of lung inflammation, alveolar macrophages produce cytokines, such as IL-1 and TNF- α , which affect endothelial cells that express adhesion molecules and release chemokines.¹⁹ Expression of adhesion molecules is a crucial step for leukocyte mobilization, as they are responsible for the firm adhesion and transmigration of leukocytes from the capillary endothelium into the lung parenchyma.¹⁹ E2 regulates adhesion molecule expression, and there is a correlation between low levels of estrogen and higher expression of soluble ICAM-1 and

FIGURE 4. Photomicrographs of immunohistochemical reaction showing ICAM-1 and VCAM on lung tissue (original magnification $\times 40$). The female rats were subjected to BD. In the treated groups, animals were submitted to BD and continuously treated with E2 immediately after BD confirmation (E2-T0) or after 3 h of BD (E2-T3). Controls were false operated animals (Sham). BD, brain death; E2, 17 β -estradiol; ICAM, intercellular adhesion molecule; VCAM, vascular cell adhesion molecule.

FIGURE 5. IL-1 β and TNF- α gene expression (A, B) and concentrations in lung homogenates (C, D). The female rats were subjected to BD. In the treated groups, animals were submitted to BD and continuously treated with E2 immediately after BD confirmation (E2-T0) or after 3 h of BD (E2-T3). Controls were false operated animals (Sham). Data are expressed as mean $\pm$ SEM from 8 animals. $^{\ast}P \leq 0.05$ in relative to BD. BD, brain death; E2, 17 β -estradiol; IL, interleukin; TNF, tumor necrosis factor.

VCAM, in females,²⁰ and E2 is able to inhibit cytokine-stimulated VCAM-1 and ICAM-1 expression by endothelial cells.²¹ In this regard, the acute reduction of E2 after the BD model would contribute to the increase in adhesion molecule expression. BD increased both molecules on gene expression and VCAM on protein expression. Our data confirmed the effect of E2 on VCAM and ICAM-1 at the gene and protein levels. This indicates that in the

BD model, E2 leads to a reduction in adhesion molecule expression in the lungs, similar to what is observed in females after ischemia and reperfusion or trauma—hemorrhage model.^{22,23}

Cytokine production plays a crucial role in mediating inflammatory injury. IL-1 β and TNF- α cytokines are produced early in the inflammatory response by alveolar macrophages via activation of NF- κ B and are released in

FIGURE 6. MMP-9 protein (A) and gene (B) expression, MMP activity (C), and lung photomicrography of MMP-9 protein expression. The female rats were subjected to BD. In the treated groups, animals were submitted to BD and continuously treated with E2 immediately after BD confirmation (E2-T0) or after 3 h of BD (E2-T3). Controls were false operated animals (Sham). Data are expressed as mean $\pm$ SEM from 5 photos/sample, 2 samples/animals, and 5 animals/group (A), and 8 animals on (B, C). (B) $^{\gamma}P \leq 0.05$ in relation to BD. BD, brain death; E2, 17 β -estradiol; MMP, matrix metalloproteinase.

the serum.²⁴ Our data show that BD elicits lung synthesis and release of IL-1 β , which is involved in enhanced expression of both ICAM-1 and VCAM on endothelial cells.^{25,26} More importantly, E2 treatment could have indirectly affected the expression of adhesion molecules in the lungs by reducing the local release of IL-1 β and TNF- α , thereby contributing to lower leukocyte infiltration.

In addition to cytokines and chemokines during lung inflammation, MMPs are released to facilitate the clearance of foreign and harmful agents; however, when in excess, these enzymes may destroy the extracellular matrix, disrupt resident cells, and further stimulate the inflammatory process.²⁷ Here, MMP-9, which plays an important role in lung injury,²⁷ was increased by means of genetic and protein expression as well as MMP activity after BD. MMP-9 is released from stimuli of TNF- α , IL-8, and IL-1 β mainly on inflammatory cells, such as neutrophils, monocytes, macrophages, and even endothelial cells.²⁸⁻³⁰ Lung MMP-9 was attenuated by E2 treatment, observed after both treatment time points on the MMP activity assay and gene expression. Other studies have confirmed the role of E2 in suppressing MMP-9 gene expression and attenuating hemorrhage, permeability, and leukocyte infiltration.^{31,32}

Additionally, the effect of E2 on lung edema formation after BD, parameters of vascular permeability, and nitric oxide synthases were assessed. In pathological situations, there is an overexpression of iNOS resulting in an excess of NO and imbalanced NO production.³³ Proinflammatory agents trigger iNOS expression in different cells, such as macrophages and smooth muscle cells.³⁴ In our model, we observed that females registered increased total nitrites on lung homogenates accompanied by increased lung microvascular permeability after BD, which could be a result of increased iNOS expression. iNOS has proinflammatory properties, influencing the secretion of proinflammatory cytokines, endothelial expression of P-selectin and ICAM-1, increasing neutrophil infiltration and related oxidative and nitrosative stress.³⁵

It is known that apart from E2 levels decrease, corticosterone levels also decrease after BD,^{7,36} which could contribute to edema formation. Acutely, glucocorticoids inhibit vasodilation and increase vascular permeability that occurs following inflammatory insult.³⁷ E2 regulates the release of glucocorticoids in females during stress conditions, and females have higher basal levels of corticosterone and a faster secretion rate.³⁸ Our data show that E2 controlled the increase in parenchymal edema but not alveolar edema. Thus, further

FIGURE 7. Permeability assessment with EB dye extravasation (A), NO_x in lung homogenates (B), eNOS (C), and iNOS (D) protein expression in the lungs. The female rats were subjected to BD. In the treated groups, animals were submitted to BD and continuously treated with E2 immediately after BD confirmation (E2-T0) or after 3 h of BD (E2-T3). Controls were false operated animals (Sham). Data are expressed as mean ± SEM from 8 animals in (A, B), and 5 photos/samples, 2 samples/animal, and 5 animals/group in (C, D). **P* ≤ 0.05 in relation to BD. BD, brain death; E2, 17β-estradiol; EB, Evans blue; eNOS, endothelial nitric oxide synthase; iNOS, inducible nitric oxide synthase.

studies should evaluate the association of E2 and corticosterone after BD, focusing on the control the edema formation and decrease the microvascular permeability.

Taken together the results point to E2 effects in reducing leukocyte infiltration, release of inflammatory mediators, adhesion molecules, and MMP activity in the lungs. E2 effects depend on several factors, including the estrogen receptors (ERs) involved in the response. Despite the fact that both classic ERs are required for pulmonary alveolar formation, lung cells were shown to express higher levels of ERβ than ERα mRNAs.³⁹ Conversely, ERα, and not ERβ, is expressed in alveolar macrophages and endothelial cells of lung vessels.⁴⁰ In parallel, E2 can also act through nongenomic mechanisms to rapidly activate signaling pathways and modulate protein expression, function, and distribution. Here, we could suggest that E2 treatment was able to elicit rapid effects by intracellular signaling pathways and to inhibit inflammatory gene expression by modulating transcription factors as NF-κB.^{21,41}

This study has some limitations, such as the possibility that the chosen time point for analysis at 6 hours after BD limits the results that take effect before that period or after it. The study used E2, which is not a receptor-selective agonist. Lung function was not specifically addressed; nonetheless, our study confirmed the important protective effects of E2 in the lungs after BD in females. The study adds to the important role of female sex hormones in

the inflammatory response of females to BD, indicating that E2 may be a significant element in the improvement of donor treatment. Our results suggest that E2 exerts a direct control, decreasing the release of chemokines and restraining the cell traffic into the lungs. Considering these factors, there is a therapeutic potential of E2 that should be considered to improve donor lung quality and be further explored.

ACKNOWLEDGMENTS

Luiz Felipe Pinho Moreira is a fellow researcher of the CNPq. This study was presented at the 18th Congress of the European Society for Organ Transplantation, September 2017, Barcelona, Spain; American Thoracic Society 2018 International Conference, 2018, San Diego, United States; 19th Congress of the European Society for Organ Transplantation, September 2019, Copenhagen, Denmark.

REFERENCES

1. Miñambres E, Coll E, Duerto J, et al. Effect of an intensive lung donor-management protocol on lung transplantation outcomes. *J Heart Lung Transplant*. 2014;33:178–184.
2. Zitur LJ, Chlebeck PJ, Odorico SK, et al. Brain death enhances activation of the innate immune system and leads to reduced renal metabolic gene expression. *Transplantation*. 2019;103:1821–1833.

3. Avlonitis VS, Fisher AJ, Kirby JA, et al. Pulmonary transplantation: the role of brain death in donor lung injury. *Transplantation*. 2003;75:1928–1933.
4. Anwar ASMT, Lee JM. Medical management of brain-dead organ donors. *Acute Crit Care*. 2019;34:14–29.
5. International Society of Heart and Lung Transplantation Registry, Sato M, Gutierrez C, et al. The effect of gender combinations on outcome in human lung transplantation: the International Society of Heart and Lung Transplantation Registry experience. *J Heart Lung Transplant*. 2006;25:634–637.
6. Demir A, Coosemans W, Decaluwé H, et al. Donor-recipient matching in lung transplantation: which variables are important? *Eur J Cardiothorac Surg*. 2015;47:974–983.
7. Breithaupt-Faloppa AC, Ferreira SG, Kudo GK, et al. Sex-related differences in lung inflammation after brain death. *J Surg Res*. 2016;200:714–721.
8. Castardo-de-Paula JC, de Campos BH, Amorim EDT, et al. Cardiovascular risk and the effect of nitric oxide synthase inhibition in female rats: the role of estrogen. *Exp Gerontol*. 2017;97:38–48.
9. Imahara SD, Jelacic S, Junker CE, et al. The influence of gender on human innate immunity. *Surgery*. 2005;138:275–282.
10. Fantozzi ET, Breithaupt-Faloppa AC, Ricardo-da-Silva FY, et al. Estradiol mediates the long-lasting lung inflammation induced by intestinal ischemia and reperfusion. *J Surg Res*. 2018;221:1–7.
11. Schmidt EP, Yang Y, Janssen WJ, et al. The pulmonary endothelial glycocalyx regulates neutrophil adhesion and lung injury during experimental sepsis. *Nat Med*. 2012;18:1217–1223.
12. Baatz H, Steinbauer M, Harris AG, et al. Kinetics of white blood cell staining by intravascular administration of rhodamine 6G. *Int J Microcirc Clin Exp*. 1995;15:85–91.
13. Breithaupt-Faloppa AC, Breithaupt-Faloppa AC, Fantozzi ET, et al. Protective effect of estradiol on acute lung inflammation induced by an intestinal ischemic insult is dependent on nitric oxide. *Shock*. 2013;40:203–209.
14. Rostron AJ, Avlonitis VS, Cork DM, et al. Hemodynamic resuscitation with arginine vasopressin reduces lung injury after brain death in the transplant donor. *Transplantation*. 2008;85:597–606.
15. Bonnano Abib ALdO, Correia CdJ, Armstrong-Jr R, et al. The influence of female sex hormones on lung inflammation after brain death—an experimental study. *Transpl Int*. 2020;33:279–287.
16. Armstrong-Jr R, Ricardo-da-Silva FY, de Jesus Correia C, et al. Treatment with 17 β -estradiol protects donor heart against brain death effects in female rat. *Transpl Int*. 2020.
17. Suzuki H, Suematsu M, Miura S, et al. Rat CINC/gro: a novel mediator for locomotive and secretagogue activation of neutrophils in vivo. *J Leukoc Biol*. 1994;55:652–657.
18. Wolpe SD, Cerami A. Macrophage inflammatory proteins 1 and 2: members of a novel superfamily of cytokines. *FASEB J*. 1989;3:2565–2573.
19. Ward PA. Recruitment of inflammatory cells into lung: roles of cytokines, adhesion molecules, and complement. *J Lab Clin Med*. 1997;129:400–404.
20. Bonello N, Norman RJ. Soluble adhesion molecules in serum throughout the menstrual cycle. *Hum Reprod*. 2002;17:2272–2278.
21. Hou X, Pei F. Estradiol inhibits cytokine-induced expression of VCAM-1 and ICAM-1 in cultured human endothelial cells via AMPK/PPAR α Activation. *Cell Biochem Biophys*. 2015;72:709–717.
22. Yu HP, Hsieh YC, Suzuki T, et al. Salutary effects of estrogen receptor-beta agonist on lung injury after trauma-hemorrhage. *Am J Physiol Lung Cell Mol Physiol*. 2006;290:L1004–L1009.
23. Kang KP, Lee JE, Lee AS, et al. Effect of gender differences on the regulation of renal ischemia-reperfusion-induced inflammation in mice. *Mol Med Rep*. 2014;9:2061–2068.
24. Lentsch AB, Ward PA. Regulation of experimental lung inflammation. *Respir Physiol*. 2001;128:17–22.
25. Kim H, Hwang J-S, Woo C-H, et al. TNF- α -induced up-regulation of intercellular adhesion molecule-1 is regulated by a Rac-ROS-dependent cascade in human airway epithelial cells. *Exp Mol Med*. 2008;40:167–175.
26. Lin CC, Lee CW, Chu TH, et al. Transactivation of Src, PDGF receptor, and Akt is involved in IL-1 β -induced ICAM-1 expression in A549 cells. *J Cell Physiol*. 2007;211:771–780.
27. Tetley TD. New perspectives on basic mechanisms in lung disease. 6. Proteinase imbalance: its role in lung disease. *Thorax*. 1993;48:560–565.
28. Partridge CA, Jeffrey JJ, Malik AB. A 96-kDa gelatinase induced by TNF- α contributes to increased microvascular endothelial permeability. *Am J Physiol*. 1993;265(5 Pt 1):L438–L447.
29. Masure P, Proost P, Van Damme J, et al. Purification and identification of 91-kDa neutrophil gelatinase release by the activating peptide interleukin-8. *Eur J Biochem*. 1991;198:391–398.
30. Shapiro SD, Senior RM. Matrix metalloproteinases. Matrix degradation and more. *Am J Respir Cell Mol Biol*. 1999;20:1100–1102.
31. Lee JY, Choi HY, Na WH, et al. 17 β -estradiol inhibits MMP-9 and SUR1/TrpM4 expression and activation and thereby attenuates BSCB disruption/hemorrhage after spinal cord injury in male rats. *Endocrinology*. 2015;156:1838–1850.
32. Na W, Lee JY, Kim WS, et al. 17 β -estradiol ameliorates tight junction disruption via repression of MMP transcription. *Mol Endocrinol*. 2015;29:1347–1361.
33. Boughton-Smith NK, Evans SM, Hawkey CJ, et al. Nitric oxide synthase activity in ulcerative colitis and Crohn's disease. *Lancet*. 1993;342:338–340.
34. Nathan C. Nitric oxide as a secretory product of mammalian cells. *Faseb J*. 1992;6:3051–3064.
35. Cuzzocrea S, Chatterjee PK, Mazzon E, et al. Role of induced nitric oxide in the initiation of the inflammatory response after postischemic injury. *Shock*. 2002;18:169–176.
36. Ferreira SG, Armstrong-Jr R, Kudo GK, et al. Differential effects of brain death on rat microcirculation and intestinal inflammation: female versus male. *Inflammation*. 2018;41:1488–1497.
37. Perretti M, Ahluwalia A. The microcirculation and inflammation: site of action for glucocorticoids. *Microcirculation*. 2000;7:147–161.
38. Kalil B, Leite CM, Carvalho-Lima M, et al. Role of sex steroids in progesterone and corticosterone response to acute restraint stress in rats: sex differences. *Stress*. 2013;16:452–460.
39. Couse JF, Lindzey J, Grandien K, et al. Tissue distribution and quantitative analysis of estrogen receptor-alpha (ERalpha) and estrogen receptor-beta (ERbeta) messenger ribonucleic acid in the wild-type and ERalpha-knockout mouse. *Endocrinology*. 1997;138:4613–4621.
40. Vegeto E, Cuzzocrea S, Crisafulli C, et al. Estrogen receptor-alpha as a drug target candidate for preventing lung inflammation. *Endocrinology*. 2010;151:174–184.
41. Moriarty K, Kim KH, Bender JR. Minireview: estrogen receptor-mediated rapid signaling. *Endocrinology*. 2006;147:5557–5563.