University of Groningen

Diuretic therapy as prognostic enrichment factor for clinical trials in patients with heart failure with reduced ejection fraction

Coiro, Stefano; Girerd, Nicolas; McMurray, John J. V.; Pitt, Bertram; Swedberg, Karl; van Veldhuisen, Dirk J.; Lamiral, Zohra; Rossignol, Patrick; Zannad, Faiez

Published in:
Clinical Research in Cardiology

DOI:
[10.1007/s00392-021-01851-w](https://doi.org/10.1007/s00392-021-01851-w)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Coiro, S., Girerd, N., McMurray, J. J. V., Pitt, B., Swedberg, K., van Veldhuisen, D. J., Lamiral, Z., Rossignol, P., & Zannad, F. (2021). Diuretic therapy as prognostic enrichment factor for clinical trials in patients with heart failure with reduced ejection fraction. *Clinical Research in Cardiology*, 110(8), 1308-1320. <https://doi.org/10.1007/s00392-021-01851-w>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Diuretic therapy as prognostic enrichment factor for clinical trials in patients with heart failure with reduced ejection fraction

Stefano Coiro^{1,2} · Nicolas Girerd^{1,3} · John J. V. McMurray⁴ · Bertram Pitt⁵ · Karl Swedberg⁶ · Dirk J. van Veldhuisen⁷ · Zohra Lamiral^{1,3} · Patrick Rossignol^{1,3} · Faiez Zannad^{1,3}

Received: 10 November 2020 / Accepted: 26 March 2021
© Springer-Verlag GmbH Germany, part of Springer Nature 2021

Abstract

Background Loop diuretics are the mainstay of congestion treatment in patients with heart failure (HF). We assessed the association between baseline loop diuretic use and outcome. We also compared the increment in risk related to diuretic dose with conventional prognostic enrichment criteria used in the EMPHASIS-HF trial, which included patients with systolic HF and mild symptoms, such as prior hospitalization and elevated natriuretic peptides.

Methods Individual analyses were performed according to baseline loop diuretic usage (furosemide-equivalent dose > 40 mg, 1–40 mg, and no furosemide), and according to enrichment criteria adopted in the trial [i.e. recent hospitalization (< 30 days or 30 to 180 days prior to randomization) due to HF or other cardiovascular cause, or elevated natriuretic peptides]. The primary endpoint was a composite of cardiovascular death or HF hospitalization.

Results Loop diuretic usage at baseline (HR for > 40 mg furosemide-equivalent dose = 3.16, 95% CI 2.43–4.11; HR for 1–40 mg = 2.06, 95% CI 1.60–2.65) was significantly associated with a higher risk for the primary endpoint in a stepwise manner when compared to no baseline loop diuretic usage. In contrast, the differences in outcome rates were more modest when using history of hospitalization and/or BNP: all HR ranged from 1 (reference, non-HF related CV hospitalization > 30 days) to 2.04 (HF hospitalization < 30 days). The effect of eplerenone on the primary endpoint was consistent across subgroups in both analyses (P for interaction ≥ 0.2 for all).

Conclusions Loop diuretic usage (especially at doses > 40 mg) identified patients at higher risk than history of HF hospitalization and/or high BNP blood concentrations.

✉ Patrick Rossignol
p.rossignol@chru-nancy.fr

Stefano Coiro
stefano.coiro@hotmail.it

Nicolas Girerd
n.girerd@chru-nancy.fr

John J. V. McMurray
john.mcmurray@glasgow.ac.uk

Bertram Pitt
bpitt@med.umich.edu

Karl Swedberg
karl.swedberg@gu.se

Dirk J. van Veldhuisen
d.j.van.veldhuisen@umcg.nl

Zohra Lamiral
z.lamiral@chru-nancy.fr

Faiez Zannad
f.zannad@chru-nancy.fr

¹ Université de Lorraine, Centre D'Investigations Cliniques-INSERM CHRU de Nancy, Institut Lorrain du Coeur Et Des Vaisseaux Louis Mathieu, 4 rue du Morvan, 54500 Vandoeuvre lès Nancy, France

² Cardiology Department, Santa Maria della Misericordia University Hospital, Perugia, Italy

³ INI-CRCT (Cardiovascular and Renal Clinical Trialists) F-CRIN Network, Nancy, France

⁴ The British Heart Foundation Cardiovascular Research Centre, University of Glasgow, Glasgow, UK

⁵ University of Michigan School of Medicine, Ann Arbor, MI, USA

⁶ Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden

⁷ University of Groningen, University Medical Center Groningen, Groningen, The Netherlands

Graphic abstract

Diuretic therapy as prognostic enrichment factor for clinical trials in patients with heart failure with reduced ejection fraction

Aims

Compare the increment in risk related to **diuretic dose vs. conventional prognostic enrichment criteria** (prior hospitalization and elevated natriuretic peptides) used in the Eplerenone in Patients With Systolic Heart Failure and Mild Symptoms (EMPHASIS-HF).

Primary endpoint

Composite of CV death or HF hospitalization.

Association of loop diuretic dose and conventional prognostic enrichment criteria with CV death or HF hospitalization

Loop diuretic usage at baseline is more associated with the primary endpoint (HR 2.0 to 3.2) than history of hospitalization and/or BNP: (HR ranging from 1.3 to 2)

Interplay of loop diuretic dose and thiazide use with regards to CV death or HF hospitalization

Thiazide use does not seem to modify the risk of adverse outcome conferred by loop diuretic alone

Effect of eplerenone according to diuretic use

Eplerenone effect is overall consistent across subgroups. The estimated effect in patients without diuretics was neutral (HR 1.15)

Loop diuretic usage identified patients at higher risk than history of HF hospitalization and/or high BNP blood concentrations.

Including diuretic dose as an additional enrichment criterion can be useful in designing future HF trials.

Keywords Heart failure with reduced ejection fraction · Eplerenone · Loop diuretics · Heart failure hospitalization · Prognostic enrichment criteria

Abbreviations

HF	Heart failure
LVEF	Left ventricular ejection fraction
BNP	B-type natriuretic peptide and
NT-proBNP	N-terminal pro-BNP
EMPHASIS-HF	Eplerenone in mild patients hospitaliza- tion and survival study in heart failure

HFrEF	HF with reduced ejection fraction
HR	Hazard ratio
95% CI	95% Confidence interval
CHARM	Candesartan in heart failure: reduction in mortality and morbidity
ReBIC	Rede Brasileira de Estudos em Insu- ficiência Cardíaca

Introduction

In event-driven trials in heart failure (HF), it has been common to enroll patients at heightened risk of disease-related outcomes by enriching for selected baseline characteristics associated with these outcomes e.g. demographics (such as age), medical history (e.g. of prior HF hospitalization) and laboratory data (e.g. natriuretic peptide blood concentrations). Prognostic enrichment strategies aims at identifying patient subgroups with high event-rate in the course of the clinical study; those procedures considerably reduce the sample sizes needed to show a treatment effect, also increasing the power of the study [1].

Hospitalization for HF is associated with high event-rates in terms of mortality and HF readmissions, independently of left ventricular ejection fraction (LVEF) [2, 3]; additionally, an inverse relationship between time from HF hospitalization to randomization and event-rates was also observed.

Higher concentrations of B-type natriuretic peptide (BNP) and N-terminal pro-BNP (NT-proBNP) [4] are associated with higher rates of fatal and non-fatal events during follow-up [5, 6], with a steeper increase in cardiovascular compared to non-cardiovascular events [7]. On the other hand, there is no agreed threshold concentration for using these peptides in this way and some comorbidities (e.g. atrial fibrillation) also influence their concentration [7].

Diuretics (especially loop diuretics) play a pivotal role in relieving congestion and related symptoms in patients with HF [8], with a class I recommendation for this indication according to the latest HF guidelines [9, 10]. However, many studies have demonstrated that loop diuretic use, especially at high doses, is independently associated with reduced survival in HF with reduced ejection fraction (HFrEF) patients [11–17]. Thus, higher loop diuretic dosages identify patients with HF at particularly high risk of mortality and dose of diuretic could be considered as another approach to augmenting event rates in clinical trials. However diuretic dose has not been compared to alternative approaches to enriching for risk (such as requiring history of hospitalization/elevated natriuretic peptides).

In the present study we examined the association between loop diuretic use (and diuretic dose) and outcomes and compared these to other current enrichment criteria, using data from the Eplerenone in Mild Patients Hospitalization and Survival Study in Heart Failure (EMPHASIS-HF) [18].

Methods

Study design and participants

The design, patient eligibility criteria, study procedures and main results of EMPHASIS-HF have been reported [18]. In brief, in this randomized double-blind trial, patients with NYHA functional class II HF and LVEF no greater than 30% (or, if > 30 to 35%, a QRS duration of > 130 ms on electrocardiography) were randomly assigned to receive eplerenone (up to 50 mg daily) or placebo, in addition to recommended therapy.

Enrollment criteria in EMPHASIS-HF

In EMPHASIS-HF, all patients either had to have a hospitalization for a cardiovascular reason within 6 months or documentation of elevated plasma natriuretic peptide concentrations (i.e. BNP concentration of at least 250 pg/ml or NT-proBNP of at least 500 pg/ml for males and 750 pg/ml for females, as assessed within 15 days of randomization) [18]. Cardiovascular hospitalization was defined as follows: heart failure (first or subsequent HF hospitalization), acute myocardial infarction, angina pectoris (unstable), cardiac arrhythmia (atrial fibrillation, atrial flutter, supraventricular arrhythmias, or ventricular arrhythmias), stroke/cerebral vascular accident, or other cardiovascular causes (e.g. hypotension, peripheral vascular disease).

Assessment of diuretic use

Use of diuretics was considered part of standard therapy in EMPHASIS-HF, if clinically indicated to minimize fluid retention [19]. Drug name and dose were collected at baseline and during follow-up on case report forms. Furosemide-equivalent doses were calculated, with bumetanide 1 mg or torsemide 20 mg considered equivalent to 40 mg of furosemide [20].

Study outcomes

The primary outcome in the trial was the composite of time to first occurrence of death from cardiovascular causes or hospitalization for HF [18]. Secondary outcomes were the individual components of the primary outcome, all-cause death, cardiovascular hospitalization, and the composite of death or hospitalization for a cardiovascular reason. Adjudication of the outcomes was carried out by an independent committee according to pre-specified criteria [18].

Statistical analysis

For the present analysis, patients were grouped into 3 subgroups according to the main enrichment criterion met at baseline: (i) HF-related or, (ii) non-HF related (i.e. due to other cardiovascular causes) prior cardiovascular hospitalization, (iii) elevated natriuretic peptide blood concentrations. Timing of prior hospitalization for HF or for other cardiovascular causes before enrollment (i.e. within 6 months according to inclusion criteria) was categorized as follows: (i) < 30 days or (ii) 30–180 days prior to randomization. Additionally, patients were grouped according to furosemide-equivalent dose as follows: (i) no baseline loop diuretic, (ii) furosemide-equivalent 1–40 mg, (iii) furosemide-equivalent > 40 mg [16, 21]. Patients were also grouped according to loop and thiazide diuretic use [i.e. (1) no baseline diuretic, (2) thiazide use, (3) furosemide 1–40 mg, (4) furosemide 1–40 mg + thiazide use, (5) furosemide > 40 mg, and (6) furosemide > 40 mg + thiazide use], and to additional furosemide dosage subgroups (i.e. no loop diuretic, 1–20 mg, 21–40 mg, 41–80, and > 80 mg).

Continuous variables are expressed as mean ($\pm$ standard deviation) or median (25–75th percentiles); categorical variables as numbers and percentages. Baseline characteristics were defined in the whole cohort according to the respective enrichment criteria at baseline (i.e. prior HF or cardiovascular hospitalization, elevated natriuretic peptides), and furosemide-equivalent dose. Characteristics were compared among enrichment criteria subgroups, between timing categories within hospitalization subgroups, and among furosemide-equivalent subgroups. Additionally, proportions of furosemide-equivalent dose categories were defined according to enrichment criteria at baseline and their timing. Comparisons of baseline characteristics were performed using Student's *t* test, Mann–Whitney, ANOVA, and chi-square test for trend, as appropriate.

Risk probabilities for the primary outcome according to enrichment criteria, furosemide-equivalent categories, and furosemide-thiazide combinations were calculated in distinct analyses using the Kaplan–Meier method and plotted as survival curves; respective associations with the primary outcome were assessed by univariable Cox proportional hazards models and the subgroup with the lowest event rate were taken as reference. Additionally, associations of enrichment criteria and loop diuretic use with primary and secondary outcomes were also assessed. Multivariable Cox analyses were also performed. Multicollinearity was checked using R function “check_multicollinearity” from Performance package, and obtained variance inflation factors indicated no significant collinear relationship. The treatment effect of eplerenone according to enrichment criteria (and their timing), furosemide-equivalent dose, and loop-thiazide diuretic combinations on primary and secondary outcomes were

assessed in separate analyses by introducing an interaction term within univariable models. Harrell's C-statistics for time-to-event data were used to assess the ability of furosemide to predict the primary outcome, in the total cohort, and in subgroups according to study enrichment criteria.

Associations were expressed as hazard ratios and their 95% confidence interval (HR 95% CI). A *P*-value < 0.05 was considered statistically significant. Statistical analyses were performed using SAS software, version 9.4 (SAS Institute, Cary, North Carolina) the R software (R Foundation for Statistical Computing).

Results

Patient characteristics

Of the 2737 patients randomized in EMPHASIS-HF, 2722 (99.5%) and 2715 (99.2%) had information available for the prespecified enrichment criteria and furosemide-equivalent dose, respectively, at baseline. The main baseline characteristics are outlined in Tables 1 and Table 2.

The overlap between furosemide subgroups and selected prognostic enrichment criteria are illustrated in Fig. S1 in Supplementary material.

When compared to patients with a prior hospitalization (HF-related or for other cardiovascular causes), the subgroup enrolled on the basis of elevated natriuretic peptide blood concentrations were more likely to be male, and had a worse clinical profile, including lower diastolic blood pressure and LVEF at randomization (all *P* values < 0.03) (Table 1). Patients in the subgroup enrolled on the basis of a prior HF hospitalization were more likely to be on a loop diuretic (furosemide-equivalent dose 1–40, > 40 mg, and > 40 mg + thiazide/thiazide-like diuretic), and less likely to be on a beta-blocker compared to the other patients (all *P* values < 0.01). Finally, the subgroup with a prior non-HF related cardiovascular hospitalization was more likely to have HF of an ischemic etiology, to be on an thiazide/thiazide-like diuretic, or to have worse renal function as compared to other subgroups (all *P* values < 0.03).

In the subgroup with a prior HF hospitalization, patients with HF hospitalization < 30 days were more likely to have HF due to an ischemic cause, to be on an angiotensin-converting enzyme inhibitors, an angiotensin II receptor blockers or a thiazide/thiazide-like diuretic, to have lower systolic blood pressure and higher LVEF as compared to those with a less recent HF hospitalization; they were also less likely to be on a loop diuretic (furosemide-equivalent dose 1–40 mg) compared to the other HF hospitalization subgroup (all *P* values < 0.04). There were few imbalances in baseline risk

Table 1 Patient characteristics according to enrichment criteria (history of HFH within 30 days or between 30 and 180 days prior to randomization, history of CVH within 30 days or between 30 and 180 days prior to randomization, and elevated natriuretic peptide concentrations)

Characteristic	History of HFH (1)			History of non-HF related CVH (2)			Elevated natriuretic peptides (3)	P-value for trend (1) vs. (2) vs. (3)
	Within 180 days (n = 1496)		P (1 b vs. 1 a)	Within 30 days (n = 689) (1 a)		P (2 a vs. 2 b)		
	Within 180 days (n = 1496)	Between 30 and 180 (n = 807) (1 b)		Within 30 days (n = 246) (2 a)	Between 30 and 180 (n = 596) (2 b)			
Age (years)	68.7 ± 7.7	68.8 ± 7.3	0.73	68.7 ± 8.0	67.8 ± 6.8	68.7 ± 7.6	68.8 ± 8.1	0.68
Male sex	1141 (76.3%)	535 (77.6%)	0.25	606 (75.1%)	185 (75.2%)	473 (79.4%)	318 (82.8%)	0.022
SBP (mmHg)	124 ± 17	122 ± 16	0.001	125 ± 17	125 ± 16	125 ± 17	124 ± 1	0.11
DBP (mmHg)	75 ± 10	75 ± 10	0.92	75 ± 10	75 ± 10	75 ± 10	72 ± 11	<0.0001
HR (bpm)	71 (64–80)	7 (64–80)	0.18	70 (64–80)	70 (63–79)	70 (62–78)	68 (60–76)	<0.0001
LVEF (%)	27 (23–30)	27 (24–30)	0.01	26 (23–30)	28 (25–30)	28 (25–30)	25 (21–30)	<0.0001
eGFR < 60ml/min/1.73 m ²	979 (65.4%)	433 (62.8%)	0.053	546 (67.6%)	165 (67.1%)	433 (72.7%)	238 (62.0%)	0.002
Non ischemic HF	556 (37.3%)	218 (31.6%)	<0.0001	338 (42.1%)	52 (21.1%)	103 (17.3%)	128 (33.3%)	<0.0001
Ischemic HF	936 (62.7%)	471 (68.4%)	<0.0001	465 (57.9%)	194 (78.9%)	493 (82.7%)	256 (66.7%)	<0.0001
BMI > 30 kg/m ²	390 (26.2%)	173 (25.2%)	0.42	217 (27.1%)	78 (31.7%)	158 (26.7%)	109 (28.6%)	0.47
Previous MI	636 (42.5%)	314 (45.6%)	0.027	322 (39.9%)	147 (59.8%)	386 (64.8%)	206 (53.6%)	<0.0001
Previous HF Hosp.	890 (59.5%)	409 (59.4%)	0.92	481 (59.6%)	136 (55.3%)	225 (37.8%)	180 (46.9%)	<0.0001
Diabetes	507 (33.9%)	232 (33.7%)	0.87	275 (34.1%)	62 (25.2%)	165 (27.7%)	118 (30.7%)	0.002
ICD	94 (6.3%)	40 (5.8%)	0.48	54 (6.7%)	33 (13.4%)	124 (20.8%)	111 (28.9%)	<0.0001
Pacemaker	165 (11.0%)	78 (11.3%)	0.74	87 (10.8%)	28 (11.4%)	108 (18.1%)	92 (24.0%)	<0.0001
Thiazide/thiazide like diuretic	275 (18.4%)	147 (21.3%)	0.006	128 (15.9%)	61 (24.8%)	121 (20.3%)	59 (15.4%)	0.024
ACE inhibitor	1163 (78.2%)	546 (79.7%)	0.2	617 (76.9%)	212 (86.5%)	458 (77.2%)	279 (73.0%)	0.025
ARB	276 (18.6%)	125 (18.2%)	0.77	151 (18.8%)	28 (11.4%)	121 (20.4%)	99 (25.9%)	0.002
ACE inhibitor or ARB	1401 (94.2%)	655 (95.6%)	0.032	746 (93.0%)	236 (96.3%)	553 (93.3%)	354 (92.7%)	0.51
Beta-blocker	1264 (85.0%)	579 (84.5%)	0.63	685 (85.4%)	219 (89.4%)	532 (89.7%)	345 (90.3%)	0.0009
No furosemide	221 (14.9%)	119 (17.4%)	0.013	102 (12.8%)	91 (37.3%)	226 (38.0%)	102 (26.9%)	<0.0001
Furosemide 1–40 mg	824 (55.6%)	356 (52.0%)	0.01	468 (58.6%)	116 (47.5%)	284 (47.8%)	181 (47.8%)	0.0003
Furosemide 1–40 mg + thiazide	103 (7.0%)	49 (7.2%)	0.77	54 (6.8%)	17 (7.0%)	29 (4.9%)	20 (5.3%)	0.26
Furosemide > 40 mg	438 (29.5%)	210 (30.7%)	0.38	228 (28.6%)	37 (15.2%)	84 (14.1%)	96 (25.3%)	<0.0001
Furosemide > 40 mg + thiazide	76 (5.1%)	42 (6.1%)	0.10	34 (4.3%)	7 (2.9%)	11 (1.9%)	15 (4.0%)	0.002

HF heart failure, CVH cardiovascular hospitalization, HFH heart failure hospitalization; BNP B-type natriuretic peptide, SBP systolic blood pressure, DBP diastolic blood pressure, HR heart rate, LVEF left ventricular ejection fraction, eGFR estimated glomerular filtration rate, HF heart failure, ICD implantable cardioverter defibrillator, BMI body mass index, MI myocardial infarction, ACE angiotensin converting enzyme, ARB angiotensin II receptor blockers

Table 2 Baseline patient characteristics according to furosemide-equivalent dose

Furosemide-equivalent dose	None		1–40 mg		> 40 mg		P-value for trend
Characteristic	N	m ± SD/median (Q1–Q3)/n(%) (n = 647)	n	m ± SD/median (Q1–Q3)/n(%) (n = 1411)	n	m ± SD/median (Q1–Q3)/n(%) (n = 657)	
Age (Years)	647	68.0 ± 7.4	1411	69.0 ± 7.8	657	68.5 ± 7.4	0.037
Male gender	647	507 (78.4%)	1411	1083 (76.8%)	657	519 (79.0%)	0.47
SBP (mmHg)	646	128 ± 16	1411	124 ± 17	657	121 ± 17	< 0.0001
DBP (mmHg)	646	76 ± 10	1411	75 ± 10	657	73 ± 10	< 0.0001
HR (bpm)	646	68.0 (62.0–76.0)	1410	70.0 (63.0–80.0)	657	72.0 (64.0–80.0)	< 0.0001
LVEF (%)	612	28.0 (25.0–30.0)	1342	27.0 (24.0–30.0)	614	25.0 (21.0–30.0)	< 0.0001
eGFR < 60 ml/min/1.73 m ²	647	513 (79.3%)	1409	926 (65.7%)	655	369 (56.3) %	< 0.0001
Non ischemic HF	646	160 (24.8%)	1409	454 (32.2%)	655	223 (34.0%)	0.0004
Ischemic HF	646	486 (75.2%)	1409	955 (67.8%)	655	432 (66.0%)	0.0004
BMI > 30 kg/m ²	647	180 (27.8%)	1401	349 (24.9%)	652	205 (31.4%)	0.008
Previous MI	647	362 (56.0%)	1411	709 (50.2%)	657	299 (45.5%)	0.0008
Previous HFH	647	248 (38.3%)	1411	755 (53.5%)	657	420 (63.9%)	< 0.0001
Diabetes	647	153 (23.6%)	1411	438 (31.0%)	657	262 (39.9%)	< 0.0001
ICD	647	89 (13.8%)	1411	162 (11.5%)	657	106 (16.1%)	0.012
Pacemaker	647	75 (11.6%)	1411	208 (14.7%)	657	108 (16.4%)	0.039
Thiazide/thiazide like diuretic	647	236 (36.5%)	1411	169 (12.0%)	657	110 (16.7%)	< 0.0001
ACE-inhibitor	631	500 (79.2%)	1411	1118 (79.2%)	657	487 (74.1%)	0.023
ARB	631	108 (17.1%)	1411	277 (19.6%)	657	141 (21.5%)	0.14
ACE-inhibitor or ARB	631	594 (94.1%)	1411	1337 (94.8%)	657	608 (92.5%)	0.14
Beta-blocker	631	553 (87.6%)	1411	1235 (87.5%)	657	567 (86.3%)	0.7
Non-HF CVH < = 180	647	317/647 (58.8%)	1411	400/1411 (32.7%)	657	121/657 (18.4%)	< 0.0001
Non-HF CVH 30–180	647	226/647 (34.9%)	1411	284/1411 (20.1%)	657	84/657 (12.8%)	0.93
Non-HF CVH < 30	647	91/647 (14.1%)	1411	116/1411 (8.2%)	657	37/657 (5.6%)	0.93
HFH < = 180	647	222/647 (34.3%)	1411	824/1411 (58.4%)	657	438/657 (66.7%)	< 0.0001
HFH 30–180	647	102/647 (15.8%)	1411	468/1411 (33.2%)	657	228/657 (34.7%)	0.013
HFH < 30	647	119/647 (18.4%)	1411	356/1411 (25.2%)	657	210/657 (32.0%)	0.013
BNP	647	102/647 (15.8%)	1411	182/1411 (12.9%)	657	96/657 (14.7%)	0.2

See Table 1 for abbreviations

factor relative to the timing of non-HF related cardiovascular hospitalization before randomization (Table 1).

Patients treated with a loop diuretic at baseline (especially those with furosemide-equivalent dose > 40 mg) were older, more likely to be obese, diabetic, and have an implantable device, but less likely to have an ischemic etiology or to be on a thiazide/thiazide-like diuretic, compared to patients not treated with a loop diuretic at baseline. Patients treated with a loop diuretic also had lower blood pressure and LVEF, and a higher heart rate, compared to those not treated with a loop diuretic (all *P* values < 0.04) (Table 2).

Patients with loop diuretic at baseline were more likely to have had a previous HF hospitalization when compared to patients not treated with loop diuretic [66.7% (438/657) for furosemide > 40 mg, and 58.4% (824/1411) for furosemide 1–40 mg, versus 34.3% (222/647) in patients without diuretics, *P*-value < 0.0001] (Table 2, Fig. S1 in Supplementary Material). With regard to non-HF related cardiovascular

hospitalization, the opposite trend was observed [18.4% (121/657) for furosemide > 40 mg, and 32.7% (400/1411) for furosemide 1–40 mg, versus 58.8% (317/647) in patients without diuretics, *P*-value < 0.0001]. Differences in terms of elevated natriuretic peptide at baseline were not statistically significant across furosemide subgroups (*P*-value = 0.2) (Table 2).

Impact of conventional prognostic enrichment criteria and loop diuretic use on outcomes

After 2 years of follow-up, the risk of cardiovascular death or HF hospitalization was highest among patients with a prior HF hospitalization: 28.5% [95% CI 24.7–32.1%] in those with hospitalization < 30 days and 27.5% (95% CI 23.9–30.9%) in those with hospitalization 30–180 days before randomization (Fig. 1a). Lower event rates were observed in patients enrolled using natriuretic peptide

Fig. 1 a–d Kaplan Meier curves for enrichment criteria—Survival curves for cardiovascular death or HF hospitalization according to enrichment criteria (non-HF related CVH within 30 days or 30–180 days prior to randomization, HFH within 30 days or 30–180 days prior to randomization or elevated natriuretic peptides;

panel **a**), to furosemide subgroups (3 subgroups, panel **b**), to furosemide-thiazide combination subgroups (panel **c**), and to 5 furosemide subgroups (panel **d**). *HF* heart failure, *CVH* cardiovascular hospitalization, *HFH* heart failure hospitalization

(24.0%, 95% CI 17.0–30.5%) or non-HF related cardiovascular hospitalization criteria [20.0% (95% CI 14.2–25.4%) and 14.8% (95% CI 11.5–17.9%) for hospitalization < 30 days and 30–180 days before randomization, respectively] (Fig. 1a). When taking patients randomized within 30–180 days of non-HF related cardiovascular hospitalization as reference, a prior HF hospitalization within 30 days (HR 2.04, 95% CI 1.60–2.62), or within 30–180 days (HR 1.82, 95% CI 1.42–2.33) were both

associated with significantly higher risks of the primary outcome (Fig. 2a).

Similarly, taking the subgroup without loop-diuretic at baseline as reference, both patients with a furosemide-equivalent dose > 40 mg (HR 3.16, 95% CI 2.43–4.11) and 1–40 mg (HR 2.06, 95% CI 1.60–2.65) displayed a significantly higher risk of the primary outcome (Figs. 1b and 2a). Importantly, patients with a furosemide-equivalent dose > 40 mg had a very high-risk for event (32.9%

Fig. 2 a, b Enrichment factors and risk of primary outcome—Association of enrichment criteria (CVH within 30 days or 30–180 days prior to randomization, HFH within 30 days or 30–180 days prior to randomization or elevated natriuretic peptides) and loop diuretic use (Panel a), and of furosemide-thiazide combination subgroups (Panel b) with rates of cardiovascular death or HF hospitalization (unadjusted hazard ratios). See Fig. 1 for abbreviations

at 2 years) which was comparable to or higher than that of patients with a HF hospitalization <30 days (28.5% at 2 years). Similar results were observed by multivariable analyses for both original study inclusion criteria and loop-diuretic subgroups (Fig. S2 in Supplementary material). Additionally, the prognostic accuracy of furosemide use as assessed by Harrell's C index is statically significant in the total cohort as well as in risk subgroups according to original study enrichment inclusion criteria, with superimposable results (Table 1 in Supplementary material).

When taking into account subgroups according to loop and thiazide diuretic combinations, thiazide/thiazide-like diuretic use does not seem to modify the risk of adverse outcome conferred by loop diuretic alone, for both furosemide >40 mg (thiazide = 32.7% vs. no thiazide 33.0%,

HR $\approx$ 3.3), and furosemide 1–40 mg (thiazide = 27.9% vs. no thiazide 24.4%, HR $\approx$ 2.2). The risk associated with thiazide diuretic use alone is comparable to that of patients taking no diuretics (Figs. 1c and 2b). Finally, a fair stepwise increase in the risk of primary outcome was observed when considering furosemide 1–20, 21–40, 41–80, and >80 mg subgroups (Fig. 1d).

Without taking into account timing of hospitalization or loop-diuretic dose, a prior HF hospitalization (non-HF related cardiovascular hospitalization subgroup as reference; HRs > 1.15, all P values < 0.001) and loop diuretic use (no baseline loop diuretic as reference; HRs > 1.70, all P values < 0.0001) were both significantly associated with the primary and all secondary outcomes (Fig. S3 in Supplementary material). Recent (versus less recent—considered

as reference) non-HF related cardiovascular or HF hospitalization were overall not significantly associated with worse prognosis. (Fig. S3 in Supplementary material).

Effect of eplerenone according to conventional prognostic enrichment criteria and baseline loop-diuretic use

According to enrichment criteria categories, the treatment effect of eplerenone on the primary outcome was significant in both patients randomized within 30 days of an HF hospitalization (HR 0.53, 95% CI 0.40–0.70) as well as in patients with natriuretic peptide criteria (HR 0.54, 95% CI 0.30–0.95) (Fig. 3a); no significant interaction between treatment effect of eplerenone and enrichment criteria was observed (interaction P -value = 0.20) (Fig. 3a).

Both patients with a furosemide-equivalent dose > 40 mg (HR 0.65, 95% CI 0.50–0.86) and 1–40 mg (HR 0.63, 95% CI 0.51–0.79) significantly benefitted from eplerenone

treatment with respect to the primary outcome; as for the enrichment criteria analysis, the interaction did not reach statistical significance (interaction P = 0.64) (Fig. 3a). Similar results were observed when considering loop-diuretic/thiazide diuretic combinations (interaction P = 0.51; Fig. 3b).

Overall, interaction analyses between treatment effect of eplerenone and enrichment criteria, loop diuretic (use and dosage), or timing of prior cardiovascular hospitalization (non-HF-related) were not significant for all studied primary and secondary outcomes (Table 2 in Supplementary materials).

Discussion

In this post hoc analysis of the EMPHASIS-HF study, we found that, among patients with HFrEF and mild symptoms, loop diuretics usage (especially at high doses) identified participants with a two–three fold greater risk

Fig. 3 a, b Effect of eplerenone according to enrichment criteria—Forest plot for treatment effect on cardiovascular death or hospitalization for heart failure according to enrichment criteria and loop diuretic use (Panel a), and to furosemide-thiazide combination subgroups (Panel b)

of cardiovascular death or HF hospitalization, which is a similar or greater increment in risk than that conferred by a well-established risk-enrichment criteria such as prior HF hospitalization. The higher risk related to loop diuretic use was consistent for all secondary endpoints, and the treatment effect of eplerenone was not influenced by this background diuretic use or dose.

Prognostic enrichment strategies in heart failure clinical trials

For event-driven studies, prognostic enrichment strategies usually involve identification of patients with a higher probability of having a disease-related endpoint. This approach is intended to permit a smaller sample size as a result of the higher event rate, but assumes that the proportional reduction in events with treatment is maintained [1]. These risk-enhancing variables usually include clinical and laboratory measurements, medical history, genomic and proteomic measurements.

The concept that a recent hospitalization for HF can identify a high-risk patient population is well-described [2, 3], and confirmed in recent HF studies [22]. Findings from the Candesartan in heart failure: reduction in mortality and morbidity (CHARM) program (enrolling 7599 HF patients with or without systolic dysfunction) showed that the magnitude of increased risk of cardiovascular death or HF hospitalization in HFrEF with a prior HF hospitalization before randomization was approximately 60%, and that the time interval between hospitalization and randomization was inversely associated with adverse outcome [2]. Another post hoc analysis on the same cohort taking into account HF hospitalizations occurring during study follow-up found similar results in terms of mortality [3].

Due to differences in statistical methodology between our study and the above post hoc analyses from the CHARM Program (i.e. different reference category, comparison among multiple enrichment criteria), a direct comparison cannot be made. However, even in a worst-case scenario as in the present study (i.e. a prior cardiovascular but not HF-related hospitalization as reference category), our results demonstrated that HF hospitalization within 6 months prior to randomization was associated with a 75% higher risk for the combined outcome, with an inverse relationship between timing of hospitalization and risk of adverse outcome (Fig. 3 in Supplementary material).

On the other hand, the assumption that a previous HF hospitalization would carry higher risk in HF trials was not confirmed in the TOPCAT trial, showing marked regional differences in terms of primary combined endpoint event-rates between patients randomized from Eastern Europe compared with those enrolled from the Americas (respectively, 14.7 vs. 2.4 per 100 patient-years) [23]. This suggest

that, although HF hospitalization is generally seen as an objective enrollment criterion, non-adjudicated qualifying hospitalization can lead to inclusion of different patient populations, and subsequent treatment effect heterogeneity [23].

With regard to eplerenone treatment effect, a previous post hoc analysis of EMPHASIS-HF showed that the beneficial effect of eplerenone on outcomes was unaffected by the timing of the qualifying cardiovascular hospitalization [24]. Similarly, we found no significant overall interaction between eplerenone treatment effect and enrichment criteria.

Diuretic therapy as enrichment factor in heart failure with reduced ejection fraction clinical trials

Loop diuretics are used for relieving congestive HF symptoms and signs, with 80% of HF patients being prescribed this medications [25]. Although diuretics have long been acknowledged as having positive effects on chronic HF patients, several retrospective studies have highlighted that their use, especially in high doses, is associated with adverse outcome [25]. As suggested by the current European Guidelines [10], diuretics might be temporarily discontinued in selected asymptomatic patients without signs of congestion. To this purpose, the Rede Brasileira de Estudos em Insuficiência Cardíaca (ReBIC trial), a multicentre double-blind randomized study which enrolled a cohort of 188 apparently euvolaemic HF outpatients (LVEF $\leq 45\%$) on stable low dose of furosemide (40 up to 80 mg/day), recently demonstrated that diuretic withdrawal do not significantly impact on the self-perception of dyspnea and use of additional diuretics up to 90 after randomization [26]. However, the study was underpowered to assess the effect of such a strategy on hard clinical endpoints, and larger trials with long-term follow-up are needed.

Despite growing data, the causal relationship between diuretics use and adverse outcome have been challenged on the grounds that a high diuretic dose more likely reflects the severity of the disease rather than a true risk factor among patients with HF and systolic dysfunction (i.e. “the sicker the patient, the higher the diuretic dose”) [27].

In this respect, Pellicori et al., in a cohort of 1190 HF outpatient with mixed LVEF, demonstrated that loop diuretics did not independently predict adverse outcome at long-term follow-up by multivariable analyses, but rather identifies patients with more advanced HF and level of congestion, which may account for their worse prognosis [28].

In any case, the primary aim of our study was not focused on adding further evidence on the prognostic impact of diuretic therapy (which was already demonstrated in several studies as reported above), but rather to directly compare the baseline risk associated with loop-diuretic therapy with that of other baseline factors usually adopted as prognostic

enrichment criteria (i.e. prior HF hospitalization or natriuretic peptide blood concentrations).

Although diuretic dose may be a powerful, simple prognostic enrichment criteria in HFrEF trials, care should be taken to avoid investigators deliberately prescribing the dose required to fit the inclusion criteria, or at least reporting that they have done so. This is also the case for history of HF hospitalization as highlighted above. This may potentially be mitigated by a policy of quality control and data ascertainment. Our results demonstrate that the risk associated with loop-diuretic at least matches or even exceeds that associated with history of HF hospitalization.

On the other hand, it should be noted that “greater risk” does not equal “greater modifiable risk”, and a proportion of patients with more advanced HF may be refractory to the benefits of a particular therapy. Clinical trialists should keep from enrolling patients with very high event rates, but rather target intermediate risk populations (the so-called “sweet-spot”), associated with larger treatment effects [6, 29].

Finally, we showed how the stepwise increase in the risk of adverse outcome associated with furosemide use appears to be unaffected by the concomitant use of thiazide; indeed, the risk associated with a given dosage of furosemide (i.e. furosemide > 40 and 1–40 mg) is quite comparable to that of respective subgroups contemporarily taking thiazide diuretics (Fig. 1c).

Of note, the risk-rate of adverse outcome associated with thiazide use alone is comparable to that of patients without diuretics at baseline (about 12.0% at 2-year follow-up); that risk is, respectively, about half and one third of that observed in subgroups with furosemide 1–40 and > 40 mg (Fig. 1c). This HFrEF subgroup, not taking loop diuretics regularly (or taking only thiazide diuretics) represents about 25% of our total cohort, and similar proportions are observed in more contemporary study [30]. Further studies are needed to assess whether is still suitable to include that low-risk subgroup in forthcoming HFrEF trial design.

Limitations

This is a post hoc analysis of a randomized clinical trial that was not powered to identify differences in terms of outcomes according loop diuretic use. Additionally, in EMPHASIS-HF, elevated natriuretic peptides were requested only in the absence of a recent cardiovascular hospitalization [18], otherwise their concentration was not routinely assessed.

Conclusions

Dose of loop diuretics (especially at high doses) identifies a subset of HF patients with high risk of adverse clinical outcomes, a risk similar or greater in magnitude than the risk

conferred by other well-established prognostic enrichment factors, namely HF hospitalization and raised natriuretic peptide concentrations. We believe that including diuretic dose as an additional enrichment criterion can be useful in designing future HF trials.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00392-021-01851-w>.

Acknowledgements The EMPHASIS HF trial was sponsored by Pfizer. SC, NG, PR, FZ are supported by the RHU Fight- HF, a public grant overseen by the French National Research Agency (ANR) as part of the second “Investissements d’Avenir” program (reference: ANR-15-RHUS-0004).

Funding None.

Declarations

Conflict of interest Dr. Girerd received board fees from Novartis and honoraria from Servier. Dr. Rossignol received personal fees (consulting) from Novartis, Relypsa, AstraZeneca, Grünenthal, Stealth Peptides, Fresenius, Vifor Fresenius Medical Care Renal Pharma, Vifor, and CTMA; lecture fees from Bayer and CVRx; and is cofounder of CardioRenal. Dr Zannad received steering committee or advisory board fees from Amgen, AstraZeneca, Bayer, Boehringer, Boston Scientific, Cardior, CVRx, Janssen, Livanova, Merck, Mundipharma, Novartis, NovoNordisk and Vifor Fresenius. Dr Pitt received personal fees from Bayer, KDP pharmaceuticals, Astra Zeneca, Sanofi, Relypsa/Vifor, scPharmaceuticals, stealth peptides, tricida, personal fees from G3 pharmaceuticals. Dr McMurray received fees for his consulting or trial committee work with Abbvie, Amgen, AstraZeneca/Medimmune, Bayer, Bristol Myers Squibb, DalCor, GlaxoSmithKline, Merck, Novartis, Resverlogix, Sanofi-Aventis and Stealth Therapeutics. Dr Swedberg received personal fees from Novartis, AstraZeneca, Pfizer, and Novartis. Dr. Van Veldhuisen received board membership fees or travel expenses from Novartis, Corvia Medical, ARCA Biopharma and Johnson and Johnson. All other authors have no conflicts of interest to declare.

References

1. (FDA). FaDA (2019) Enrichment strategies for clinical trials to support determination of effectiveness of human drugs and biological products guidance for industry [Internet]. Silver Spring, MD: FDA. <https://www.fda.gov/media/121320/download>
2. Bello NA, Claggett B, Desai AS, McMurray JJ, Granger CB, Yusuf S, Swedberg K, Pfeffer MA, Solomon SD (2014) Influence of previous heart failure hospitalization on cardiovascular events in patients with reduced and preserved ejection fraction. *Circ Heart Fail* 7(4):590–595. <https://doi.org/10.1161/circheartfailure.113.001281>
3. Solomon SD, Dobson J, Pocock S, Skali H, McMurray JJ, Granger CB, Yusuf S, Swedberg K, Young JB, Michelson EL, Pfeffer MA (2007) Influence of nonfatal hospitalization for heart failure on subsequent mortality in patients with chronic heart failure. *Circulation* 116(13):1482–1487. <https://doi.org/10.1161/circulationaha.107.696906>
4. Ibrahim NE, Burnett JC Jr, Butler J, Camacho A, Felker GM, Fiuzat M, O’Connor C, Solomon SD, Vaduganathan M, Zile MR, Januzzi JL Jr (2020) Natriuretic peptides as inclusion criteria in

- clinical trials: a JACC heart failure position paper. *JACC Heart Fail*. <https://doi.org/10.1016/j.jchf.2019.12.010>
5. Hartmann F, Packer M, Coats AJ, Fowler MB, Krum H, Mohacs P, Rouleau JL, Tendera M, Castaigne A, Anker SD, Amann-Zalan I, Hoersch S, Katus HA (2004) Prognostic impact of plasma N-terminal pro-brain natriuretic peptide in severe chronic congestive heart failure: a substudy of the carvedilol prospective randomized cumulative survival (COPERNICUS) trial. *Circulation* 110(13):1780–1786. <https://doi.org/10.1161/01.Cir.0000143059.68996.A7>
 6. Anand IS, Rector TS, Cleland JG, Kuskowski M, McKelvie RS, Persson H, McMurray JJ, Zile MR, Komajda M, Massie BM, Carson PE (2011) Prognostic value of baseline plasma amino-terminal pro-brain natriuretic peptide and its interactions with irbesartan treatment effects in patients with heart failure and preserved ejection fraction: findings from the I-PRESERVE trial. *Circ Heart Fail* 4(5):569–577. <https://doi.org/10.1161/circheartfailure.111.962654>
 7. Savarese G, Orsini N, Hage C, Vedin O, Cosentino F, Rosano GMC, Dahlström U, Lund LH (2018) Utilizing NT-proBNP for eligibility and enrichment in trials in HFpEF, HFmrEF, and HFrEF. *JACC Heart Fail* 6(3):246–256. <https://doi.org/10.1016/j.jchf.2017.12.014>
 8. Faris RF, Flather M, Purcell H, Poole-Wilson PA, Coats AJS (2012) Diuretics for heart failure. *Cochrane Database Syst Rev*. <https://doi.org/10.1002/14651858.CD003838.pub3>
 9. Yancy CW, Jessup M, Bozkurt B, Butler J, Casey DE, Drazner MH, Fonarow GC, Geraci SA, Horwich T, Januzzi JL, Johnson MR, Kasper EK, Levy WC, Masoudi FA, McBride PE, McMurray JJV, Mitchell JE, Peterson PN, Riegel B, Sam F, Stevenson LW, Tang WHW, Tsai EJ, Wilkoff BL (2013) 2013 ACCF/AHA guideline for the management of heart failure. *Circulation* 128(16):e240–e327. <https://doi.org/10.1161/CIR.0b013e31829e8776>
 10. Ponikowski P, Voors AA, Anker SD, Bueno H, Cleland JG, Coats AJ, Falk V, Gonzalez-Juanatey JR, Harjola VP, Jankowska EA, Jessup M, Linde C, Nihoyannopoulos P, Parissis JT, Pieske B, Riley JP, Rosano GM, Ruilope LM, Ruschitzka F, Rutten FH, van der Meer P (2016) ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: the task force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC). Developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur J Heart Fail* 18(8):891–975. <https://doi.org/10.1002/ehf.592>
 11. Eshaghian S, Horwich TB, Fonarow GC (2006) Relation of loop diuretic dose to mortality in advanced heart failure. *Am J Cardiol* 97(12):1759–1764. <https://doi.org/10.1016/j.amjcard.2005.12.072>
 12. Martins J, Lourenco P, Araujo JP, Mascarenhas J, Lopes R, Azevedo A, Bettencourt P (2011) Prognostic implications of diuretic dose in chronic heart failure. *J Cardiovasc Pharmacol Ther* 16(2):185–191. <https://doi.org/10.1177/1074248410388807>
 13. Testani JM, Cappola TP, Brensinger CM, Shannon RP, Kimmel SE (2011) Interaction between loop diuretic-associated mortality and blood urea nitrogen concentration in chronic heart failure. *J Am Coll Cardiol* 58(4):375–382. <https://doi.org/10.1016/j.jacc.2011.01.052>
 14. Dini FL, Ghio S, Klersy C, Rossi A, Simioniuc A, Scelsi L, Genta FT, Ciccoira M, Tavazzi L, Temporelli PL (2013) Effects on survival of loop diuretic dosing in ambulatory patients with chronic heart failure using a propensity score analysis. *Int J Clin Pract* 67(7):656–664. <https://doi.org/10.1111/ijcp.12144>
 15. Cubbon RM, Woolston A, Adams B, Gale CP, Githorpe MS, Baxter PM, Kearney LC, Mercer B, Rajwani A, Batin PD, Kahn M, Sapsford RJ, Witte KK, Kearney MT (2014) Prospective development and validation of a model to predict heart failure hospitalisation. *Heart* 100(12):923–929. <https://doi.org/10.1136/heartjnl-2013-305294>
 16. Damman K, Kjekshus J, Wikstrand J, Cleland JG, Komajda M, Wedel H, Waagstein F, McMurray JJ (2016) Loop diuretics, renal function and clinical outcome in patients with heart failure and reduced ejection fraction. *Eur J Heart Fail* 18(3):328–336. <https://doi.org/10.1002/ehf.462>
 17. Ahmed A, Husain A, Love TE, Gambassi G, Dell'Italia LJ, Francis GS, Gheorghiade M, Allman RM, Meleth S, Bourge RC (2006) Heart failure, chronic diuretic use, and increase in mortality and hospitalization: an observational study using propensity score methods. *Eur Heart J* 27(12):1431–1439. <https://doi.org/10.1093/eurheartj/ehi890>
 18. Zannad F, McMurray JJV, Krum H, van Veldhuisen DJ, Swedberg K, Shi H, Vincent J, Pocock SJ, Pitt B (2010) Eplerenone in patients with systolic heart failure and mild symptoms. *N Engl J Med* 364(1):11–21. <https://doi.org/10.1056/NEJMoa1009492>
 19. Zannad F, McMurray JJ, Drexler H, Krum H, van Veldhuisen DJ, Swedberg K, Shi H, Vincent J, Pitt B (2010) Rationale and design of the eplerenone in mild patients hospitalization and survival study in heart failure (EMPHASIS-HF). *Eur J Heart Fail* 12(6):617–622. <https://doi.org/10.1093/eurjhf/hfq049>
 20. DeVore AD, Hasselblad V, Mentz RJ, O'Connor CM, Armstrong PW, McMurray JJ, Ezekowitz JA, Tang WH, Starling RC, Voors AA, Califf RM, Hernandez AF (2015) Loop diuretic dose adjustments after a hospitalization for heart failure: insights from ASCEND-HF. *Eur J Heart Fail* 17(3):340–346. <https://doi.org/10.1002/ehf.235>
 21. Ferreira JP, Eschalier R, Duarte K, Damman K, Gustafsson F, Schou M, Girerd N, Fay R, Tala S, Pitt B, Zannad F, Rossignol P (2020) Reduced diuretic dose in patients treated with eplerenone: data from the EPHEsus trial. *Circ Heart Fail* 13(5):e006597. <https://doi.org/10.1161/circheartfailure.119.006597>
 22. Vaduganathan M, Claggett BL, Desai AS, Anker SD, Perrone SV, Janssens S, Milicic D, Arango JL, Packer M, Shi VC, Lefkowitz MP, McMurray JJV, Solomon SD (2020) Prior heart failure hospitalization, clinical outcomes, and response to sacubitril/valsartan compared with valsartan in HFpEF. *J Am Coll Cardiol* 75(3):245–254. <https://doi.org/10.1016/j.jacc.2019.11.003>
 23. Pfeffer MA, Claggett B, Assmann SF, Boineau R, Anand IS, Clausell N, Desai AS, Diaz R, Fleg JL, Gordeev I, Heitner JF, Lewis EF, O'Meara E, Rouleau JL, Probstfield JL, Shaburishvili T, Shah SJ, Solomon SD, Sweitzer NK, McKinlay SM, Pitt B (2015) Regional variation in patients and outcomes in the treatment of preserved cardiac function heart failure with an aldosterone antagonist (TOPCAT) trial. *Circulation* 131(1):34–42. <https://doi.org/10.1161/circulationaha.114.013255>
 24. Girerd N, Collier T, Pocock S, Krum H, McMurray JJ, Swedberg K, Van Veldhuisen DJ, Vincent J, Pitt B, Zannad F (2015) Clinical benefits of eplerenone in patients with systolic heart failure and mild symptoms when initiated shortly after hospital discharge: analysis from the EMPHASIS-HF trial. *Eur Heart J* 36(34):2310–2317. <https://doi.org/10.1093/eurheartj/ehv273>
 25. Kapelios CJ, Malliaras K, Kaldara E, Vakrou S, Nanas JN (2018) Loop diuretics for chronic heart failure: a foe in disguise of a friend? *Eur Heart J Cardiovasc Pharmacother* 4(1):54–63. <https://doi.org/10.1093/ehjcvp/pvx020>
 26. Rohde LE, Rover MM, Figueiredo Neto JA, Danzmann LC, Bertoldi EG, Simões MV, Silvestre OM, Ribeiro ALP, Moura LZ, Beck-da-Silva L, Prado D, Sant'Anna RT, Bridi LH, Zimmerman A, Raupp da Rosa P, Biolo A (2019) Short-term diuretic withdrawal in stable outpatients with mild heart failure and no fluid retention receiving optimal therapy: a double-blind, multicentre, randomized trial. *Eur Heart J* 40(44):3605–3612. <https://doi.org/10.1093/eurheartj/ehz554>

27. Mielniczuk LM, Tsang SW, Desai AS, Nohria A, Lewis EF, Fang JC, Baughman KL, Stevenson LW, Givertz MM (2008) The association between high-dose diuretics and clinical stability in ambulatory chronic heart failure patients. *J Card Fail* 14(5):388–393. <https://doi.org/10.1016/j.cardfail.2008.01.015>
28. Pellicori P, Cleland JGF, Zhang J, Kallvikbacka-Bennett A, Urbinati A, Shah P, Kazmi S, Clark AL (2016) Cardiac dysfunction, congestion and loop diuretics: their relationship to prognosis in heart failure. *Cardiovasc Drugs Ther* 30(6):599–609. <https://doi.org/10.1007/s10557-016-6697-7>
29. Cleland JGF, Tavazzi L, Daubert JC, Tageldien A, Freemantle N (2009) Cardiac resynchronization therapy: are modern myths preventing appropriate use? *J Am Coll Cardiol* 53(7):608–611. <https://doi.org/10.1016/j.jacc.2008.10.040>
30. Vardeny O, Claggett B, Kachadourian J, Desai AS, Packer M, Rouleau J, Zile MR, Swedberg K, Lefkowitz M, Shi V, McMurray JJV, Solomon SD (2019) Reduced loop diuretic use in patients taking sacubitril/valsartan compared with enalapril: the PARADIGM-HF trial. *Eur J Heart Fail* 21(3):337–341. <https://doi.org/10.1002/ejhf.1402>