University of Groningen

Exercise capacity in patients with repaired Tetralogy of Fallot aged 6 to 63 years

Eshuis, Graziella; Hock, Julia; Marchie du Sarvaas, Gideon; van Duinen, Hiske; Neidenbach, Rhoia; van den Heuvel, Freek; Hillege, Hans; Berger, Rolf Mf; Hager, Alfred

Published in:
Heart

DOI:
[10.1136/heartjnl-2020-318928](https://doi.org/10.1136/heartjnl-2020-318928)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2022

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Eshuis, G., Hock, J., Marchie du Sarvaas, G., van Duinen, H., Neidenbach, R., van den Heuvel, F., Hillege, H., Berger, R. M., & Hager, A. (2022). Exercise capacity in patients with repaired Tetralogy of Fallot aged 6 to 63 years. *Heart*, 108(3), 186-193. <https://doi.org/10.1136/heartjnl-2020-318928>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Original research

Exercise capacity in patients with repaired Tetralogy of Fallot aged 6 to 63 years

Graziella Eshuis ¹, Julia Hock,² Gideon Marchie du Sarvaas,¹ Hiske van Duinen,³ Rhoia Neidenbach,² Freek van den Heuvel,¹ Hans Hillege,⁴ Rolf MF Berger,¹ Alfred Hager²

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/heartjnl-2020-318928>).

For numbered affiliations see end of article.

Correspondence to

Graziella Eshuis, Center of Congenital Heart Disease, Department of Paediatric Cardiology, Beatrix Children's Hospital, University of Groningen, University Medical Centre Groningen, Groningen, The Netherlands; g.eshuis01@umcg.nl

Received 31 December 2020

Revised 2 April 2021

Accepted 16 April 2021

ABSTRACT

Objectives This study aimed to provide a perspective for the interpretation of exercise capacity (peakVO₂) in patients with repaired Tetralogy of Fallot (patients with rTOF) by describing the course of peakVO₂ from patients aged 6–63 years.

Methods A retrospective study was performed between September 2001 and December 2016 in the German Heart Centre Munich, Germany, and in the University Medical Centre Groningen, the Netherlands. A total of 1175 cardiopulmonary exercise tests (CPETs) were collected from 586 patients with rTOF, 46% female. Maximal exertion was verified using a respiratory exchange ratio ≥ 1.00 . PeakVO₂ was modelled using time-dependent multilevel models for repeated measurements (n=889 in 300 patients), and compared with subject-specific reference values calculated by the models of Bongers *et al* and Mylius *et al*.

Results The peakVO₂ of patients with rTOF was reduced at all ages. At the age of 6, the peakVO₂ was 614 mL/min (70% of predicted (95% CI 67 to 73)). The reduced increase in peakVO₂ during adolescence resulted in a significant lower maximum peakVO₂ of 1209 mL/min at 25 years (65% predicted, p<0.001). A linear decline after 25 years was observed in patients and references, although patients showed an accelerated decline, with a -0.24% point of predicted (95% CI 0.11 to 0.38) per year without differences between sexes (p=0.263).

Conclusions This study provides a context for peakVO₂ across ages in patients with rTOF under contemporary treatment strategies. It showed that the reduction in peakVO₂ originates from childhood and declines over time. Sex differences in patients with rTOF were similar to natural existing sex differences.

cross-sectional studies, whereas cohort studies with repeated measures over time are scarce.

Ageing has been shown to affect peakVO₂.⁸ In healthy children, peakVO₂ increases associated with a growth-related increase in height and weight.^{9–11} During adolescence, weight gain is predominantly determined by increased skeletal muscle mass, most pronounced in boys.^{10–12} After reaching adulthood, peakVO₂ gradually declines during life.¹³ This decline is partially explained by a loss of muscle mass and a decrease of chronotropic response.^{8 12 14}

Previous cross-sectional studies in patients with rTOF reported a reduced peakVO₂ of 60%–80% of predicted.^{15–18} Longitudinal studies assessing the peakVO₂ course in patients with rTOF are rare, especially in children. These studies are mainly small studies in adults.^{19 20} A slow attrition of peakVO₂, about 1% point per year, is described under contemporary treatment strategies.^{19 20}

To adequately interpret serial CPETs in individual patients, knowledge of the peakVO₂ course across a wide age range is essential. The peakVO₂ course, and its variation between sexes, will put an individual's result in a more proper perspective. It will provide better insight into when the deviations of patients with rTOF from the healthy references occur. This allows for customised and more optimally timed interventions, including lifestyle advice and physical training programmes. This study aimed to show the peakVO₂ course from childhood to the ageing adult.

METHODS

Design, setting and patients

This multicentre cohort study included consecutive CPETs of patients with rTOF followed in the German Heart Centre Munich, Germany, or in the University Medical Center Groningen, the Netherlands. Patients were retrospectively included between September 2001 and December 2016. Clinical and demographic data were collected from medical records. Patients or the Public were not involved in the design, conduct, reporting or dissemination plans of our research.

Cardiopulmonary exercise test

All patients performed symptom-limited upright sitting CPET with an incremental protocol on a bicycle ergometer with breath-by-breath analysis, according to international guidelines and recommendations.^{21 22} The CPETs performed on

INTRODUCTION

Serial exercise testing using cardiopulmonary exercise tests (CPETs) is an emerging strategy to monitor patients with congenital heart disease (CHD). Measuring exercise capacity (maximal oxygen consumption, peakVO₂) is increasingly advocated as a diagnostic and prognostic tool to recognise clinical deterioration early in the course of the disease.^{1–3} The CPET has been adopted in international guidelines to support treatment decisions, including valve replacement interventions in patients with repaired Tetralogy of Fallot (rTOF).^{3–7} Such recommendations are predominantly based on

© Author(s) (or their employer(s)) 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Eshuis G, Hock J, Marchie du Sarvaas G, *et al*. Heart Epub ahead of print: [please include Day Month Year]. doi:10.1136/heartjnl-2020-318928

a treadmill were excluded, since treadmill exercise is a weight-bearing form of physical activity, while cycling is not. By using only cycling tests, the effects of weight fluctuations over time were minimised.²³

In short, a patient started the CPET with a resting and warming-up phase, followed by a progressive incremental workload protocol, adapted to the individual patient, aiming for an exercise duration of 8–12 min for both children and adults. The workload protocol was determined based on age, height and subjective fitness of the patient. The test ended when the patient could not maintain the pedal cycle rhythm, or if any arrhythmias occurred. CPETs were considered valid if the VO_2 reached a plateau (>1 min) despite increasing workload (only scarce in adults) and/or a complementary respiratory exchange ratio (RER) ≥ 1.00 combined with the presence of subjective signs of maximal effort (facial flushing, sweating, unwillingness to continue despite encouragement).^{21 22} Additional analyses of the peak VO_2 course in adults who reached RER ≥ 1.00 but RER < 1.1 , vs those with a RER ≥ 1.1 showed no significant difference in slope nor intercept. Therefore, all were included.

Outcome

The primary outcome value was peak VO_2 , either expressed in absolute mL/min, or as percentage of predicted.^{13 24} Peak VO_2 was defined as the mean VO_2 of the last 30 s of the incremental protocol. The absolute peak VO_2 was compared with subject-specific references based on two prediction models. The model of Bongers *et al* was used in girls until the age of 13, and boys until the age of 15.²⁴ For participants above those ages, weight and height contributed to their peak VO_2 and therefore reference values reported by Mylius *et al* were used.^{9–13}

Statistical analyses

Patient characteristics were described using means and SD or median with IQR depending on the distribution. All variables were visually checked and tested for normality using histograms, PP and QQ plots and the Kolmogorov-Smirnov test. To depict the age-related distribution of the population, the population was divided into age-based quartiles at initial measurement. Differences between sexes were tested using t-test for normally distributed variables, Mann-Whitney U test for skewed variables and χ^2 test was used for dichotomous variables. Deviations from the reference values were tested by combining subject-specific references and patients measurements into one time-dependent model.

Modelling peak VO_2 course

The peak VO_2 of all initial measurements were plotted in a box plot with age-based quartiles to show the distribution across ages. Scatterplots of the repeated measurements were used to visualise the peak VO_2 course, for the individual within-person variation (see online supplemental file). The models' lines including 95% CIs (dotted lines) were plotted over the scatterplots. This was used to visually assess the fit of the model.

Time-dependent multilevel models for repeated measures using an unstructured covariance structure with maximum likelihood estimation were used to calculate the course over time using repeated measures ($n=889$, in 300 patients).²⁵ This model was suitable for an unbalanced dataset, including various waves (eg, visits), non-identical spacing and different starting points. Assumptions of the model were: (1) linearity checked by empirical growth plots and scatterplots, and (2) normality and homoscedasticity of the residuals checked by normal probability

plots for residuals and scatterplots of the residuals against age. The statistical best fit was checked by using the Bayesian Information Criterion (BIC) and Akaike Information Criterion (AIC). The best model had the lowest BIC and AIC. For detailed information about the statistical methods and model building methods, see online supplemental file. Statistical analyses were performed using SPSS version 26 and STATA version 15. A p value of <0.05 was considered significant for determinants.

RESULTS

Baseline characteristics

A total of 586 individual patients (46% female patients) were included in the study, and 1175 valid CPETs were available for analyses. In 300 patients (51%), at least two measurements were available for analyses, and in 141 patients (24%) three or more measurements were available ($n=889$). Characteristics at initial measurement of all patients divided based on age quartiles are shown in figure 1 and table 1. At initial measurement, the median age was 21 (IQR 15–30) years. The youngest patient was 6 and the oldest 63 years of age. Almost half of the patients (42%) were 18 years or younger ($n=247$) at initial measurement. The median interval between the initial and follow-up CPET was 24 (IQR 12–38) months, with a maximal interval of 11 years. There was no significant difference in baseline characteristics between the single measurement sample and the repeated measurement sample, besides the morphology (table 2).

Peak VO_2 course in patients with rTOF

The absolute peak VO_2 course (mL/min) in the 300 patients with repeated measures and their subject-specific references^{13 24} is depicted in figure 2A. Additionally, the relative peak VO_2 course (%-predicted) is shown in figure 2B. The indexed peak VO_2 course (mL/min/kg) is shown in figure 3 (AIC 14.725; BIC 14.771) and figure 4 stratified for sex (AIC 14.235; BIC 14.344).

The cubic prediction model was the best fit to describe the absolute peak VO_2 course (AIC 26.054; BIC 26.098). The absolute peak VO_2 at age 6 (intercept of the model) seemed lower in patients (614 mL/min) compared with subject-specific references (637 mL/min), however, not significant, $p=0.764$. Thereafter, the peak VO_2 increased faster in references (280 mL/min per year) compared with children with rTOF (167 mL/min per year), $p<0.001$. The course was described by the model for patients' peak VO_2 (mL/min) = $614 + 167 \times (\text{age}-6) - 6 \times (\text{age}-6)^2 + 0.06 \times (\text{age}-6)^3$ vs the model for references' peak VO_2 (mL/min) = $637 + 280 \times (\text{age}-6) - 10 \times (\text{age}-6)^2 + 0.10 \times (\text{age}-6)^3$; $p<0.001$. The maximal peak VO_2 was reached around 20–25 years of age in both patients and references. Thereafter, there was a gradual decline over the years. This is also visible in the relative peak VO_2 course (figure 2).

The linear model was used to describe the relative peak VO_2 course (BIC 6.910; AIC 6.882). The relative peak VO_2 of patients with rTOF at age 6 was 70% (95% CI 67 to 73). Patients showed a reduced peak VO_2 at all ages, and an accelerated decline compared with references with 0.24% points of predicted (95% CI 0.11 to 0.38) per year.

Sex-related differences in exercise capacity in patients with rTOF

The absolute peak VO_2 (mL/min) modelled and stratified for sex is depicted in figure 5. Again, the cubic model was the best fit (AIC 25.547; BIC 25.652). No significant difference between male and female patients at the age of 6 ($p<0.906$) was observed. Both sexes showed a lower peak VO_2 than the references at age 6

Figure 1 Box plot of the absolute and relative peakVO₂ at initial cardiopulmonary exercise test of all 586 individual patient, divided by age quartiles to show the age distribution and effect of age. The left boxes represent the absolute peakVO₂ (mL/min) corresponding to the left y-axis. The right boxes represent the relative peakVO₂ (%-predicted) corresponding to the right y-axis.

($p < 0.001$). The steep and linear increase in peakVO₂ is accelerated in boys compared with girls ($p < 0.001$). The model for female patients' peakVO₂ (mL/min) = $922 + 167 \times (\text{age}-6) - 5.3 \times (\text{age}-6)^2 + 0.05 \times (\text{age}-6)^3$ vs the model for male patients' peakVO₂ (mL/min) = $935 + 322 \times (\text{age}-6) - 9.7 \times (\text{age}-6)^2 + 0.09 \times (\text{age}-6)^3$; $p < 0.001$. All effects for male and female patients are significantly lower compared with the subject-specific references (figure 5, $p < 0.001$), also appreciated in figure 6, which shows the percentage of predicted.

The linear model was the best fit to model relative peakVO₂ (AIC 6.885; BIC 6.923). The relative peakVO₂ at age 6 was not

significantly different between boys and girls ($p = 0.389$). The annual decline in comparison to the subject-specific references was not significantly different between sexes ($p = 0.263$).

DISCUSSION

This study showed the peakVO₂ course in 300 patients with rTOF aged 6 to 63 years ($n = 889$ CPETs). The peakVO₂ was reduced at all ages and deteriorated significantly faster in patients compared with healthy reference values with -0.24% of predicted point per year. Female patients had a lower peakVO₂ than male patients,

Table 1 Overview of 586 patients with rTOF divided into age-based quartiles at initial visit

	Q1		Q2		Q3		Q4	
Patient characteristics	n=146		n=146		n=147		n=147	
Age range, years	6.4–14.7		14.7–20.5		20.5–30.1		30.1–62.8	
Age, years	11.6	(9.8–13.4)	16.8	(15.6–18.4)	25.2	(23.0–27.4)	37.1	(33.6–43.5)
Female sex, n	53	36%	63	43%	81	55%	72	49%
Morphology								
TOF	104	71%	110	75%	104	71%	125	85%
PAVSD	36	25%	27	19%	27	18%	19	13%
DORV	6	4%	9	6%	16	11%	3	2%
Functional assessment								
CPET, n	1.44	±0.8	1.85	±1.2	2.2	±1.7	2.2	±1.4
Body weight, kg	41	(30–51)	61	(54–72)	65	(57–76)	69	(58–81)
Body height, cm	150	(138–161)	169	(162–178)	170	(163–175)	170	(163–177)
BMI, kg/m ²	18.3	±3.6	21.7	±3.2	23.2	±4.0	24.7	±4.4
PeakVO ₂ , mL/min	1358	(1082–1696)	1828	(1513–2342)	1813	(1377–2227)	1584	(1233–1976)
PeakVO ₂ , %-predicted*	73	±16	69	±16	63	±13	59	±15
PeakVO ₂ , mL/min/kg	35.8	(29.3–42.1)	31.7	(26.4–37.4)	26.8	(22.2–31.4)	22.0	(18.4–28.1)

Baseline characteristics for all patients at initial visit. Values are expressed as median (IQR), counts (percentage) or mean ± SD.

*Predicted values were based on the model by Bongers *et al* for girls until 13 years, for boys until 15 years and the model by Mylius *et al* for girls >13 years and boys >15 years. Age quartiles were constructed to show the age distribution in the population Q1 (first 25%, $n = 146$), Q2 (25%–50%, $n = 146$), Q3 (50%–75%, $n = 147$) and Q4 (last 25%, $n = 147$). BMI, body mass index; CPET, cardiopulmonary exercise test; DORV, double outlet right ventricle; PAVSD, pulmonary atresia/ventricular septal defect; TOF, Tetralogy of Fallot.

Table 2 Baseline characteristics of 586 patients with rTOF

	Single measurements		Repeated measurements		P value
Patient characteristics	n=286		n=300		
Age, years	20.0	(14.7–31.5)	21.5	(14.7–28.7)	0.815*
Female sex, n	128	45%	141	47%	0.099¥
Morphology					
TOF	229	80%	214	71%	0.019¥
PAVSD	40	14%	69	23%	
DORV	17	6%	17	6%	
Functional assessment	n=286		n=300		
CPET, n	1	±0	2.3	±1.4	<0.001\$
Body weight, kg	60	(49–73)	60	(51–72)	0.512*
Body height, cm	165	(158–174)	167	(158–175)	0.261*
BMI, kg/m ²	22.0	±4.9	21.9	±4.1	0.690\$
PeakVO ₂ , mL/min	1633	(1307–2069)	1628	(1251–2008)	0.501*
PeakVO ₂ , %-predicted†	67	±17	65	±15	0.153\$
PeakVO ₂ , mL/min/kg	29.2	(22.0–36.5)	28.7	(23.1–33.8)	0.493*

Baseline characteristics for all patients at initial visit. Values are expressed as median (IQR), counts (percentage) or mean±SD.

*P<0.05 was considered significant between the populations, based on Mann-Whitney (*) or Student's t-test (\$) or χ^2 test (¥).

†Predicted values were based on the model by Bongers *et al* for girls until 13 years, for boys until 15 years and on the model by Mylius *et al* for girls >13 years and boys >15 years.

BMI, body mass index; CPET, cardiopulmonary exercise test; DORV, Double Outlet Right Ventricle; PAVSD, Pulmonary Atresia/Ventricular Septal Defect; TOF, Tetralogy of Fallot.

as is expected compared with the references (and so, the healthy population). Compared with male patients, female patients did not show significant different rates of decline in peakVO₂.

The physiological effect of age on peakVO₂

The growth and maturation of children and adolescents, and the gradual decline of the cardiopulmonary system and muscle mass in adulthood have a substantial effect on physical fitness.^{10–12}

In early childhood, body compositions are similar in boys and girls; however, during adolescence this changes dramatically.¹² As adolescence proceeds, the weight gain is predominantly determined by an increase in skeletal muscle mass, which is most pronounced in boys.¹² Therefore, it is only logical to incorporate more than just age as a predictor of peakVO₂ in older children, healthy or diseased. A linear increase related to age is assumed in boys until the age of 15 and in girls until the age of 13.^{10–12} After

Figure 2 (A) The scatterplot represents repeated peakVO₂ measurements (n=889) (orange) and the subject-specific reference peakVO₂ (green) in the dataset. The time-dependent model is shown by the line including the 95% CI (dotted line) (AIC 26.054; BIC 26.098). Model 1A—orange (*patients*): peakVO₂ (mL/min)=614+167×(age-6)-6×(age-6)²+0.06×(age-6)³. Model 1A—green (*references*): peakVO₂ (mL/min)=637+280×(age-6)-10×(age-6)²+0.10×(age-6)³. (B) The scatterplot represents calculated percentage of predicted based on the subject-specific references (purple). The time-dependent model is shown by the line including the 95% CI (dotted line) (AIC 6.882; BIC 6.910). Model 1B—purple (*patients*): peakVO₂ (%-predicted)=70-0.24×(age-6). AIC, Akaike Information Criterion; BIC, Bayesian Information Criterion.

Figure 3 The scatterplot represents repeated indexed peakVO₂ measurements (n=889) (orange) and the subject-specific reference indexed peakVO₂ (green) in the dataset. The time-dependent model is shown by the line including the 95% CI (dotted line) (AIC 14.725; BIC 14.771). Model 2—orange (*patients*): peakVO₂ (mL/min/kg)=38.9–0.9×(age-6)+0.02×(age-6)²–0.0002×(age-6)³. Model 2—green (*references*): peakVO₂ (mL/min)=42.7+0.5×(age-6)–0.03×(age-6)²+0.0002×(age-6)³. AIC, Akaike Information Criterion; BIC, Bayesian Information Criterion.

these ages, body composition influences exercise performance.¹² Figure 2 shows a combination of the predictions of Bongers *et al*²⁴ and Mylius *et al*.¹³ The linear increase suggested by Bongers *et al* for the first 18 years of a child's life is not necessarily true, especially not for girls.^{10 24} In girls, from 13 years of age body weight and height are important factors in addition to age. From this age onwards, girls accumulate more fat, whereas boys do

increase their muscle mass.^{12 26} This partly explains the natural difference in peakVO₂ between sexes.^{12 26 27}

Cross-sectional studies in healthy, active adult people described a percentage rate of change of about 1% points per year.^{28 29} The deterioration of exercise capacity is related to ageing, reduced physical activity, natural weight gain and increased fat mass during life with subsequent sarcopenia starting at 40 years of age.²⁹

The peakVO₂ course with a cardiac lesion

While growing and ageing, physiological changes should be considered when assessing and interpreting peakVO₂ of patients with rTOF.^{5–7} We found a peakVO₂ at age 6 of 70% (95% CI 67 to 73) of predicted in the current study. This result is comparable to the cross-sectional studies in children with rTOF in the literature.^{15–18} The models combined with the scatterplots showed that there is a significant variation in exercise capacity. Roughly, a range from 25% until 120% of predicted at a young age, and 25% until 80% of predicted at older age, also shown in the indexed peakVO₂ course (mL/min/kg).

At a young age, there is no significant difference between sexes yet. Probably because the natural difference between sexes is not yet expressed too.^{10–12} Both male and female patients show a similar relative peakVO₂ compared with the subject-specific reference values. The peakVO₂ in young patients with rTOF does not rise as fast as in references (figure 2A). This is also shown in figure 2B, where the predicted percentage declined linearly from the beginning. The reduced increase during childhood and adolescence might be the reason why patients reach a critical point of exercise limitation, at which daily life is affected, faster than a healthy peer. This peakVO₂ course in patients with rTOF suggests that the limitation in exercise performance described in adults with rTOF originates mainly in childhood and adolescence. In this period of life, the musculoskeletal

Figure 4 The scatterplot represents repeated indexed peakVO₂ measurements (n=889) (red for female patients, blue for male patients) and the subject-specific reference indexed peakVO₂ (green) in the dataset. The time-dependent model is shown by the line, including the 95% CI (dotted line) (AIC 14.235; BIC 14.344). Model 5—blue (*male patients*): peakVO₂ (mL/min/kg)=40.8–0.8×(age-6)+0.01×(age-6)²–0.0001×(age-6)³. Model 5—red (*female patients*): peakVO₂ (mL/min/kg)=35.9–2.7×(age-6)+0.07×(age-6)²–0.0007×(age-6)³. AIC, Akaike Information Criterion; BIC, Bayesian Information Criterion.

Figure 5 The scatterplot represents repeated peakVO₂ measurements (n=889) (red for female patients, blue for male patients) and the subject-specific reference peakVO₂ (green) in the dataset. The time-dependent model is shown by the line, including the 95% CI (dotted line) (AIC 25.547; BIC 25.652). Model 3—blue (male patients): peakVO₂ (mL/min)=935+322×(age-6)−9.7×(age-6)²+0.09×(age-6)³. Model 3—red (female patients): peakVO₂ (mL/min)=922+167×(age-6)−5.3×(age-6)²+0.05×(age-6)³. AIC, Akaike Information Criterion; BIC, Bayesian Information Criterion.

system normally grows and matures.^{13 14} These children do have residual lesions; however, they do not (yet) suffer from major cardiac problems, such as right ventricular dysfunction or dilatation, requiring pulmonary valve replacement.¹⁸ Besides affecting the heart, muscle development is failing in a young patient with rTOF.³⁰ Future assessment of patients should also implement more unconventional variables, such as muscle function.

Two studies in the literature do show a longitudinal assessment of peakVO₂ in patients with CHD. Müller *et al* described

a slow decline of −1.01% point predicted per year in peakVO₂ in patients with CHD, based on the references of Gläser *et al*.³¹ A total of 116 patients with rTOF were included in the group of 522 patients with CHD. Patients with simple and more complex CHD were also included, no differences in the rate of change of the relative peakVO₂ were observed between CHD subgroups. There was no distinction based on sex.¹⁹ The rate of decline was slower in our population, with 0.24% of predicted points per year. We did not observe a difference between sexes, except for the natural existing difference between boys and girls. Our study showed the peakVO₂ course, which provides more insight in the overall peakVO₂ of a patient with rTOF than just a difference between two measures.

Kipps *et al* also described peakVO₂ over time in sole patients with rTOF, and included 70 patients with at least two measures (n=179). They showed a detailed change in peakVO₂ as percentage of predicted for the individual patient, based on the references of Cooper *et al*.³² The individual trajectories provide a detailed overview of the peakVO₂ of the patient with rTOF; however, it also shows its diversity.²⁰ By using a statistical model to average the peakVO₂ course, we were able to zoom out. Our study provides a stronger context for individual patients by acknowledging and addressing the natural existing difference between sexes concerning peakVO₂.

In the light of explaining peakVO₂, there were no associations between clinical or demographic variables at baseline and the deterioration of peakVO₂.²⁰ Furthermore, the cardiac function at rest, measured by cardio magnetic resonance, was not associated with a decline in peakVO₂. Thus, it seems to be hard to predict which patients would deteriorate faster than the other with conventional methods.²⁰ To explain the differences between a well-performing patient with rTOF and a deteriorating patient with rTOF, future studies and assessment of patients should implement objective measurements of fitness and daily activities or leisure-time sports. Besides cardiopulmonary fitness,

Figure 6 The scatterplot represents calculated percentage of predicted based on the subject-specific predictions (red for female patients, blue for male patients). The time-dependent model is shown by the line, including the 95% CI (dotted line) (AIC 6.885; BIC 6.923). Model 4—blue (male patients): peakVO₂ (%-predicted)=71−0.32×(age-6). Model 4—red (female patients): peakVO₂ (%-predicted)=69−0.16×(age-6). AIC, Akaike Information Criterion; BIC, Bayesian Information Criterion.

the muscle status and lung function at a young age should be addressed as well.

Strengths and limitations

To our knowledge, this is the largest study to describe the peakVO₂ of sole patients with rTOF in a wide age range. Our study included a relatively young cohort with at least 42% children and adolescents (<18 years) at baseline, from two centres. The statistical model made it possible to create a representative model of the peakVO₂ course. All CPETs included were performed on a cycle ergometer, which eliminates the fluctuations in weight over time on the modality of exercise.²³ The peakVO₂ course of reference values and patients provide clinicians with a clear context for their interpretation of CPETs. Furthermore, up-to-date prediction models were used to address the difference with the references.

A limitation of this study is the retrospective design. Our dataset was unbalanced, it did not include follow-up measurements for every patient. We chose to compute the model with repeated measurements, which ensured a modelled course over time. Selection bias was minimal, since there was no difference between the single measurement sample and the repeated sample. In addition, the policy in both hospitals was not to perform the CPET only on indication, but as a part of routine follow-up. Furthermore, there was a discrepancy between the absolute versus the predicted models due to averaging by the mathematical model. This happened in the crude model (with only peakVO₂ and age), not when stratified for sex. Therefore, this discrepancy was explained by averaging out the sex effect in the patient group. Individual cardiac functional imaging, such as echocardiography or cardio magnetic resonance, was not available for analysis. Our mathematical models describing the course of peakVO₂ did not include explanatory factors (such as re-operations, sports activities, etc). Including these factors would have made the mathematical model too complex to interpret. Further

studies are warranted to investigate whether re-operations affect the lifetime trajectories of peakVO₂ in patients with rTOF.

CONCLUSION

This multicentre study showed that the reduction in peakVO₂ in patients with rTOF originates already from childhood. The peakVO₂ is reduced at all ages. Particularly, the magnitude of the rise in peakVO₂ during adolescence is smaller, while the decline in adulthood is accelerated compared with healthy references.

Author affiliations

¹Center of Congenital Heart Disease, Department of Paediatric Cardiology, Beatrix Children's Hospital, University of Groningen, University Medical Centre Groningen, Groningen, The Netherlands

²Department of Paediatric Cardiology and Congenital Heart Disease, Technical University Munich, German Heart Centre Munich, München, Germany

³Department of Biomedical Sciences of Cells & Systems, Section of Anatomy & Medical Physiology, University of Groningen, University Medical Centre Groningen, Groningen, The Netherlands

⁴Center for Congenital Heart Disease, Department of Epidemiology, University of Groningen, University Medical Centre Groningen, Groningen, The Netherlands

Twitter Graziella Eshuis @Graziell36647415

Acknowledgements The authors would like to thank all colleagues from the Paediatric Cardiology Department and Exercise Laboratory of the German Heart Centre Munich and the University Medical Centre Groningen.

Contributors All authors contributed to the manuscript. GE and RMFB are responsible for the overall content as guarantors in addition to planning, conduct and reporting of the work. JH, RN and FvdH: conduct. GMDs: reporting of the work. HvD: reporting of the work. HH: reporting of the work. AH: conduct and report of the work.

Funding GE received a student grant for the research abroad from the Dutch Heart Foundation (td/dekker/2755).

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The institutional medical ethical committees approved the research protocol and waived the need for informed consent due to the retrospective nature of the study (METc2018/345). The investigation conformed to the principles outlined in the Declaration of Helsinki.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. Please contact the corresponding author for questions about the availability of the data.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

ORCID iD

Graziella Eshuis <http://orcid.org/0000-0001-6469-1566>

REFERENCES

- Guazzi M, Bandera F, Ozemek C, *et al.* Cardiopulmonary exercise testing: what is its value? *J Am Coll Cardiol* 2017;70:1618–36.
- Egbe AC, Driscoll DJ, Khan AR, *et al.* Cardiopulmonary exercise test in adults with prior Fontan operation: the prognostic value of serial testing. *Int J Cardiol* 2017;235:6–10.
- Paolillo S, Agostoni P. Prognostic role of cardiopulmonary exercise testing in clinical practice. *Ann Am Thorac Soc* 2017;14:S53–8.
- Ritt LE, Myers J, Stein R, *et al.* Additive prognostic value of a cardiopulmonary exercise test score in patients with heart failure and intermediate risk. *Int J Cardiol* 2015;178:262–4.
- Legendre A, Richard R, Pontnau F, *et al.* Usefulness of maximal oxygen pulse in timing of pulmonary valve replacement in patients with isolated pulmonary regurgitation. *Cardiol Young* 2016;26:1310–8.
- Sabate Rotes A, Johnson JN, Burkhart HM, *et al.* Cardiorespiratory response to exercise before and after pulmonary valve replacement in patients with repaired

Key messages

What is already known on this subject?

- Cross-sectional studies in children and adults show reduced peakVO₂ in patients with repaired tetralogy of Fallot (rTOF).
- Two longitudinal studies including patients with rTOF (n=70, n=116) show the same trend.
- Both studies describe the exercise capacity either as a difference between two points in time or as differences that occur in the individual patient.

What might this study add?

- Our study is the largest study to date that encompasses 300 patients with rTOF and describe the course of peakVO₂ across a wide age range from 6 to 63 years.
- Also, this study for the first time includes a young population with 42% aged 18 years or younger and addresses sex differences.

How might this impact on clinical practice?

- This study provides a context for interpreting peakVO₂ in patients with rTOF.
- A clinician can apply this to compare the patient with healthy reference values and a contemporary cohort of patients with rTOF.

- tetralogy of Fallot: a retrospective study and systematic review of the literature. *Congenit Heart Dis* 2015;10:263–70.
- 7 Babu-Narayan SV, Diller G-P, Gheta RR, *et al.* Clinical outcomes of surgical pulmonary valve replacement after repair of tetralogy of Fallot and potential prognostic value of preoperative cardiopulmonary exercise testing. *Circulation* 2014;129:18–27.
  - 8 Tanaka H, Monahan KD, Seals DR. Age-predicted maximal heart rate revisited. *J Am Coll Cardiol* 2001;37:153–6.
  - 9 Van Brussel M, Bongers BC, Hulzebos EHJ, *et al.* A systematic approach to interpreting the cardiopulmonary exercise test in pediatrics. *Pediatr Exerc Sci* 2019;31:194–203.
  - 10 Armstrong N, Welsman JR. Assessment and interpretation of aerobic fitness in children and adolescents. *Exerc Sport Sci Rev* 1994;22:435–76.
  - 11 Fawcner S, Armstrong N. Oxygen uptake kinetic response to exercise in children. *Sports Med* 2003;33:651–69.
  - 12 Rutenfranz J, Andersen KL, Seliger V, *et al.* Maximum aerobic power and body composition during the puberty growth period: similarities and differences between children of two European countries. *Eur J Pediatr* 1981;136:123–33.
  - 13 Mylius CF, Krijnen WP, van der Schans CP, *et al.* Peak oxygen uptake reference values for cycle ergometry for the healthy Dutch population: data from the LowLands fitness registry. *ERJ Open Res* 2019;5. doi:10.1183/23120541.00056-2018. [Epub ahead of print: 01 Apr 2019].
  - 14 Roh J, Rhee J, Chaudhari V, *et al.* The role of exercise in cardiac aging: from physiology to molecular mechanisms. *Circ Res* 2016;118:279–95.
  - 15 Powell AW, Mays WA, Knecht SK, *et al.* Pulmonary effects on exercise testing in tetralogy of Fallot patients repaired with a transannular patch. *Cardiol Young* 2019;29:133–9.
  - 16 Hock J, Häcker A-L, Reiner B, *et al.* Functional outcome in contemporary children and young adults with tetralogy of Fallot after repair. *Arch Dis Child* 2019;104:129–33.
  - 17 Friedberg MK, Fernandes FP, Roche SL, *et al.* Relation of right ventricular mechanics to exercise tolerance in children after tetralogy of Fallot repair. *Am Heart J* 2013;165:551–7.
  - 18 Bhatt SM, Elci OU, Wang Y, *et al.* Determinants of exercise performance in children and adolescents with repaired tetralogy of Fallot using stress echocardiography. *Pediatr Cardiol* 2019;40:71–8.
  - 19 Müller J, Ewert P, Hager A. Only slow decline in exercise capacity in the natural history of patients with congenital heart disease: a longitudinal study in 522 patients. *Eur J Prev Cardiol* 2015;22:113–8.
  - 20 Kipps AK, Graham DA, Harrild DM, *et al.* Longitudinal exercise capacity of patients with repaired tetralogy of Fallot. *Am J Cardiol* 2011;108:99–105.
  - 21 Takken T, Bongers BC, van Brussel M, *et al.* Cardiopulmonary exercise testing in pediatrics. *Ann Am Thorac Soc* 2017;14:S123–8.
  - 22 American Thoracic Society, American College of Chest Physicians. ATS/ACCP statement on cardiopulmonary exercise testing. *Am J Respir Crit Care Med* 2003;167:211–77.
  - 23 Bar-Yoseph R, Porszasz J, Radom-Aizik S. The effect of test modality on dynamic exercise biomarkers in children, adolescents, and young adults. *Physiol Rep* 2019;7.
  - 24 Bongers BC, Brussel van M, Hulzebos EHJ. Paediatric norms for cardiopulmonary exercise testing. In: *BoxPress 's-Hertogenbosch*. 2nd edn, 2014.
  - 25 Singer JD, Willett JB. *Applied longitudinal data analysis modeling change and event occurrence*. 1st edn. Oxford: Oxford University Press, 2003: 29.
  - 26 Cureton K, Bishop P, Hutchinson P, *et al.* Sex difference in maximal oxygen uptake. *Eur J Appl Physiol Occup Physiol* 1986;54:656–60.
  - 27 Hagdorn QAJ, Beurskens NEG, Gorter TM, *et al.* Sex differences in patients with repaired tetralogy of Fallot support a tailored approach for males and females: a cardiac magnetic resonance study. *Int J Cardiovasc Imaging* 2020;36:1997–2005.
  - 28 Hawkins SA, Marcell TJ, Victoria Jaque S, *et al.* A longitudinal assessment of change in VO2max and maximal heart rate in master athletes. *Med Sci Sports Exerc* 2001;33:1744–50.
  - 29 Fitzgerald MD, Tanaka H, Tran ZV, *et al.* Age-related declines in maximal aerobic capacity in regularly exercising vs. sedentary women: a meta-analysis. *J Appl Physiol* 1997;83:160–5.
  - 30 Greutmann M, Le TL, Tobler D, *et al.* Generalised muscle weakness in young adults with congenital heart disease. *Heart* 2011;97:1164–8.
  - 31 Koch B, Schäper C, Ittermann T, *et al.* Reference values for cardiopulmonary exercise testing in healthy volunteers: the SHIP study. *Eur Respir J* 2009;33:389–97.
  - 32 Cooper DM, Weiler-Ravell D. Gas exchange response to exercise in children. *Am Rev Respir Dis* 1984;129:S47–8.