University of Groningen

Heart failure re-hospitalizations and subsequent fatal events in coronary artery disease

Ferreira, Joao Pedro; Cleland, John G.; Lam, Carolyn S. P.; Anker, Stefan D.; Mehra, Mandeep R.; van Veldhuisen, Dirk J.; Byra, William M.; La Police, David A.; Pitt, Bertram; Greenberg, Barry

Published in:
Clinical Research in Cardiology

DOI:
[10.1007/s00392-021-01830-1](https://doi.org/10.1007/s00392-021-01830-1)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Ferreira, J. P., Cleland, J. G., Lam, C. S. P., Anker, S. D., Mehra, M. R., van Veldhuisen, D. J., Byra, W. M., La Police, D. A., Pitt, B., Greenberg, B., & Zannad, F. (2021). Heart failure re-hospitalizations and subsequent fatal events in coronary artery disease: insights from COMMANDER-HF, EPHEsus, and EXAMINE. *Clinical Research in Cardiology*, 110, 1554-1563. <https://doi.org/10.1007/s00392-021-01830-1>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Heart failure re-hospitalizations and subsequent fatal events in coronary artery disease: insights from COMMANDER-HF, EPHEBUS, and EXAMINE

João Pedro Ferreira^{1,2,3} · John G. Cleland⁴ · Carolyn S. P. Lam^{5,6} · Stefan D. Anker^{7,8} · Mandeep R. Mehra⁹ · Dirk J. van Veldhuisen⁶ · William M. Byra¹⁰ · David A. La Police¹⁰ · Bertram Pitt¹¹ · Barry Greenberg¹² · Faiez Zannad^{1,2,3}

Received: 3 January 2021 / Accepted: 24 February 2021
© Springer-Verlag GmbH, DE part of Springer Nature 2021

Abstract

Background Patients with coronary artery disease (CAD) are at increased risk of developing and being hospitalised for heart failure (HFH). However, the risk of HFH versus ischemic events may vary among patients with CAD, depending on whether acute myocardial infarction (MI), left ventricular dysfunction or decompensated HF is present at baseline.

Aims We aim to explore the risk of non-fatal events (HFH, MI, stroke) and subsequent death in 3 landmark trials, COMMANDER-HF, EPHEBUS and EXAMINE that, together, included patients with CAD with and without reduced ejection fraction and acute MI.

Methods Events, person-time metrics and time-updated Cox models.

Results In COMMANDER-HF the event-rate for the composite of AMI, stroke or all-cause death was 13.5 (12.8–14.3) events/100 py. Rates for AMI and stroke were much lower (2.2 [2.0–2.6] and 1.3 [1.1–1.6] events/100 py, respectively) than the rate of HFH (16.9 [16.1–17.9] events/100 py). In EPHEBUS, the rates of MI and stroke were also lower than the rate of HFH: 7.2 (6.7–7.8), 1.9 (1.7–2.3), and 10.6 (9.9–11.3) events/100 py, but this was not true for EXAMINE with 4.4 (4.0–4.9), 0.7 (0.6–0.9), and 2.4 (2.0–2.7) events/100 py, respectively. In all 3 trials, a non-fatal event (HFH, MI or stroke) during follow-up doubled the risk of subsequent mortality. This most commonly followed a HFH.

Conclusions A first or recurrent HFH is common in patients with CAD and AMI or HFrEF and indicates a poor prognosis. Preventing the development of heart failure after AMI and control of congestion in patients with CAD and HFrEF are key unmet needs and therapeutic targets.

Registration ClinicalTrials.gov Identifier: NCT01877915. URL: <https://clinicaltrials.gov/ct2/show/NCT01877915>.

Keywords Even type · Even-rates · Heart failure · Myocardial infarction · Stroke

Introduction

Strategies to prevent and treat coronary artery disease (CAD) have led to reductions in the incidence of myocardial infarction (MI) and subsequent mortality, driven by early restoration of myocardial blood flow and better secondary prevention by controlling cardiovascular risk factors

and pharmacological interventions [1, 2]. In contrast, the incidence and prevalence of heart failure have increased, especially amongst older people, due (at least partially) to better survival after MI [3]. The development of heart failure often leads to substantial disability and impaired quality of life and is associated with high rates of hospitalizations and a poor prognosis [4]. Hospitalisations are often for worsening heart failure (HFH) that may be triggered by ischemia or many other factors [5, 6].

In the COMMANDER-HF trial, which included patients with a left ventricular ejection fraction (LVEF) < 40%, heart failure with reduced ejection fraction (HFrEF) and coronary artery disease (CAD) in sinus rhythm plus a recent hospitalisation for worsening congestion, rivaroxaban 2.5 mg bd

✉ João Pedro Ferreira
j.ferreira@chru-nancy.fr

✉ Faiez Zannad
f.zannad@chru-nancy.fr

Extended author information available on the last page of the article

added to background anti-platelet therapy did not reduce the rate of hospitalisation or death due to worsening heart failure but may have reduced vascular occlusive events [7–9]. These findings suggest that myocardial dysfunction rather than coronary events drive the progression of HFrEF with CAD [10]. Accordingly, we compared event-rates for HFH, MI and stroke as well as the risk of subsequent death after a non-fatal event (HFH, MI, and stroke) in COMMANDER-HF compared to EPHESUS (Eplerenone, a Selective Aldosterone Blocker, in Patients with Left Ventricular Dysfunction after Myocardial Infarction) [11] and EXAMINE (Alogliptin after Acute Coronary Syndrome in Patients with Type 2 Diabetes) [12].

Methods

Study populations

COMMANDER-HF was an international, double-blind, randomized trial comparing the factor-Xa inhibitor rivaroxaban (2.5 mg twice daily) vs. placebo. Key inclusion criteria included the history of chronic HF for 3 or more months, left ventricular ejection fraction (LVEF) of 40% or less, history of coronary artery disease, absence of atrial fibrillation or other indication for chronic anticoagulation, and treatment for an episode of decompensated heart failure (i.e., the index event) within the previous 21 days. Decompensated HF was defined by symptoms of worsening dyspnea or fatigue, objective signs of congestion, and/or adjustment of HF medications requiring hospital admission or unscheduled parenteral diuretic [7, 13].

EPHESUS was an international, double-blind, randomized trial comparing eplerenone vs. placebo. Patients were eligible for enrolment if they had an acute MI, and a LVEF $\leq$ 40% plus HF or diabetes within 3–14 days before randomization [11].

EXAMINE was an international, double-blind, randomized trial comparing the DPP-IV inhibitor alogliptin vs. placebo. Patients were eligible for enrolment if they had type-2 diabetes mellitus and had experienced an acute coronary syndrome within 15–90 days before randomization [12, 14].

The 3 trials were conducted in accordance with the Declaration of Helsinki and approved by the site ethics committees. All participants gave written informed consent to participate in the studies.

Study outcomes

The primary efficacy outcome in COMMANDER-HF was the composite of death from any cause, MI, or stroke. In EPHESUS the primary efficacy outcomes were a composite

of death from cardiovascular causes or hospitalization for cardiovascular causes (including heart failure, acute myocardial infarction, stroke, or ventricular arrhythmia), and death from any cause. In EXAMINE the primary endpoint was a composite of death from cardiovascular causes, nonfatal MI, or nonfatal stroke. HFH was a main secondary end-point of both the COMMANDER-HF and EXAMINE trials.

Statistical analysis

Similar methods were applied across the studied trials. Number and proportion of events, person-time exposures, and time-to-first event-rates (per 100 person-years) were computed. A “smoothed” hazard function was plotted for each time-to-first event to assess the event pattern throughout the follow-up. To make the comparisons easier to interpret we plotted all-cause death, HFH, MI, and stroke over time using the same *x*-scale and *y*-scale for all trials. Cumulative incidence rates were computed using the Nelson-Aalen method in the placebo arm of the respective trials to ease the between-trial comparison without the effect of treatments. To assess the risk of death after the first non-fatal events; time-updated Cox models were performed in the placebo groups of the respective trials adjusting for potential baseline confounders, as described in previous analysis [15]. In COMMANDER-HF the adjustment was performed on age, sex, race, diabetes, prior MI, prior stroke, history of percutaneous coronary intervention or coronary artery bypass grafting (PCI/CABG), NYHA class, body mass index, systolic blood pressure, anemia, estimated glomerular filtration rate (eGFR), dual anti-platelet therapy, rivaroxaban or placebo allocation, and geographical region stratification. The treatment effect of rivaroxaban vs. placebo was tested by the intention-to-treat principle at different follow-up times with stratification by region and “right censored” times (6, 12 months, and/or last censor). The proportional hazards assumption was tested by plotting the Schoenfeld residuals over time; the residuals were independent of time and no evidence of proportional hazards violation was found. A 2-sided *p* value of < 0.05 was considered significant. Stata® version 16 (StataCorp. 2019. College Station, TX: StataCorp LLC) was used for the analyses.

Results

Baseline characteristics

In COMMANDER-HF a total of 5022 patients were included with a median follow-up duration of 21.1 months (percentile_{25–75}, 12.9–32.8), the mean age was 66 ± 10 years, 77% were male, 82% white, and 41% had diabetes. In EPHESUS a total of 6632 patients were included with a

median follow-up duration of 15.6 months (percentile_{25–75}, 11.9–20.6), the mean age was 64 ± 11 years, 71% were male, 90% white, and 32% had diabetes. In EXAMINE a total of 5380 patients were included with a median follow-up duration of 19.2 months (percentile_{25–75}, 12.0–26.4), the mean age was 61 ± 10 years, 68% were male, 73% white, and all patients had diabetes (by inclusion criteria). The characteristics of the patients enrolled in the respective trials are described in the Supplemental Table 1.

Event rates

In COMMANDER-HF the primary outcome (a composite of MI, stroke or all-cause death) overall event rate was 13.5 (12.8–14.3) events per 100 person-years; the event rate was higher in the first 6 months and decreased thereafter: 16.8 (15.2–18.5) in the first 6 months vs. 11.9 (11.0–12.9) events per 100 person-years beyond the 12th month of follow-up. Similar findings were observed for the composite of MI, stroke or cardiovascular death. The major contributor for the observed event rate was all-cause death with an overall event rate of 11.2 (10.6–11.9) with similar findings observed for cardiovascular death (Table 1). The overall rate of MI was low: 2.2 (2.0–2.6) events per 100 person-years, but slightly higher within the first 6 months of follow-up: 3.2 (2.6–4.0) in the first 6 months vs. 1.6 (1.3–2.0) events per 100 person-years beyond the 12th month of follow-up. The rate of stroke was even lower and more homogeneous over time with an overall event rate of 1.3 (1.1–1.6) events per 100 person-years (Table 1). The rate of HFH was the highest among all outcomes with an overall rate of 16.9 (16.1–17.9) events per 100 person-years that was much higher in the first 6 months: 29.3 (27.1–31.6) vs. 9.3 (8.4–10.3) after 12 months of follow-up (Table 1; Figs. 1, 2). Bleeding rates were overall low: ≤ 2 events per 100 person-years (Supplemental Table 2). Details on multiple other events (e.g., sudden death, “pump failure” death, non-HFH) are also presented in the Supplemental Table 2. No major variations of the treatment effect (rivaroxaban vs. placebo) over time were found (Supplemental Table 3).

In EPHEUS the overall event rate of the composite outcome of MI, stroke or cardiovascular death was 16.2 (15.4–17.1) events per 100 person-years; the event rate was higher in the first 6 months and decreased thereafter: 28.6 (26.7–30.6) in the first 6 months vs. 7.8 (6.8–9) events per 100 person-years beyond the 12th month of follow-up. In EPHEUS, the MI rates were more than three-fold higher than in COMMANDER-HF, with an overall event rate of 7.2 (6.7–7.8) events per 100 person-years, also highest in the first 6 months. The rate of stroke was similar to that found in COMMANDER-HF with an overall value of 1.9 (1.7–2.3) events per 100 person-years (Table 1). The HFH rates were lower than those found in COMMANDER-HF. The overall rate was

Table 1 Event rates per 100 person-years by intervals of follow-up time

Time (months)	Person-years	Events	Rate (per 100 py)
COMMANDER-HF (<i>n</i> = 5022)			
MI, stroke, ACM			
0–6	2403	404	16.8 (15.2–18.5)
6–12	2151	290	13.5 (12–15.1)
Beyond 12	4946	590	11.9 (11–12.9)
Total	9500	1284	13.5 (12.8–14.3)
MI, stroke, CVD			
0–6	2403	367	15.3 (13.8–16.9)
6–12	2151	263	12.2 (10.8–13.8)
Beyond 12	4946	491	9.9 (9.1–10.8)
Total	9500	1121	11.8 (11.1–12.5)
MI			
0–6	2412	78	3.2 (2.6–4)
6–12	2169	59	2.7 (2.1–3.5)
Beyond 12	5048	79	1.6 (1.3–2)
Total	9629	216	2.2 (2–2.6)
Stroke			
0–6	2420	41	1.7 (1.2–2.3)
6–12	2183	30	1.4 (1–2)
Beyond 12	5075	56	1.1 (0.8–1.4)
Total	9679	127	1.3 (1.1–1.6)
ACM			
0–6	2429	320	13.2 (11.8–14.7)
6–12	2202	239	10.9 (9.6–12.3)
Beyond 12	5180	543	10.5 (9.6–11.4)
Total	9810	1102	11.2 (10.6–11.9)
CVD			
0–6	2429	282	11.6 (10.3–13)
6–12	2202	212	9.6 (8.4–11)
Beyond 12	5180	435	8.4 (7.6–9.2)
Total	9810	929	9.5 (8.9–10.1)
HFH			
0–6	2265	663	29.3 (27.1–31.6)
6–12	1872	344	18.4 (16.5–20.4)
Beyond 12	4009	373	9.3 (8.4–10.3)
Total	8146	1380	16.9 (16.1–17.9)
EPHESUS (<i>n</i> = 6632)			
MI, stroke, CVD			
0–6	2998	857	28.6 (26.7–30.6)
6–12	2676	280	10.5 (9.3–11.8)
Beyond 12	2586	202	7.8 (6.8–9)
Total	8259	1339	16.2 (15.4–17.1)
MI			
0–6	3016	427	14.2 (12.9–15.6)
6–12	2703	106	3.9 (3.2–4.7)
Beyond 12	2632	71	2.7 (2.1–3.4)
Total	8351	604	7.2 (6.7–7.8)
Stroke			
0–6	3091	86	2.8 (2.3–3.4)

Table 1 (continued)

Time (months)	Person-years	Events	Rate (per 100 py)
6–12	2822	44	1.6 (1.2–2.1)
Beyond 12	2758	38	1.4 (1–1.9)
Total	8671	168	1.9 (1.7–2.3)
ACM			
0–6	3111	590	19 (17.5–20.6)
6–12	2851	243	8.5 (7.5–9.7)
Beyond 12	2808	199	7.1 (6.2–8.1)
Total	8769	1032	11.8 (11.1–12.5)
CVD			
0–6	3111	534	17.2 (15.8–18.7)
6–12	2851	196	6.9 (6–7.9)
Beyond 12	2808	160	5.7 (4.9–6.7)
Total	8769	890	10.1 (9.5–10.8)
HFH			
0–6	2952	594	20.1 (18.6–21.8)
6–12	2610	160	6.1 (5.2–7.2)
Beyond 12	2536	101	4 (3.3–4.8)
Total	8098	855	10.6 (9.9–11.3)
EXAMINE (n = 5380)			
MI, stroke, CVD			
0–6	2504	283	11.3 (10.1–12.7)
6–12	2100	160	7.6 (6.5–8.9)
Beyond 12	3429	178	5.2 (4.5–6)
Total	8033	621	7.7 (7.1–8.4)
MI			
0–6	2511	168	6.7 (5.8–7.8)
6–12	2114	95	4.5 (3.7–5.5)
Beyond 12	3470	97	2.8 (2.3–3.4)
Total	8094	360	4.4 (4–4.9)
Stroke			
0–6	2551	23	0.9 (0.6–1.4)
6–12	2190	15	0.7 (0.4–1.1)
Beyond 12	3667	23	0.6 (0.4–0.9)
Total	8409	61	0.7 (0.6–0.9)
ACM			
0–6	2558	119	4.7 (3.9–5.6)
6–12	2204	81	3.7 (3–4.6)
Beyond 12	3708	126	3.4 (2.9–4)
Total	8470	326	3.8 (3.5–4.3)
CVD			
0–6	2558	97	3.8 (3.1–4.6)
6–12	2204	61	2.8 (2.2–3.6)
Beyond 12	3708	84	2.3 (1.8–2.8)
Total	8470	242	2.9 (2.5–3.2)
HFH			
0–6	2531	105	4.1 (3.4–5)
6–12	2156	42	1.9 (1.4–2.6)
Beyond 12	3587	48	1.3 (1–1.8)
Total	8273	195	2.4 (2–2.7)

MI myocardial infarction, CVD cardiovascular death, ACM all-cause mortality, HFH heart failure re-hospitalization

10.6 (9.9–11.3) events per 100 person-years, but also highest early in the follow-up (Table 1; Figs. 1, 2).

In EXAMINE the overall event rate of the composite outcome of MI, stroke or cardiovascular death was 7.7 (7.1–8.4) events per 100 person-years; the event rate was higher in the first 6 months and decreased thereafter. In EXAMINE, the MI rates were twofold higher than in COMMANDER-HF, with an overall event rate of 4.4 (4.0–4.9) events per 100 person-years, also highest in the first 6 months. The rate of stroke was lower than in COMMANDER-HF with an overall value of 0.7 (0.6–0.9) events per 100 person-years (Table 1). The HFH rates were much lower than those found in COMMANDER-HF and EPHEUS. The overall rate was 2.4 (2.0–2.7) events per 100 person-years, but also highest early in the follow-up (Table 1; Figs. 1, 2).

Deaths after a non-fatal event

In COMMANDER-HF, having a non-fatal event during the follow-up increased the risk of subsequent death by twofold. The risk was similar for all non-fatal events, but slightly higher after an MI: MI adjusted HR (95% CI) = 2.31 (1.73–2.61), $p < 0.001$; HFH adjusted HR (95% CI) = 1.81 (1.59–2.05), $p < 0.001$; stroke adjusted HR (95% CI) = 1.75 (1.31–2.33), $p < 0.001$. The absolute numbers and proportion of HFH events preceding death was much higher than the proportion of MI or stroke: HFH 517 (46.9%), MI 106 (9.6%), Stroke 53 (4.8%) (Table 2; Figs. 3, 4).

In EPHEUS, having a non-fatal event during the follow-up increased the risk of subsequent death by twofold to threefold. The risk was higher after an MI or stroke, than after a HFH: MI adjusted HR (95% CI) = 3.16 (2.72–3.67), $p < 0.001$; stroke adjusted HR (95% CI) = 3.01 (2.38–3.81), $p < 0.001$; HFH adjusted HR (95% CI) = 2.23 (1.94–2.58), $p < 0.001$. The absolute numbers and proportion of HFH events preceding death was also higher than the proportion of MI or stroke: HFH 316 (30.6%), MI 261 (25.3%), Stroke 82 (8.0%) (Table 2; Figs. 3, 4).

In EXAMINE, having a non-fatal event during the follow-up increased the risk of subsequent death by 1.5–2.5-fold. In EXAMINE the risk was higher after a HFH than after a MI or stroke: HFH adjusted HR (95% CI) = 2.50 (1.80–3.48), $p < 0.001$; MI adjusted HR (95% CI) = 1.59 (1.15–2.19), $p = 0.005$; stroke adjusted HR (95% CI) = 1.49 (0.70–3.18), $p = 0.30$. The absolute numbers and proportion of HFH events preceding death were similar to MI and higher than stroke: HFH 48 (14.7%), MI 46 (14.1%), Stroke 7 (2.2%) (Table 2; Figs. 3, 4).

Fig. 1 Smoothed hazard function over time. **a** COMMANDER-HF. **b** EPHESUS. **c** EXAMINE. The cumulative hazard function is on the y-axis and the follow-up time in years is displayed in the x-axis

Discussion

Our study shows that HFHs are frequent and associated with poor outcomes in patients with CAD. However, depending on the population studied the type of events and their rate may vary widely. Patients with decompensated HF but without an acute MI (as in COMMANDER-HF), present a high rate of HF re-hospitalizations frontloaded early within the follow-up; the HFH rates are much superior (eightfold or even higher) to those of MI or stroke (despite the presence of underlying CAD) and greatly increase the risk for subsequent death ($\approx$ twofold higher). In EPHESUS, with high-risk acute MI patients, HF re-hospitalizations were more frequent than recurrent MI and also independently associated with a poor prognosis. In EXAMINE the recurrent MI rates were higher than HFH, but HFH was also frequent and associated with a poorer prognosis than MI. Together, these data show

that a large window of opportunity to reduce events still exists and that targeting HF is key in this endeavour.

COMMANDER-HF was designed with the hypothesis that thrombin would play a central role in the HF progression of patients with CAD [13]. Inhibiting the factor Xa with rivaroxaban could lead to the reduction of thrombin and related pathways associated with myocyte injury, inflammation, endothelial dysfunction and microvascular thrombosis, which could then result in a reduction of ischemic/thrombotic events and also HF hospitalizations and mortality via thrombin-related pathways [16, 17]. The inclusion of patients early (within 21 days) of a decompensated HF episode would enrich the trial for outcome events. This was indeed what happened, with the vast majority of these events being HFH on which rivaroxaban had no effect [8, 18]. By computing the risk of death after the non-fatal events, one may observe that the risk of dying after a HFH was similar

Fig. 2 Cumulative incidence of events in the placebo groups of the respective trials. The individual events are displayed on the y-axis and the follow-up time in years is displayed in the x-axis

to the risk of dying after a stroke or MI. However, HFH were a much more frequent event than MI or stroke, making HF re-hospitalizations as a main target for improvement. Even in patients who had an acute MI (as in EPHESUS and EXAMINE), HF re-hospitalizations represent a major burden with deleterious consequences. Similarly, in a post-hoc analysis of the ALTITUDE trial (Aliskiren Trial in Type 2 Diabetes Using Cardio-Renal Endpoints) that included patients with type 2 diabetes plus chronic kidney disease, cardiovascular disease or both, the proportion of patients experiencing a

HFH was higher than that of MI or stroke, and the greatest number of deaths occurred in the group who had a HFH first [19].

Up-to-date HFrEF treatment including an angiotensin-receptor neprilysin inhibitor (ARNI), a sodium glucose co-transporter inhibitor (SGLT2i), and a mineralocorticoid receptor antagonist (MRA) may lead to a 50% event reduction compared with treatment with angiotensin-converting enzyme/angiotensin receptor blocker (ACEi/ARB) and a beta-blocker [20]; however, it should be noted that the

Table 2 Death after a non-fatal event

Events	Event and death	Event and survival	Adj. HR (95% CI)	<i>p</i>
COMMANDER-HF				
HFH				
<i>n</i> = 1380	517 (46.9%)	863 (22.0%)	1.81 (1.59–2.05)	<0.001
MI				
<i>n</i> = 216	106 (9.6%)	110 (2.8%)	2.13 (1.73–2.64)	<0.001
Stroke				
<i>n</i> = 127	53 (4.8%)	74 (1.9%)	1.75 (1.31–2.33)	<0.001
Total	1102	3920		
EPHESUS				
HFH				
<i>n</i> = 855	316 (30.6%)	539 (9.6%)	2.23 (1.94–2.58)	<0.001
MI				
<i>n</i> = 604	261 (25.3%)	343 (6.1%)	3.16 (2.72–3.67)	<0.001
Stroke				
<i>n</i> = 168	82 (8.0%)	86 (1.5%)	3.01 (2.38–3.81)	<0.001
Total	5600	1032		
EXAMINE				
HFH				
<i>n</i> = 195	48 (14.7%)	147 (2.9%)	2.50 (1.80–3.48)	<0.001
MI				
<i>n</i> = 360	46 (14.1%)	314 (6.2%)	1.59 (1.15–2.19)	0.005
Stroke				
<i>n</i> = 61	7 (2.2%)	54 (1.1%)	1.49 (0.70–3.18)	0.30
Total	5054	326		

MI myocardial infarction, HFH heart failure re-hospitalization

Fig. 3 Proportion of deaths following a non-fatal event in each trial. y-scale, percentage (%). The figure represents the proportion of all deaths that occurred after a non-fatal event in each trial. For example, in COMMANDER-HF 47% of all deaths occurred after a HFH. HFH heart failure hospitalization, MI myocardial infarction

implementation of HF-modifying therapies remains an issue even in developed countries [21–25]. Notwithstanding, while the therapeutic armamentarium for chronic HFrEF may sharply reduce events rates, if implemented; post worsening HF remains an unmet need with high recurrent HFH and mortality, and offers an opportunity to test

and implement novel strategies. Such opportunity was tested in the PIONEER-HF (Angiotensin–Neprilysin Inhibition in Acute Decompensated Heart Failure) trial, where initiation of sacubitril/valsartan upon stabilization of a decompensated HF episode led to a greater reduction in the NT-pro BNP than enalapril without increasing adverse events [26]. In

Fig. 4 Cumulative incidence of death after a non-fatal event. **a** COMMANDER-HF. **b** EPHESUS. **c** EXAMINE. *HFH* heart failure hospitalization, *MI* myocardial infarction

VICTORIA-HF (Vericiguat in Patients with Heart Failure and Reduced Ejection Fraction) patients had to have evidence of worsening HF and most patients (67%) had been hospitalized for HF in the previous 3 months which resulted in a high rate of HFH and death in this trial, slightly reduced with vericiguat treatment [23]. In COMPASS (Rivaroxaban with or without Aspirin in Stable Cardiovascular Disease) patients with a HF history also had a higher rate of all the studied events, including HFH [27, 28].

These examples show that the opportunity for improvement in populations that had a recent HF hospitalization, MI or both is considerable and optimizing the use of therapies that have been shown to improving outcomes is key to achieve this goal [29].

Limitations

The findings here depicted may only be generalized to patients with similar characteristics of those included in these trials, due to the relatively young, white and male populations included, our findings may not be applicable to patients with different characteristics. Furthermore, the inclusion/exclusion criteria and level of monitoring performed in a trial context may further limit the generalization of these results to patients in the community, but allows a detailed assessment of adjudicated cause-specific events. The timing of enrollment after the acute event was similar in COMMANDER-HF and EPHESUS (less than 21 and 3–14 days, respectively), but more widespread in EXAMINE (15–90 days) which might have contributed to the lower overall event rate in EXAMINE. The follow-up times were

slightly different between trials (median in months of 21 in COMMANDER-HF, 16 in EPHEUS, and 19 in EXAMINE), which might have influenced the event rates and the assessment of fatal events after non-fatal ones. EPHEUS was published almost 20 years ago; the MI therapeutics have improved much since and the observed high death risk after MI could be lower in a contemporary cohort.

Conclusions

The burden of HF re-hospitalizations is enormous in patients with CAD and is associated with a high risk of subsequent death. Reducing HF in CAD represents an unmet need that should be addressed in future studies.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00392-021-01830-1>.

Funding The COMMANDER-HF trial was supported by Janssen Research and Development.

Declarations

Conflict of interest Dr Ferreira is a consultant for Boehringer-Ingelheim. Dr Zannad reports steering committee personal fees from Applied Therapeutics, Amgen, Bayer, Boehringer, Novartis, Janssen, Cellprothera and CVRx, advisory board personal fees from, AstraZeneca, Vifor Fresenius, Cardior, Cereno pharmaceutical, Corvidia, Merck, Myokardia, NovoNordisk and Owkin, stock options at Cereno and G3Pharmaceutical, and being the founder of the Global Cardiovascular Clinical Trialist Forum. Dr Lam has received research support from Boston Scientific, Bayer, Roche Diagnostics, AstraZeneca, Medtronic, and Vifor Pharma; has served as consultant or on the Advisory Board/ Steering Committee/ Executive Committee for Abbott Diagnostics, Amgen, Applied Therapeutics, AstraZeneca, Bayer, Biofourmis, Boehringer Ingelheim, Boston Scientific, Corvia Medical, Cytokinetics, Darma Inc., Us2.ai, JanaCare, Janssen Research & Development LLC, Medtronic, Menarini Group, Merck, MyoKardia, Novartis, Novo Nordisk, Radcliffe Group Ltd., Roche Diagnostics, Sanofi, Stealth BioTherapeutics, The Corpus, Vifor Pharma and WebMD Global LLC; and serves as co-founder & non-executive director of Us2.ai. Dr Greenberg reports consulting fees from Amgen, Cytokinetics, EBR Systems, Impulse Dynamics, Ionis, Jaan, Janssen, Merck, Myokardia, Sanofi, Vifor, Viking, Windtree and Zensun. All the other authors report not having relevant conflicts of interest regarding the content of this manuscript.

References

- Smolina K, Wright FL, Rayner M, Goldacre MJ (2012) Determinants of the decline in mortality from acute myocardial infarction in England between 2002 and 2010: linked national database study. *BMJ* 344:d8059
- Bandosz P, O'Flaherty M, Drygas W, Rutkowski M, Koziarek J, Wyrzykowski B, Bennett K, Zdrojewski T, Capewell S (2012) Decline in mortality from coronary heart disease in Poland after socioeconomic transformation: modelling study. *BMJ* 344:d8136
- Conrad N, Judge A, Tran J, Mohseni H, Hedgecott D, Crespiello AP, Allison M, Hemingway H, Cleland JG, McMurray JJV, Rahimi K (2017) Temporal trends and patterns in heart failure incidence: a population-based study of 4 million individuals. *Lancet* 391:572–580
- Conrad N, Judge A, Canoy D, Tran J, Pinho-Gomes AC, Millett ERC, Salimi-Khorshidi G, Cleland JG, McMurray JJV, Rahimi K (2019) Temporal trends and patterns in mortality after incident heart failure: a longitudinal analysis of 86000 individuals. *JAMA Cardiol* 4:1102–1111
- Kobayashi M, Voors AA, Girerd N, Billotte M, Anker SD, Cleland JG, Lang CC, Ng LL, van Veldhuisen DJ, Dickstein K, Metra M, Duarte K, Rossignol P, Zannad F, Ferreira JP (2020) Heart failure etiologies and clinical factors precipitating for worsening heart failure: findings from BIOSTAT-CHF. *Eur J Intern Med* 71:62–69
- Arrigo M, Gayat E, Parenica J, Ishihara S, Zhang J, Choi DJ, Park JJ, Alhabib KF, Sato N, Miro O, Maggioni AP, Zhang Y, Spinar J, Cohen-Solal A, Iwashyna TJ, Mebazaa A (2017) Precipitating factors and 90-day outcome of acute heart failure: a report from the intercontinental GREAT registry. *Eur J Heart Fail* 19(2):201–208
- Zannad F, Anker SD, Byra WM, Cleland JGF, Fu M, Gheorghiade M, Lam CSP, Mehra MR, Neaton JD, Nessel CC, Spiro TE, van Veldhuisen DJ, Greenberg B (2018) Rivaroxaban in patients with heart failure, sinus rhythm, and coronary disease. *N Engl J Med* 379:1332–1342
- Mehra MR, Vaduganathan M, Fu M, Ferreira JP, Anker SD, Cleland JGF, Lam CSP, van Veldhuisen DJ, Byra WM, Spiro TE, Deng H, Zannad F, Greenberg B (2019) A comprehensive analysis of the effects of rivaroxaban on stroke or transient ischaemic attack in patients with heart failure, coronary artery disease, and sinus rhythm: the COMMANDER HF trial. *Eur Heart J* 40(44):3593–3602
- Greenberg B, Neaton JD, Anker SD, Byra WM, Cleland JGF, Deng H, Fu M, La Police DA, Lam CSP, Mehra MR, Nessel CC, Spiro TE, van Veldhuisen DJ, Vanden Boom CM, Zannad F (2019) Association of rivaroxaban with thromboembolic events in patients with heart failure, coronary disease, and sinus rhythm: a post hoc analysis of the COMMANDER HF trial. *JAMA Cardiol* 4(6):515–523
- Cleland JGF, Pellicori P (2019) Myocardial dysfunction and coronary artery disease as therapeutic targets in heart failure. *Circulation* 140(7):538–541
- Pitt B, Williams G, Remme W, Martinez F, Lopez-Sendon J, Zannad F, Neaton J, Roniker B, Hurley S, Burns D, Bittman R, Kleiman J (2001) The EPHEUS trial: eplerenone in patients with heart failure due to systolic dysfunction complicating acute myocardial infarction. Eplerenone post-AMI heart failure efficacy and survival study. *Cardiovasc Drugs Ther* 15(1):79–87
- White WB, Cannon CP, Heller SR, Nissen SE, Bergenstal RM, Bakris GL, Perez AT, Fleck PR, Mehta CR, Kupfer S, Wilson C, Cushman WC, Zannad F (2013) Alogliptin after acute coronary syndrome in patients with type 2 diabetes. *N Engl J Med* 369(14):1327–1335
- Zannad F, Greenberg B, Cleland JG, Gheorghiade M, van Veldhuisen DJ, Mehra MR, Anker SD, Byra WM, Fu M, Mills RM (2015) Rationale and design of a randomized, double-blind, event-driven, multicentre study comparing the efficacy and safety of oral rivaroxaban with placebo for reducing the risk of death, myocardial infarction or stroke in subjects with heart failure and significant coronary artery disease following an exacerbation of heart failure: the COMMANDER HF trial. *Eur J Heart Fail* 17(7):735–742
- White WB, Bakris GL, Bergenstal RM, Cannon CP, Cushman WC, Fleck P, Heller S, Mehta C, Nissen SE, Perez A, Wilson C, Zannad F (2011) EXamination of cArdiovascular outcoMes with alogliptIN versus standard of carE in patients with type 2 diabetes

- mellitus and acute coronary syndrome (EXAMINE): a cardiovascular safety study of the dipeptidyl peptidase 4 inhibitor alogliptin in patients with type 2 diabetes with acute coronary syndrome. *Am Heart J* 162(4):620–626.e1
15. Ferreira JP, Rossignol P, Dewan P, Lamiral Z, White WB, Pitt B, McMurray JJV, Zannad F (2019) Income level and inequality as complement to geographical differences in cardiovascular trials. *Am Heart J* 218:66–74
 16. Zannad F, Stough WG, Regnault V, Gheorghiade M, Deliargyris E, Gibson CM, Agewall S, Berkowitz SD, Burton P, Calvo G, Goldstein S, Verheugt FW, Koglin J, O'Connor CM (2013) Is thrombosis a contributor to heart failure pathophysiology? Possible mechanisms, therapeutic opportunities, and clinical investigation challenges. *Int J Cardiol* 167(5):1772–1782
 17. Ferreira JP, Girerd N, Alshalash S, Konstam MA, Zannad F (2016) Antithrombotic therapy in heart failure patients with and without atrial fibrillation: update and future challenges. *Eur Heart J* 37:2455–2464
 18. Ferreira JP, Girerd N, Gregson J, Latar I, Sharma A, Pfeffer MA, McMurray JJV, Abdul-Rahim AH, Pitt B, Dickstein K, Rossignol P, Zannad F (2018) Stroke risk in patients with reduced ejection fraction after myocardial infarction without atrial fibrillation. *J Am Coll Cardiol* 71(7):727–735
 19. Jhund PS, McMurray JJ, Chaturvedi N, Brunel P, Desai AS, Finn PV, Haffner SM, Solomon SD, Weinrauch LA, Claggett BL, Pfeffer MA (2015) Mortality following a cardiovascular or renal event in patients with type 2 diabetes in the ALTITUDE trial. *Eur Heart J* 36(36):2463–2469
 20. Vaduganathan M (2020) Estimating lifetime benefits of comprehensive disease-modifying pharmacological therapies in patients with heart failure with reduced ejection fraction: a comparative analysis of three randomised controlled trials. *Lancet* 396:121–128
 21. Tromp J, Ferreira JP, Janwanishstaporn S, Shah M, Greenberg B, Zannad F, Lam CSP (2020) Heart failure around the world. *Eur J Heart Fail* 21:1187–1196
 22. Greene SJ, Fonarow GC, DeVore AD, Sharma PP, Vaduganathan M, Albert NM, Duffy CI, Hill CL, McCague K, Patterson JH, Spertus JA, Thomas L, Williams FB, Hernandez AF, Butler J (2019) Titration of medical therapy for heart failure with reduced ejection fraction. *J Am Coll Cardiol* 73(19):2365–2383
 23. Armstrong PW, Pieske B, Anstrom KJ, Ezekowitz J, Hernandez AF, Butler J, Lam CSP, Ponikowski P, Voors AA, Jia G, McNulty SE, Patel MJ, Roessig L, Koglin J, O'Connor CM (2020) Vericiguat in patients with heart failure and reduced ejection fraction. *N Engl J Med* 382:1883–1893
 24. McMurray JJV, Solomon SD, Inzucchi SE, Kober L, Kosiborod MN, Martinez FA, Ponikowski P, Sabatine MS, Anand IS, Belohlavek J, Bohm M, Chiang CE, Chopra VK, de Boer RA, Desai AS, Diez M, Drozd J, Dukar A, Ge J, Howlett JG, Katova T, Kitakaze M, Ljungman CEA, Merkely B, Nicolau JC, O'Meara E, Petrie MC, Vinh PN, Schou M, Tereshchenko S, Verma S, Held C, DeMets DL, Docherty KF, Jhund PS, Bengtsson O, Sjostrand M, Langkilde AM (2019) Dapagliflozin in patients with heart failure and reduced ejection fraction. *N Engl J Med* 381:1995–2008
 25. McMurray JJ, Packer M, Desai AS, Gong J, Lefkowitz MP, Rizkala AR, Rouleau JL, Shi VC, Solomon SD, Swedberg K, Zile MR (2014) Angiotensin-neprilysin inhibition versus enalapril in heart failure. *N Engl J Med* 371(11):993–1004
 26. Velazquez EJ, Morrow DA, DeVore AD, Duffy CI, Ambrosy AP, McCague K, Rocha R, Braunwald E (2019) Angiotensin-neprilysin inhibition in acute decompensated heart failure. *N Engl J Med* 380(6):539–548
 27. Branch KR, Probstfield JL, Eikelboom JW, Bosch J, Maggioni AP, Cheng RK, Bhatt DL, Avezum A, Fox KAA, Connolly SJ, Shestakovska O, Yusuf S (2019) Rivaroxaban with or without aspirin in patients with heart failure and chronic coronary or peripheral artery disease. *Circulation* 140(7):529–537
 28. Eikelboom JW, Connolly SJ, Bosch J, Dagenais GR, Hart RG, Shestakovska O, Diaz R, Alings M, Lonn EM, Anand SS, Widimsky P, Hori M, Avezum A, Piegas LS, Branch KRH, Probstfield J, Bhatt DL, Zhu J, Liang Y, Maggioni AP, Lopez-Jaramillo P, O'Donnell M, Kakkar A, Fox KAA, Parkhomenko AN, Ertl G, Stork S, Keltai M, Ryden L, Pogossova N, Dans AL, Lanus F, Commerford PJ, Torp-Pedersen C, Guzik TJ, Verhamme PB, Vinereanu D, Kim JH, Tonkin AM, Lewis BS, Felix C, Yusuf K, Steg PG, Metsarinne KP, Cook Bruns N, Misselwitz F, Chen E, Leong D, Yusuf S (2017) Rivaroxaban with or without aspirin in stable cardiovascular disease. *N Engl J Med* 377:1319–1330
 29. Zannad F (2017) Rising incidence of heart failure demands action. *Lancet* 391:518–519

Authors and Affiliations

João Pedro Ferreira^{1,2,3} · John G. Cleland⁴ · Carolyn S. P. Lam^{5,6} · Stefan D. Anker^{7,8} · Mandeep R. Mehra⁹ · Dirk J. van Veldhuisen⁶ · William M. Byra¹⁰ · David A. La Police¹⁰ · Bertram Pitt¹¹ · Barry Greenberg¹² · Faiez Zannad^{1,2,3}

¹ Centre d'investigations Cliniques Plurithématique, Inserm 1433, Université de Lorraine, Institut Lorrain du Coeur et des Vaisseaux Louis Mathieu, CHRU Nancy–Hopitaux de Brabois, 4 rue du Morvan, 54500 Nancy, Vandoeuvre les Nancy, France

² CHRU de Nancy, Inserm U1116, Nancy, France

³ FCRIN INI-CRCT, Nancy, France

⁴ Robertson Centre for Biostatistics and Clinical Trials, University of Glasgow, Glasgow, Scotland

⁵ National Heart Centre Singapore, Duke-National University of Singapore, Singapore, Singapore

⁶ Department of Cardiology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands

⁷ Berlin–Brandenburg Center for Regenerative Therapies, Berlin, Germany

⁸ Department of Cardiology, German Center for Cardiovascular Research Partner Site Berlin, Charité Universitätsmedizin Berlin, Berlin, Germany

⁹ Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

¹⁰ Janssen Research and Development, Raritan, NJ, USA

¹¹ Department of Medicine, University of Michigan School of Medicine, Ann Arbor, USA

¹² Cardiology Division, Department of Medicine, University of California, La Jolla, San Diego, USA