University of Groningen

A presumed Lazarus coral

Tilstra, Arjen; Hoeksema, Bert W.; Wild, Christian

Published in:
Bulletin of Marine Science

DOI:
[10.5343/bms.2021.0007](https://doi.org/10.5343/bms.2021.0007)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Tilstra, A., Hoeksema, B. W., & Wild, C. (2021). A presumed Lazarus coral: Outstanding regeneration capacity of a *Goniopora* coral exposed to air over several months . *Bulletin of Marine Science*, 97(3), 473-474. <https://doi.org/10.5343/bms.2021.0007>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

A presumed Lazarus coral: outstanding regeneration capacity of a *Goniopora* coral exposed to air over several months

Arjen Tilstra ^{1*}, Bert W Hoeksema ^{2,3}, Christian Wild ¹

¹ Marine Ecology Department, Faculty of Biology and Chemistry, University of Bremen, Leobener Str. 6 (UFT), 28359 Bremen, Germany. ² Taxonomy, Systematics and Geodiversity Group, Naturalis Biodiversity Center, 2300 RA Leiden, The Netherlands. ³ Groningen Institute for Evolutionary Life Sciences, University of Groningen, 9700 CC Groningen, The Netherlands. * Corresponding author email: <tilstra@uni-bremen.de>

It is a common and well-known phenomenon that reef corals harboring shallow intertidal reefs regularly get exposed to air during low tide. This can cause temporary stress, visible as tissue retraction (Brown et al. 1994), bleaching (Anthony and Kerswell 2007), and temperature shifts (visualized by thermal imagery) of air exposed colonies (Heyward and Negri 2020), with the latter two depending on factors such as coral morphology and wave exposure. In the long term, these recurrent aerial exposures are not necessarily lethal, but they can lead to significant changes in coral holobiont physiology (e.g., by shifts in the microbiome; Sweet et al. 2017). However, long-lasting aerial exposure caused by extreme low tides can be deleterious for certain coral species (Anthony and Kerswell 2007, Schoepf et al. 2015, Mejía-Rentería et al. 2020). Here we present an observation of nondeleterious prolonged exposure to air in an ex situ cultured fleshy reef-building coral.

In May 2020, a colony of the reef-building scleractinian *Goniopora* sp., which had been in stable culture for several years (salinity 33.0, temperature 26 °C, nitrate ± 2.0 mg L⁻¹, phosphate ± 0.05 mg L⁻¹, calcium 440 mg L⁻¹, magnesium 1350 mg L⁻¹, alkalinity 3.0 mEq L⁻¹), retracted its polyps after extended low light exposure caused by shading from faster growing coral colonies. As a result, the coral lost all visible tissue and was discarded from the holding tank at the beginning of June after it was presumed dead. It was subsequently left outside to face the Dutch weather in the city of Rotterdam on a pile of limestone rocks facing south. The skeleton was introduced back into the holding tank approximately 3 mo later to support the substrate for a colony of another coral. During these 3 mo outside the water, the skeleton was exposed to a mean (SD) temperature of 18.4 (3.5) °C (range = 6.2–33.8 °C), mean (SD) global radiation of 1801 (641) J cm⁻² (range = 517–3050 J cm⁻²), and precipitation ranging between 0 and 30.6 mm d⁻¹ (KNMI 2021).

Within 2 wks after reintroduction, the skeleton generated tissue that was first assessed as dead tissue being slowly discarded from the skeleton (Panel A). Surprisingly, soon after this initial observation, the layer of tissue appeared to consist of living coral polyps (Panel B). In the following weeks, the coral showed a further improvement of its health by regaining its natural color (partially due to recolonization of zooxanthellae), while polyp expansion increased (Panel C). Finally, after about 4 mo, the colony fully regained its original color, while the coenosarc between all individual polyps appeared to be fully fused (Panel D).

Previous documented cases of extreme polyp regeneration from apparently dead scleractinian coral tissue did not involve such long periods of draught and concerned only a few scleractinian taxa, such as fungiid and dendrophyllid corals (e.g., Hoeksema 2004, Luz et al. 2018, Luz et al. 2021). While coral reef literature describes the effects of aerial exposure for minutes to several hours per day (e.g., Rosser and Veron 2011, Castrillón-Cifuentes et al. 2017, Sweet et al. 2017), to our knowledge, this is the longest period described in the literature of a reef coral that has survived out of water conditions. We hypothesize that this remarkable ability to regenerate may be due to the depth to which the coral retracted its polyps within the skeleton (Kitano et al. 2013), potentially linked to production of persistent mucus that efficiently protected the retracted coral tissue against desiccation (Krupp 1984), and to the regrowth of possible tissue remnants (DeFilippo et al. 2016). The remaining retracted tissue may have been in a state of quiescence (dormancy), a survival mechanism often demonstrated by temperate scleractinian corals as a response to cold water stress (Grace 2017). Our observation may lead to new ways to investigate the regenerative capacity of reef-building corals and may help coral reef conservation by identifying more resilient coral species.

ACKNOWLEDGMENTS

We thank Glenn Fong of Dutch Synthetic Reefing (DSR) for sharing this observation and providing the information and photographs. We also thank Kevin Favier for his help with photo editing.

LITERATURE CITED

- Anthony KRN, Kerswell AP. 2007. Coral mortality following extreme low tides and high solar radiation. *Mar Biol.* 151(5):1623–1631. <https://doi.org/10.1007/s00227-006-0573-0>
- Brown BE, Le Tissier MDA, Dunne RP. 1994. Tissue retraction in the scleractinian coral *Coeloseris mayeri*, its effect upon coral pigmentation, and preliminary implications for heat balance. *Mar Ecol Prog Ser.* 105:209–218. <https://doi.org/10.3354/meps105209>
- Castrillón-Cifuentes AL, Lozano-Cortés DF, Zapata FA. 2017. Effect of short-term subaerial exposure on the cauliflower coral, *Pocillopora damicornis*, during a simulated extreme low-tide event. *Coral Reefs.* 36(2):401–414. <https://doi.org/10.1007/s00338-017-1552-2>
- DeFilippo L, Burmester EM, Kaufman L, Rotjan RD. 2016. Patterns of surface lesion recovery in the Northern Star Coral, *Astrangia poculata*. *J Exp Mar Biol Ecol.* 481:15–24. <https://doi.org/10.1016/j.jembe.2016.03.016>
- Grace S. 2017. Winter quiescence, growth rate, and the release from competition in the temperate scleractinian coral *Astrangia poculata* (Ellis & Solander 1786). *Northeast Nat (Steuben).* 24(7):B119–B134. <https://doi.org/10.1656/045.024.s715>
- Heyward A, Negri A. 2020. Distinct thermal signatures of coral morphologies above the waterline. *Bull Mar Sci.* 96(4):765–766. <https://doi.org/10.5343/bms.2019.0112>
- Hoeksema BW. 2004. Impact of budding on free-living corals at East Kalimantan, Indonesia. *Coral Reefs.* 23:492. <https://doi.org/10.1007/s00338-004-0402-1>
- Kitano YF, Obuchi M, Uyeno D, Miyazaki K, Fukami H. 2013. Phylogenetic and taxonomic status of the coral *Goniopora stokesi* and related species (Scleractinia: Poritidae) in Japan based on molecular and morphological data. *Zool Stud.* 52:25. <https://doi.org/10.1186/1810-522X-52-25>
- KNMI. 2021. Daily data for Rotterdam (no. 344) between 1 June, 2020 and 31 August, 2020. R Netherlands Meteorol Inst. Accessed 3 February, 2020. Available from: <https://www.knmi.nl/nederland-nu/klimatologie/daggegevens>
- Krupp DA. 1984. Mucus production by corals exposed during an extreme low tide. *Pac Sci.* 38(1):1–11.
- Luz BLP, Capel KCC, Zilberberg C, Flores AAV, Migotto AE, Kitahara MV. 2018. A polyp from nothing: the extreme regeneration capacity of the Atlantic invasive sun corals *Tubastraea coccinea* and *T. tagusensis* (Anthozoa, Scleractinia). *J Exp Mar Biol Ecol.* 503:60–65. <https://doi.org/10.1016/j.jembe.2018.02.002>
- Luz BLP, Miller DJ, Kitahara MV. 2021. High regenerative capacity is a general feature within colonial dendrophyllid corals (Anthozoa, Scleractinia). *J Exp Zool B Mol Dev Evol.* 336:281–292. <https://doi.org/10.1002/jez.b.23021>
- Mejía-Rentería JC, Castellanos-Galindo GA, Osorio-Cano JD, Casella E. 2020. Subaerial exposure of coral reefs during spring low tides in the eastern Pacific. *Bull Mar Sci.* 96(1):219–220. <https://doi.org/10.5343/bms.2019.0085>
- Rosser NL, Veron JEN. 2011. Australian corals thriving out of water in an extreme environment. *Coral Reefs.* 30(1):21. <https://doi.org/10.1007/s00338-010-0689-z>
- Schoepf V, Stat M, Falter JL, McCulloch MT. 2015. Limits to the thermal tolerance of corals adapted to a highly fluctuating, naturally extreme temperature environment. *Sci Rep.* 5:17639. <https://doi.org/10.1038/srep17639>
- Sweet MJ, Brown BE, Dunne RP, Singleton I, Bulling M. 2017. Evidence for rapid, tide-related shifts in the microbiome of the coral *Coelastrea aspera*. *Coral Reefs.* 36(3):815–828. <https://doi.org/10.1007/s00338-017-1572-y>

Date Submitted: 10 February, 2021.

Date Accepted: 3 May, 2021.

Available Online: 3 May, 2021.