University of Groningen

Decadal shift in foraging strategy of a migratory southern ocean predator

van der Berg, Gideon; Vermeulen, Els; Valenzuela, Luciano O.; Bérubé, Martine; Ganswindt, Andre; Grocke, Darren R.; Hall, Grant; Hulva, Pavel; Neveceralova, Petra; Palsboll, Per

Published in:
Global Change Biology

DOI:
[10.1111/gcb.15465](https://doi.org/10.1111/gcb.15465)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

van der Berg, G., Vermeulen, E., Valenzuela, L. O., Bérubé, M., Ganswindt, A., Grocke, D. R., Hall, G., Hulva, P., Neveceralova, P., Palsboll, P., & Carroll, E. (2021). Decadal shift in foraging strategy of a migratory southern ocean predator. *Global Change Biology*, 27(5), 1052-1067.
<https://doi.org/10.1111/gcb.15465>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Decadal shift in foraging strategy of a migratory southern ocean predator

Gideon L. van den Berg¹ | Els Vermeulen¹ | Luciano O. Valenzuela^{2,3,4} |
Martine Bérubé^{5,6} | Andre Ganswindt¹ | Darren R. Gröcke⁷ | Grant Hall¹ |
Pavel Hulva^{8,9} | Petra Neveceralova^{8,10,11} | Per J. Palsbøll^{5,6} | Emma L. Carroll¹²

¹Department of Zoology and Entomology, Mammal Research Institute, University of Pretoria, Pretoria, South Africa

²Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Laboratorio de Ecología Evolutiva Humana (LEEH, Facultad de Ciencias Sociales, Unidad de Enseñanza Universitaria Quequén, UNCPBA, Buenos Aires, Argentina

³School of Biological Sciences, University of Utah, Salt Lake City, UT, USA

⁴Instituto de Conservación de Ballenas, Buenos Aires, Argentina

⁵Marine Evolution and Conservation Group, Groningen Institute of Evolutionary Life Sciences, University of Groningen, Groningen, The Netherlands

⁶Centre for Coastal Studies, Provincetown, MA, USA

⁷Stable Isotope Biogeochemistry Laboratory (SIBL), Department of Earth Sciences, Durham University, Durham, UK

⁸Department of Zoology, Faculty of Science, Charles University, Prague, Czech Republic

⁹Department of Biology and Ecology, Faculty of Science, University of Ostrava, Ostrava, Czech Republic

¹⁰Ivanhoe Sea Safaris, Gansbaai, South Africa

¹¹Dyer Island Conservation Trust, Great White House, Kleinbaai, South Africa

¹²School of Biological Sciences, University of Auckland, Auckland, New Zealand

Correspondence

Emma L. Carroll, School of Biological Sciences, University of Auckland, Auckland, New Zealand.
Email: e.carroll@auckland.ac.nz

Funding information

Royal Society; Royal Society of New Zealand Te Apārangi; ENI; Total; Exxon Mobil Corporation; Charles University, Grant/Award Number: 1140217; Newton International Fellowship

ABSTRACT

Rapid anthropogenic environmental change is expected to impact a host of ecological parameters in Southern Ocean ecosystems. Of critical concern are the consequences of these changes on the range of species that show fidelity to migratory destinations, as philopatry is hypothesized to help or hinder adaptation to climate change depending on the circumstances. Many baleen whales show philopatry to feeding grounds and are also capital breeders that meet migratory and reproductive costs through seasonal energy intake. Southern right whales (*Eubalaena australis*, SRWs) are capital breeders that have a strong relationship between reproductive output and foraging success. The population dynamics of South Africa's population of SRWs are characterized by two distinct periods: the 1990s, a period of high calving rates; and the late 2010s, a period associated with lowered calving rates. Here we use analyses of stable carbon ($\delta^{13}\text{C}$) and nitrogen ($\delta^{15}\text{N}$) isotope values from SRW biopsy samples ($n = 122$) collected during these two distinct periods to investigate foraging ecology of the South African population of SRWs over a time period coincident with the demographic shift. We show that South African SRWs underwent a dramatic northward shift, and diversification, in foraging strategy from 1990s to 2010s. Bayesian mixing model results suggest that during the 1990s, South African SRWs foraged on prey isotopically similar to South Georgia/Islands Georgias del Sur krill. In contrast, in the

2010s, South African SRWs foraged on prey isotopically consistent with the waters of the Subtropical Convergence, Polar Front and Marion Island. We hypothesize that this shift represents a response to changes in preferred habitat or prey, for example, the decrease in abundance and southward range contraction of Antarctic krill. By linking reproductive decline to changing foraging strategies for the first time in SRWs, we show that altering foraging strategies may not be sufficient to adapt to a changing ocean.

KEYWORDS

Bayesian mixing model, cetacean, climate change, foraging ecology, indicator species, reproductive success, SIBER, Southern Ocean, southern right whale, stable isotopes

1 | INTRODUCTION

Oceanic ecosystems are changing rapidly on a global scale due to a host of anthropogenic impacts (Bindoff et al., 2019). The most pronounced physical changes include increasing ocean temperatures, the southward shift of fronts in the Southern Ocean, ocean acidification and changes in seasonal sea-ice extents (Böning et al., 2008; Bracegirdle et al., 2008; Orr et al., 2005; Sokolov & Rintoul, 2009). The consequences of these physical changes on Southern Ocean food webs are complex (Constable et al., 2014), and are likely to have multifaceted ecological impacts on marine species, including physiology, phenology, behaviour, population densities and structure including migratory patterns, food web architecture and dynamics, and species ranges (Trathan et al., 2007). Of particular concern is the impact that such physical change will have on Antarctic krill (*Euphausia superba*), the dominant mid-trophic species in Southern Ocean food webs (Atkinson et al., 2004; Reid & Croxall, 2001). The southwest Atlantic sector of the Southern Ocean, which contains over 50% of Antarctic krill stocks, has seen a major reduction in Antarctic krill density from 1976 to 2003 (Atkinson et al., 2004) as well as a significant southward contraction in Antarctic krill over the past 90 years (Atkinson et al., 2019). Population models constructed by Murphy et al. (2007) indicate that a regional warming of 1°Celsius during the coming century could lead to a 95% reduction in the abundance and biomass of Antarctic krill in the southwest Atlantic Ocean.

Changes in krill availability can have profound negative impacts on krill predators in the Southern Ocean (Barbosa et al., 2012; Forcada & Trathan, 2009; Forcada et al., 2005; Fraser & Hofmann, 2003). Particularly disconcerting are the potential impacts on capital breeders; that is, species relying on seasonal energy reserves for reproduction (see Jönsson, 1997), such as southern right whales (*Eubalaena australis*, hereafter SRW). SRWs consume large amounts of krill and copepods during the austral summer on high-productivity foraging grounds which are stored as blubber; sustaining migration, mating and calving during the remainder of the year (Lockyer, 2007). SRW mothers rely on their energy reserves to sustain themselves and their calves during lactation (Lockyer, 2007), losing up to 25% of their body volume during a few months (Christiansen et al., 2018).

Sufficient energy reserves are thus crucial for both mother and calf survival (Lockyer, 2007).

The reproductive output of SRWs in Argentina (expected calving rate) and Brazil (observed calving rate) are correlated with krill abundance at a summer foraging ground off South Georgia (Islas Georgias del Sur), which, in turn, is linked to large-scale global climate drivers such as the El Niño–Southern Oscillation and the Southern Annular Mode (Leaper et al., 2006; Seyboth et al., 2016). Several studies have shown that reproduction is suppressed in cetaceans during periods of nutritional stress (Greene et al., 2003; Hlista et al., 2009; Lockyer, 1986; Reeves et al., 2001; Ward et al., 2009; Williams et al., 2013), potentially the cause of lower reproductive output in Argentinean SRWs (Rowntree et al., 2013). If summer foraging and reproductive success are correlated in SRWs, then it follows that a decrease in SRW reproductive rates may be caused by reduced biological productivity at offshore foraging grounds.

SRWs show migratory fidelity to destinations, with long-term photo-identification and genotype studies showing philopatry to natal wintering grounds (i.e. Bannister, 2001; Best, 1990; Best et al., 2001; Carroll et al., 2016; Patenaude & Baker, 2001; Payne, 1986). Furthermore, correlations between isotopic profiles ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$), indicative of summer foraging grounds, and maternally inherited mitochondrial DNA haplotypes support the hypothesis that SRW calves learn annual summer foraging destinations from their mothers during a prolonged period of parental care (Carroll et al., 2015; Valenzuela et al., 2009). This transmission of information alludes to the presence of migratory culture in SRWs, where culture is ‘information or behaviour – shared by a population or subpopulation – which is acquired from conspecifics through some form of social learning’ (Rendell & Whitehead, 2001, p. 364). Culture is believed to have evolved as a highly adaptive strategy to pass valuable information between conspecifics (Keith & Bull, 2017). Such transmission of information can occur within generations (horizontally), or in this case, across generations (vertically; Whitehead et al., 2004). It is important to note, however, that all behaviour is the result of interactions between genetics and the environment—a concept well illustrated by a network-based diffusion analysis by Allen et al. (2013), which allowed for the simultaneous consideration of ecological, social and genetic factors as drivers of learned behaviour in humpback whales (*Megaptera novaeangliae*).

Philopatry can be particularly advantageous in relatively stable environments (Keith & Bull, 2017), for example, in aiding individuals to find suitable foraging grounds in a vast ocean. However, on a species level, philopatry (assuming a degree of diversity) can also allow species to track changes in the environment (Kokko & Sutherland, 2001). For example, SRWs in the southwest Atlantic have been found to exhibit diverse foraging strategies (Valenzuela et al., 2018) which may allow the species to cope with potential productivity changes at some of their foraging grounds.

However, in the face of extremely rapid anthropogenic climate change (Bindoff et al., 2019), conserved decisions in SRWs brought about by migratory culture may prevent the exploitation of new feeding areas, and may even become maladaptive if conditions in many of the established foraging grounds become less optimal (Keith & Bull, 2017; Whitehead et al., 2004). In these instances, rapid anthropogenic climate change can lead to ecological traps (as defined by Keith & Bull, 2017; Schlaepfer et al., 2002), wherein environmental cues which individuals use to assess habitat quality become decoupled from the habitat's true quality, causing individuals to make maladaptive habitat choices. Nonetheless, innovation from individual SRWs, followed by horizontal cultural transmission, may provide a highly effective method to facilitate adaptive behavioural changes in SRW populations (Keith & Bull, 2017; Whitehead et al., 2004).

Previous studies have inferred changes in South American SRW foraging success via inference from changes in their calving output and correlations with global climate indices (Leaper et al., 2006; Seyboth et al., 2016). More recently, the SRW population from Península Valdés in Argentina has been shown to have a decreased growth rate between 2007 and 2016 (Crespo et al., 2019). Interestingly, the reproductive output of the South African SRW population is characterized by an initial period (1971–2006) with high calving rates (Brandão et al., 2010) followed by a second period during the late 2010s with reduced calving rates (Brandão et al., 2018; Vermeulen et al., 2018), and an associated decreased population growth rate (Brandão et al., 2018). Being capital breeders, an associated reduction in foraging success in South African SRWs may be assumed. We therefore aim to investigate temporal trends in their foraging strategies, using stable isotope analyses.

Stable carbon ($\delta^{13}\text{C}$) and nitrogen ($\delta^{15}\text{N}$) isotope ratios of skin biopsy samples have been widely used in the study of cetacean foraging ecology (Hooker et al., 2001; Ruiz-Cookey et al., 2004; Todd, 1997; Valenzuela et al., 2018; Witteveen et al., 2011), likely indicating where whales have foraged in the few months prior to sampling (Busquets-Vass et al., 2017). Stable isotope values (e.g. $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) of tissue samples can thus serve as indicators of foraging grounds, as their signatures are incorporated directly from food sources into consumer tissues, with varying degrees of discrimination (Newsome et al., 2010), and their values have predictable patterns of change across terrestrial and marine systems (Graham et al., 2010; Hobson, 1999).

The isotopic composition of an organism can be used to quantify its 'isotopic niche' or 'δ-space' and can be used as a powerful tool to investigate the ecological niche of that organism (Newsome et al., 2007). Isotopic niche space (range of isotopic values, defined as parts per mil or ‰) and niche area (area occupied in space characterized by

two or more isotopes, ‰²) can reveal patterns of resource partitioning or variation in foraging strategies (e.g. Buelow et al., 2018; Das et al., 2017; Dehnhard et al., 2020; Mendes et al., 2018; Ryan et al., 2013). Additionally, assuming the isotopic composition of a system's baseline is known, isotopic mixing models can take these observations further by reconstructing the proportional diet of the consumer (Phillips et al., 2014; Stock et al., 2018), thus allowing further inference to be made into variation and specialization of cetacean foraging strategies and foraging ground location (e.g. Witteveen et al., 2011).

Importantly in the context of marine megafauna foraging ecology, a significant latitudinal gradient in $\delta^{13}\text{C}$ values of phytoplankton and zooplankton is found in the world's oceans, a pattern that is particularly accentuated in the Southern Ocean (Cherel & Hobson, 2007; Espinasse et al., 2019; Magozzi et al., 2017; McMahon et al., 2013a). This gradient is partly due to processes driven by temperature variations (McMahon et al., 2013a), CO_2 concentrations (Peterson & Fry, 1987) and variations in the metabolism of the primary producer with latitude (Hobson et al., 2010). The latitudinal gradient is a well-established principle in top marine predator foraging ecology, it shows temporal stability across decades (Logan et al., 2020), and has proven an effective indicator of broad foraging habitats in both short (i.e. ≤ 4 years; e.g. Cherel & Hobson, 2007; Lübcker et al., 2017; Valenzuela et al., 2018) and long-term studies (i.e. ≥ 14 years; e.g. Logan et al., 2020; Mestre et al., 2020).

Here, we take this approach and assess the isotopic niche space and diet composition of South African SRWs sampled during a period with high calving rates (the 1990s) and during a period with reduced calving rates (the late 2010s). Should the stable isotope values of South African SRWs remain constant across time, then this would be indicative of consistent foraging strategies. Here, the population could be subject to an ecological trap, driven by the combination of being highly philopatric capital breeders showing fidelity to a foraging ground that has become suboptimal in an ocean experiencing change. Alternatively, should there be substantial changes in the isotopic values of South African SRWs across time, then we infer that the changes reflect alterations in foraging behaviour. This could be an indication of adaptation to changing oceanic conditions, but given the recent declines in reproductive output, the shift in foraging may have led to SRWs feeding in less productive regions or on less nutritious prey. Regardless, as long-lived, wide-ranging ocean predators, the response of SRWs to changing oceanic conditions will likely be indicative of broad-scale productivity changes in their foraging grounds in general.

2 | METHODS

2.1 | Sample collection

South African SRW skin biopsy samples (Table 1) analysed in this study are a mixture of previously collected (1990s, 2015 and 2016) and recently collected samples (2019). The recently collected samples were obtained using small stainless steel biopsy darts deployed from a crossbow (Lambertsen, 1987) on SRWs in Walker Bay, Hermanus (34°26'S, 19°18'E) and in San Sebastian Bay, Witsand

TABLE 1 Southern right whale skin samples used for isotopic analyses, shown by sampling year and age class: cows (adult-nursing females) and unaccompanied adults (males, non-nursing females, pregnant females, juveniles). Unknown age classes are non-calves

Decade	Year	Sample size (n)	Sampling month					Age class		
			July	August	September	October	November	Cows	Unaccompanied adults	Unknown
1990	1995	13	3	6	0	4	0	1	11	1
	1996	13	7	0	0	6	0	6	2	5
	1997	18	0	0	13	0	5	17	1	0
	Total	44	10	6	13	10	5	24	14	6
2010	2015	2	0	0	0	2	0	0	0	2
	2016	27	2	5	14	6	0	1	9	17
	2019	49	0	0	49	0	0	41	8	0
	Total	78	2	5	63	8	0	42	17	19
Summary		122	12	11	76	18	5	66	31	25

(34°23'S 20°52'E; Figure S1) in September 2019 (close to the August SRW calving peak; Best, 1994). Samples were stored frozen until stable isotope analyses. Prior to obtaining a biopsy sample, individual whales were photo-identified based on natural markings (Payne et al., 1983) using a DJI phantom 4 pro drone (35 cm diameter, 1388 g, www.dji.com), thus avoiding re-sampling of whales. Age classes (unaccompanied adults/juveniles) were visually determined based on the size of individuals, and adult-nursing females (cows) were identified by the presence of calves. Previously collected samples include those collected from free-ranging SRWs on South African coastal waters (Figure S1) from 1995 to 1997 (biopsy samples stored in dimethyl sulfoxide, DMSO; Best et al., 2003) and 2015–2016 (sloughed skin stored in ethanol; Carroll et al., 2020). The effect of DMSO storage is removed by lipid extraction (Burrows et al., 2014; Busquets-Vass et al., 2017; Newsome et al., 2018; Todd et al., 1997) and storage in ethanol has no effect on the isotopic composition of animal tissues (Hobson et al., 1997). Nevertheless, we assessed the C/N ratios of all samples to ensure that lipid extraction had normalized the C/N ratio across samples which were preserved differently.

2.2 | Stable isotope analysis

All skin samples underwent freeze-drying and lipid extraction following Todd et al. (1997). Approximately 0.5–0.6 mg of dried skin was weighed into tin capsules pre-cleaned with toluene. For the 1990s and 2015/2016 samples, carbon ($\delta^{13}\text{C}$) and nitrogen ($\delta^{15}\text{N}$) isotope ratios were measured using a Costech 4010 elemental analyzer coupled to a Thermo Scientific Delta V isotope ratio mass spectrometer at the Stable Isotope Biogeochemistry Laboratory (SIBL, Durham University, UK). The $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ ratios of the 2019 samples were measured using an elemental analyser (Flash EA 1112 Series) coupled to a Delta V Plus stable light isotope ratio mass spectrometer via a ConFlo IV system, housed at the Stable Isotope Laboratory, Mammal Research Institute, University of Pretoria (Pretoria, South Africa). Analyses in all laboratories used

the internationally accepted standards Vienna Pee Dee Belemnite (VPDB) for $\delta^{13}\text{C}$, and atmospheric nitrogen for $\delta^{15}\text{N}$ (AIR); units are expressed as parts per thousand, or per mil (‰) to ensure comparability. Analytical precision for $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ of the 1990 and 2015/2016 samples were assessed via analyses of in-house and international reference materials for each daily run, which were stringently calibrated against international standards (e.g. USGS 40, IAEA 600, IAEA N2), and was measured to be $\pm 0.1\text{‰}$ (SD; Carroll et al., 2015). Analytical precision for $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ of the 2019 samples was monitored via routine analyses of in-house standards (Merck Gel: $\delta^{13}\text{C} = -20.26\text{‰}$, $\delta^{15}\text{N} = 7.89\text{‰}$, C% = 41.28, N% = 15.29 and DL-Valine: $\delta^{13}\text{C} = -10.57\text{‰}$, $\delta^{15}\text{N} = -6.15\text{‰}$, C% = 55.50, N% = 11.86), which were calibrated against international standards (NBS 22, IAEA-CH-3, IAEA-CH-6, IAEA-CH-7, IAEA N-1, IAEA N-2, IAEA NO-3), and was measured to be $\pm 0.08\text{‰}$ (SD) for both carbon and nitrogen, respectively.

2.3 | Statistical analyses

$\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values for South African SRW skin samples ($n = 122$) were plotted in R v3.6.3 (R Core Team, 2020) to visually inspect the data for any trends. Distributions of stable isotope values were assessed for normality using the Shapiro–Wilk test. Differences in stable isotope values of SRW skin samples grouped into (a) decade; (b) year and (c) month of sampling were assessed via Kruskal–Wallis tests and post hoc Dunn's multiple comparisons tests (Dunn, 1964). Statistical analyses were conducted in R v3.6.3 (R Core Team, 2020).

2.4 | Southern right whale isotopic niche space

We estimated, and examined trends in, the isotopic niche space (‰) of South African SRWs sampled during the two decades via the Stable Isotope Bayesian Ellipses (SIBER) package (version 2.1.4) in R (Jackson et al., 2011). Bivariate ellipses were used to delineate

isotopic niche space ($\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ value 95% confidence interval ellipses) for SRWs sampled in the 1990s and 2010s, respectively, and the total isotopic niche space for the sampled population. Bayesian standard ellipses areas (SEA_B) were plotted using the SIBER routine to show niche overlap. Niche area and overlap ($\%^2$) were estimated based on 1,000,000 Markov chain Monte Carlo (MCMC) iterations, after 100,000 burn-ins, implemented in rjags (Plummer, 2018). SEA_B were fitted to the data using a vague normal prior on the means and a vague Inverse Wishart prior on the covariance matrix of the sample (Jackson et al., 2011). Convergence was assessed using the Gelman and Rubin (1992) scale reduction factor with values below 1.1 indicating convergence.

2.5 | Dietary reconstruction

To further our understanding of South African SRW foraging ecology, we estimated the proportional composition of South African SRW diet using isotopic mixing models. This was done in two parts: the compilation of a reference zooplankton dataset and the running of mixing models.

2.5.1 | Zooplankton stable isotope values

SRWs feed on copepods and krill (Tormosov et al., 1998), and those that winter in South African waters are thought to have at least three

summer foraging grounds (Figure 1; Best, 2007). However, these locations are based largely on illegal Soviet catch data (Tormosov et al., 1998) and Townsend's (1935) charts of open boat whaling catches, and their use by the contemporary population is not certain (Best, 2007). Contemporary information from satellite tags deployed on SRWs in South Africa in 2001 revealed that the Subtropical Convergence (STC), the Polar Front (PF) and the west coast of South Africa are likely important for South African SRW foraging (Mate et al., 2011).

Given the remote location of foraging grounds of South African SRWs (see Figure 1), preventing the direct sampling of prey, we obtained $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of likely prey taxa from the SRWs' assumed historical and contemporary foraging grounds (Figure 1; Best, 2007) from the literature (Table S1), as done in other studies using mixing models to estimate diet composition in marine mammals (i.e. Lübcker et al., 2017; Valenzuela et al., 2018). However, to account for the uncertainty in contemporary foraging grounds, additional published data were acquired for areas known to serve as feeding grounds for SRWs in the Southwest Atlantic Ocean (acquired from Valenzuela et al., 2018). The geographical locations (Table S1) correspond to sampling locations in the original study. All zooplankton samples underwent lipid extraction or had a lipid correction model applied to their $\delta^{13}\text{C}$ values (Table S1). Zooplankton stable isotope data from broad overlapping oceanographic regions were grouped together to aid the data visualization and analyses. For example, the area code 'Far South'

FIGURE 1 Map of the South Atlantic, Southwest Indian and Southern Ocean, with blue shading indicating the summer foraging grounds of the South African population of southern right whales, as suggested by Best (2007); yellow shading: the major winter breeding grounds of southern right whales within the map extent; circles, squares and triangles: the locations of zooplankton samples acquired from the literature and whose stable isotope profiles are used in this study, organized by area and prey source codes: Far South (FS), Polar Front (PF), Scotia Sea (SS), Subtropical Convergence (STC), west and south coasts of South Africa (SA), west coast of Namibia (Nam), South Georgia (SG), Marion Island (MI); copepods (C), euphausiids (E); positions of the Subtropical Front (STF), Subantarctic Front (SAF), Polar Front (PF; Orsi & Harris, 2019); and the minimum and maximum sea ice extents for the period of 1989–1999 (Smith & Jacka, 2003)

(FS) consists of zooplankton collected at 55°S, 65°S, the Lazarev Sea and the Weddell Gyre (Figure 1). The remaining prey sources were grouped to represent the *Polar Front* (PF), *Scotia Sea* (SS), *Subtropical Convergence* (STC), *west coast of Namibia* (Nam), the *west and south coasts of South Africa* (SA), *South Georgia* (SG) and *Marion Island* (MI; Figure 1; Table S1).

2.5.2 | Bayesian mixing models

Bayesian stable isotope mixing models, implemented in R using the MixSIAR package (version 3.1.10; Stock & Semmens, 2016a), were used to estimate the proportional composition of prey (i.e. sources) to the diet of South African SRWs. Adopting a Bayesian mixing model approach overcomes certain limitations of linear mixing models, and allows for the estimation of proportional contributions of diet in undetermined systems (i.e. more sources than isotopes) while incorporating uncertainties in the isotopic values of sources and trophic enrichment factors (TEFs; Parnell et al., 2010).

Among several other important considerations (see Phillips et al., 2014), mixing models with more than seven prey sources are unlikely to yield precise and interpretable results (Stock et al., 2018), and mixing models cannot differentiate between sources with similar isotopic signatures (Phillips et al., 2014). Thus, isotopically similar copepod and euphausiid sources from the same region (see area codes in Figure 1) were grouped (Phillips et al., 2014). $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ value means and SD of the various source groups were then acquired from source distributions which we calculated via weighting the means of each taxon by sample size (Phillips et al., 2014). We scrutinized isotope mixing polygons (isospaces) to ensure that the sources from each geographical region (see area codes in Figure 1) were sufficiently isotopically dissimilar, and as a result, we assessed two mixing models with slightly different combinations of sources. The mixing models assessed employed multiple data for the consumer (individual whales, $n = 122$), mean $\pm$ SD and sample size for each potential source (Table S2), and TEFs with their associated isotopic variability (i.e. mean $\pm$ SD).

The proportional contribution of each potential source to the diet of South African SRWs was estimated by running three MCMC chains, each comprising 3,000,000 iterations of which the first 1,500,000 iterations were discarded. Sampling (during the last 1,500,000 iterations) was conducted at every 500th iteration. Potential sources had uniform (i.e. assumed non-informative) prior distributions. As suggested by Stock and Semmens (2016b), a process $\times$ residual error structure was implemented. TEFs used were $1.28 \pm 0.38\text{‰}$ for $\delta^{13}\text{C}$ values and $2.82 \pm 0.3\text{‰}$ for $\delta^{15}\text{N}$ values (Borrell et al., 2012). Model convergence was assessed via the Gelman and Rubin (1992) scale reduction factor with values less than 1.1 indicating convergence and the Geweke test, a two-sided z-test comparing the mean of the first part of the chain with the mean of the second part (Stock & Semmens, 2016a). Finally, different mixing models were assessed using the widely applicable information criterion (WAIC) and approximate leave-one-out cross-validation (LOO; Stock et al., 2018; Vehtari et al., 2017).

3 | RESULTS

3.1 | Stable isotope ratios of skin samples

SRW skin biopsy samples collected in South Africa over the period of 1995–2019 had an overall mean $\delta^{13}\text{C}$ of $-21.98 \pm 1.89\text{‰}$ (range -25.95 to -16.28‰ , $n = 122$) and a mean $\delta^{15}\text{N}$ of $7.12 \pm 1.03\text{‰}$ (range: 5.01 – 12.19‰ , $n = 122$). Samples collected in the 1990s had a mean $\delta^{13}\text{C}$ of $-23.76 \pm 1.17\text{‰}$ (range -25.95 to -20.46‰ , $n = 44$) and a mean $\delta^{15}\text{N}$ of $7.14 \pm 0.43\text{‰}$ (range 6.46 – 7.94‰ , $n = 44$). Samples collected in 2010s had a mean $\delta^{13}\text{C}$ of $-20.98 \pm 1.43\text{‰}$ (range -23.61 to -16.28‰ , $n = 78$) and a mean $\delta^{15}\text{N}$ of $7.12 \pm 1.25\text{‰}$ (range 5.01 – 12.19‰ , $n = 78$; see Supporting Information 1 for full dataset). The $\delta^{13}\text{C}$ distribution of the combined dataset did not significantly violate the assumption of normality (Shapiro–Wilk W-test: $n = 122$; $p > 0.05$; Figure 2), while the $\delta^{15}\text{N}$ distribution of the combined dataset was significantly non-normal (Shapiro–Wilk W-test: $n = 122$; $p < 0.001$; Figure 2). Visual inspection of isotope values suggests substantial isotopic differentiation between SRW skin biopsies collected in 1995, 1996 and 1997 compared to 2015, 2016 and 2019 (Figure 2).

Skin samples collected in different decades (1990 and 2010) show significant differences in $\delta^{13}\text{C}$ values (Kruskal–Wallis [K-W] $\chi^2 = 66.3$, $p < 0.001$; Figure S2: see Supporting Information 2 for all R code used in the analysis) but not in $\delta^{15}\text{N}$ values (K-W $\chi^2 = 1.79$, $p > 0.05$; Figure S2). Skin samples collected in different years show significant differences in $\delta^{13}\text{C}$ (K-W $\chi^2 = 66.5$, $p < 0.001$; Figure 3) and in $\delta^{15}\text{N}$ (K-W $\chi^2 = 12.7$, $p < 0.05$; Figure 3). Post hoc Dunn's

FIGURE 2 $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values from 122 southern right whale skin biopsy samples collected over the period of 1995–2019. Values are grouped by the year (colour) and decade (1990s are open circles, 2010s are filled circles) of sample collection. Frequency distributions of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ are shown as marginal histograms. $\delta^{13}\text{C}$ values are normally distributed (Shapiro–Wilk test: $n = 122$; $p > 0.05$), while $\delta^{15}\text{N}$ values are not normally distributed (Shapiro–Wilk test: $n = 122$; $p < 0.001$)

multiple comparisons tests indicated that $\delta^{13}\text{C}$ values for the years 2019, 2016 and 2015 were all significantly higher than the $\delta^{13}\text{C}$ values for the years 1995, 1996 and 1997 (Figure 3; Tables S3 and S4). Post hoc Dunn's multiple comparisons tests indicated that $\delta^{15}\text{N}$ values for the year 1995 were significantly higher than the $\delta^{15}\text{N}$ values for the years 1996, 1997, 2016 and 2019 (Figure 3; Tables S3 and S5).

Skin samples collected in different months within the decade of 1990 show significant differences in $\delta^{15}\text{N}$ (K-W $\chi^2 = 11.2$, $p < 0.05$; Figure S3) but not in $\delta^{13}\text{C}$ values (K-W $\chi^2 = 1.9$, $p > 0.05$; Figure S3). Post hoc Dunn's multiple comparisons tests indicated that $\delta^{15}\text{N}$ values for the month of September were lower than the $\delta^{15}\text{N}$ values for the months of July and August (Figure S3; Tables S6 and S7). Skin samples collected in different months within the decade of 2010 show no significant differences in $\delta^{15}\text{N}$ (K-W $\chi^2 = 6.7$, $p > 0.05$; Figure S4) nor in $\delta^{13}\text{C}$ values (K-W $\chi^2 = 3.1$, $p > 0.05$; Figure S4).

3.2 | Southern right whale isotopic niche space

The SIBER assessment of the South African SRW samples collected in two different decades indicated a substantial expansion and shift

FIGURE 3 Stable isotope values ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) from 122 southern right whale skin biopsy samples grouped by year of collection: 1995 ($n = 13$), 1996 ($n = 13$), 1997 ($n = 18$), 2015 ($n = 2$), 2016 ($n = 27$) and 2019 ($n = 49$). Boxplots represent the median, 25th and 75th percentiles, minimum and maximum values

FIGURE 4 Variation in $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values in southern right whale skin biopsy samples ($n = 122$) categorized by decade of collection. (a) 95% CI bivariate ellipses of southern right whale skin biopsy samples collected in 1995–1997 (black dots) and in 2015–2019 (red dots), demonstrating a significant isotopic niche shift and expansion in South African southern right whales in the 2010 decade. (b) Central tendency and measures of uncertainty (50%, 75% and 95% credible intervals, black dot is the mode of each distribution) of Bayesian standard ellipses areas (SEA_b) for southern right whale skin samples collected in 1995–1997 and in 2015–2019

in the isotopic niche space of SRWs sampled in the 2010 decade (Figure 4). SEA_b modes of the 1990 and 2010 SRW samples were 1.69‰^2 and 4.84‰^2 , respectively (Figure 4).

3.3 | Dietary estimations

Overall, zooplankton from coastal areas (Nam & SA) and the STC tended to have higher $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values than those from colder waters further south (Figure S5). When corrected for trophic enrichment (TEF 1.28‰ for $\delta^{13}\text{C}$ and 2.82‰ for $\delta^{15}\text{N}$; Borrell et al., 2012), SRW skin isotope ratios overlapped with ratios for zooplankton from several locations, including South Georgia, the PF, Marion Island, the STC and coastal Africa (Figure S5).

FIGURE 5 MixSIAR isotope mixing polygon (isospace) for mixing model 1 (M1) illustrating source (prey groups) and southern right whale isotope signatures. $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of southern right whale skin biopsy samples are grouped by decade (1990: grey dots, 2010: black dots). Each source is corrected for trophic level. The error bars in the sources correspond to the standard deviations calculated from data in Table S1. Sources are colour coded by region and prey source composition is indicated by euphausiid and copepod clipart. See Table S2 for uncorrected prey source statistical summaries

Sources from the west coast of Namibia and sources from the STC could not be differentiated by mixing models, and thus, given the lack of evidence for South African SRWs making use of coastal Namibia as a foraging ground (Best, 2007; Mate et al., 2011), we excluded this source from the mixing model source pool. All sampled areas in which both prey types were present, apart from the PF, had isotopically similar signatures for copepods and euphausiids (Figure S5). Thus, to account for the isotopic dissimilarity between copepods and euphausiids from the PF, we ran two different mixing models in MixSIAR. The first mixing model (M1) had seven sources, each source being a combination, where possible, of copepods and euphausiids from the same region (Figure 5). The second model (M2) also had seven sources, but here with copepods

and euphausiids from the PF separated (Figure S6). Mixing model M2 could not differentiate between copepods from the PF and euphausiids from South Georgia (see Figure S5), so euphausiids from South Georgia were excluded from mixing model M2 (Figure S6).

LOO statistics provide a strong weighting preference for mixing model M1 over M2; thus, the graphical results for mixing model M2 can be found in the supplement (Figures S6 and S7; Table S9). Mixing model M1 estimated that euphausiids from South Georgia contribute 43% (median, with 95% Bayesian credible intervals: 25%–59%, see Table S8) to the diet of SRWs sampled in the 1990 decade (Figure 6). Smaller dietary contributions to the 1990 decade were estimated with copepods and euphausiids from Far South (FS) contributing 16% (5%–27%) and euphausiids from Marion Island contributing 14% (1%–27%; Figure 6).

FIGURE 6 MixSIAR model outputs for mixing model M1 showing estimates (median, 50% and 95% credibility intervals) of diet composition for South African southern right whales (SRW) sampled in 1995–1997 and 2015–2019. FS is Far South, MI is Marion Island, PF is Polar Front, SG is South Georgia, SS is Scotia Sea, STC is Subtropical Convergence, SA is west and south coasts of South Africa, and euphausiids and copepods are represented by clipart. The two distinct mixing model potential outcomes for SRWs sampled in 2015–2019 are indicated by S1 and S2

Dietary contributions estimated for SRWs sampled in the 2010 decade by M1 were complex, with the presence of three bimodal posterior distributions representing two distinct potential outcomes (Figure 6). Specifically, the diets of the SRWs sampled in the 2010 decade either comprised primarily of euphausiids from Marion Island (78%, 0–91%) or both copepods and euphausiids from the PF (7%, 0%–34%) and STC (4%, 0%–63%). All remaining sources used in model M1 had potential contributions of <10%.

Mixing model M2 estimated that copepods and euphausiids from Far South (FS) contribute 33% (24%–42%) to the diet of SRWs sampled in the 1990 decade. Smaller dietary contributions to the 1990 decade were estimated with euphausiids from Marion Island contributing 21% (11%–32%) and copepods from the PF contributing 17% (3%–31%; Figure S7; Table S8). For SRWs sampled in the 2010 decade, only a single source, euphausiids from Marion Island, were estimated to substantially contribute to SRW diet, with a contribution of 84% (72%–93%; Figure S7). All remaining sources used in model M2 had potential contributions of <10%.

4 | DISCUSSION

The detection of changes in marine megafauna population dynamics and foraging ecology provides insight into how anthropogenic change can influence natural systems (e.g. Davis et al., 2017, 2020; Fleming et al., 2016). Here, we used stable isotope analyses of skin biopsy samples to investigate long-term shifts in foraging ecology of South African SRWs. Results show that the population has recently shifted and diversified its foraging strategies, suggesting that their previously productive foraging grounds have changed over time. Considering the vast oceanic range of SRW foraging grounds, our findings point towards large-scale ecosystem changes in the Southern Ocean. This is perhaps not surprising, given the unprecedented impacts of recent climate change on Southern Ocean physical features, which, in turn, have driven regional changes on all levels of Antarctic marine food webs (see Rogers et al., 2020 and references therein). The shift in foraging strategies is also concurrent with a time period associated with lowered reproductive output in the South African SRW population (Brandão et al., 2018; Vermeulen et al., 2018). As highly migratory capital breeders, SRWs have a strong relationship between foraging success and their reproductive output (Leaper et al., 2006; Lockyer, 2007; Seyboth et al., 2016); therefore, questions arise on their recent success in foraging.

4.1 | Decadal shift in southern right whale isotope values

Results of this study demonstrate a significant shift in $\delta^{13}\text{C}$ and a broadening of $\delta^{15}\text{N}$ values (Figure 3) of SRW skin biopsy samples over the period 1995–2019. SRWs sampled in the 1990-decade have lower $\delta^{13}\text{C}$ values than the 2010-decade, and their

reconstructed diet consists predominantly of prey sources from regions with isotopic signatures similar to euphausiids from South Georgia. Our findings for the 1990-decade are supported by isotopic data from 11 South African SRW baleen plates collected from 1963 to 1994 (Best & Schell, 1996), which suggested foraging occurred predominantly south of the STC. In contrast, SRWs sampled in the 2010-decade, with higher $\delta^{13}\text{C}$ values in skin, likely foraged further north, in regions with isotopic signatures similar to those found at the STC, PF and Marion Island. While the shift in diet is clear, the mixing model found two possible outcomes for the 2010-decade between which it could not distinguish (1) euphausiids from Marion Island or (2) copepods and euphausiids from the PF and STC. Despite the model converging, these two scenarios are potentially confounded as Marion Island is located between the PF and STC along an isotopic gradient, rather than being discrete isotopic regions. Prey sources from the STC are isotopically discrete from those collected at Marion Island and the PF (Figure 5); and despite post hoc tests indicating isotopic similarity between sources from the PF and Marion Island, the mixing model results do not suggest that the two regions are isotopically equivalent. Additionally, whaling records, photo-identification, and satellite tagging data reveal that both Marion Island and the PF represent likely foraging grounds used by South African SRWs (IWC, 2013; Mate et al., 2011; Smith et al., 2012). Therefore, prey sources from Marion Island, the PF and STC were retained in the mixing model.

The mixing models presented here reveal that a northward shift in the foraging of South African SRWs has likely occurred recently. The findings of Tormosov et al. (1998), which show that SRWs mostly feed on krill when south of 50°S, and copepods when north of 40°S, suggest that the northward shift in foraging presented here is likely accompanied by an increasing proportion of copepods in the diet of SRWs. Further work using microchemical markers or information from offshore movements of satellite-tagged SRWs would allow further inference to be made into South African SRW foraging ecology and diet composition. Fatty acids are widely used to assess the foraging ecology of marine mammals (e.g. Budge et al., 2008; Haug et al., 2017; Herman et al., 2005; Hooker et al., 2001; Marón et al., 2020; Olsen & Grahl-Nielsen, 2003) and they are known to undergo little biochemical change when passed up the food chain (Iverson, 2009). Similarly, compound-specific isotope analyses of amino acids have been proven a valuable tool in top predator foraging ecology (e.g. Nelson et al., 2018; Pomerleau et al., 2017; Zupcic-Moore et al., 2017) as it provides information about essential amino acids which are preserved unchanged through the food chain (Larsen et al., 2013). Incorporating compound-specific analyses and *a priori* narrowing likely foraging grounds with satellite tag work into the Bayesian mixing model approach would further amplify the ecosystem baseline signals, inform model prior distributions (see Moore & Semmens, 2008) and would ultimately permit more accurate estimation of diet composition in SRWs.

The evidence for a shift in foraging strategy in South African SRWs presented here, occurred between a period characterized by

rapid population growth (1990-decade; Brandão et al., 2010) and one characterized by an increase in calving intervals and decreased growth rate (2010-decade; Brandão et al., 2018; Vermeulen et al., 2018). It is unlikely that this suboptimal reproductive output and the shift in foraging strategy we present could have occurred coincidentally, given (1) SRWs are capital breeders, which rely on accumulated energy reserves for successful reproduction (Lockyer, 2007); (2) periods of suppressed reproduction have been shown to occur during periods of nutritional stress in cetaceans (Greene et al., 2003; Hlista et al., 2009; Lockyer, 1986; Reeves et al., 2001; Ward et al., 2009; Williams et al., 2013) and (3) the reproductive output of SRWs in Argentina and Brazil has been found to respond to fluctuations in krill abundance (Leaper et al., 2006; Seyboth et al., 2016). These concurrent events thus suggest not only a shift in foraging strategy but also a decreased foraging success in the 2010-decade.

Alternatively, the shift in foraging strategy in South Africa's population of SRWs may be related to SRW population recovery. A higher density of SRWs in the last decade could lead to more intra-specific competition for potentially shrinking prey sources, thus resulting in a fraction of the population having suboptimal foraging success, and potentially explaining the lowered reproductive output seen in South Africa. However, the mechanisms by which SRWs will respond to approaching their carrying capacity are not yet fully understood and the data to test this idea are not available. Furthermore, the pre-exploitation, global historical SRW population size has been estimated to consist of 101,882 individuals (95% posterior probability 88,282–123,780; Jackson et al., 2008), while the most recent estimate of global abundance indicates a population size of approximately 13,611 SRWs (IWC, 2013), thus, the global population does not appear close to full recovery.

Had the foraging strategy of South African SRWs remained constant during the 1990s and 2010s, then further investigation into the hypothesis of an ecological trap may be warranted, whereby SRWs may be making maladaptive habitat choices based on former reliable environmental information (Schlaepfer et al., 2002), leading to poor body condition and suboptimal reproductive output. However, we provide evidence for the inverse: SRWs appear to have shifted and potentially diversified their foraging strategies, possibly making use of anomalously high productivity at the STC (Del Castillo et al., 2019), yet still displaying suboptimal reproductive output. Although it is intuitive to suggest that diversification in foraging may provide some degree of resilience to large-scale oceanic changes (see Cartwright et al., 2019)—our results, combined with the increase in calving intervals and decreased population growth rate (Brandão et al., 2018; Vermeulen et al., 2018), suggest that the potential foraging plasticity provided by diversifying foraging strategies may in fact not be sufficient to ensure continued foraging success, and therefore successful reproduction, in South African SRWs.

The shift in foraging strategy could be linked to an increase in productivity in the region between the STC and PF: the Sub Antarctic Zone; hereafter SAAZ. Over the period 1987–2018, Del Castillo et al. (2019) found strong increases in surface chlorophyll concentrations in the Pacific, Atlantic and Indian sectors of the

Southern Ocean, particularly in the SAAZ. Furthermore, since approximately 2010, winter months in the Atlantic region of the SAAZ have seen extremely high chlorophyll concentrations, characteristic of summer month concentrations (Del Castillo et al., 2019). We hypothesize that this expansion of productivity in the SAAZ into winter has resulted in year-round high productivity at specific SRW foraging grounds. Given the high productivity in the SAAZ, South African SRWs may not need to migrate to foraging grounds south of the PF. Alternatively, they may still migrate to foraging grounds south of the PF but also take advantage of high productivity in the SAAZ on their northwards migration. How this may have led to an apparent decreased foraging success remains to be determined.

4.2 | Cetaceans as indicators of large-scale oceanic changes

Identifying species that may be useful for elucidating the links between physical environmental change and the ecological responses of top marine predators can provide insight into ecological dynamics and inform predictions for future responses of marine top predators in the face of rapid anthropogenic climate change (Fleming et al., 2016). However, the foraging and migratory responses of cetaceans to climate change appear context-dependant.

Spatial and/or temporal mismatches between the life history or phenology of krill-dependent species and their prey availability, and thus their migratory habits, may be key to understanding the effects of climate change on the survival and reproduction of baleen whales (Tulloch et al., 2019). There is some evidence suggesting substantial flexibility in baleen whale migration. For example, grey whales (*Eschrichtius robustus*) have begun to respond to warming by remaining in Arctic areas over winter (Moore et al., 2007), and Spitsbergen's bowhead whales (*Balaena mysticetus*), unlike larger bowhead whale populations, have recently been found to forage and overwinter in cold, sea-ice associated waters. Dietary flexibility has also been documented; Fleming et al. (2016) found that the diet of humpback whales between 1993 and 2012 in the California Current System (CCS), inferred from isotope data, shifted with available prey resources in response to climate variability, indicating that the species is a promising indicator for ecosystem dynamics in the CCS. Indeed, the North Atlantic right whale (*E. glacialis*), closely related to the SRW, has shown changes in its seasonal distribution in recent years, likely linked to changing prey resources (Davis et al., 2017). This acoustic study, and a more recent one studying North Atlantic right whales, humpback, sei (*Balaenoptera borealis*), fin (*Balaenoptera physalus*) and blue whales (*Balaenoptera musculus*), showed distributional changes in all species that suggested they were keeping pace with environmental changes in the North Atlantic (Davis et al., 2020).

However, historic whaling records and more than 15 years of satellite-derived data have shown that humpback whale migrations in the southwest Atlantic have not changed during dynamic oceanographic and geomagnetic conditions (Horton et al., 2020).

Furthermore, Abrahms et al. (2019) found that both long-term memory and resource tracking play vital roles in blue whale migrations, suggesting that other highly migratory species may struggle to adapt to climate change. What is clear from these examples, as well as the results we present, is that we are seeing differences in baleen whale responses to climate change, and these differences tend to be correlated to poor population recovery. Indeed, although capital breeders like right whales may have some advantage given that their life history strategy decouples environmental productivity at their wintering breeding grounds from reproductive success, their reliance on foraging in some of the most rapidly changing parts of the world (Jansen et al., 2020; Rogers et al., 2020) seems to limit this benefit.

4.3 | Caveats

Our study has several assumptions common to stable isotope analyses: first, stable isotope values in the ecosystem (isotope baseline) have not changed enough over the time period of our study to influence the results; second, the prey dataset and TEF lead to an accurate reconstruction of diet; and third, we are only detecting shifts over the period of integration of isotopes into the whales' skin.

The accurate interpretation of stable isotope values of a consumer requires an understanding of the spatial and temporal variation in isotopic compositions of food web baselines (Magozzi et al., 2017). Disentangling whether a change in a consumer's stable isotope values represents a shift in foraging behaviour or a shift in isotopic baseline is challenging (McMahon et al., 2013b). However, the increase in $\delta^{13}\text{C}$ in South African SRWs is unlikely to be due to ecosystem baseline changes, as it is both substantial and in the opposite direction to the general trend of decreases in oceanic baseline $\delta^{13}\text{C}$ values (de la Vega et al., 2019; Hanson et al., 2009; Lorrain et al., 2020; Mestre et al., 2020; Newsome, Etnier, et al., 2007). Specifically, $\delta^{13}\text{C}$ values are known to decline by approximately -0.1 to -0.2‰ per decade (Sonnerup et al., 1999) due to ocean absorption of anthropogenic carbon emissions, that is, the Suess (1955) effect. As a result, several studies have detected decreases in marine predator $\delta^{13}\text{C}$ values that could not be attributed to shifts in foraging strategies, but rather to changes in abiotic conditions and/or other biotic components of the Southern Ocean (i.e. Suess effect or changes in biological productivity; de la Vega et al., 2019; Hanson et al., 2009; Lorrain et al., 2020; Mestre et al., 2020; Newsome, Etnier, et al., 2007; Quillfeldt et al., 2010). However, these studies have shown a small magnitude of change in $\delta^{13}\text{C}$ (e.g. -1.4‰ from 2004 to 2017 for elephant seals in Mestre et al., 2020; and -1.8‰ for three tuna species from 2000 to 2015 in Lorrain et al., 2020) in the direction of the Suess effect, compared with the mean increase in $\delta^{13}\text{C}$ we present for South African SRWs from -23.76‰ in the 1990s to -20.98‰ in the 2010 (i.e. $+2.78\text{‰}$). Finally, recent work (e.g. Logan et al., 2020) suggests temporal stability in isotopic latitudinal gradients across decades in the Southern Ocean isoscape, thus supporting this supposition.

The isotopic dietary reconstruction presented here relies on the accurate representation of South African SRW prey isotopic values and that the TEF used accurately reflects the isotopic enrichment from SRW prey to SRW skin. We included a carefully vetted copepod and euphausiid dataset (i.e. data were lipid-extracted or lipid normalized) available for regions of the South Atlantic Ocean, the southwest Indian Ocean and the Southern Ocean which South African SRWs may utilize as foraging grounds (Table S1). This approach has some limitations and sources of variance (for more details, please see Valenzuela et al., 2018). Nevertheless, this approach is commonly used in mixing models, as it represents the best available data (Lübcker et al., 2017; Valenzuela et al., 2018). Similarly, we used a TEF calculated for fin whale skin (Borrell et al., 2012) as to the best of our knowledge, a species-specific TEF has not been calculated for SRWs. This TEF ($1.28 \pm 0.38\text{‰}$ for $\delta^{13}\text{C}$ and $2.82 \pm 0.3\text{‰}$ for $\delta^{15}\text{N}$) is similar to the average values for whole animals ($\approx 3\text{‰}$ for nitrogen and 0 to 1‰ for carbon; Peterson & Fry, 1987), and has previously been used in mixing models to reconstruct blue, sei, fin (Silva et al., 2019) and SRW diet (Valenzuela et al., 2018).

The timing of isotopic integration is also an important consideration in this and other isotope ecology studies. For whale skin, integration is thought to be approximately 3–5 months (Busquets-Vass et al., 2017), and so samples collected in the austral winter represent foraging in autumn or late summer. The 1990-decade data presented here, baleen plate analysis (Best & Schell, 1996) and historical whaling data (Smith et al., 2012) all suggest that higher latitude foraging grounds were visited later in summer and into autumn, prior to the northward migration. The work presented here suggests that the STC and PF, rather than foraging grounds further south, are now visited by SRWs prior to the northward migration. However, we cannot rule out that higher latitude foraging grounds are visited in spring or early summer, as the isotopic signatures would no longer be present in the whale skin; modern baleen whale plates would provide further information on this. Furthermore, it is important to acknowledge that SRWs may have subtly different isotopic signatures in their skin despite having visited the same foraging grounds. This can be a result of visiting foraging grounds in a different order, having different levels of foraging effort at each location or by spending different amounts of time at each foraging ground. This is a limitation of many isotope studies on marine mammals and other vagile marine predators.

5 | CONCLUSION

Stable isotope analyses reveal that the South African population of SRWs underwent a significant northward shift, and diversification, in foraging strategy from the 1990s to 2010s. This shift could represent a strategy to cope with changes in preferred prey or habitat, a positive sign that the species could potentially adapt to changing resources. However, the concurrent decline in reproductive success suggests the shift could also be a suboptimal foraging strategy.

This contributes to the growing body of evidence that responses to climate change is context- and species-dependent in cetaceans but highlights how shifting foraging strategies may not be sufficient to cope with a changing environment.

ACKNOWLEDGEMENTS

We wish to thank ExxonMobil, Total and ENI for their support in the collection of the 2019 biopsy samples. These biopsy samples were collected under South African Department of Environmental Affairs research permit (RES2019/19), Cape Nature sample collection permit (CN44-28-5255) and ethical clearance from the University of Pretoria Faculty of Natural and Agricultural Sciences Ethics Committee (NAS271/2019). Permits and ethics for the 1990s samples can be found in Carroll et al. (2015) and for the 2015/16 samples in Carroll et al. (2020). Collection of samples from 2015 to 2016 was funded by Charles University Grant Agency (1140217). Stable isotope analysis of the 1990s and 2015/16 samples was funded by a Newton International Fellowship to E.L.C. and E.L.C. was supported by a Rutherford Discovery Fellowship from the Royal Society of New Zealand Te Apārangi. Copepod, krill, and whale clipart used in this manuscript were provided by Phylopic. Copepod and krill clipart were used under the Public Domain Dedication 1.0 license and the Public Domain Mark 1.0 license, respectively. Whale clipart was provided by Chris Huh, used under the Creative Commons Attribution-ShareAlike 3.0 Unported license, and only clipart colour was edited. The authors have no conflict of interest to declare.

DATA AVAILABILITY STATEMENT

The whale skin isotope data are available in Supporting Information 1 and R code used in the analysis is available in Supporting Information 2. All Supplementary Figures and Tables are in Supplementary Material.

ORCID

Gideon L. van den Berg <https://orcid.org/0000-0002-1789-7382>

Emma L. Carroll <https://orcid.org/0000-0003-3193-7288>

REFERENCES

- Abrahms, B., Hazen, E. L., Aikens, E. O., Savoca, M. S., Goldbogen, J. A., Bograd, S. J., Jacox, M. G., Irvine, L. M., Palacios, D. M., & Mate, B. R. (2019). Memory and resource tracking drive blue whale migrations. *Proceedings of the National Academy of Sciences of the United States of America*, 116(12), 5582–5587. <https://doi.org/10.1073/pnas.1819031116>
- Allen, J., Weinrich, M., Hoppitt, W., & Rendell, L. (2013). Network-based diffusion analysis reveals cultural transmission of lobe feeding in humpback whales. *Science*, 340(6131), 485–488. <https://doi.org/10.1126/science.1231976>
- Atkinson, A., Hill, S. L., Pakhomov, E. A., Siegel, V., Reiss, C. S., Loeb, V. J., Steinberg, D. K., Schmidt, K., Tarling, G. A., Gerrish, L., & Sailley, S. F. (2019). Krill (*Euphausia superba*) distribution contracts southward during rapid regional warming. *Nature Climate Change*, 9(2), 142–147. <https://doi.org/10.1038/s41558-018-0370-z>
- Atkinson, A., Siegel, V., Pakhomov, E., & Rothery, P. (2004). Long-term decline in krill stock and increase in salps within the Southern Ocean. *Nature*, 432(7013), 100–103. <https://doi.org/10.1038/nature02996>
- Bannister, J. (2001). Status of southern right whales (*Eubalaena australis*) off Australia. *Journal of Cetacean Research & Management*, 2, 103–110.
- Barbosa, A., Benzal, J., de León, A., & Moreno, J. (2012). Population decline of chinstrap penguins (*Pygoscelis antarctica*) on Deception Island, South Shetlands, Antarctica. *Polar Biology*, 35(9), 1453–1457. <https://doi.org/10.1007/s00300-012-1196-1>
- Best, P. B. (1990). Trends in the inshore right whale population off South Africa, 1969–1987. *Marine Mammal Science*, 6(2), 93–108. <https://doi.org/10.1111/j.1748-7692.1990.tb00232.x>
- Best, P. B. (1994). Seasonality of reproduction and the length of gestation in southern right whales *Eubalaena australis*. *Journal of Zoology*, 232(2), 175–189. <https://doi.org/10.1111/j.1469-7998.1994.tb01567.x>
- Best, P. B. (2007). *Whales and dolphins of the southern African subregion*. Cambridge University Press.
- Best, P. B., Brandão, A., & Butterworth, D. S. (2001). Demographic parameters of southern right whales off South Africa. *Journal of Cetacean Research and Management Special Issue*, 2(2), 161–169.
- Best, P. B., Schaeff, C. M., Reeb, D., & Palsbøl, P. J. (2003). Composition and possible function of social groupings of southern right whales in south African waters. *Behaviour*, 140(11/12), 1469–1494. <https://www.jstor.org/stable/4536102>
- Best, P. B., & Schell, D. (1996). Stable isotopes in southern right whale (*Eubalaena australis*) baleen as indicators of seasonal movements, feeding and growth. *Marine Biology*, 124(4), 483–494. <https://doi.org/10.1007/BF00351030>
- Bindoff, N. L., Cheung, W. W. L., Kairo, J., Aristegui, J., Guinder, V. A., Hallberg, R., Hilmi, N., Jiao, N., Karim, M., Levin, L., O'Donoghue, S., Purca Cuicapusa, S. R., Rinkevich, B., Suga, T., Tagliabue, A., & Williamson, P. (2019). Changing ocean, marine ecosystems, and dependent communities. *IPCC special report on the ocean and cryosphere in a changing climate* (pp. 447–588). United Nations.
- Böning, C. W., Dispert, A., Visbeck, M., Rintoul, S. R., & Schwarzkopf, F. U. (2008). The response of the Antarctic circumpolar current to recent climate change. *Nature Geoscience*, 1(12), 864–869. <https://doi.org/10.1038/ngeo362>
- Borrell, A., Abad-Oliva, N., Gómez-Campos, E., Giménez, J., & Aguilar, A. (2012). Discrimination of stable isotopes in fin whale tissues and application to diet assessment in cetaceans. *Rapid Communications in Mass Spectrometry*, 26(14), 1596–1602. <https://doi.org/10.1002/rcm.6267>
- Bracegirdle, T. J., Connolley, W. M., & Turner, J. (2008). Antarctic climate change over the twenty first century. *Journal of Geophysical Research Atmospheres*, 113(3), 1–13. <https://doi.org/10.1029/2007JD008933>
- Brandão, A., Best, P., & Butterworth, D. S. (2010). Estimates of demographic parameters for southern right whales. *IWC Scientific Committee Report, SC/62/BRG3*, 1–17.
- Brandão, A., Vermeulen, E., Ross-gillespie, A., Findlay, K., & Butterworth, D. S. (2018). Updated application of a photo-identification based assessment model to southern right whales in South African waters, focussing on inferences to be drawn from a series of appreciably lower counts of calving females over 2015 to 2017. *IWC Scientific Committee Report, SC/67B/SH2*. <https://doi.org/10.13140/RG.2.2.14456.88325>
- Budge, S. M., Springer, A. M., Iverson, S. J., Sheffield, G., & Rosa, C. (2008). Blubber fatty acid composition of bowhead whales, *Balaena mysticetus*: Implications for diet assessment and ecosystem monitoring. *Journal of Experimental Marine Biology and Ecology*, 359(1), 40–46. <https://doi.org/10.1016/j.jembe.2008.02.014>
- Buelow, C. A., Reside, A. E., Baker, R., & Sheaves, M. (2018). Stable isotopes reveal opportunistic foraging in a spatiotemporally heterogeneous environment: Bird assemblages in mangrove forests. *PLoS One*, 13(11), <https://doi.org/10.1371/journal.pone.0206145>

- Burrows, D. G., Reichert, W. L., & Bradley Hanson, M. (2014). Effects of decomposition and storage conditions on the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ isotope values of killer whale (*Orcinus orca*) skin and blubber tissues. *Marine Mammal Science*, 30(2), 747–762. <https://doi.org/10.1111/mms.12076>
- Busquets-Vass, G., Newsome, S. D., Calambokidis, J., Serra-Valente, G., Jacobsen, J. K., Aguiñiga-García, S., & Gendron, D. (2017). Estimating blue whale skin isotopic incorporation rates and baleen growth rates: Implications for assessing diet and movement patterns in mysticetes. *PLoS One*, 12(5), 1–25. <https://doi.org/10.1371/journal.pone.0177880>
- Carroll, E. L., Baker, C. S., Watson, M., Alderman, R., Bannister, J., Gaggiotti, O. E., Gröcke, D. R., Patenaude, N., & Harcourt, R. (2015). Cultural traditions across a migratory network shape the genetic structure of southern right whales around Australia and New Zealand. *Scientific Reports*, 5(November), 1–12. <https://doi.org/10.1038/srep16182>
- Carroll, E. L., Fewster, R. M., Childerhouse, S. J., Patenaude, N. J., Boren, L., & Baker, C. S. (2016). First direct evidence for natal wintering ground fidelity and estimate of juvenile survival in the New Zealand southern right whale *Eubalaena australis*. *PLoS One*, 11(1), 1–17. <https://doi.org/10.1371/journal.pone.0146590>
- Carroll, E. L., Ott, P. H., McMillan, L. F., Galletti Vernazzani, B., Neveceralova, P., Vermeulen, E., Gaggiotti, O. E., Andriolo, A., Baker, C. S., Bamford, C., Best, P., Cabrera, E., Calderan, S., Chirife, A., Fewster, R. M., Flores, P. A. C., Frasier, T., Freitas, T. R. O., Groch, K., ... Jackson, J. A. (2020). Genetic diversity and connectivity of southern right whales (*Eubalaena australis*) found in the Brazil and Chile-Peru wintering grounds and the south Georgia (Islas Georgias del Sur) feeding ground. *Journal of Heredity*, 111(3), 263–276. <https://doi.org/10.1093/jhered/esaa010>
- Cartwright, R., Venema, A., Hernandez, V., Wyels, C., Cesere, J., & Cesere, D. (2019). Fluctuating reproductive rates in Hawaii's humpback whales, *Megaptera novaeangliae*, reflect recent climate anomalies in the North Pacific. *Royal Society Open Science*, 6(3), 181463. <https://doi.org/10.1098/rsos.181463>
- Cherel, Y., & Hobson, K. (2007). Geographical variation in carbon stable isotope signatures of marine predators: A tool to investigate their foraging areas in the Southern Ocean. *Marine Ecology Progress Series*, 329, 281–287. <https://doi.org/10.3354/meps329281>
- Christiansen, F., Vivier, F., Charlton, C., Ward, R., Amerson, A., Burnell, S., & Bejder, L. (2018). Maternal body size and condition determine calf growth rates in southern right whales. *Marine Ecology Progress Series*, 592, 267–281. <https://doi.org/10.3354/meps12522>
- Constable, A. J., Melbourne-Thomas, J., Corney, S. P., Arrigo, K. R., Barbraud, C., Barnes, D. K. A., Bindoff, N. L., Boyd, P. W., Brandt, A., Costa, D. P., Davidson, A. T., Ducklow, H. W., Emmerson, L., Fukuchi, M., Gutt, J., Hindell, M. A., Hofmann, E. E., Hosie, G. W., Iida, T., ... Ziegler, P. (2014). Climate change and Southern Ocean ecosystems I: How changes in physical habitats directly affect marine biota. *Global Change Biology*, 20(10), 3004–3025. <https://doi.org/10.1111/gcb.12623>
- Crespo, E. A., Pedraza, S. N., Dans, S. L., Svendsen, G. M., Degradi, M., & Coscarella, M. A. (2019). The southwestern Atlantic southern right whale, *Eubalaena australis*, population is growing but at a decelerated rate. *Marine Mammal Science*, 35(1), 93–107. <https://doi.org/10.1111/mms.12526>
- Das, K., Holleville, O., Ryan, C., Berrow, S., Gilles, A., Ody, D., & Michel, L. N. (2017). Isotopic niches of fin whales from the Mediterranean Sea and the Celtic Sea (North Atlantic). *Marine Environmental Research*, 127, 75–83. <https://doi.org/10.1016/j.marenvres.2017.03.009>
- Davis, G. E., Baumgartner, M. F., Bonnell, J. M., Bell, J., Berchok, C., Bort Thornton, J., Brault, S., Buchanan, G., Charif, R. A., Cholewiak, D., Clark, C. W., Corkeron, P., Delarue, J., Dudzinski, K., Hatch, L., Hildebrand, J., Hodge, L., Klinck, H., Kraus, S., ... Van Parijs, S. M. (2017). Long-term passive acoustic recordings track the changing distribution of North Atlantic right whales (*Eubalaena glacialis*) from 2004 to 2014. *Scientific Reports*, 7(1), 1–12. <https://doi.org/10.1038/s41598-017-13359-3>
- Davis, G. E., Baumgartner, M. F., Corkeron, P. J., Bell, J., Berchok, C., Bonnell, J. M., Bort Thornton, J., Brault, S., Buchanan, G. A., Cholewiak, D. M., Clark, C. W., Delarue, J., Hatch, L. T., Klinck, H., Kraus, S. D., Martin, B., Mellinger, D. K., Moors-Murphy, H., Niekirk, S., ... Van Parijs, S. M. (2020). Exploring movement patterns and changing distributions of baleen whales in the western North Atlantic using a decade of passive acoustic data. *Global Change Biology*, 26(9), 4812–4840. <https://doi.org/10.1111/gcb.15191>
- de la Vega, C., Jeffreys, R. M., Tuerena, R., Ganeshram, R., & Mahaffey, C. (2019). Temporal and spatial trends in marine carbon isotopes in the Arctic Ocean and implications for food web studies. *Global Change Biology*, 25(12), 4116–4130. <https://doi.org/10.1111/gcb.14832>
- Dehnhard, N., Achurch, H., Clarke, J., Michel, L. N., Southwell, C., Sumner, M. D., Eens, M., & Emmerson, L. (2020). High inter- and intraspecific niche overlap among three sympatrically breeding, closely related seabird species: Generalist foraging as an adaptation to a highly variable environment? *Journal of Animal Ecology*, 89(1), 104–119. <https://doi.org/10.1111/1365-2656.13078>
- Del Castillo, C. E., Signorini, S. R., Karaköylü, E. M., & Rivero-Calle, S. (2019). Is the southern ocean getting greener? *Geophysical Research Letters*, 46(11), 6034–6040. <https://doi.org/10.1029/2019GL083163>
- Dunn, O. J. (1964). Multiple comparisons using rank sums. *Technometrics*, 6(3), 241–252. <https://doi.org/10.1080/00401706.1964.10490181>
- Espinasse, B., Pakhomov, E. A., Hunt, B. P. V., & Bury, S. J. (2019). Latitudinal gradient consistency in carbon and nitrogen stable isotopes of particulate organic matter in the Southern Ocean. *Marine Ecology Progress Series*, 631, 19–30. <https://doi.org/10.3354/meps13137>
- Fleming, A. H., Clark, C. T., Calambokidis, J., & Barlow, J. (2016). Humpback whale diets respond to variance in ocean climate and ecosystem conditions in the California Current. *Global Change Biology*, 22(3), 1214–1224. <https://doi.org/10.1111/gcb.13171>
- Forcada, J., & Trathan, P. N. (2009). Penguin responses to climate change in the Southern Ocean. *Global Change Biology*, 15(7), 1618–1630. <https://doi.org/10.1111/j.1365-2486.2009.01909.x>
- Forcada, J., Trathan, P. N., Reid, K., & Murphy, E. J. (2005). The effects of global climate variability in pup production of Antarctic fur seals. *Ecology*, 86(9), 2408–2417. <https://doi.org/10.1890/04-1153>
- Fraser, W., & Hofmann, E. (2003). A predator's perspective on causal links between climate change, physical forcing and ecosystem response. *Marine Ecology Progress Series*, 265, 1–15. <https://doi.org/10.3354/meps265001>
- Gelman, A., & Rubin, D. B. (1992). Inference from iterative simulation using multiple sequences. *Statistical Science*, 7(4), 457–472. <https://doi.org/10.1214/ss/1177011136>
- Graham, B. S., Koch, P. L., Newsome, S. D., McMahon, K. W., & Aurioles, D. (2010). Using isoscapes to trace the movements and foraging behavior of top predators in oceanic ecosystems. In J. B. West, G. J. Bowen, T. E. Dawson, & K. P. Tu (Eds.), *Isoscapes* (pp. 299–318). Springer. https://doi.org/10.1007/978-90-481-3354-3_14
- Greene, C. H., Pershing, A. J., Kenney, R. D., & Jossi, J. W. (2003). Impact of climate variability on the recovery of endangered north Atlantic right whales. *Oceanography*, 16(4), 98–103. <https://doi.org/10.5670/oceanog.2003.16>
- Hanson, N. N., Wurster, C. M., Bird, M. I., Reid, K., & Boyd, I. L. (2009). Intrinsic and extrinsic forcing in life histories: Patterns of growth and stable isotopes in male Antarctic fur seal teeth. *Marine Ecology Progress Series*, 388, 263–272. <https://doi.org/10.3354/meps08158>
- Haug, T., Falk-Petersen, S., Greenacre, M., Hop, H., Lindstrøm, U., Meier, S., Nilssen, K. T., & Wold, A. (2017). Trophic level and fatty acids in harp seals compared with common minke whales in the

- Barents Sea. *Marine Biology Research*, 13(9), 919–932. <https://doi.org/10.1080/17451000.2017.1313988>
- Herman, D. P., Burrows, D. G., Wade, P. R., Durban, J. W., Matkin, C. O., Leduc, R. G., Barrett-Lennard, L. G., & Krahn, M. M. (2005). Feeding ecology of eastern North Pacific killer whales *Orcinus orca* from fatty acid, stable isotope, and organochlorine analyses of blubber biopsies. *Marine Ecology Progress Series*, 302, 275–291. <https://doi.org/10.3354/meps302275>
- Hlista, B. L., Sosik, H. M., Traykovski, L. V. M., Kenney, R. D., & Moore, M. J. (2009). Seasonal and interannual correlations between right-whale distribution and calving success and chlorophyll concentrations in the Gulf of Maine, USA. *Marine Ecology Progress Series*, 394, 289–302. <https://doi.org/10.3354/meps08267>
- Hobson, K. A. (1999). Tracing origins and migration of wildlife using stable isotopes: A review. *Oecologia*, 120(3), 314–326. <https://doi.org/10.1007/s004420050865>
- Hobson, K. A., Barnett-Johnson, R., & Cerling, T. (2010). Using isoscapes to track animal migration. In *Isoscapes* (Issue November, pp. 273–298). Springer. https://doi.org/10.1007/978-90-481-3354-3_13
- Hobson, K. A., Gibbs, H. L., & Gloutney, M. L. (1997). Preservation of blood and tissue samples for stable-carbon and stable-nitrogen isotope analysis. *Canadian Journal of Zoology*, 75(10), 1720–1723. <https://doi.org/10.1139/z97-799>
- Hooker, S. K., Iverson, S. J., Ostrom, P., & Smith, S. C. (2001). Diet of northern bottlenose whales inferred from fatty-acid and stable-isotope analyses of biopsy samples. *Canadian Journal of Zoology*, 79(8), 1442–1454. <https://doi.org/10.1139/cjz-79-8-1442>
- Horton, T. W., Zerbini, A. N., Andriolo, A., Danilewicz, D., & Sucunza, F. (2020). Multi-decadal humpback whale migratory route fidelity despite oceanographic and geomagnetic change. *Frontiers in Marine Science*, 7(June), 1–19. <https://doi.org/10.3389/fmars.2020.00414>
- International Whaling Commission. (2013). Report of the IWC workshop on the assessment of southern right whales. *Journal of Cetacean Research & Management*, 14(Supplement), 442–451.
- Iverson, S. J. (2009). Tracing aquatic food webs using fatty acids: From qualitative indicators to quantitative determination. In *Lipids in aquatic ecosystems* (pp. 281–308). Springer. <https://doi.org/10.1007/978-0-387-89366-2>
- Jackson, A. L., Inger, R., Parnell, A. C., & Bearhop, S. (2011). Comparing isotopic niche widths among and within communities: SIBER – Stable Isotope Bayesian Ellipses in R. *Journal of Animal Ecology*, 80, 595–602. <https://doi.org/10.1111/j.1365-2656.2011.01806.x>
- Jackson, J. A., Patenaude, N. J., Carroll, E. L., & Baker, C. S. (2008). How few whales were there after whaling? Inference from contemporary mtDNA diversity. *Molecular Ecology*, 17, 236–251. <https://doi.org/10.1111/j.1365-294X.2007.03497.x>
- Jansen, E., Christensen, J. H., Dokken, T., Nisancioglu, K. H., Vinther, B. M., Capron, E., Guo, C., Jensen, M. F., Langen, P. L., Pedersen, R. A., Yang, S., Bentsen, M., Kjær, H. A., Sadatzki, H., Sessford, E., & Stendel, M. (2020). Past perspectives on the present era of abrupt Arctic climate change. *Nature Climate Change*, 10(8), 714–721. <https://doi.org/10.1038/s41558-020-0860-7>
- Jönsson, K. I., & Jonsson, K. I. (1997). Capital and income breeding as alternative tactics of resource use in reproduction. *Oikos*, 78(1), 57. <https://doi.org/10.2307/3545800>
- Keith, S. A., & Bull, J. W. (2017). Animal culture impacts species' capacity to realise climate-driven range shifts. *Ecography*, 40(2), 296–304. <https://doi.org/10.1111/ecog.02481>
- Kokko, H., & Sutherland, W. J. (2001). Ecological traps in changing environments: Ecological and evolutionary consequences of a behaviourally mediated Allee effect. *Evolutionary Ecology Research*, 3, 537–551.
- Lambertsen, R. H. (1987). A biopsy system for large whales and its use for cytogenetics. *Journal of Mammalogy*, 68(2), 443–445. <https://doi.org/10.2307/1381495>
- Larsen, T., Ventura, M., Andersen, N., O'Brien, D. M., Piatkowski, U., & McCarthy, M. D. (2013). Tracing carbon sources through aquatic and terrestrial food webs using amino acid stable isotope fingerprinting. *PLoS One*, 8(9). <https://doi.org/10.1371/journal.pone.0073441>
- Leaper, R., Cooke, J., Trathan, P., Reid, K., Rowntree, V., & Payne, R. (2006). Global climate drives southern right whale (*Eubalaena australis*) population dynamics. *Biology Letters*, 2(2), 289–292. <https://doi.org/10.1098/rsbl.2005.0431>
- Lockyer, C. (1986). Body fat condition in northeast Atlantic fin whales, *Balaenoptera physalus*, and its relationship with reproduction and food resource. *Canadian Journal of Fisheries and Aquatic Sciences*, 43(1), 142–147. <https://doi.org/10.1139/f86-015>
- Lockyer, C. (2007). All creatures great and smaller: A study in cetacean life history energetics. *Journal of the Marine Biological Association of the United Kingdom*, 87(4), 1035–1045. <https://doi.org/10.1017/S0025315407054720>
- Logan, J. M., Pethybridge, H., Lorrain, A., Somes, C. J., Allain, V., Bodin, N., Choy, C. A., Duffy, L., Goñi, N., Graham, B., Langlais, C., Ménard, F., Olson, R., & Young, J. (2020). Global patterns and inferences of tuna movements and trophodynamics from stable isotope analysis. *Deep-Sea Research Part II: Topical Studies in Oceanography*, 175(May, 2019). <https://doi.org/10.1016/j.dsr2.2020.104775>
- Lorrain, A., Pethybridge, H., Cassar, N., Receveur, A., Allain, V., Bodin, N., Bopp, L., Choy, C. A., Duffy, L., Fry, B., Goñi, N., Graham, B. S., Hobday, A. J., Logan, J. M., Ménard, F., Menkes, C. E., Olson, R. J., Pagendam, D. E., Point, D., ... Young, J. W. (2020). Trends in tuna carbon isotopes suggest global changes in pelagic phytoplankton communities. *Global Change Biology*, 26(2), 458–470. <https://doi.org/10.1111/gcb.14858>
- Lübcker, N., Reisinger, R., Oosthuizen, W., de Bruyn, P., van Tonder, A., Pistorius, P., & Bester, M. (2017). Low trophic level diet of juvenile southern elephant seals *Mirounga leonina* from Marion Island: A stable isotope investigation using vibrissal regrowths. *Marine Ecology Progress Series*, 577(June), 237–250. <https://doi.org/10.3354/meps12240>
- Magozzi, S., Yool, A., Vander Zanden, H. B., Wunder, M. B., & Trueman, C. N. (2017). Using ocean models to predict spatial and temporal variation in marine carbon isotopes. *Ecosphere*, 8(5). <https://doi.org/10.1002/ecs2.1763>
- Marón, C., Budge, S., Ward, R., Valenzuela, L., Di Martino, M., Ricciardi, M., Sironi, M., Uhart, M., Seger, J., & Rowntree, V. (2020). Fatty acids and stable isotopes ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$) in southern right whale *Eubalaena australis* calves in relation to age and mortality at Peninsula Valdés, Argentina. *Marine Ecology Progress Series*, 646, 189–200. <https://doi.org/10.3354/meps13387>
- Mate, B. R., Best, P. B., Lagerquist, B. A., & Winsor, M. H. (2011). Coastal, offshore, and migratory movements of South African right whales revealed by satellite telemetry. *Marine Mammal Science*, 27(3), 455–476. <https://doi.org/10.1111/j.1748-7692.2010.00412.x>
- McMahon, K. W., Hamady, L. L., & Thorrold, S. R. (2013a). A review of ecogeochemistry approaches to estimating movements of marine animals. *Limnology and Oceanography*, 58(2), 697–714. <https://doi.org/10.4319/lo.2013.58.2.0697>
- McMahon, K. W., Hamady, L. L., & Thorrold, S. R. (2013b). Ocean ecogeochemistry: A review. *Oceanography and Marine Biology: An Annual Review*, 51. <https://doi.org/10.1201/b15406>
- Mendes, R. F., Ramos, J. A., Paiva, V. H., Calado, J. G., Matos, D. M., & Ceia, F. R. (2018). Foraging strategies of a generalist seabird species, the yellow-legged gull, from GPS tracking and stable isotope analyses. *Marine Biology*, 165(10). <https://doi.org/10.1007/s00227-018-3421-0>
- Mestre, J., Authier, M., Cherel, Y., Harcourt, R., McMahon, C. R., Hindell, M. A., Charrassin, J.-B., & Guinet, C. (2020). Decadal changes in blood $\delta^{13}\text{C}$ values, at-sea distribution, and weaning mass of southern elephant seals from Kerguelen Islands. *Proceedings of the Royal Society B: Biological Sciences*, 287(1933), 20201544. <https://doi.org/10.1098/rspb.2020.1544>

- Moore, J., & Semmens, B. X. (2008). Incorporating uncertainty and prior information into stable isotope mixing models. *Ecology Letters*, 11(5), 470–480. <https://doi.org/10.1111/j.1461-0248.2008.01163.x>
- Moore, S. E., Wynne, K. M., Kinney, J. C., & Grebmeier, J. M. (2007). Gray whale occurrence and forage southeast of Kodiak, Island. *Alaska. Marine Mammal Science*, 23(2), 419–428. <https://doi.org/10.1111/j.1748-7692.2007.00102.x>
- Murphy, E. J., Trathan, P. N., Watkins, J. L., Reid, K., Meredith, M. P., Forcada, J., Thorpe, S. E., Johnston, N. M., & Rothery, P. (2007). Climatically driven fluctuations in Southern Ocean ecosystems. *Proceedings of the Royal Society B: Biological Sciences*, 274(1629), 3057–3067. <https://doi.org/10.1098/rspb.2007.1180>
- Nelson, M. A., Quakenbush, L. T., Mahoney, B. A., Taras, B. D., & Wooller, M. J. (2018). Fifty years of Cook Inlet beluga whale feeding ecology from isotopes in bone and teeth. *Endangered Species Research*, 36, 77–87. <https://doi.org/10.3354/esr00890>
- Newsome, S. D., Chivers, S. J., & Berman Kowalewski, M. (2018). The influence of lipid-extraction and long-term DMSO preservation on carbon ($\delta^{13}\text{C}$) and nitrogen ($\delta^{15}\text{N}$) isotope values in cetacean skin. *Marine Mammal Science*, 34(2), 277–293. <https://doi.org/10.1111/mms.12454>
- Newsome, S. D., Clementz, M. T., & Koch, P. L. (2010). Using stable isotope biogeochemistry to study marine mammal ecology. *Marine Mammal Science*, 26(3), 509–572. <https://doi.org/10.1111/j.1748-7692.2009.00354.x>
- Newsome, S. D., Etnier, M., Kurlle, C., Waldbauer, J., Chamberlain, C., & Koch, P. (2007). Historic decline in primary productivity in western Gulf of Alaska and eastern Bering Sea: isotopic analysis of northern fur seal teeth. *Marine Ecology Progress Series*, 332, 211–224. <https://doi.org/10.3354/meps332211>
- Newsome, S. D., Martinez del Rio, C., Bearhop, S., & Phillips, D. L. (2007). A niche for stable isotope ecology. *Frontiers in Ecology and the Environment*, 5(8), 429–436. <https://doi.org/10.1890/060150.01>
- Olsen, E., & Grahl-Nielsen, O. (2003). Blubber fatty acids of minke whales: Stratification, population identification and relation to diet. *Marine Biology*, 142(1), 13–24. <https://doi.org/10.1007/s00227-002-0934-2>
- Orr, J. C., Fabry, V. J., Aumont, O., Bopp, L., Doney, S. C., Feely, R. A., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., Key, R. M., Lindsay, K., Maier-Reimer, E., Matear, R., Monfray, P., Mouchet, A., Najjar, R. G., Plattner, G.-K., Rodgers, K. B., ... Yool, A. (2005). Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature*. <https://doi.org/10.1038/nature04095>
- Orsi, A. H., & Harris, U. (2019). *Fronts of the Antarctic circumpolar current – GIS data, ver. 1*. Australian Antarctic Data Centre. https://data.aad.gov.au/metadata/records/antarctic_circumpolar_current_fronts
- Parnell, A. C., Inger, R., Bearhop, S., & Jackson, A. L. (2010). Source partitioning using stable isotopes: Coping with too much variation. *PLoS One*, 5(3), e9672. <https://doi.org/10.1371/journal.pone.0009672>
- Patenaude, N. J., & Baker, C. S. (2001). Population status and habitat use of southern right whales in the sub-Antarctic Auckland Islands of New Zealand. *Journal of Cetacean Research & Management Special Issue*, 2(July, 1993), 111–116.
- Payne, R. (1986). Long term behavioral studies of the southern right whale (*Eubalaena australis*). *Reports – International Whaling Commission, Special Issue*, 10, 161–167.
- Payne, R., Brazier, O., Dorsey, E. M., Perkins, J. S., Rowntree, V. J., & Titus, A. (1983). External features in southern right whales, *Eubalaena australis*, and their use in identifying individuals. In R. Payne (Ed.), *Communication and behavior of whales* (1st ed., pp. 295–328). Westview Press.
- Peterson, B. J., & Fry, B. (1987). Stable isotopes in ecosystem studies. *Annual Review of Ecology and Systematics*, 18, 293–320. <https://doi.org/10.1146/annurev.es.18.110187.001453>
- Phillips, D. L., Inger, R., Bearhop, S., Jackson, A. L., Moore, J. W., Parnell, A. C., Semmens, B. X., & Ward, E. J. (2014). Best practices for use of stable isotope mixing models in food-web studies. *Canadian Journal of Zoology*, 92(10), 823–835. <https://doi.org/10.1139/cjz-2014-0127>
- Plummer, M. (2018). *rjags: Bayesian graphical models using MCMC. R Package Version, 4-8*. <https://CRAN.R-Project.Org/Package=rjags>
- Pomerleau, C., Heide-Jørgensen, M. P., Ferguson, S. H., Stern, H. L., Høyer, J. L., & Stern, G. A. (2017). Reconstructing variability in West Greenland ocean biogeochemistry and bowhead whale (*Balaena mysticetus*) food web structure using amino acid isotope ratios. *Polar Biology*, 40(11), 2225–2238. <https://doi.org/10.1007/s00300-017-2136-x>
- Quillfeldt, P., Masello, J. F., McGill, R. A. R., Adams, M., & Furness, R. W. (2010). Moving polewards in winter: A recent change in the migratory strategy of a pelagic seabird? *Frontiers in Zoology*, 7, 1–11. <https://doi.org/10.1186/1742-9994-7-15>
- R Core Team. (2020). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. <https://www.r-project.org/>
- Reeves, R., Rolland, R., & Clapham, P. (2001). Causes of reproductive failure in North Atlantic right whales: New avenues of research. *Northeast Fisheries Science Center Report of a Workshop Held 26-28 April 2000, Falmouth, Massachusetts*, 2–17. <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.318.8720&rep=rep1&type=pdf>
- Reid, K., & Croxall, J. P. (2001). Environmental response of upper trophic-level predators reveals a system change in an Antarctic marine ecosystem. *Proceedings of the Royal Society B: Biological Sciences*, 268(1465), 377–384. <https://doi.org/10.1098/rspb.2000.1371>
- Rendell, L., & Whitehead, H. (2001). Culture in whales and dolphins. *Behavioral and Brain Sciences*, 24(2), 309–382. <https://doi.org/10.1017/S0140525X0100396X>
- Rogers, A. D., Frinault, B. A. V., Barnes, D. K. A., Bindoff, N. L., Downie, R., Ducklow, H. W., Friedlaender, A. S., Hart, T., Hill, S. L., Hofmann, E. E., Linse, K., McMahon, C. R., Murphy, E. J., Pakhomov, E. A., Reygondeau, G., Staniland, I. J., Wolf-Gladrow, D. A., & Wright, R. M. (2020). Antarctic futures: An assessment of climate-driven changes in ecosystem structure, function, and service provisioning in the southern ocean. *Annual Review of Marine Science*, 12, 87–120. <https://doi.org/10.1146/annurev-marine-010419-011028>
- Rowntree, V. J., Uhart, M. M., Sironi, M., Chirife, A., Di Martino, M., La Sala, L., Musmeci, L., Mohamed, N., Andrejuk, J., McAlouse, D., Sala, J. E., Carribero, A., Rally, H., Franco, M., Adler, F. R., Brownell, R. L., Seger, J., & Rowles, T. (2013). Unexplained recurring high mortality of southern right whale *Eubalaena australis* calves at Península Valdés, Argentina. *Marine Ecology Progress Series*, 493, 275–289. <https://doi.org/10.3354/meps10506>
- Ruiz-Cooley, R. I., Gendron, D., Aguiñiga, S., Mesnick, S., & Carriquiry, J. D. (2004). Trophic relationships between sperm whales and jumbo squid using stable isotopes of C and N. *Marine Ecology Progress Series*, 277, 275–283. <https://doi.org/10.3354/meps277275>
- Ryan, C., McHugh, B., Trueman, C. N., Sabin, R., Deaville, R., Harrod, C., Berrow, S. D., & O'Connor, I. (2013). Stable isotope analysis of baleen reveals resource partitioning among sympatric rorquals and population structure in fin whales. *Marine Ecology Progress Series*, 479, 251–261. <https://doi.org/10.3354/meps10231>
- Schlaepfer, M. A., Runge, M. C., & Sherman, P. W. (2002). Ecological and evolutionary traps. *Trends in Ecology & Evolution*, 17(10), 474–480. [https://doi.org/10.1016/S0169-5347\(02\)02580-6](https://doi.org/10.1016/S0169-5347(02)02580-6)
- Seyboth, E., Groch, K. R., Dalla Rosa, L., Reid, K., Flores, P. A. C., & Secchi, E. R. (2016). Southern right whale (*Eubalaena australis*) reproductive success is influenced by krill (*Euphausia superba*) density and climate. *Scientific Reports*, 6. <https://doi.org/10.1038/srep28205>
- Silva, M. A., Borrell, A., Prieto, R., Gauffier, P., Bérubé, M., Palsbøl, P. J., & Colaço, A. (2019). Stable isotopes reveal winter feeding in different habitats in blue, fin and sei whales migrating through the Azores. *Royal Society Open Science*, 6(8). <https://doi.org/10.1098/rsos.181800>

- Smith, D. T., & Jacka, J. (2003). *Extents of Antarctic sea ice – GIS data – 1973–1999, Ver. 1*. Australian Antarctic Data Centre. https://data.aad.gov.au/metadata/records/sea_ice_extent_gis
- Smith, T. D., Reeves, R. R., Josephson, E. A., & Lund, J. N. (2012). Spatial and seasonal distribution of American whaling and whales in the age of sail. *PLoS One*, 7(4). <https://doi.org/10.1371/journal.pone.0034905>
- Sokolov, S., & Rintoul, S. R. (2009). Circumpolar structure and distribution of the antarctic circumpolar current fronts: 2. Variability and relationship to sea surface height. *Journal of Geophysical Research: Oceans*, 114(11), 1–15. <https://doi.org/10.1029/2008JC005248>
- Sonnerup, R. E., Quay, P. D., McNichol, A. P., Bullister, J. L., Westby, T. A., & Anderson, H. L. (1999). Reconstructing the oceanic ^{13}C Suess Effect. *Global Biogeochemical Cycles*, 13(4), 857–872. <https://doi.org/10.1029/1999GB900027>
- Stock, B. C., Jackson, A. L., Ward, E. J., Parnell, A. C., Phillips, D. L., & Semmens, B. X. (2018). Analyzing mixing systems using a new generation of Bayesian tracer mixing models. *PeerJ*, 2018(6), 1–27. <https://doi.org/10.7717/peerj.5096>
- Stock, B. C., & Semmens, B. X. (2016a). MixSIAR GUI user manual. Version 3.1, March, 1–42. <https://doi.org/10.5281/zenodo.47719>
- Stock, B. C., & Semmens, B. X. (2016b). Unifying error structures in commonly used biotracer mixing models. *Ecology*, 97(10), 2562–2569. <https://doi.org/10.1002/ecs.1517>
- Suess, H. E. (1955). Radiocarbon concentration in modern wood. *Science*, 122(3166), 415–417. <https://doi.org/10.1126/science.122.3166.415-a>
- Todd, S. K. (1997). Dietary patterns of humpback whales (*Megaptera novaeangliae*) in the Northwest Atlantic: Evidence from ^{13}C and ^{15}N stable isotopes. *Doctoral (PhD) Thesis, Memorial University of Newfoundland*.
- Todd, S., Ostrom, P., Lien, J., & Abrajano, J. (1997). Use of biopsy samples of humpback whale (*Megaptera novaeangliae*) skin for stable isotope ($\delta^{13}\text{C}$) determination. *Journal of Northwest Atlantic Fishery Science*, 22(July), 71–76. <https://doi.org/10.2960/J.v22.a6>
- Tormosov, D. D., Mikhalev, Y. A., Best, P. B., Zemsky, V. A., Sekiguchi, K., & Brownell, R. L. (1998). Soviet catches of southern right whales *Eubalaena australis*, 1951–1971. Biological data and conservation implications. *Biological Conservation*, 86(2), 185–197. [https://doi.org/10.1016/S0006-3207\(98\)00008-1](https://doi.org/10.1016/S0006-3207(98)00008-1)
- Townsend, C. H. (1935). The distribution of certain whales as shown by logbook records of American whaleships. *Zoologica*, 19, 3–50.
- Trathan, P. N., Forcada, J., & Murphy, E. J. (2007). Environmental forcing and Southern Ocean marine predator populations: Effects of climate change and variability. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 362(1488), 2351–2365. <https://doi.org/10.1098/rstb.2006.1953>
- Tulloch, V. J. D., Plagányi, É. E., Brown, C., Richardson, A. J., & Matear, R. (2019). Future recovery of baleen whales is imperiled by climate change. *Global Change Biology*, 25(4), 1263–1281. <https://doi.org/10.1111/gcb.14573>
- Valenzuela, L. O., Rowntree, V., Sironi, M., & Seger, J. (2018). Stable isotopes ($\delta^{15}\text{N}$, $\delta^{13}\text{C}$, $\delta^{34}\text{S}$) in skin reveal diverse food sources used by southern right whales *Eubalaena australis*. *Marine Ecology Progress Series*, 603, 243–255. <https://doi.org/10.3354/meps12722>
- Valenzuela, L. O., Sironi, M., Rowntree, V. J., & Seger, J. (2009). Isotopic and genetic evidence for culturally inherited site fidelity to feeding grounds in southern right whales (*Eubalaena australis*). *Molecular Ecology*, 18(5), 782–791. <https://doi.org/10.1111/j.1365-294X.2008.04069.x>
- Vehtari, A., Gelman, A., & Gabry, J. (2017). Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. *Statistics and Computing*, 27(5), 1413–1432. <https://doi.org/10.1007/s11222-016-9696-4>
- Vermeulen, E., Wilkinson, C., Thornton, M., Peters, I. T., & Findlay, K. (2018). Report on the mammal research institute whale unit southern right whale survey - 2017. *IWC Scientific Committee Report, SC/67B/SH/*. <https://doi.org/10.13140/RG.2.2.33382.45125>
- Ward, E. J., Holmes, E. E., & Balcomb, K. C. (2009). Quantifying the effects of prey abundance on killer whale reproduction. *Journal of Applied Ecology*, 46(3), 632–640. <https://doi.org/10.1111/j.1365-2664.2009.01647.x>
- Whitehead, H., Rendell, L., Osborne, R. W., & Würsig, B. (2004). Culture and conservation of non-humans with reference to whales and dolphins: Review and new directions. *Biological Conservation*, 120(3), 427–437. <https://doi.org/10.1016/j.biocon.2004.03.017>
- Williams, R., Vikingsson, G. A., Gislason, A., Lockyer, C., New, L., Thomas, L., & Hammond, P. S. (2013). Evidence for density-dependent changes in body condition and pregnancy rate of North Atlantic fin whales over four decades of varying environmental conditions. *ICES Journal of Marine Science*, 70(6), 1273–1280. <https://doi.org/10.1093/icesjms/fst059>
- Witteveen, B. H., Worthy, G. A. J., Foy, R. J., & Wynne, K. M. (2011). Modeling the diet of humpback whales: An approach using stable carbon and nitrogen isotopes in a Bayesian mixing model. *Marine Mammal Science*, 28(3), 1–18. <https://doi.org/10.1111/j.1748-7692.2011.00508.x>
- Zupcic-Moore, J. R., Ruiz-Cooley, R. I., Paliza, O., Koch, P. L., & McCarthy, M. D. (2017). Using stable isotopes to investigate foraging variation and habitat use of sperm whales from northern Peru. *Marine Ecology Progress Series*, 579, 201–212. <https://doi.org/10.3354/meps12281>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: van den Berg GL, Vermeulen E, Valenzuela LO, et al. Decadal shift in foraging strategy of a migratory southern ocean predator. *Glob Change Biol*. 2021;27:1052–1067. <https://doi.org/10.1111/gcb.15465>