University of Groningen

Don't throw the baby out with the bathwater

Riese, Harriette; Ormel, Johan; Aleman, Andre; Servaas, Michelle N.; Jeronimus, Bertus F.

Published in:
Neuroimage

DOI:
[10.1016/j.neuroimage.2015.11.012](https://doi.org/10.1016/j.neuroimage.2015.11.012)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2016

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Riese, H., Ormel, J., Aleman, A., Servaas, M. N., & Jeronimus, B. F. (2016). Don't throw the baby out with the bathwater: Depressive traits are part and parcel of neuroticism. *Neuroimage*, 15(125), 1103. <https://doi.org/10.1016/j.neuroimage.2015.11.012>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Comments and Controversies

Don't throw the baby out with the bathwater: Depressive traits are part and parcel of neuroticism

H. Riese^{a,*}, J. Ormel^a, A. Aleman^b, M.N. Servaas^{a,b}, B.F. Jeronimus^a

^a University of Groningen, University Medical Center Groningen, Department of Psychiatry, Interdisciplinary Center for Psychopathology and Emotion regulation, PO Box 30.001 (CC72), 9700 RB, Groningen, The Netherlands

^b University of Groningen, University Medical Center Groningen, Neuroimaging Center, Department of Neuroscience, PO Box 196, 9700 AD, Groningen, The Netherlands

In their comment (Bianchi and Laurent, 2015) on a recent publication in NeuroImage (Everaerd et al., 2015), Bianchi and Laurent argue that “depressive symptomatology should be systematically controlled for in neuroticism research” in order to be able to interpret the “true effect” of neuroticism. Unfortunately, their uncommon (see for example Table 2 in the meta-analysis of Servaas et al. (2013), where depression was specifically not mentioned as a covariate) advice lacks a theoretical or strong empirical basis. On the contrary, the high correlation of $\rho = 0.73$ between depressive symptoms (as measured with the Beck Depression Inventory) and neuroticism reveals that these cannot be easily disentangled, but rather suggests a significant degree of convergence, which may be based on shared (genetic) etiology (Genetics of Personality Consortium et al., 2015). We therefore argue that neuroticism, a personality trait known to be a robust predictor of vulnerability for physical and common mental health disorders, including depression (Hakulinen et al., 2015; Lahey, 2009), deserves a more nuanced approach than being reduced to variance left after systematic adjustment for depressive symptoms.

We are concerned that Bianchi and Laurent's advice could be potentially misleading to researchers, since controlling for third factors should be grounded in a specific research question. At least a distinction should be made between trait depressive symptoms (i.e. the depressive symptoms one usually experiences) and state depressive symptoms, because a temporary episode of (subthreshold) depression can inflate neuroticism reports (Ormel et al., 2013b). Depending on the research question, it can be defended to adjust the neuroticism personality trait for temporary increases in depressive symptoms (state depression), although more elegant alternatives would be to either postpone personality assessment until the episode has remitted or use other-reports (e.g. parent, partner) to assess an individual's trait scores.

Moreover, there is substantial consensus about the multifaceted nature of neuroticism. Depression is one of these facets, next to anxiety, hostility, and self-consciousness, among others (John et al., 2008; Ormel et al., 2013b). A systematic control for symptoms of depression seems therefore quite arbitrary, as one could also control for symptoms of e.g. anxiety ($r = .65$ with neuroticism, and $r = .77$ with depression;

Jeronimus et al., 2013), or both. In our opinion, future research could benefit from systematically acknowledging the heterogeneity of neuroticism. Revealing facet-specific neurobiological underpinnings of neuroticism may help to better understand the association between neuroticism, stress-reactivity, and mental disorders (Haas et al., 2007; Ormel et al., 2013a). In sum, we feel that a systematic control for a person's usual depressive symptoms results in a mutilated neuroticism construct and may yield uninterpretable results.

References

- Bianchi, R., Laurent, E., 2015. Depressive symptomatology should be systematically controlled for in neuroticism research. NeuroImage. <http://dx.doi.org/10.1016/j.neuroimage.2015.07.088> (Epub ahead of print, pii: S1053-8119(15)00732-6).
- Everaerd, D., Klumpers, F., van Wingen, G., Tendolcar, I., Fernandez, G., 2015. Association between neuroticism and amygdala responsivity emerges under stressful conditions. NeuroImage 112, 218–224.
- Genetics of Personality Consortium, de Moor, M.H., van den Berg, S.M., Verweij, K.J., Krueger, R.F., Luciano, M., et al., 2015. Meta-analysis of genome-wide association studies for neuroticism, and the polygenic association with major depressive disorder. JAMA Psychiatry 72 (7), 642–650.
- Haas, B.W., Omura, K., Constable, R.T., Canli, T., 2007. Emotional conflict and neuroticism: personality-dependent activation in the amygdala and subgenual anterior cingulate. Behav. Neurosci. 121 (0735-7044; 2), 249–256.
- Hakulinen, C., Elovainio, M., Pulkki-Raback, L., Virtanen, M., Kivimäki, M., Jokela, M., 2015. Personality and depressive symptoms: individual participant meta-analysis of 10 cohort studies. Depress. Anxiety 32 (7), 461–470.
- Jeronimus, B.F., Ormel, J., Aleman, A., Penninx, B.W., Riese, H., 2013. Negative and positive life events are associated with small but lasting change in neuroticism. Psychol. Med. 43 (11), 2403–2415.
- John, O.P., Robins, R., Pervin, L.A., 2008. Handbook of Personality: Theory and Research. third ed. Guilford, New York.
- Lahey, B.B., 2009. Public health significance of neuroticism. Am. Psychol. 64, 241–256.
- Ormel, J., Bastiaansen, A., Riese, H., Bos, E.H., Servaas, M., Ellenbogen, M., et al., 2013a. The biological and psychological basis of neuroticism: current status and future directions. Neurosci. Biobehav. Rev. 37 (1), 59–72.
- Ormel, J., Jeronimus, B.F., Kotov, R., Riese, H., Bos, E.H., Hankin, B., et al., 2013b. Neuroticism and common mental disorders: meaning and utility of a complex relationship. Clin. Psychol. Rev. 33 (5), 686–697.
- Servaas, M.N., van der Velde, J., Costafreda, S.G., Horton, P., Ormel, J., Riese, H., et al., 2013. Neuroticism and the brain: a quantitative meta-analysis of neuroimaging studies investigating emotion processing. Neurosci. Biobehav. Rev. 37 (8), 1518–1529.

* Corresponding author.