University of Groningen

Ischemia-Induced Neural Stem/Progenitor Cells in the Pia Mater Following Cortical Infarction

Nakagomi, Takayuki; Molnar, Zoltan; Nakano-Doi, Akiko; Taguchi, Akihiko; Saino, Orië; Kubo, Shuji; Clausen, Martijn; Yoshikawa, Hiroo; Nakagomi, Nami; Matsuyama, Tomohiro

Published in:
Stem cells and development

DOI:
[10.1089/scd.2011.0279](https://doi.org/10.1089/scd.2011.0279)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2011

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Nakagomi, T., Molnar, Z., Nakano-Doi, A., Taguchi, A., Saino, O., Kubo, S., Clausen, M., Yoshikawa, H., Nakagomi, N., & Matsuyama, T. (2011). Ischemia-Induced Neural Stem/Progenitor Cells in the Pia Mater Following Cortical Infarction. *Stem cells and development*, 20(12), 2037-2051.
<https://doi.org/10.1089/scd.2011.0279>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Ischemia-Induced Neural Stem/Progenitor Cells in the Pia Mater Following Cortical Infarction

Takayuki Nakagomi,^{1,2} Zoltán Molnár,² Akiko Nakano-Doi,¹ Akihiko Taguchi,³ Orié Saino,¹ Shuji Kubo,¹ Martijn Clausen,^{1,4} Hiroo Yoshikawa,⁵ Nami Nakagomi,^{1,6,7} and Tomohiro Matsuyama¹

Increasing evidence shows that neural stem/progenitor cells (NSPCs) can be activated in the nonconventional neurogenic zones such as the cortex following ischemic stroke. However, the precise origin, identity, and subtypes of the ischemia-induced NSPCs (iNSPCs), which can contribute to cortical neurogenesis, is currently still unclear. In our present study, using an adult mouse cortical infarction model, we found that the leptomeninges (pia mater), which is widely distributed within and closely associated with blood vessels as microvascular pericytes/perivascular cells throughout central nervous system (CNS), have NSPC activity in response to ischemia and can generate neurons. These observations indicate that microvascular pericytes residing near blood vessels that are distributed from the leptomeninges to the cortex are potential sources of iNSPCs for neurogenesis following cortical infarction. In addition, our results propose a novel concept that the leptomeninges, which cover the entire brain, have an important role in CNS restoration following brain injury such as stroke.

Introduction

STROKE IS ONE OF THE MAJOR causes of death and disability in developed countries. The central nervous system (CNS) is known for its limited regenerative capacity. However, several studies have demonstrated that the CNS shows some regenerative potential since cerebral ischemia is followed by activation of endogenous neurogenesis. Although adult cerebral cortical neurogenesis in adults remains controversial, accumulating evidence indicates that new neurons are generated in the cerebral cortex of adult mammals under pathological conditions [1–4]. This suggests that neural stem/progenitor cells (NSPCs) are activated in the cortex during brain injuries such as hypoxic/ischemic stroke. In support of this idea, we have previously shown that NSPCs develop in the poststroke cortex of adult mice and refer to these as ischemia-induced NSPCs (iNSPCs) [5–8]. iNSPCs form nestin-positive neurospheres in vitro that can self-renew and differentiate into electrophysiologically functional neurons, astrocytes, and myelin-producing oligodendrocytes [5]. In addition, we have shown that the nestin-positive iNSPCs develop in the perivascular regions of the poststroke cortex near endothelial cells (ECs) after stroke [6,9]. These observations suggest that iNSPCs may originate, at least in part,

from perivascular cells/microvascular pericytes widely distributed in CNS. However, the origin and identity of the iNSPCs that contribute to cortical neurogenesis [5–8] remain unclear.

Increasing evidence indicates that NSPCs [10,11] and neuronal progenitors [12] are induced at the subpial/cortical layer 1 regions in response to cortical ischemia/injury, suggesting that NSPCs can be activated in the cortical surface adjacent to an injured region. Leptomeninges (including pia mater and arachnoid mater), which cover the entire brain cortex, regulate NSPC development [13] and cortical neurogenesis [14]. In addition, a recent study has shown that stem/progenitor cells with neuronal differentiation potential reside in the leptomeninges during development [15]. However, no study has examined pia mater as a source of iNSPCs following cortical infarction.

In the present study we investigated whether the leptomeninges (pia mater), which is widely distributed within and closely associated with blood vessels throughout CNS including the cortex, have NSPC activity in response to ischemia and can generate neurons. Further, we investigated whether microvascular pericytes that are distributed from the pia mater to the cortex are the origin of iNSPCs following cortical infarction.

¹Institute for Advanced Medical Sciences, Hyogo College of Medicine, Hyogo, Japan.

²Department of Physiology, Anatomy and Genetics, University of Oxford, Oxford, United Kingdom.

³Department of Cerebrovascular Disease, National Cardiovascular Center, Osaka, Japan.

⁴Department of Molecular Neurobiology, University of Groningen, Groningen, The Netherlands.

Departments of ⁵Internal Medicine and ⁶Pathology, Hyogo College of Medicine, Hyogo, Japan.

⁷Department of Tumour Pathology, Nuffield Department of Clinical Laboratory Sciences, University of Oxford, Oxford, United Kingdom.

Materials and Methods

Animal studies

All procedures were approved by the Animal Care Committee of Hyogo College of Medicine and were in accordance with the criteria outlined in the "Guide for the Care and Use of Laboratory Animals" prepared by the National Academy of Science of the USA. Quantitative analyses were performed by investigators who were blinded to the experimental protocol and identity of the samples under study.

Induction of focal cerebral ischemia

Cerebral ischemia was induced in 6-week-old male CB-17/1cr-+/+Jcl mice (Clea Japan Inc., Tokyo, Japan). Permanent focal cerebral ischemia was produced by ligation and disconnection of the distal portion of the left middle cerebral artery (MCA) [5–8,16]. In brief, the left MCA was isolated, electrocauterized, and disconnected just distal to its crossing of the olfactory tract (distal M1 portion) under halothane inhalation. The infarct area in mice of this background has been shown to be highly reproducible and limited to the ipsilateral cerebral cortex [5–8,16].

Histological analysis

Immunohistochemistry was performed as described previously [5,6,8]. In brief, mice were deeply anesthetized with sodium pentobarbital (50 mg/kg) and perfused transcardially with 4% paraformaldehyde. Next, coronal brain sections (20 μ m) were prepared and they were stained with antibodies to nestin (Chemicon, Temecula, CA), Ki67 (BD Pharmingen, San Diego, CA), 5-bromo-2'-deoxyuridine (BrdU; Abcam, Cambridge, United Kingdom), CD31 (Santa Cruz Biotechnology, Santa Cruz, CA), NG2 (Chemicon), platelet-derived growth factor receptor α (PDGFR α ; Santa Cruz Biotechnology), PDGFR β (Santa Cruz Biotechnology), glial fibrillary acidic protein (GFAP; DAKO, Glostrup, Denmark), A2B5 (Chemicon), Olig2 (Chemicon), Sox2 (Chemicon), Sox10 (R&D Systems, Minneapolis, MN), and glutamic acid decarboxylase 67 (GAD67; Sigma, St. Louis, MO). To quantify cell proliferation *in vivo*, 50 mg/kg of BrdU (Sigma) was administered subcutaneously daily for 3 days before sacrifice. Primary antibodies were observed using Alexa Fluor 488- or 555-conjugated secondary antibodies. Nuclei were stained with 4',6-diamino-2-phenylindole (DAPI; Kirkegaard & Perry Laboratories, Gaithersburg, MD). Images of sections were captured using a confocal laser microscope (LSM510; Carl Zeiss, Jena, Germany). The number of nestin-positive cells was counted at the poststroke pia mater and cortex, the nonischemic ipsilateral area, and regions within or around the subventricular zone (SVZ; at a distance of 1–100 and 101–200 μ m from the ventricular wall, respectively). Each area (10 sections/mouse, $n=5$) was measured using ImageJ software from NIH Image as described previously [5–7]. Results were expressed as the number of cells/mm². In another set of experiments, mice were anesthetized and placed in a stereotactic apparatus 2 days before induction of ischemia. A lentiviral vector encoding green fluorescent protein (GFP; 5.2×10^5 transducing unit (TU)/ μ L) [17] was injected into the pia mater as described previously [5].

Cell culture

To examine whether nestin-positive cells developed in the poststroke pia mater were NSPCs, we cultured pial cells under the conditions that promoted the formation of neurospheres as described previously [5,6,8]. In brief, mice were deeply anesthetized with sodium pentobarbital on poststroke day 3, and the pia mater was carefully removed under a microscope (Carton, Pathumthani, Thailand). Absence of cortical contamination was confirmed by histological analysis. The removed tissues were then mechanically dissociated by passage through 18-, 23-, and 27-gauge needles to prepare a single-cell suspension. The resulting cell suspensions were incubated in Dulbecco's modified Eagle's medium (DMEM) containing 20 ng/mL of basic fibroblast growth factor (bFGF; Peprotech, Rocky Hill, NJ), 20 ng/mL of epidermal growth factor (EGF; Peprotech), and 1% N2 supplement (Invitrogen, Carlsbad, CA). Total number of neurosphere-like cell clusters exceeding 80 μ m in diameter was counted every day for 30 days using a phase-contrast microscope (Olympus, Tokyo, Japan). On day 7 of incubation, cell clusters were fixed, cut on a cryostat (8- μ m thick sections), and stained with antibody to nestin, NG2 and GFAP as described previously [5,6,8]. In another set of experiments, whole brains were removed on poststroke day 3, and coronal brain slices (6 mm thickness) were incubated in DMEM containing bFGF and EGF. On day 7 of incubation, brain sections were fixed, cut on a cryostat into 20- μ m-thick sections, and subjected to immunohistochemistry.

Differentiation of neurospheres

To induce cell differentiation, neurosphere-like cell clusters were incubated on poly-L-lysine-coated glass coverslips for 6 days in neurobasal medium (Invitrogen) containing the B-27 supplement (Invitrogen) and all-*trans* retinoic acid (0.2 μ M; Sigma) [6]. Differentiated cells were stained with antibodies to microtubule-associated protein-2 (MAP2; Chemicon), neurofilament (Chemicon), Tuj-1 (Chemicon), GFAP (Chemicon), O4 (Chemicon), and myelin-associated glycoprotein (MAG; gift from Prof. Shinobu Inagaki, University of Osaka, Osaka, Japan).

Electron microscopy

Brains at poststroke day 3 were sequentially fixed in 1% glutaraldehyde, 1% OsO₄, ethanol, and propylene oxide, and embedded in Epon as described previously [18]. Sections were stained with uranyl acetate/lead citrate and examined under a HITACHI 7100 electron microscope (Tokyo, Japan) operated at 75 kV.

Reverse transcription-polymerase chain reaction

Total RNA was extracted from embryonic stem (ES) cells as well as pia mater- and SVZ-derived spheres. cDNA was amplified under the following conditions: 15 s at 94°C, 30 s at 56°C, and 1 min at 68°C (40 cycles). Primer sequences were as follows: GFAP forward, TCGGCCAGTTACCAGGAGG and GFAP reverse, ATGGTGATGCGGTTTCTTCG (amplicon size, 176 bp); aquaporin-4 (AQP4) forward, AACTGGGCAA ACCACTGGATA and AQP4 reverse, TCTCCCGAAGAGTC TTTCCCC (amplicon size, 278 bp); glyceraldehyde-3-phosphate dehydrogenase (GAPDH) forward, TGGTATCGTGAAG

GACTCATGAC and GAPDH reverse, ATGCCAGTGAGC TTCCCGTTCAGC (amplicon size, 180 bp); Snail forward, CTTGTGTCTGCACGACCTGT and reverse, CTTACATC CGAGTGGGTTT (amplicon size, 161 bp); Slug forward, AACATTTCAACGCCTCCAAG and Slug reverse, GCCGAC GATGTCCATACAGT (amplicon size, 162 bp); Twist forward, ACGCAGTCGCTGAACGAGGC and Twist reverse, GTA CAGGAAGTCGATGTACC (amplicon size, 119 bp); Sox9 forward, AAGAGGCCACGGAACAGACTCA and Sox9 reverse, GACCCTGAGATTGCCCAGAGTG (amplicon size, 132 bp); NG2 forward, ATGCTTCTCAGCCCGGGACA and NG2 reverse, GGTTGCGGCCATTGAGAATG (amplicon size, 424 bp); PDGFR β forward, ACAATTCCTGCGGAGTGACAG and PDGFR β reverse, AAAAGTACCAGTGAAACCTCGCTG (amplicon size, 114 bp); Olig2 forward, ACGCCAGCCTGGTG TCTAGT and Olig2 reverse, TCGGGCTCAGTCATCTGCTT (amplicon size, 304 bp); nestin forward, CACTAGAAAGCA GGAACCAG and nestin reverse, AGATGGTTCACAATCC TCTG (amplicon size, 307 bp); Sox2 forward, TTGGGAGGGG TGCAAAAAGA and Sox2 reverse, CCTGCGAAGCGCC TAACGTA (amplicon size, 312 bp); Oct4 forward, AGCTGCT GAAGCAGAAGAGG and Oct4 reverse, GGTTCTCATTGTT GTCGGCT (amplicon size, 138 bp); Klf4 forward, GTGCCC CGACTAACCGTTG and Klf4 reverse, GTCGTTGAACTCC TCGGTCT (amplicon size, 185 bp); c-myc forward, ATGCC CCTCAACGTGAACTTC and c-myc reverse, CGCAACATAG GATGGAGAGCA (amplicon size, 228 bp); and β -actin forward, GCTCGTCGTCGACAAGGGCTC and β -actin reverse, CAAACATGATCTGGGTCATCTTCTC (amplicon size, 353 bp).

Western blot analysis

Samples (10 μ g) were separated by 10% sodium dodecyl sulfate–polyacrylamide gel electrophoresis. Separated proteins were transferred electrophoretically onto nitrocellulose membranes. The membranes were incubated with antibodies to Sox2 (Chemicon), Oct4 (Chemicon), Klf4 (Santa Cruz Biotechnology), or c-myc-antibody (Santa Cruz Biotechnology) and then with peroxidase-labeled secondary antibodies. Antibody labeling of protein band was detected with enhanced chemiluminescence reagents (Chemi-Lumi One, Nacalai Tesque, Kyoto, Japan), according to the manufacturer's instructions.

Statistical analysis

Results are reported as the mean $\pm$ standard deviation. Statistical comparisons among groups were determined by 1-way analysis of variance. Individual comparisons were performed using Student's *t*-test when appropriate. *P* < 0.05 was considered to be statistically significant.

Results

Nestin-positive cells develop in the poststroke pia mater after ischemia

We reported previously that nestin-positive iNSPCs develop in the poststroke cortex in the brains of adult mice [5–8]. Thus, we examined the location of nestin-positive cells on poststroke day 3 by immunohistochemistry to determine their precise origin. Consistent with our previous study [5], nestin-positive cells were observed in the poststroke cortex (Fig. 1A, B), whereas they were rarely detectable in the

nonischemic cortex (Fig. 1A, C). Nestin-positive cells were expressed consistently in the conventional neurogenic zones such as SVZ (Fig. 1A, D). In addition to these areas, we found that nestin-positive cells developed in poststroke, but not in nonischemic pia mater (Fig. 1A–C). To evaluate cell proliferation in the poststroke pia mater, Ki67 was used as a marker of dividing cells [19]. On day 3 after stroke, extensive expression of Ki67 was observed in the poststroke pia mater/cortex, whereas few Ki67-positive cells were observed in the nonischemic ipsilateral pia mater/cortex (Fig. 2A–C). Further, cells that emerged in the poststroke pia mater/cortex could be birthdated with daily BrdU injections for 3 days, indicating that the nestin/BrdU double-positive cells in these regions were proliferative. In contrast, proliferative cells were rarely observed in the nonischemic ipsilateral pia mater/cortex (Fig. 2A, D–G).

We also investigated the time course of nestin expression after stroke (Fig. 1E–H). On poststroke day 1, some weakly immunoreactive cells were observed in the poststroke pia mater and cortex (Fig. 1F, Q). The number of nestin-positive cells in these areas increased on poststroke day 3 (Fig. 1G, Q). On poststroke day 7, these nestin-positive cells decreased in number (Fig. 1H, Q). These findings were consistent with our previous study [7]. Nestin-positive cells were rarely observed in the nonischemic ipsilateral pia mater or cortex for any of the ages studied (poststroke days 1–7; Fig. 1I–L, Q). Nestin-positive cells were observed in the ipsilateral SVZ (Fig. 1M–P) as early as poststroke day 1 (Fig. 1N, R), followed by an increase in number by day 3 (Fig. 1O, R). However, these cells were still restricted to the area close to SVZ (also see Fig. 1D). On poststroke day 7, some nestin-positive cells migrated out of SVZ, but they did not reach the poststroke cortex at that time (Fig. 1P, R). These findings are consistent with our previous study showing that GFP-labeled SVZ cells migrate out of the dorsal horn of the rostral migratory stream on poststroke day 7, but they do not yet reach the poststroke cortex at that time [5].

Pial nestin-positive cells spread into the cortical parenchyma as microvascular pericytes/perivascular cells

Until now, our data indicate that nestin-positive cells appear in the poststroke pia mater [5]. Pia mater is histologically continuous with the cortical parenchyma along the blood vessels [20,21]. Therefore, we further investigated the relationship between the pial nestin-positive cells and parenchymal blood vessels by double staining for nestin and CD31, a marker for ECs. We observed that nestin-positive cells in the poststroke pia mater developed in proximity to CD31-positive ECs and that some pial nestin-positive cells spread into the cortex through cortical layer I along the blood vessels (Fig. 3A–C). Whole-mount specimens [22] of poststroke pia mater also showed that nestin-positive cells were present near CD31-positive ECs (Fig. 3D–F). We further investigated whether nestin-positive cells developing in these regions coexpressed NG2, a marker of microvascular pericytes [23]. Double labeling for nestin and NG2 showed that some nestin-positive cells in the poststroke pia mater (~60%) as well as those in the poststroke cortex [5] coexpressed NG2 lying next to ECs (Fig. 3G–I). Further, a large proportion of nestin-positive pial cells (~95%) coexpressed

FIG. 1. Nestin-positive cells develop in the poststroke pia mater. Immunohistochemistry for nestin using the diaminobenzidine (DAB) reaction 3 days after stroke (**A–D**) showed that nestin-positive cells developed in the poststroke pia mater (**B**, arrows) and cortex (**A**, **B**) and were rarely observed in the nonischemic ipsilateral cortex (**A**, **C**). Nestin-positive cells were also observed in SVZ (**A**, **D**). Immunohistochemistry of the poststroke pia mater/cortex (**E–H**) showed that these cells developed on poststroke day 1 (**F**), 3 (**G**), and 7 (**H**) [nestin (**F–H**, green), DAPI (**F–H**, blue)]. In contrast, nestin-positive cells were rarely observed in the nonischemic ipsilateral cortex (**I–L**) on poststroke day 1 (**J**), 3 (**K**), and 7 (**L**) [nestin (**J–L**, green), DAPI (**J–L**, blue)]. In SVZ (**M–P**), nestin-positive cells were found on poststroke day 1 (**N**), 3 (**O**), and 7 (**P**) [nestin (**N–P**, green), DAPI (**N–P**, blue)], although they did not reach the poststroke cortex until day 7 after stroke. Semiquantitative analysis of nestin-positive cells for each area is shown (**Q**, **R**). Insets (**B**, **G**) show higher magnification of nestin-positive cells within the post-stroke pia mater. Panels (**B**), (**C**), and (**D**) show higher magnification of insets in panel (**A**) delineated by the black rectangle. Scale bars = 500 μ m (**A**) and 100 μ m (**B**, **F**, **J**, **N**). Results displayed are representative of 5 repetitions of the experimental protocol. DAPI, 4',6-diamino-2-phenylindole; SVZ, subventricular zone.

PDGFR β , another marker for microvascular pericytes [24] (Fig. 3J–L).

Pial nestin-positive cells have NSPCs activity to generate neurons

Previous studies have demonstrated that nestin/NG2-positive CNS microvascular pericytes [25] and PDGFR β -expressing pericytes [26] have NSPC activity and can generate neurons. In addition, we have recently shown that nestin-positive iNSPCs develop in the perivascular region of the postischemic cortex in proximity to ECs [6,9]. Based on these findings, we investigated whether pial nestin-positive cells developing near ECs have NSPC activity. To study this, we cultured pial cells as mentioned above. Neurosphere-like cell clusters appeared on day 4 of the incubation (Fig. 4A). Their number then gradually increased at later time points (day 4, 1.2 ± 0.4 ; day 10, 6.0 ± 1.2 ; day 20, 9.5 ± 2.5). Thirty days after incubation, the number of cell clusters reached ~ 14 (13.6 ± 3.8). In contrast, we did not obtain neurosphere-

like cell clusters from nonischemic pia mater. The pial iNSPCs also possessed self-renewal capacity after being incubated in the same medium at a clonal density of 5 cells/ μ L [5,6], but cluster formation was limited to between 3 and 5 cell passages. These observations were consistent with the traits of the iNSPCs obtained from the poststroke cortex [5] and suggest that pial iNSPCs are more likely to include neural progenitors than neural stem cells (NSCs) similar to neurospheres derived from other brain subregions such as the hippocampus [27] and subcortical white matter [28]. Immunohistochemistry revealed that virtually almost all cells forming spheres expressed nestin, and some coexpressed NG2 (Fig. 4B–D). The population of nestin- and NG2-positive cells was counted in an adherent monolayer culture [8] of pial cells. Similar to the neurospheres, $\sim 95\%$ of the cells expressed nestin, and $\sim 60\%$ of them expressed NG2 (Fig. 4E–G).

We used organotypic cortical slice cultures to obtain additional independent evidence that nestin-positive cells isolated from poststroke pia mater could expand in response to bFGF and EGF. Although nestin-positive cells were rarely

FIG. 2. Nestin-positive cells in the poststroke pia mater proliferate after ischemia. Brain sections on poststroke day 3 (A–F) showed that many Ki67-positive cells (B, arrows) were present within the poststroke pia mater (B). In contrast, Ki67-positive cells were rarely observed in the nonischemic ipsilateral pia mater (C) [Ki67 (B, C, red), DAPI (B, C, blue)]. Some nestin-positive cells developing in the poststroke pia mater expressed BrdU (F, arrows), whereas this expression was observed less often in the nonischemic ipsilateral pia mater [nestin (D, F, green), BrdU (E, F, red)]. The number of pial nestin/BrdU-positive cells increased significantly in the ischemic area compared to that in the nonischemic ipsilateral area (G). * $P < 0.05$ versus nonischemic pia mater ($n = 5$). Scale bar = 20 μm (B) and 200 μm (D). Results displayed are representative of 5 repetitions of the experimental protocol. BrdU, 5-bromo-2'-deoxyuridine.

observed in normal pia mater/cortex (data not shown) or the nonischemic ipsilateral pia mater (Fig. 4H, L; 3.2 ± 1.3 cells/ mm^2) and cortex (Fig. 4H), many nestin-positive cells were observed in the poststroke pia mater (Fig. 4I, L; 3100.8 ± 189.4 cells/ mm^2) and cortex (Fig. 4I). However, in the absence of bFGF and EGF, fewer nestin-positive cells were observed, even in the poststroke pia mater (Fig. 4J, L; 393.8 ± 55.5 cells/ mm^2) and cortex (Fig. 4J). After removing the poststroke pia mater, few nestin-positive cells were observed in the cortex (Fig. 4K). These observations strongly suggest that nestin-positive cells in the poststroke pia mater proliferate in response to bFGF and EGF and they are responsible for forming neurosphere-like cell clusters in vitro.

Double staining for nestin and CD31 was performed because nestin has also been shown to be expressed in cells with endothelial phenotype [29]. Pial nestin-positive cells proliferated in response to bFGF and EGF but did not co-express CD31 (Supplementary Fig. S1A–C; Supplementary Data are available online at www.liebertonline.com/scd). These observations indicate that the nestin-positive cells were not ECs, consistent with the traits of nestin-positive cells within the poststroke cortex [6].

Next, we investigated whether pial nestin-positive neurosphere-like cell clusters could differentiate into mature neural cells, including neurons, astrocytes, and oligodendrocytes. Cell clusters were plated on poly-L-lysine-coated glass coverslips for 7 days under conditions conducive to differentiation [6]. After differentiation, the cells expressed neuronal (Fig. 4M: MAP2, 4N: neurofilament, 4O: Tuj-1), astrocyte (Fig. 4O: GFAP), and oligodendrocyte (Fig. 4P: O4, 4Q: MAG) markers. Cell population analysis for differenti-

ated spheres showed that these cells predominantly differentiated into neuronal cells and oligodendrocytes rather than astrocytes (Tuj-1-positive neuronal cells $36.7\% \pm 6.6\%$; GFAP-positive astrocytes $4.8\% \pm 2.3\%$; and O4-positive oligodendrocytes $30.0\% \pm 6.1\%$). To confirm that these nestin-positive cells originated from pial cells, pia mater was labeled with a GFP-expressing vector. Then, the poststroke pia mater isolated 3 days after stroke was incubated in DMEM containing bFGF and EGF. GFP-positive neurospheres were obtained (Fig. 4R) and were found to have the same properties as shown above (Fig. 4A–D, M–Q).

To further explore the stem-cell nature of pial iNSPCs in vivo, we examined Sox2 expression, a marker of persistent NSPCs [30]. Sox2 expression was observed in nestin-positive cells located in SVZ and ependyma (data not shown), as described previously [31,32]. In contrast, Sox2-positive cells were rarely detectable in the nonischemic pia mater/cortex (Fig. 4S). However, Sox2-positive cells were observed in the poststroke pia mater and cortex, and some of them ($\sim 60\%$) coexpressed nestin (Fig. 4T–W). Although the pial cells revealed several NSPC characteristics following ischemia, recent studies have demonstrated that NSPCs derived from other regions (e.g., SVZ-derived NSCs [33] and cortex-derived NSPCs [5]) also possess pluripotent/undifferentiated cell markers. Thus, these markers, including Sox2, Oct4, Klf4, and c-myc, which are expressed in induced pluripotent stem (iPS) cells [34], were examined by western blot analysis. As a control, we used ES cells [5], which express these 4 factors as described previously [34], in our western blot analysis. Although none of these 4 factors were observed in nonischemic pia mater, Sox2, Klf4, and c-myc, but not Oct4, expression

FIG. 3. Nestin-positive cells observed in the perivascular area by microvascular pericyte marker expression. Coronal brain sections on poststroke day 3 showed that nestin-positive cells developed in close association with CD31-positive ECs from the poststroke pia mater to the cortex [nestin (A, C, green), CD31 (B, C, red), DAPI (A–C, blue)]. Note that some nestin-positive cells in the pia mater were connected to those within in the poststroke cortex through cortical layer I (A, arrowheads). Similarly, whole-mount samples obtained from the poststroke pia mater showed that nestin-positive cells developed around CD31-positive ECs [nestin (D, F, green); CD31 (E, F, red); and DAPI (D–F, blue)]. Immunohistochemistry on poststroke day 3 showed that nestin-positive cells in the poststroke pia mater/cortex coexpressed the microvascular pericyte markers such as NG2 (I, arrows) [nestin (G, I, green); NG2 (H, I, red); and DAPI (G–I, blue)] and PDGFR β (L, arrows) [nestin (J, L, green); PDGFR β (K, L, red); and DAPI (J–L, blue)]. Scale bars = 50 μ m (A, D) and 100 μ m (G, J). Results displayed are representative of 5 repetitions of the experimental protocol. ECs, endothelial cells; L1, cortical layer I; P, pia mater; PDGFR β , platelet-derived growth factor receptor beta.

was observed in pial iNSPCs (Fig. 4X), consistent with previous studies [5,33].

Pial nestin-positive iNSPCs originate, at least in part, from brain pericytes

Thus far, our data indicate that pial nestin-positive iNSPCs spread into the cortical parenchyma as perivascular cells and express pericyte markers. These results strongly suggest that these cells originate, at least in part, from microvascular pericytes. To confirm this, we isolated PDGFR β -positive cells from poststroke mice by magnetic cell sorting [26]. Immunohistochemistry showed that PDGFR β -positive cells were positive for nestin (Fig. 5A–C) and NG2 (Fig. 5D–F). These cells formed cell clusters in the medium that promoted neurosphere formation, as mentioned above (Fig. 5G). After dif-

ferentiation (Fig. 5H), they generated MAP2- and Tuj-1-positive neurons (Fig. 5I, J), GFAP-positive astrocytes (Fig. 5J), and O4-positive oligodendrocytes (Fig. 5K). Because brain pericytes are considered to have a neural crest origin, similar to the pia mater [35,36], we further investigated whether the pial iNSPCs possess neural crest markers in addition to those of pericytes. The pial nestin-positive iNSPCs largely (~95%) expressed Sox10, a neural crest marker (Fig. 5L–N). Polymerase chain reaction (PCR) analysis showed that iNSPCs, which expressed a NSPC marker (nestin) and pericytic markers (NG2 and PDGFR β), displayed various neural crest markers such as Snail, Slug, Twist, and Sox9 (Fig. 5O), although these markers were rarely observed in ES cells, as described previously [37]. These observations support our present finding that pial iNSPCs are derived from brain pericytes, which are neural crest derivatives [36].

FIG. 4. Pial nestin-positive cells show NSPC activity. Neurosphere-like cell clusters were obtained from the poststroke pia mater cultured in Dulbecco's modified Eagle's medium (DMEM) containing bFGF and EGF (A). Immunohistochemistry revealed that virtually almost all cells forming spheres expressed nestin, and some coexpressed NG2 [nestin (B, D, green); NG2 (C, D, red); and DAPI (B, C, blue)]. Similarly, pial cells incubated as adherent monolayer cultures largely expressed nestin and some coexpressed NG2 [nestin (E, G, green); NG2 (F, G, red); and DAPI (E–G, blue)]. To obtain additional independent evidence that nestin-positive cells isolated from poststroke pia mater can expand in response to bFGF and EGF, coronal brain samples, including the poststroke area, were incubated in the same medium that promoted sphere formation (DMEM containing bFGF and EGF; H–K). Although nestin-positive cells were rarely observed in the nonischemic ipsilateral pia mater/cortex [nestin (H, green) and DAPI (H, blue)], many nestin-positive cells were confirmed in the poststroke pia mater/cortex [nestin (I, green) and DAPI (I, blue)]. However, when brain samples were incubated without bFGF and EGF, fewer nestin-positive cells were observed in the poststroke pia mater/cortex [nestin (J, green) and DAPI (J, blue)]. Further, the expansion of pial nestin-positive cells was not detectable after removing the pia mater, even when brain samples were incubated in DMEM containing bFGF and EGF [nestin (K, green) and DAPI (K, blue)]. Semiquantitative analysis for the number of pial nestin-positive cells is shown (L). After differentiation (M–Q), the cells forming the neurospheres expressed neuronal [MAP2 (M, green); neurofilament (N, green); Tuj-1 (O, green); and DAPI (M–O, blue)], astrocyte [GFAP (O, red)], and oligodendrocyte markers [O4 (P, green); MAG (Q, green); and DAPI (P, Q, blue)] markers. GFP-positive neurospheres were obtained from cultures of GFP-labeled pia mater [GFP (R, green)]. Immunohistochemistry on poststroke day 3 for Sox2 showed a marker of persistent NSPCs (S–W). Although Sox2-positive cells were rarely detectable in the nonischemic pia mater/cortex [Sox2 (S, red), and DAPI (S, blue)], they were confirmed in poststroke pia mater and cortex, and some Sox2-positive cells in these regions coexpressed nestin [nestin (T, V, W, green); Sox2 (U, V, W, red); and DAPI (T–W, blue)]. Expression of several pluripotent/undifferentiated cell markers in pial iNSPCs (X). Western blot analysis showed that ES cells possessed Yamanaka's 4 factors (Sox2, Oct4, Klf4, and c-myc) needed for iPS cell induction. Although all 4 of these factors were not observed in cells isolated from the nonischemic pia mater, pial iNSPCs expressed 3 of them, viz., Sox2, Klf4, and c-myc. * $P < 0.05$ versus nonischemic pia mater with bFGF plus EGF, and # $P < 0.05$ versus ischemic pia mater with bFGF plus EGF group ($n = 5$) (L). Panel (W) shows higher magnification of the insets in panel (V), delineated by the white box. Scale bars = 100 μ m (A, B, E, H, M, R, S, T), and 50 μ m (W). The results displayed are representative of 5 repetitions of the experimental protocol. bFGF, basic fibroblast growth factor-basic; EGF, epidermal growth factor; ES cells, embryonic stem cells; GFAP, glial fibrillary acidic protein; GFP, green fluorescent protein; iNSPCs, ischemia-induced neural stem/progenitor cells; iPS cells, induced pluripotent stem cells; MAG, myelin-associated glycoprotein; MAP-2, microtubule-associated protein-2; P, pia mater.

FIG. 5. Pial nestin-positive iNSPCs are, at least in part, derived from brain pericytes. PDGFR β -positive cells were isolated by MACS. Immunohistochemistry showed that these cells coexpressed nestin [PDGFR β (A, C, red); nestin (B, C, green); and DAPI (A–C)] and NG2 [PDGFR β (D, F, red); NG2 (E, F, green); and DAPI (D–F)]. PDGFR β -positive cells formed neurosphere-like cell clusters (G). These spheres differentiated into (H) neuronal cells [MAP2 (I, green); Tuj-1 (J, green); and DAPI (I, J, blue)], astrocytes [GFAP (J, red), and DAPI (J, blue)], and oligodendrocytes [O4 (K, green), and DAPI (K, blue)]. Immunohistochemistry on poststroke day 3 (L–N) showed that the neural crest marker Sox10 was expressed in the nuclei of nestin-positive iNSPCs isolated from the poststroke pia mater [nestin (L, N, green); Sox10 (M, N, red); and DAPI (L–N)]. PCR analysis showed that the pial iNSPCs expressed various neural crest markers, including Snail, Slug, Twist, and Sox9, as well as markers for NSPCs (nestin) and pericytes (NG2 and PDGFR β), whereas expression of neural crest markers was rarely observed in ES cells (O). Scale bars = 10 μ m (A, D), 100 μ m (G–I), and 50 μ m (L). Results displayed are representative of 5 repetitions of the experimental protocol.

Comparison of pial iNSPCs to other types of CNS stem/progenitor cells

Radial glia cells are derived from neuroepithelial cells during development and give rise to several types of CNS stem/progenitor cells such as SVZ astrocytes [38,39], ependymal cells [40,41], and oligodendrocyte precursor cells (OPCs) [42,43] in adults [44]. Although pial iNSPCs expressed pericyte markers, such as NG2 and PDGFR β , the precise characteristics of NG2-expressing cells in poststroke pia mater remain unclear. Previous studies have reported that NG2 is also expressed in resident glia/glia precursors with NSPC activity [45,46], and in ependymal cells (a specialized glia), which acquire NSPC activity after CNS injury [41]. Anatomically, the pia mater spreads into the ventricular wall where it connects with ependymal cells to form the choroid plexus, which produces cerebrospinal fluid [14]. Ependymal cells were originally considered to be the resident stem cell population in the perivascular zone [47]. Although it is controversial whether ependymal cells have NSPC activity or not, recent studies confirm that ependymal cells do not play a role in adult neurogenesis under normal conditions, but they do possess NSPC activity and can differentiate into neurons, astrocytes, and oligodendrocytes in

response to CNS injuries including ischemic stroke [40,41]. Further, ependymal cells strongly express PDGFR α [48] and are penetrated by the apical process of NSCs in SVZ (type B cells), which express GFAP [38,39]. Based on these findings, we investigated whether pial iNSPCs are similar to periventricular NSPCs such as ependymal cells and/or SVZ NSCs (astrocytes). Consistent with previous studies, PDGFR α (Fig. 6A) and GFAP (Fig. 6B) expression was confirmed in the ependymal [48] and SVZ cells [38,39], respectively. PDGFR α -positive cells were present in the poststroke pia mater, and some nestin-positive cells in the poststroke pia mater co-expressed PDGFR α (Fig. 6C–E). As expected, nestin and GFAP double-positive reactive astrocyte-like cells were observed in the peri-stroke area (Fig. 7A, C, F) [49,50]. However, cells neither in the poststroke pia mater nor in the poststroke cortex expressed GFAP (Figs. 6F and 7A, B, D). In contrast, many GFAP-positive astrocytes were observed within the nonischemic ipsilateral pia mater/cortex (Fig. 7A, C, E). Similarly, most (90%) of the cultured pial nestin-positive cells were PDGFR α positive (Fig. 6G–I), but did not express GFAP (Fig. 6J–L), indicating that pial iNSPCs are not reactive astrocytes. Because the pial NSPCs did not express GFAP in vivo (Figs. 6F and 7A, B, D) and differentiated into astrocytes at a very low frequency (Figs. 4M–Q and 5I–K), it

FIG. 6. Comparison of pial iNSPCs to previously reported CNS stem/progenitor cells. The traits of pial iNSPCs were analyzed and compared with those of SVZ astrocytes, ependymal cells, and OPCs (A–Y). Immunohistochemistry on post-stroke day 3 showed that PDGFR α - and GFAP-positive cells were located in the ependyma [PDGFR α (A, red) and DAPI (A, blue)] and SVZ cells [GFAP (B, red), and DAPI (B, blue)], respectively. At the same time, PDGFR α -positive cells were also confirmed in the poststroke pia mater, and they partially coexpressed nestin [nestin (C, E, green); PDGFR α (D, E, red); and DAPI (C–E, blue)]. Cells in the poststroke pia mater and cortex did not express GFAP on poststroke day 3 [GFAP (F, red) and DAPI (B, blue)]. Nestin-positive cells isolated from the poststroke pia mater coexpressed PDGFR α [nestin (G, I, green); PDGFR α (H, I, red); and DAPI (G–I, blue)] but not GFAP [nestin (J, L, green); GFAP (K, L, red); and DAPI (J–L, blue)]. Similar to the *in vivo* findings, cell clusters from poststroke pia mater largely expressed nestin but not GFAP [nestin (M, O, green); GFAP (N, O, red); and DAPI (M–O, blue)]. In contrast, virtually almost all cells forming SVZ-derived neurospheres expressed GFAP rather than nestin [nestin (P, R, green); GFAP (Q, R, red); and DAPI (P–R, blue)]. Parallel to these findings, PCR analysis showed that expression of astrocytic markers such as GFAP and AQP4 was much lower or undetectable in the pia mater-derived spheres than in SVZ-derived neurospheres (S). Microvilli-like structures were confirmed by electron microscopy in the ependymal cells (T), but these structures were not observed in cells within the poststroke pia mater (U). The poststroke pia mater/cortex did not show the OPC markers A2B5 [A2B5 (V, red) and DAPI (V, blue)] or Olig2 [Olig2 (W, red) and DAPI (W, blue)]. Olig2 was induced in pial spheres under conditions promoting OPCs (iNSPCs-OPCs) (X) and almost all of these cells differentiated into O4- and MAG-positive oligodendrocytes [O4 (Y, green); MAG (Y, red); and DAPI (Y, blue)]. Scale bars = 100 μ m (A, C, E, G, J, M, P, V, W), 2 μ m (T, U), and 50 μ m (Y). Results displayed are representative of 5 repetitions of the experimental protocol. AQP4, aquaporin4; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; OPCs, oligodendrocyte precursor cells; CNS, central nervous system.

is also possible that they are different from SVZ NSCs [38,39]. Thus, we further compared several astrocyte markers in pial neurosphere-like cell clusters with those in SVZ-derived neurospheres. Similar to the *in vivo* findings, pial neurosphere-like cell clusters largely expressed nestin but not GFAP (Fig. 6M–O). In contrast, virtually almost all cells forming SVZ-derived neurospheres expressed GFAP rather than nestin (Fig. 6P–R). In accordance with these findings, reverse transcription (RT)-PCR analysis showed that GFAP expression in the pial spheres was much weaker than that in SVZ-derived neurospheres. In addition, AQP4, which is predominantly expressed in astrocyte foot processes [51], was not expressed in pia-derived spheres, although it was rich in SVZ-derived neurospheres (Fig. 6S; GFAP, 176 bp; AQP4, 278 bp; GAPDH, 180 bp). These observations are consistent with previous studies showing that ependyma with NSPC activity display weak [41] or no GFAP expression [47]. Taken together, these results suggest that pial iNSPCs may have similar characteristics to ependymal cells rather than to SVZ astrocytes. To confirm this, we further investigated whether the poststroke pial cells have the structures of lipid droplets, microvilli, and cilia, which ependyma cells possess [47,48]. These structures were confirmed in the ependyma by electron microscopy (Fig. 6T). However, microvilli-like structures were not observed in cells located in the poststroke pia mater (Fig. 6U). These findings indicate that pial iNSPCs do not have completely identical characteristics to those of ependymal cells.

Because OPCs also express NG2 [52] and PDGFR α [53], and A2B5-positive OPCs have NSPC activity [42,43], we further investigated whether pial nestin-positive cells might have OPC characteristics with multipotent NSPC activity. Immunohistochemistry on poststroke day 3 showed that A2B5-positive (Fig. 6V) and Olig2-positive (another OPC marker) cells [54] (Fig. 6W) were not present in poststroke pia mater or cortex. However, following incubation in OPC-

promoting medium [55], Olig2 expression was induced in pial iNSPCs (iNSPCs-OPCs) (Fig. 6X) and almost all these cells (~98%) induced from the iNSPCs differentiated into O4- and/or MAG-positive oligodendrocytes (Fig. 6Y). These observations indicate that pial nestin-positive iNSPCs are different from reported multipotent OPCs [42,43], although they partially co-express OPC markers.

Moreover, pial iNSPCs did not express GAD67 (data not shown), which is a marker of layer I progenitors [12]. Schematic representation for the fate of iNSPCs (Fig. 8) and their distinct expression of markers are summarized in Table 1.

Discussion

We demonstrated for the first time that nestin-positive iNSPCs, which are potential sources of neurons driving cortical neurogenesis, developed in the poststroke pia mater. In addition, we showed that they developed in the cortex as has been described previously [5,6]. Mignone and colleagues have shown that in developing transgenic nestin-GFP embryos GFP expression was evident as early as day 7 of embryonic development (e7). At e8, a GFP signal was seen predominantly in the neural plate, and by e10 intense GFP fluorescence was seen throughout the neuroepithelium. At e10 to e12, GFP signals marked the entire thickness of the cerebral wall, but GFP expression became weaker near the pial surface and stronger in the ventricular zones starting from e12. Finally, in the adult brain, GFP was selectively expressed in SVZ and the subgranular zone (SGZ) within the dentate gyrus of the hippocampus in areas related to continuous neurogenesis [56]. However, after brain injury/ischemia, nestin-positive cells and/or nestin-positive NSPCs were observed in nonconventional neurogenic zones such as the cortex [5,9] and striatum [57], although they were rarely observed in the absence of brain ischemia/injury [5]. These

FIG. 8. Schematic representation for the fate of iNSPCs following cortical infarction. Following cortical infarction, pial iNSPCs, which are presumably derived from the neural crest-brain pericyte lineage, develop in the post-stroke pia mater in close association with blood vessels throughout the brains, including the cortex. These cells reveal markers for NSPCs and pluripotent/undifferentiated cells in addition to those for pericytes and the neural crest. Pial iNSPCs do not have identical characteristics to those of other types of proposed CNS stem/progenitor cells, such as SVZ NSCs (astrocytes), ependymal cells, OPCs, reactive astrocytes, or layer I progenitors. NSCs, neural stem cells.

findings suggest that NSPC activation and neuronal homeostasis are maintained under physiological conditions, at least in part, in specific brain regions (such as SVZ and SGZ). However, after brain injury/ischemia, it would appear that regional NSPCs are mobilized to accelerate tissue repair in adult mammalian brains, in addition to activating neurogenesis in SVZ and SVG [58].

Previous studies have demonstrated that stem/progenitor cells in SVZ have the capacity to migrate toward injured lesions, including those in the cortex [59]. However, our previous study failed to demonstrate cell migration from SVZ to the cortex after cerebral infarction *in vivo* but showed that iNSPCs in the poststroke cortex originated, at least in part, from within the cerebral cortex [5]. Consistent with that study, subsequent studies have shown that NSPCs developing within and around the poststroke cortex are derived from locally activated stem/progenitor cells and not from SVZ cells [12,49], indicating that it is more

likely that iNSPCs from the poststroke pia mater/cortex rather than NSCs from SVZ are involved in cortical repair after stroke, particularly from the acute phase. Previous studies have indicated that NSPCs express an astrocyte marker, GFAP, after brain injury; thus, reactive astrocytes are proposed as a potential origin of injury-iNSPCs [49]. However, we could not find GFAP-positive cells within the poststroke area itself, and isolated pial iNSPCs showed few astrocytic traits compared with SVZ NSCs, suggesting that pial iNSPCs are a distinct population with a different origin. In the present study, we further demonstrated that pial iNSPCs after cortical infarction originate, at least in part, from the microvascular pericytes/perivascular cells. Pericytes with multipotent progenitor activity have been identified in various organs [60] in addition to CNS [25]. Although a previous study showed that pericyte-derived NSPCs can be isolated from CNS of noninjured animals [25], we could not obtain iNSPCs from nonischemic pia

TABLE 1. EXPRESSION MARKERS OF ISCHEMIA-INDUCED NEURAL STEM/PROGENITOR CELLS AND COMPARISON TO OTHER TYPES OF CENTRAL NERVOUS SYSTEM STEM/PROGENITOR CELLS

Expression markers	<i>Nestin</i>	<i>Sox2</i>	<i>NG2</i>	<i>PDGFRβ</i>	<i>GFAP</i>	<i>PDGFRα</i>	<i>A2B5</i>	<i>Olig2</i>	<i>GAD67</i>
iNSPCs	+	+	+	+	ND (in vivo) ND~weak (in vitro)	+	ND	ND	ND
SVZ astrocytes		+			+				
Ependymal cells		+	+		ND~weak	+			
Reactive astrocytes	+				+				
OPCs			+			+	+	+	
Layer1 progenitors	ND								+
Resident glia/glia precursors			+			+			

+, positive expression; ND, nondetectable; iNSPC, ischemia-induced neural stem/progenitor cells; SVZ, subventricular zone; NSC, neural stem cells; OPC, oligodendrocyte precursor cells; GFAP, glial fibrillary acidic protein; PDGFRα, platelet-derived growth factor receptor alpha; PDGFRβ, platelet-derived growth factor receptor beta.

mater and cortex [5]. Consistent with these findings, only pericytes located within ischemic pia mater/cortex but not within nonischemic pia mater/cortex expressed NSPC markers such as nestin and Sox2. These observations suggest that ischemia/hypoxia may enhance the stem-like nature of brain pericytes, although we do not yet know which signaling and/or factors are essential for their induction. Interestingly, it has been demonstrated that some CNS pericytes originate from the neural crest [36] as well as the pia mater [35], and our current study showed that pial iNSPCs expressed various neural crest markers in addition to pericytic markers. These observations also support that pial iNSPCs originated from brain pericytes, which are neural crest derivatives, providing a novel concept that neural crest-derived cells may play a crucial role in CNS repair following cortical infarction.

It is still highly controversial whether periventricular NSPCs are SVZ astrocytes (type 2 cells), ependymal cells, or both [61]. However, our combined immunohistochemical, RT-PCR, electron microscopic, and cell culture analyses demonstrated that pial iNSPCs do not have characteristics identical to those of SVZ astrocytes or ependymal cells. Although pial nestin-positive iNSPCs expressed NG2, a pericyte marker [23], it is also expressed in resident glial cells/glia progenitors [62], which have NSPC activity after brain injury [46,63]. Combined with the present results that the pial iNSPCs expressed other glia markers such as PDGFRα, and that they have the potential to convert into OPCs under certain conditions [55], it is possible that iNSPCs may also have glial cell characteristics. However, neuron production from NG2- and/or PDGFRα-positive glial cells is rare [63,64], suggesting that iNSPCs may be different from these glial cells. The precise origin, lineage, and traits of iNSPCs warrant further investigation; nevertheless, our current study revealed a novel NSPC population.

Neuronal progenitors have been reported to be induced in layer I of the cerebral cortex in response to ischemia. Such neuronal progenitors express GAD67 but not nestin [12]. In contrast, pial iNSPCs did not express GAD67 but revealed strong nestin expression. Therefore, these neuronal progenitors may not be identical to iNSPCs that we showed in the present study. Alternatively, Bifari and colleagues have shown that novel stem/progenitor cells with neuronal differentiation potential reside in the leptomeninges during

development [15]. In the present study, we further demonstrated that pial iNSPCs with neuronal differentiation potential were induced in the poststroke pia mater. These observations suggest that neuronal progenitors in the poststroke cortex including neocortical layer I may originate, in part, from pial iNSPCs that migrate toward the injured cortex in addition to NSPCs that originate in the cortex (e.g., cortical pericytes-derived iNSPCs and/or other types of proposed NSPCs such as resident glia, reactive astrocytes, and layer I progenitors).

In the present study, we found that iNSPCs developed in close association with blood vessels of the brain from the pia mater to the cortex in the poststroke area. These results suggest that environmental controls surrounding iNSPCs such as ECs, which are an important niche for NSPCs [65], and perivascular (Virchow-Robin) spaces, in which many inflammatory cells infiltrate after brain injury [21], are also essential for cortical neurogenesis after ischemic stroke. In support of these view points, our previous studies showed that adequate support of iNSPCs in the in vivo environment of the poststroke cortex (e.g., regulation of ECs [6,8] and of inflammatory cells [7]) could promote iNSPCs proliferation and enhance neurogenesis.

In conclusion, we have demonstrated that iNSPCs, which are potential sources for neocortical neurogenesis, developed in the poststroke pia mater, suggesting that pia mater that covers the entire brain cortex is an important target for cortical neurogenesis following stroke. Thus, deposition of several materials, including cell sheets [66], on poststroke pia mater may make it possible to promote cortical repair/neurogenesis without cerebral parenchymal damage by intracerebral cell transplantation in the future.

Acknowledgments

This work was partially supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology (21700363 to T. Nakagomi and 21500359 to T. Matsuyama), Hyogo Science and Technology Association (to T. Nakagomi), and Takeda Science Foundation, 2009 (to T. Nakagomi). We would like to thank Y. Okinaka, Y. Tanaka, and Y. Tatsumi for technical assistance, and F.M. Oeschger, C.E. Turnquist-Held, F.G. Szele, H.B. Stolp, F. García-Moreno, W.Z. Wang, C.C. Young, and E.H. Chang for helpful discussions.

Author Disclosure Statement

The authors declare that no conflict of interest exists.

References

- Magavi SS, BR Leavitt and JD Macklis. (2000). Induction of neurogenesis in the neocortex of adult mice. *Nature* 405: 951–955.
- Jiang W, W Gu, T Brannstrom, R Rosqvist and P Wester. (2001). Cortical neurogenesis in adult rats after transient middle cerebral artery occlusion. *Stroke* 32:1201–1207.
- Yang Z, MV Covey, CL Bitel, L Ni, GM Jonakait and SW Levison. (2007). Sustained neocortical neurogenesis after neonatal hypoxic/ischemic injury. *Ann Neurol* 61:199–208.
- Jin K, X Wang, L Xie, XO Mao, W Zhu, Y Wang, J Shen, Y Mao, S Banwait and DA Greenberg. (2006). Evidence for stroke-induced neurogenesis in the human brain. *Proc Natl Acad Sci U S A* 103:13198–13202.
- Nakagomi T, A Taguchi, Y Fujimori, O Saino, A Nakano-Doi, S Kubo, A Gotoh, T Soma, H Yoshikawa, et al. (2009). Isolation and characterization of neural stem/progenitor cells from post-stroke cerebral cortex in mice. *Eur J Neurosci* 29:1842–1852.
- Nakano-Doi A, T Nakagomi, M Fujikawa, N Nakagomi, S Kubo, S Lu, H Yoshikawa, T Soma, A Taguchi and T Matsuyama. (2010). Bone marrow mononuclear cells promote proliferation of endogenous neural stem cells through vascular niches after cerebral infarction. *Stem Cells* 28:1292–1302.
- Saino O, A Taguchi, T Nakagomi, A Nakano-Doi, S Kashiwamura, N Doe, N Nakagomi, T Soma, H Yoshikawa, et al. (2010). Immunodeficiency reduces neural stem/progenitor cell apoptosis and enhances neurogenesis in the cerebral cortex after stroke. *J Neurosci Res* 88:2385–2397.
- Nakagomi N, T Nakagomi, S Kubo, A Nakano-Doi, O Saino, M Takata, H Yoshikawa, DM Stern, T Matsuyama and A Taguchi. (2009). Endothelial cells support survival, proliferation, and neuronal differentiation of transplanted adult ischemia-induced neural stem/progenitor cells after cerebral infarction. *Stem Cells* 27:2185–2195.
- Nakayama D, T Matsuyama, H Ishibashi-Ueda, T Nakagomi, Y Kasahara, H Hirose, A Kikuchi-Taura, DM Stern, H Mori and A Taguchi. (2010). Injury-induced neural stem/progenitor cells in post-stroke human cerebral cortex. *Eur J Neurosci* 31:90–98.
- Sirko S, A Neitz, T Mittmann, A Horvat-Brocker, A von Holst, UT Eysel and A Faissner. (2009). Focal laser-lesions activate an endogenous population of neural stem/progenitor cells in the adult visual cortex. *Brain* 132:2252–2264.
- Xue JH, H Yanamoto, Y Nakajo, N Tohnai, Y Nakano, T Hori, K Iihara and S Miyamoto. (2009). Induced spreading depression evokes cell division of astrocytes in the subpial zone, generating neural precursor-like cells and new immature neurons in the adult cerebral cortex. *Stroke* 40:e606–e613.
- Ohira K, T Furuta, H Hioki, KC Nakamura, E Kuramoto, Y Tanaka, N Funatsu, K Shimizu, T Oishi, et al. (2010). Ischemia-induced neurogenesis of neocortical layer 1 progenitor cells. *Nat Neurosci* 13:173–179.
- Sockanathan S and N Gaiano. (2009). Preview. Meninges play a radical role in embryonic neural stem cell regulation. *Cell Stem Cell* 5:455–456.
- Siegenthaler JA, AM Ashique, K Zarbalis, KP Patterson, JH Hecht, MA Kane, AE Folias, Y Choe, SR May, et al. (2009). Retinoic acid from the meninges regulates cortical neuron generation. *Cell* 139:597–609.
- Bifari F, I Decimo, C Chiamulera, E Bersan, G Malpeli, J Johansson, V Lisi, B Bonetti, G Fumagalli, G Pizzolo and M Krampera. (2009). Novel stem/progenitor cells with neuronal differentiation potential reside in the leptomeningeal niche. *J Cell Mol Med* 13:3195–3208.
- Taguchi A, T Soma, H Tanaka, T Kanda, H Nishimura, H Yoshikawa, Y Tsukamoto, H Iso, Y Fujimori, et al. (2004). Administration of CD34+ cells after stroke enhances neurogenesis via angiogenesis in a mouse model. *J Clin Invest* 114:330–338.
- Kubo S, MC Seleme, HS Soifer, JL Perez, JV Moran, HH Kazazian Jr. and N Kasahara. (2006). L1 retrotransposition in nondividing and primary human somatic cells. *Proc Natl Acad Sci U S A* 103:8036–8041.
- Yamamoto K, T Matsuyama, S Shiosaka, S Inagaki, E Senba, Y Shimizu, I Ishimoto, T Hayakawa, M Matsumoto and M Tohyama. (1983). Overall distribution of substance P-containing nerves in the wall of the cerebral arteries of the guinea pig and its origins. *J Comp Neurol* 215:421–426.
- Gatter KC, MS Dunnill, J Gerdes, H Stein and DY Mason. (1986). New approach to assessing lung tumours in man. *J Clin Pathol* 39:590–593.
- Zhang ET, CB Inman and RO Weller. (1990). Interrelationships of the pia mater and the perivascular (Virchow-Robin) spaces in the human cerebrum. *J Anat* 170:111–123.
- Hutchings M and RO Weller. (1986). Anatomical relationships of the pia mater to cerebral blood vessels in man. *J Neurosurg* 65:316–325.
- Fricke B, M von During and KH Andres. (1997). Topography and immunocytochemical characterization of nerve fibers in the leptomeningeal compartments of the rat. A light- and electron-microscopical study. *Cell Tissue Res* 287:11–22.
- Ozerdem U, E Monosov and WB Stallcup. (2002). NG2 proteoglycan expression by pericytes in pathological microvasculature. *Microvasc Res* 63:129–134.
- Hellstrom M, M Kalen, P Lindahl, A Abramsson and C Betsholtz. (1999). Role of PDGF-B and PDGFR-beta in recruitment of vascular smooth muscle cells and pericytes during embryonic blood vessel formation in the mouse. *Development* 126:3047–3055.
- Dore-Duffy P, A Katychiev, X Wang and E Van Buren. (2006). CNS microvascular pericytes exhibit multipotential stem cell activity. *J Cereb Blood Flow Metab* 26:613–624.
- Jung KH, K Chu, ST Lee, JJ Bahn, D Jeon, JH Kim, S Kim, CH Won, M Kim, SK Lee and JK Roh. (2011). Multipotent PDGFRbeta-expressing cells in the circulation of stroke patients. *Neurobiol Dis* 41:489–497.
- Bull ND and PF Bartlett. (2005). The adult mouse hippocampal progenitor is neurogenic but not a stem cell. *J Neurosci* 25:10815–10821.
- Nunes MC, NS Roy, HM Keyoung, RR Goodman, G McKhann 2nd, L Jiang, J Kang, M Nedergaard and SA Goldman. (2003). Identification and isolation of multipotential neural progenitor cells from the subcortical white matter of the adult human brain. *Nat Med* 9:439–447.
- Gravdal K, OJ Halvorsen, SA Haukaas and LA Akslen. (2009). Proliferation of immature tumor vessels is a novel marker of clinical progression in prostate cancer. *Cancer Res* 69:4708–4715.

30. Ellis P, BM Fagan, ST Magness, S Hutton, O Taranova, S Hayashi, A McMahon, M Rao and L Pevny. (2004). SOX2, a persistent marker for multipotential neural stem cells derived from embryonic stem cells, the embryo or the adult. *Dev Neurosci* 26:148–165.
31. Suh H, A Consiglio, J Ray, T Sawai, KA D'Amour and FH Gage. (2007). In vivo fate analysis reveals the multipotent and self-renewal capacities of Sox2+ neural stem cells in the adult hippocampus. *Cell Stem Cell* 1:515–528.
32. Brazel CY, TL Limke, JK Osborne, T Miura, J Cai, L Pevny and MS Rao. (2005). Sox2 expression defines a heterogeneous population of neurosphere-forming cells in the adult murine brain. *Aging Cell* 4:197–207.
33. Kim JB, V Sebastiao, G Wu, MJ Arauzo-Bravo, P Sasse, L Gentile, K Ko, D Ruau, M Ehrlich, et al. (2009). Oct4-induced pluripotency in adult neural stem cells. *Cell* 136:411–419.
34. Takahashi K, K Tanabe, M Ohnuki, M Narita, T Ichisaka, K Tomoda and S Yamanaka. (2007). Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell* 131:861–872.
35. Morse DE and JL Cova. (1984). Pigmented cells in the leptomeninges of the cat. *Anat Rec* 210:125–132.
36. Etchevers HC, C Vincent, NM Le Douarin and GF Couly. (2001). The cephalic neural crest provides pericytes and smooth muscle cells to all blood vessels of the face and forebrain. *Development* 128:1059–1068.
37. Aihara Y, Y Hayashi, M Hirata, N Arikawa, S Shibata, N Nagoshi, M Nakanishi, K Ohnuma, M Warashina, et al. (2010). Induction of neural crest cells from mouse embryonic stem cells in a serum-free monolayer culture. *Int J Dev Biol* 54:1287–1294.
38. Doetsch F, I Caille, DA Lim, JM Garcia-Verdugo and A Alvarez-Buylla. (1999). Subventricular zone astrocytes are neural stem cells in the adult mammalian brain. *Cell* 97:703–716.
39. Jackson EL, JM Garcia-Verdugo, S Gil-Perotin, M Roy, A Quinones-Hinojosa, S VandenBerg and A Alvarez-Buylla. (2006). PDGFR alpha-positive B cells are neural stem cells in the adult SVZ that form glioma-like growths in response to increased PDGF signaling. *Neuron* 51:187–199.
40. Carlen M, K Meletis, C Goritz, V Darsalia, E Evergren, K Tanigaki, M Amendola, F Barnabe-Heider, MS Yeung, et al. (2009). Forebrain ependymal cells are Notch-dependent and generate neuroblasts and astrocytes after stroke. *Nat Neurosci* 12:259–267.
41. Moreno-Manzano V, FJ Rodriguez-Jimenez, M Garcia-Rosello, S Lainez, S Erceg, MT Calvo, M Ronaghi, M Lloret, R Planells-Cases, JM Sanchez-Puelles and M Stojkovic. (2009). Activated spinal cord ependymal stem cells rescue neurological function. *Stem Cells* 27:733–743.
42. Gaughwin PM, MA Caldwell, JM Anderson, CJ Schwiening, JW Fawcett, DA Compston and S Chandran. (2006). Astrocytes promote neurogenesis from oligodendrocyte precursor cells. *Eur J Neurosci* 23:945–956.
43. Kondo T and M Raff. (2000). Oligodendrocyte precursor cells reprogrammed to become multipotential CNS stem cells. *Science* 289:1754–1757.
44. Kriegstein A and A Alvarez-Buylla. (2009). The glial nature of embryonic and adult neural stem cells. *Annu Rev Neurosci* 32:149–184.
45. Belachew S, R Chittajallu, AA Aguirre, X Yuan, M Kirby, S Anderson and V Gallo. (2003). Postnatal NG2 proteoglycan-expressing progenitor cells are intrinsically multipotent and generate functional neurons. *J Cell Biol* 161:169–186.
46. Yokoyama A, A Sakamoto, K Kameda, Y Imai and J Tanaka. (2006). NG2 proteoglycan-expressing microglia as multipotent neural progenitors in normal and pathologic brains. *Glia* 53:754–768.
47. Coskun V, H Wu, B Bianchi, S Tsao, K Kim, J Zhao, JC Biancotti, L Hutnick, RC Krueger Jr., G Fan, J de Vellis and YE Sun. (2008). CD133+ neural stem cells in the ependyma of mammalian postnatal forebrain. *Proc Natl Acad Sci U S A* 105:1026–1031.
48. Danilov AI, W Gomes-Leal, H Ahlenius, Z Kokaia, E Carlemalm and O Lindvall. (2009). Ultrastructural and antigenic properties of neural stem cells and their progeny in adult rat subventricular zone. *Glia* 57:136–152.
49. Shimada IS, BM Peterson and JL Speers. (2010). Isolation of locally derived stem/progenitor cells from the peri-infarct area that do not migrate from the lateral ventricle after cortical stroke. *Stroke* 41:e552–e560.
50. Oki K, N Kaneko, H Kanki, T Imai, N Suzuki, K Sawamoto and H Okano. (2010). Musashi1 as a marker of reactive astrocytes after transient focal brain ischemia. *Neurosci Res* 66:390–395.
51. Xu M, W Su and QP Xu. (2010). Aquaporin-4 and traumatic brain edema. *Chin J Traumatol* 13:103–110.
52. Ulrich R, F Seeliger, M Kreutzer, PG Germann and W Baumgartner. (2008). Limited remyelination in Theiler's murine encephalomyelitis due to insufficient oligodendroglial differentiation of nerve/glia antigen 2 (NG2)-positive putative oligodendroglial progenitor cells. *Neuropathol Appl Neurobiol* 34:603–620.
53. Hall A, NA Giese and WD Richardson. (1996). Spinal cord oligodendrocytes develop from ventrally derived progenitor cells that express PDGF alpha-receptors. *Development* 122:4085–4094.
54. Billon N, C Jolicœur, QL Ying, A Smith and M Raff. (2002). Normal timing of oligodendrocyte development from genetically engineered, lineage-selectable mouse ES cells. *J Cell Sci* 115:3657–3665.
55. Chen Y, V Balasubramanian, J Peng, EC Hurlock, M Tallquist, J Li and QR Lu. (2007). Isolation and culture of rat and mouse oligodendrocyte precursor cells. *Nat Protoc* 2:1044–1051.
56. Mignone JL, V Kukekov, AS Chiang, D Steindler and G Enikolopov. (2004). Neural stem and progenitor cells in nestin-GFP transgenic mice. *J Comp Neurol* 469:311–324.
57. Kronenberg G, LP Wang, M Synowitz, K Gertz, J Katchanov, R Glass, C Harms, G Kempermann, H Kettenmann and M Endres. (2005). Nestin-expressing cells divide and adopt a complex electrophysiologic phenotype after transient brain ischemia. *J Cereb Blood Flow Metab* 25:1613–1624.
58. Jin K, M Minami, JQ Lan, XO Mao, S Bateur, RP Simon and DA Greenberg. (2001). Neurogenesis in dentate subgranular zone and rostral subventricular zone after focal cerebral ischemia in the rat. *Proc Natl Acad Sci U S A* 98:4710–4715.
59. Goings GE, V Sahni and FG Szele. (2004). Migration patterns of subventricular zone cells in adult mice change after cerebral cortex injury. *Brain Res* 996:213–226.
60. Crisan M, CW Chen, M Corselli, G Andriolo, L Lazzari and B Peault. (2009). Perivascular multipotent progenitor cells in human organs. *Ann N Y Acad Sci* 1176:118–123.
61. Chojnacki AK, GK Mak and S Weiss. (2009). Identity crisis for adult periventricular neural stem cells: subventricular zone astrocytes, ependymal cells or both? *Nat Rev Neurosci* 10:153–163.

62. Stallcup WB and L Beasley. (1987). Bipotential glial precursor cells of the optic nerve express the NG2 proteoglycan. *J Neurosci* 7:2737–2744.
63. Zawadzka M, LE Rivers, SP Fancy, C Zhao, R Tripathi, F Jamen, K Young, A Goncharevich, H Pohl, et al. (2010). CNS-resident glial progenitor/stem cells produce Schwann cells as well as oligodendrocytes during repair of CNS demyelination. *Cell Stem Cell* 6:578–590.
64. Richardson WD, KM Young, RB Tripathi and I McKenzie. (2011). NG2-glia as multipotent neural stem cells: fact or fantasy? *Neuron* 70:661–673.
65. Shen Q, Y Wang, E Kokovay, G Lin, SM Chuang, SK Goderie, B Roysam and S Temple. (2008). Adult SVZ stem cells lie in a vascular niche: a quantitative analysis of niche cell-cell interactions. *Cell Stem Cell* 3:289–300.
66. Derval N, L Barandon, P Dufourcq, L Leroux, JM Lamaziere, D Daret, T Couffinhal and C Duplaa. (2008). Epicardial deposition of endothelial progenitor and mesenchymal

stem cells in a coated muscle patch after myocardial infarction in a murine model. *Eur J Cardiothorac Surg* 34:248–254.

Address correspondence to:

Dr. Takayuki Nakagomi
Institute for Advanced Medical Sciences
Hyogo College of Medicine,
1-1 Mukogawacho, Nishinomiya
Hyogo 663-8501
Japan

E-mail: nakagomi@hyo-med.ac.jp

Received for publication June 4, 2011

Accepted after revision July 5, 2011

Prepublished on Liebert Instant Online July 5, 2011

This article has been cited by:

1. Ilaria Decimo, Sissi Dolci, Gabriella Panuccio, Marco Riva, Guido Fumagalli, Francesco Bifari. 2021. Meninges: A Widespread Niche of Neural Progenitors for the Brain. *The Neuroscientist* **27**:5, 506-528. [[Crossref](#)]
2. Stefania Zorzini, Andrea Corsi, Francesca Ciarpella, Emanuela Bottani, Sissi Dolci, Giorgio Malpeli, Annachiara Pino, Alessia Amenta, Guido Francesco Fumagalli, Cristiano Chiamulera, Francesco Bifari, Ilaria Decimo. 2021. Environmental Enrichment Induces Meningeal Niche Remodeling through TrkB-Mediated Signaling. *International Journal of Molecular Sciences* **22**:19, 10657. [[Crossref](#)]
3. Kazuhide Hayakawa, Evan Y. Snyder, Eng H. Lo. 2021. Meningeal Multipotent Cells: A Hidden Target for CNS Repair?. *NeuroMolecular Medicine* **23**:3, 339-343. [[Crossref](#)]
4. Koji Ohira. 2021. Cortical adult neurogenesis and its biological implication. *Clinical and Experimental Neuroimmunology* **10**. . [[Crossref](#)]
5. Milton H. Hamblin, Jean-Pyo Lee. 2021. Neural Stem Cells for Early Ischemic Stroke. *International Journal of Molecular Sciences* **22**:14, 7703. [[Crossref](#)]
6. Sofia Passera, Marta Boccazzi, Cindy Bokobza, Valerie Faivre, Fabio Mosca, Juliette Van Steenwinckel, Monica Fumagalli, Pierre Gressens, Bobbi Fleiss. 2021. Therapeutic potential of stem cells for preterm infant brain damage: Can we move from the heterogeneity of preclinical and clinical studies to established therapeutics?. *Biochemical Pharmacology* **186**, 114461. [[Crossref](#)]
7. Akhlaq A. Farooqui. Molecular aspects of regeneration and neuroprotection in neurotraumatic diseases 181-224. [[Crossref](#)]
8. Liyun Wang, Xiaoxing Xiong, Luyuan Zhang, Jian Shen. 2021. Neurovascular Unit: A critical role in ischemic stroke. *CNS Neuroscience & Therapeutics* **27**:1, 7-16. [[Crossref](#)]
9. Yoichi Morofuji, Shinsuke Nakagawa, Takashi Fujimoto, Susumu Yamaguchi, Kenta Ujifuku, Takayuki Matsuo. Biology of Pericytes – Recent Advances: Role of Pericytes in Brain Metastasis 75-98. [[Crossref](#)]
10. Jamie N. Mayo, Shawn E. Bearden. 2020. Pericytes as Cell Therapy for Locomotor Recovery. *Current Tissue Microenvironment Reports* **1**:4, 199-207. [[Crossref](#)]
11. Mikhail M. Moisenovich, Denis N. Silachev, Anastasia M. Moysenovich, Anastasia Yu. Arkhipova, Konstantin V. Shaitan, Vladimir G. Bogush, Vladimir G. Debabov, Alexander V. Latanov, Irina B. Pevzner, Ljubava D. Zorova, Valentina A. Babenko, Egor Y. Plotnikov, Dmitry B. Zorov. 2020. Effects of Recombinant Spidroin rS1/9 on Brain Neural Progenitors After Photothrombosis-Induced Ischemia. *Frontiers in Cell and Developmental Biology* **8**. . [[Crossref](#)]
12. Takayuki Nakagomi, Yasue Tanaka, Nami Nakagomi, Tomohiro Matsuyama, Shinichi Yoshimura. 2020. How Long Are Reperfusion Therapies Beneficial for Patients after Stroke Onset? Lessons from Lethal Ischemia Following Early Reperfusion in a Mouse Model of Stroke. *International Journal of Molecular Sciences* **21**:17, 6360. [[Crossref](#)]
13. Saujanya Rajbhandari, Mikiya Beppu, Toshinori Takagi, Akiko Nakano-Doi, Nami Nakagomi, Tomohiro Matsuyama, Takayuki Nakagomi, Shinichi Yoshimura. 2020. Ischemia-Induced Multipotent Stem Cells Isolated from Stroke Patients Exhibit Higher Neurogenic Differentiation Potential than Bone Marrow-Derived Mesenchymal Stem Cells. *Stem Cells and Development* **29**:15, 994-1006. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplementary Material](#)]
14. Yasue Tanaka, Nami Nakagomi, Nobutaka Doe, Akiko Nakano-Doi, Toshinori Sawano, Toshinori Takagi, Tomohiro Matsuyama, Shinichi Yoshimura, Takayuki Nakagomi. 2020. Early Reperfusion Following Ischemic Stroke Provides Beneficial Effects, Even After Lethal Ischemia with Mature Neural Cell Death. *Cells* **9**:6, 1374. [[Crossref](#)]
15. Wenting Zhang, Ling Zhu, Chengrui An, Rongrong Wang, Liqun Yang, Weifeng Yu, Peiying Li, Yanqin Gao. 2020. The blood brain barrier in cerebral ischemic injury – Disruption and repair. *Brain Hemorrhages* **1**:1, 34-53. [[Crossref](#)]
16. Tae-Ryong Riew, Xuyan Jin, Hong Lim Kim, Soojin Kim, Mun-Yong Lee. 2020. Ultrastructural and Molecular Characterization of Platelet-derived growth factor Beta-Positive Leptomeningeal Cells in the Adult Rat Brain. *Molecular Neurobiology* **57**:3, 1484-1501. [[Crossref](#)]
17. Stuart P. Atkinson. 2020. A preview of selected articles. *STEM CELLS Translational Medicine* **9**:1, 1-3. [[Crossref](#)]
18. Stephanie Bonney, Swati Mishra, Samuel J. Pleasure, Julie A. Siegenthaler. Meninges and vasculature 1037-1063. [[Crossref](#)]
19. Yuki Maeda, Nami Nakagomi, Akiko Nakano-Doi, Hiroto Ishikawa, Yoshiki Tatsumi, Yoshio Bando, Hiroo Yoshikawa, Tomohiro Matsuyama, Fumi Gomi, Takayuki Nakagomi. 2019. Potential of Adult Endogenous Neural Stem/Progenitor Cells in the Spinal Cord to Contribute to Remyelination in Experimental Autoimmune Encephalomyelitis. *Cells* **8**:9, 1025. [[Crossref](#)]
20. Takayuki Nakagomi, Toshinori Takagi, Mikiya Beppu, Shinichi Yoshimura, Tomohiro Matsuyama. 2019. Neural regeneration by regionally induced stem cells within post-stroke brains: Novel therapy perspectives for stroke patients. *World Journal of Stem Cells* **11**:8, 452-463. [[Crossref](#)]

21. Li Lin, Chu Lisheng, Ren Cuicui, Wang Jun, Sun Siqi, Li Tianyi, Yin Yuanjun. 2019. Enhanced Migration of Bone Marrow-Derived Mesenchymal Stem Cells with Tetramethylpyrazine and Its Synergistic Effect on Angiogenesis and Neurogenesis After Cerebral Ischemia in Rats. *Stem Cells and Development* **28**:13, 871-881. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
22. Nikolai M. Nemirovich-Danchenko, Marina Yu. Khodanovich. 2019. New Neurons in the Post-ischemic and Injured Brain: Migrating or Resident?. *Frontiers in Neuroscience* **13**. . [[Crossref](#)]
23. Kyosuke Yamanishi, Nobutaka Doe, Keiichiro Mukai, Kaoru Ikubo, Takuya Hashimoto, Noriko Uwa, Miho Sumida, Yosif El-Darawish, Naomi Gamachi, Wen Li, Sachi Kuwahara-Otani, Seishi Maeda, Yuko Watanabe, Tetsu Hayakawa, Hiromichi Yamanishi, Tomohiro Matsuyama, Hideshi Yagi, Haruki Okamura, Hisato Matsunaga. 2019. Interleukin-18-deficient mice develop hippocampal abnormalities related to possible depressive-like behaviors. *Neuroscience* **408**, 147-160. [[Crossref](#)]
24. Beppu Mikiya, Nakagomi Takayuki, Takagi Toshinori, Nakano-Doi Akiko, Sakuma Rika, Kuramoto Yoji, Tatebayashi Kotaro, Matsuyama Tomohiro, Yoshimura Shinichi. 2019. Isolation and Characterization of Cerebellum-Derived Stem Cells in Poststroke Human Brain. *Stem Cells and Development* **28**:8, 528-542. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplementary Material](#)]
25. Bikei Ryu, Hidekazu Sekine, Jun Homma, Tomonori Kobayashi, Eiji Kobayashi, Takakazu Kawamata, Tatsuya Shimizu. 2019. Allogeneic adipose-derived mesenchymal stem cell sheet that produces neurological improvement with angiogenesis and neurogenesis in a rat stroke model. *Journal of Neurosurgery* **7**, 1-14. [[Crossref](#)]
26. Austin C. Boese, Quan-Son Eric Le, Dylan Pham, Milton H. Hamblin, Jean-Pyo Lee. 2018. Neural stem cell therapy for subacute and chronic ischemic stroke. *Stem Cell Research & Therapy* **9**:1. . [[Crossref](#)]
27. Tae-Ryong Riew, Jeong-Heon Choi, Hong Lim Kim, Xuyan Jin, Mun-Yong Lee. 2018. PDGFR- β -Positive Perivascular Adventitial Cells Expressing Nestin Contribute to Fibrotic Scar Formation in the Striatum of 3-NP Intoxicated Rats. *Frontiers in Molecular Neuroscience* **11**. . [[Crossref](#)]
28. Sakuma Rika, Takahashi Ai, Nakano-Doi Akiko, Sawada Rikako, Kamachi Saeko, Beppu Mikiya, Takagi Toshinori, Yoshimura Shinichi, Matsuyama Tomohiro, Nakagomi Takayuki. 2018. Comparative Characterization of Ischemia-Induced Brain Multipotent Stem Cells with Mesenchymal Stem Cells: Similarities and Differences. *Stem Cells and Development* **27**:19, 1322-1338. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplementary Material](#)]
29. M.C. Rusu, V.S. Mănoiu, D. Crețoiu, S.M. Crețoiu, A.D. Vrapciu. 2018. Stromal cells/telocytes and endothelial progenitors in the perivascular niches of the trigeminal ganglion. *Annals of Anatomy - Anatomischer Anzeiger* **218**, 141-155. [[Crossref](#)]
30. G S P Santos, P H D M Prazeres, A Mintz, A Birbrair. 2018. Role of pericytes in the retina. *Eye* **32**:3, 483-486. [[Crossref](#)]
31. Toshinori Takagi, Shinichi Yoshimura. 2018. A national survey of acute thrombectomy (RESCUE-Japan Project) and possibility of regenerative therapy using injury-induced multipotent stem cells. *Cerebral Blood Flow and Metabolism (Japanese journal of cerebral blood flow and metabolism)* **30**:1, 59-64. [[Crossref](#)]
32. Toshinori Takagi, Shinichi Yoshimura, Rika Sakuma, Akiko Nakano-Doi, Tomohiro Matsuyama, Takayuki Nakagomi. 2017. Novel Regenerative Therapies Based on Regionally Induced Multipotent Stem Cells in Post-Stroke Brains: Their Origin, Characterization, and Perspective. *Translational Stroke Research* **8**:6, 515-528. [[Crossref](#)]
33. Sissi Dolci, Annachiara Pino, Valeria Berton, Pau Gonzalez, Alice Braga, Marta Fumagalli, Elisabetta Bonfanti, Giorgio Malpeli, Francesca Pari, Stefania Zorzini, Clelia Amoroso, Denny Moscon, Francisco J. Rodriguez, Guido Fumagalli, Francesco Bifari, Ilaria Decimo. 2017. High Yield of Adult Oligodendrocyte Lineage Cells Obtained from Meningeal Biopsy. *Frontiers in Pharmacology* **8**. . [[Crossref](#)]
34. Annachiara Pino, Guido Fumagalli, Francesco Bifari, Ilaria Decimo. 2017. New neurons in adult brain: distribution, molecular mechanisms and therapies. *Biochemical Pharmacology* **141**, 4-22. [[Crossref](#)]
35. Tatebayashi Kotaro, Tanaka Yasue, Nakano-Doi Akiko, Sakuma Rika, Kamachi Saeko, Shirakawa Manabu, Uchida Kazutaka, Kageyama Hiroto, Takagi Toshinori, Yoshimura Shinichi, Matsuyama Tomohiro, Nakagomi Takayuki. 2017. Identification of Multipotent Stem Cells in Human Brain Tissue Following Stroke. *Stem Cells and Development* **26**:11, 787-797. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplementary Material](#)]
36. Masayo Nakata, Takayuki Nakagomi, Mitsuyo Maeda, Akiko Nakano-Doi, Yoshihiro Momota, Tomohiro Matsuyama. 2017. Induction of Perivascular Neural Stem Cells and Possible Contribution to Neurogenesis Following Transient Brain Ischemia/Reperfusion Injury. *Translational Stroke Research* **8**:2, 131-143. [[Crossref](#)]
37. Francesco Bifari, Ilaria Decimo, Annachiara Pino, Enric Llorens-Bobadilla, Sheng Zhao, Christian Lange, Gabriella Panuccio, Bram Boeckx, Bernard Thienpont, Stefan Vinckier, Sabine Wyns, Ann Bouché, Diether Lambrechts, Michele Giugliano, Mieke Dewerchin, Ana Martin-Villalba, Peter Carmeliet. 2017. Neurogenic Radial Glia-like Cells in Meninges Migrate and Differentiate into Functionally Integrated Neurons in the Neonatal Cortex. *Cell Stem Cell* **20**:3, 360-373.e7. [[Crossref](#)]
38. Adriana Octaviana Dulamea. Role of Oligodendrocyte Dysfunction in Demyelination, Remyelination and Neurodegeneration in Multiple Sclerosis 91-127. [[Crossref](#)]

39. Yoshihiro Momota. 2017. Induction of pericyte-derived neural stem cells following transient brain ischemia/reperfusion injury. *Cerebral Blood Flow and Metabolism (Japanese journal of cerebral blood flow and metabolism)* **28**:2, 347-351. [[Crossref](#)]
40. Rika Sakuma, Maiko Kawahara, Akiko Nakano-Doi, Ai Takahashi, Yasue Tanaka, Aya Narita, Sachi Kuwahara-Otani, Tetsu Hayakawa, Hideshi Yagi, Tomohiro Matsuyama, Takayuki Nakagomi. 2016. Brain pericytes serve as microglia-generating multipotent vascular stem cells following ischemic stroke. *Journal of Neuroinflammation* **13**:1. . [[Crossref](#)]
41. Harmanvir Ghuman, Michel Modo. 2016. Biomaterial applications in neural therapy and repair. *Chinese Neurosurgical Journal* **2**:1. . [[Crossref](#)]
42. Lawrence Edelstein, Kjell Fuxe, Michael Levin, Bogdan O. Popescu, John Smythies. 2016. Telocytes in their context with other intercellular communication agents. *Seminars in Cell & Developmental Biology* **55**, 9-13. [[Crossref](#)]
43. Ana Pombero, Raquel Garcia-Lopez, Salvador Martinez. 2016. Brain mesenchymal stem cells: physiology and pathological implications. *Development, Growth & Differentiation* **58**:5, 469-480. [[Crossref](#)]
44. Jialei Yang, Xiufen Zhang, Yin Wu, Bo Zhao, Xunyuan Liu, Yuanhang Pan, Yonghong Liu, Yuqiang Ding, Mengsheng Qiu, Ya-Zhou Wang, Gang Zhao. 2016. Wnt/ β -catenin signaling mediates the seizure-facilitating effect of postischemic reactive astrocytes after pentylentetrazole-kindling. *Glia* **10**, n/a-n/a. [[Crossref](#)]
45. Mike Dragunow, Sheryl Feng, Justin Rustenhoven, Maurice Curtis, Richard Faull. 2016. Studying Human Brain Inflammation in Leptomeningeal and Choroid Plexus Explant Cultures. *Neurochemical Research* **41**:3, 579-588. [[Crossref](#)]
46. Laura Cristina Ceafalan, Bogdan Ovidiu Popescu. Juxtacerebral Tissue Regeneration Potential: Telocytes Contribution 397-402. [[Crossref](#)]
47. Li Chen, Xiaoting Wang, Jian Zhang, Chao Dang, Gang Liu, Zhijian Liang, Gelun Huang, Weijia Zhao, Jinsheng Zeng. 2016. Tongxinluo Enhances Neurogenesis and Angiogenesis in Peri-Infarct Area and Subventricular Zone and Promotes Functional Recovery after Focal Cerebral Ischemic Infarction in Hypertensive Rats. *Evidence-Based Complementary and Alternative Medicine* **2016**, 1-9. [[Crossref](#)]
48. Quentin Marlier, Sebastien Verteneuil, Renaud Vandenbosch, Brigitte Malgrange. 2015. Mechanisms and Functional Significance of Stroke-Induced Neurogenesis. *Frontiers in Neuroscience* **9**. . [[Crossref](#)]
49. A.I. Derjac-Aramă, C. Sarafoleanu, C.M. Manea, M.I. Nicolescu, A.D. Vrapciu, M.C. Rusu. 2015. Regenerative Potential of Human Schneiderian Membrane: Progenitor Cells and Epithelial-Mesenchymal Transition. *The Anatomical Record* **298**:12, 2132-2140. [[Crossref](#)]
50. Abdelhak Belmadani, Dongjun Ren, Bula J. Bhattacharyya, Katharina B. Rothwangl, Thomas J. Hope, Harris Perlman, Richard J. Miller. 2015. Identification of a sustained neurogenic zone at the dorsal surface of the adult mouse hippocampus and its regulation by the chemokine SDF-1. *Hippocampus* **25**:11, 1224-1241. [[Crossref](#)]
51. Jamie N. Mayo, Shawn E. Bearden. 2015. Driving the Hypoxia-Inducible Pathway in Human Pericytes Promotes Vascular Density in an Exosome-Dependent Manner. *Microcirculation* **22**:8, 711-723. [[Crossref](#)]
52. BIN CHEN, JING TAO, YUKUN LIN, RUHUI LIN, WEILIN LIU, LIDIAN CHEN. 2015. Electro-acupuncture exerts beneficial effects against cerebral ischemia and promotes the proliferation of neural progenitor cells in the cortical peri-infarct area through the Wnt/ β -catenin signaling pathway. *International Journal of Molecular Medicine* **36**:5, 1215-1222. [[Crossref](#)]
53. Francesco Bifari, Valeria Berton, Annachiara Pino, Marijana Kusalo, Giorgio Malpeli, Marzia Di Chio, Emanuela Bersan, Eliana Amato, Aldo Scarpa, Mauro Krampera, Guido Fumagalli, Ilaria Decimo. 2015. Meninges harbor cells expressing neural precursor markers during development and adulthood. *Frontiers in Cellular Neuroscience* **9**. . [[Crossref](#)]
54. Nakagomi Takayuki, Nakano-Doi Akiko, Kawamura Miki, Matsuyama Tomohiro. 2015. Do Vascular Pericytes Contribute to Neurovasculogenesis in the Central Nervous System as Multipotent Vascular Stem Cells?. *Stem Cells and Development* **24**:15, 1730-1739. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
55. Jeong Sook Noh, Ha-Jin Pak, Yoo-Jin Shin, Tae-Ryong Riew, Joo-Hee Park, Young Wha Moon, Mun-Yong Lee. 2015. Differential expression of the calcium-sensing receptor in the ischemic and border zones after transient focal cerebral ischemia in rats. *Journal of Chemical Neuroanatomy* **66-67**, 40-51. [[Crossref](#)]
56. Suet-Ping Wong, Jessica E. Rowley, Andia N. Redpath, Jessica D. Tilman, Tariq G. Fellous, Jill R. Johnson. 2015. Pericytes, mesenchymal stem cells and their contributions to tissue repair. *Pharmacology & Therapeutics* **151**, 107-120. [[Crossref](#)]
57. Yihua Bei, Fei Wang, Changqing Yang, Junjie Xiao. 2015. Telocytes in regenerative medicine. *Journal of Cellular and Molecular Medicine* **19**:7, 1441-1454. [[Crossref](#)]
58. Takayuki Nakagomi, Shuji Kubo, Akiko Nakano-Doi, Rika Sakuma, Shan Lu, Aya Narita, Maiko Kawahara, Akihiko Taguchi, Tomohiro Matsuyama. 2015. Brain Vascular Pericytes Following Ischemia Have Multipotential Stem Cell Activity to Differentiate Into Neural and Vascular Lineage Cells. *STEM CELLS* **33**:6, 1962-1974. [[Crossref](#)]

59. Anne Schwerk, Jennifer Altschüler, Manfred Roch, Manfred Gossen, Christine Winter, Jürgen Berg, Andreas Kurtz, Levent Akyüz, Barbara Steiner. 2015. Adipose-derived human mesenchymal stem cells induce long-term neurogenic and anti-inflammatory effects and improve cognitive but not motor performance in a rat model of Parkinson's disease. *Regenerative Medicine* **10**:4, 431-446. [[Crossref](#)]
60. Helen B. Stolp, Zoltán Molnár. 2015. Neurogenic niches in the brain: help and hindrance of the barrier systems. *Frontiers in Neuroscience* **9**. . [[Crossref](#)]
61. Alexander Birbrair, Tan Zhang, Zhong-Min Wang, Maria Laura Messi, Akiva Mintz, Osvaldo Delbono. 2015. Pericytes at the intersection between tissue regeneration and pathology: Figure 1. *Clinical Science* **128**:2, 81-93. [[Crossref](#)]
62. Takayuki Nakagomi, Akiko Nakano-Doi, Aya Narita, Tomohiro Matsuyama. 2015. Concise Review: Are Stimulated Somatic Cells Truly Reprogrammed into an ES/iPS-Like Pluripotent State? Better Understanding by Ischemia-Induced Multipotent Stem Cells in a Mouse Model of Cerebral Infarction. *Stem Cells International* **2015**, 1-6. [[Crossref](#)]
63. Tomohiro Matsuyama, Takayuki Nakagomi. 2015. Neuroprotection and neurogenesis by brain vascular pericytes. *Cerebral Blood Flow and Metabolism (Japanese journal of cerebral blood flow and metabolism)* **26**:2, 145-149. [[Crossref](#)]
64. Takayuki Nakagomi. 2015. Brain-derived ischemia-induced multipotent stem cells. *Cerebral Blood Flow and Metabolism (Japanese journal of cerebral blood flow and metabolism)* **26**:2, 203-206. [[Crossref](#)]
65. Manoj Kumar, Zsolt Csaba, Stéphane Peineau, Rupali Srivastava, Sowmyalakshmi Rasika, Shyamala Mani, Pierre Gressens, Vincent El Ghouzi. 2014. Endogenous cerebellar neurogenesis in adult mice with progressive ataxia. *Annals of Clinical and Translational Neurology* **1**:12, 968-981. [[Crossref](#)]
66. Guo-Dong Feng, Bao-Rong He, Fan Lu, Lin-Hong Liu, Lingling Zhang, Bo Chen, Zu-Ping He, Ding-Jun Hao, Hao Yang. 2014. Fibroblast Growth Factor 4 Is Required but not Sufficient for the Astrocyte Dedifferentiation. *Molecular Neurobiology* **50**:3, 997-1012. [[Crossref](#)]
67. Alexander Birbrair, Tan Zhang, Zhong-Min Wang, Maria L. Messi, Akiva Mintz, Osvaldo Delbono. 2014. Pericytes: multitasking cells in the regeneration of injured, diseased, and aged skeletal muscle. *Frontiers in Aging Neuroscience* **6**. . [[Crossref](#)]
68. Stanley Bazarek, Daniel A. Peterson. 2014. Prospects for engineering neurons from local neocortical cell populations as cell-mediated therapy for neurological disorders. *Journal of Comparative Neurology* **522**:12, 2857-2876. [[Crossref](#)]
69. N. Paolone, S. Manohar, S.H. Hayes, K.M. Wong, R.J. Salvi, J.S. Baizer. 2014. Dissociation of doublecortin expression and neurogenesis in unipolar brush cells in the vestibulocerebellum and dorsal cochlear nucleus of the adult rat. *Neuroscience* **265**, 323-331. [[Crossref](#)]
70. Sherif M. Amr, Ashraf Gouda, Wael T. Koptan, Ahmad A. Galal, Dina Sabry Abdel-Fattah, Laila A. Rashed, Hazem M. Atta, Mohammad T. Abdel-Aziz. 2014. Bridging defects in chronic spinal cord injury using peripheral nerve grafts combined with a chitosan-laminin scaffold and enhancing regeneration through them by co-transplantation with bone-marrow-derived mesenchymal stem cells: Case series of 14 patients. *The Journal of Spinal Cord Medicine* **37**:1, 54-71. [[Crossref](#)]
71. Adriano Nogueira, Mari Sogayar, Alison Colquhoun, Sheila Siqueira, Ariel Nogueira, Paulo Marchiori, Manoel Teixeira. 2014. Existence of a potential neurogenic system in the adult human brain. *Journal of Translational Medicine* **12**:1, 75. [[Crossref](#)]
72. Florence Appaix. 2014. Brain mesenchymal stem cells: The other stem cells of the brain?. *World Journal of Stem Cells* **6**:2, 134. [[Crossref](#)]
73. Alexander Birbrair, Tan Zhang, Zhong-Min Wang, Maria Laura Messi, Grigori N. Enikolopov, Akiva Mintz, Osvaldo Delbono. 2013. Role of Pericytes in Skeletal Muscle Regeneration and Fat Accumulation. *Stem Cells and Development* **22**:16, 2298-2314. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)] [[Supplementary Material](#)]
74. Frederic Mercier, Tina M. Weatherby, Daniel K. Hartline. 2013. Meningeal-like organization of neural tissues in calanoid copepods (Crustacea). *Journal of Comparative Neurology* **521**:4, 760-790. [[Crossref](#)]
75. Yoo-Jin Shin, Hong Lim Kim, Jang-Mi Park, Jeong Min Cho, Seong Yun Kim, Mun-Yong Lee. 2013. Characterization of nestin expression and vessel association in the ischemic core following focal cerebral ischemia in rats. *Cell and Tissue Research* **351**:3, 383-395. [[Crossref](#)]
76. Yoo-Jin Shin, Jang-Mi Park, Jeong Min Cho, Jung-Ho Cha, Seong Yun Kim, Mun-Yong Lee. 2013. Induction of vascular endothelial growth factor receptor-3 expression in perivascular cells of the ischemic core following focal cerebral ischemia in rats. *Acta Histochemica* **115**:2, 170-177. [[Crossref](#)]
77. Mike Dragunow. 2013. Meningeal and choroid plexus cells—Novel drug targets for CNS disorders. *Brain Research* **1501**, 32-55. [[Crossref](#)]

78. Kyeung Min Joo, Bong Gu Kang, Je Young Yeon, Yu Jin Cho, Jae Yeol An, Hyeon Suk Song, Jun Ho Won, Sang Jeong Kim, Seung-Chyul Hong, Do-Hyun Nam. 2013. Experimental and clinical factors influencing long-term stable in vitro expansion of multipotent neural cells from human adult temporal lobes. *Experimental Neurology* **240**, 168-177. [[Crossref](#)]
79. Shogo Ninomiya, Shigeyuki Esumi, Kunimasa Ohta, Takaichi Fukuda, Tetsufumi Ito, Itaru Imayoshi, Ryoichiro Kageyama, Toshio Ikeda, Shigeyoshi Itohara, Nobuaki Tamamaki. 2013. Amygdala kindling induces nestin expression in the leptomeninges of the neocortex. *Neuroscience Research* **75**:2, 121-129. [[Crossref](#)]
80. Alexander Birbrair, Tan Zhang, Zhong-Min Wang, Maria Laura Messi, Grigori N. Enikolopov, Akiva Mintz, Osvaldo Delbono. 2013. Skeletal muscle pericyte subtypes differ in their differentiation potential. *Stem Cell Research* **10**:1, 67-84. [[Crossref](#)]
81. Varun Sharma, Tina W Ling, Sarah S Rewell, David L Hare, David W Howells, Angela Kourakis, Peter J Wookey. 2012. A Novel Population of α -Smooth Muscle Actin-Positive Cells Activated in a Rat Model of Stroke: An Analysis of the Spatio-Temporal Distribution in Response to Ischemia. *Journal of Cerebral Blood Flow & Metabolism* **32**:11, 2055-2065. [[Crossref](#)]
82. Takayuki Nakagomi, Zoltán Molnár, Akihiko Taguchi, Akiko Nakano-Doi, Shan Lu, Yukiko Kasahara, Nami Nakagomi, Tomohiro Matsuyama. 2012. Leptomeningeal-Derived Doublecortin-Expressing Cells in Poststroke Brain. *Stem Cells and Development* **21**:13, 2350-2354. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
83. Arthur Giraldo-Guimarães, Helder Teixeira de Freitas, Bárbara de Paula Coelho, Hugo Macedo-Ramos, Rosalia Mendez-Otero, Leny A. Cavalcante, Wagner Baetas-da-Cruz. 2012. Bone marrow mononuclear cells and mannose receptor expression in focal cortical ischemia. *Brain Research* **1452**, 173-184. [[Crossref](#)]
84. B.O. Popescu, M. Gherghiceanu, S. Kostin, L. Ceafalan, L.M. Popescu. 2012. Telocytes in meninges and choroid plexus. *Neuroscience Letters* **516**:2, 265-269. [[Crossref](#)]