University of Groningen

Pituitary hyperplasia during primary hypothyroidism

Hoogenberg, K; van Tol, KM

Published in:
Thyroid

DOI:
[10.1089/105072503768499743](https://doi.org/10.1089/105072503768499743)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2003

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Hoogenberg, K., & van Tol, KM. (2003). Pituitary hyperplasia during primary hypothyroidism. *Thyroid*, 13(8), 831-832. <https://doi.org/10.1089/105072503768499743>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Images in Thyroidology*

Section Editor: Yaron Tomer

Pituitary Hyperplasia During Primary Hypothyroidism

K. Hoogenberg and K.M. van Tol

ELEVATED PROLACTIN LEVELS accompanying primary hypothyroidism may lead physicians to consider the presence of a concomitant pituitary disease, either a microprolactinoma or a sellar mass of any origin with pituitary stalk compression. The fact that successive magnetic resonance imaging may further increase the suspicion of a pituitary disorder is illustrated in this case. It shows enlargement of the pituitary gland in a 51-year-old man with primary hypothyroidism with an elevated thyroid stimulating hormone level (120 mU/L). Serum prolactin (580 mU/L, reference range, 50–200 mU/L) was also increased (Fig. 1A and 1B). After 6 months of levothyroxine substitution, the enlargement of the pituitary gland as well as the elevated serum thyroid stimulating hormone and prolactin levels normalized (Fig. 1C and 1D).

Address reprint requests to:
K. Hoogenberg
Department of Internal Medicine
Division of Endocrinology
Groningen University Hospital
P.O. Box 30.001
9700 RB Groningen
The Netherlands

E-mail: K.Hoogenberg@Int.azg.nl

(See images on next page →)

Department of Internal Medicine, Division of Endocrinology, Groningen University Hospital, sGroningen, The Netherlands.

*If you would like to submit an image for publication in "Images in Thyroidology," please send two copies of high-quality black-and-white images with a short legend. In special circumstances color figures will be published. Please inquire for details from the Section Editor. All material must be original and neither published nor submitted elsewhere. The legend should give relevant clinical information. The entire legend should be typed double-spaced, and should be no more than 200 words. Send all submissions to Yaron Tomer, M.D., Box 1055, Mount Sinai School of Medicine, One Gustave L. Levy Place, New York, NY 10029; E-mail: Yaron.Tomer@mssm.edu

FIG. 1. T1-weighted gadolinium enhanced magnetic resonance imaging of the pituitary gland during hypothyroidism (Fig. 1A and 1B) and 6 months after levothyroxine substitution (Fig. 1C and 1D). The coronal (A) and sagittal (B) images show an enlarged pituitary gland, 12 mm in size, with a normal gadolinium distribution. Furthermore, a small portion of the pituitary is near the optic chiasm. On similarly performed coronal (C) and sagittal (D) images, the pituitary gland enlargement resolved with normalisation of its size and contour.