University of Groningen

Hamilton cycles and perfect matchings in the KPKVB model

Fountoulakis, Nikolaos; Mitsche, Dieter; Müller, Tobias; Schepers, Markus

Published in:
Stochastic processes and their applications

DOI:
[10.1016/j.spa.2020.09.012](https://doi.org/10.1016/j.spa.2020.09.012)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Fountoulakis, N., Mitsche, D., Müller, T., & Schepers, M. (2021). Hamilton cycles and perfect matchings in the KPKVB model. *Stochastic processes and their applications*, 131, 340-358.
<https://doi.org/10.1016/j.spa.2020.09.012>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Hamilton cycles and perfect matchings in the KPKVB model

Nikolaos Fountoulakis^{a,1}, Dieter Mitsche^{b,2}, Tobias Müller^{c,3},
Markus Schepers^{c,*,4}

^a School of Mathematics, University of Birmingham, United Kingdom

^b Institut Camille Jordan, Univ. Lyon, Université Jean Monnet, France

^c Benoulli Institute, Groningen University, The Netherlands

Received 27 January 2019; received in revised form 10 June 2020; accepted 6 September 2020

Available online 17 September 2020

Abstract

In this paper we consider the existence of Hamilton cycles and perfect matchings in a random graph model proposed by Krioukov et al. in 2010. In this model, nodes are chosen randomly inside a disk in the hyperbolic plane and two nodes are connected if they are at most a certain hyperbolic distance from each other. It has been previously shown that this model has various properties associated with complex networks, including a power-law degree distribution, “short distances” and a non-vanishing clustering coefficient. The model is specified using three parameters: the number of nodes n , which we think of as going to infinity, and $\alpha, \nu > 0$, which we think of as constant. Roughly speaking α controls the power law exponent of the degree sequence and ν the average degree.

Here we show that for every $\alpha < 1/2$ and $\nu = \nu(\alpha)$ sufficiently small, the model does not contain a perfect matching with high probability, whereas for every $\alpha < 1/2$ and $\nu = \nu(\alpha)$ sufficiently large, the model contains a Hamilton cycle with high probability.

© 2020 Elsevier B.V. All rights reserved.

Keywords: Hamilton cycle; Perfect matching; Hyperbolic random graph; Poisson process; Randomized algorithm; Tiling

* Corresponding author.

E-mail addresses: n.fountoulakis@bham.ac.uk (N. Fountoulakis), dmitsche@unice.fr (D. Mitsche), tobias.muller@rug.nl (T. Müller), markus.schepers@uni-mainz.de, m.f.schepers@rug.nl (M. Schepers).

¹ Research partially supported by the Alan Turing Institute, United Kingdom, grant no. EP/N510129/1.

² Research partially supported by IDEXLYON of Université de Lyon, France (Programme Investissements d’Avenir ANR16-IDEX-0005) and by Labex MILYON, France/ANR-10-LABX-0070.

³ Research partially supported by NWO, The Netherlands grants 639.032.529 and 612.001.409.

⁴ Research partially supported by NWO, The Netherlands grant 639.032.529.

1. Introduction

A *Hamilton cycle* in a graph is a cycle which contains all vertices of the graph. A graph is called *Hamiltonian* if it contains at least one Hamilton cycle. A *matching* is a set of edges that do not share endpoints and a *perfect matching* is a matching that covers all vertices of the graph.

Hamilton cycles and perfect matchings are classic themes in graph theory. Historically the existence of Hamilton cycles and perfect matchings in a random graph has been a central theme in the theory of random graphs as well. In particular, in the theory of the Erdős–Rényi model the threshold for having a Hamilton cycle as well as the simultaneous emergence in the random graph process of a Hamilton cycle together with having minimum degree at least two are among the classic results in the field [2,7,19,20,26]. In the context of random geometric graphs in the Euclidean plane, analogous results have been obtained [4,9,23]. The emergence of Hamilton cycles was also considered in other models, including the preferential attachment model [13] and the random d -regular graph model [27].

In this paper, we will consider the problem of the existence of a Hamilton cycle and a perfect matching in a model of random graphs that involves points taken randomly in the hyperbolic plane. This model was introduced by Krioukov, Papadopoulos, Kitsak, Vahdat and Boguñá [21] in 2010 — we abbreviate it as *the KPKVB model*. We should however note that the model also goes by several other names in the literature, including *hyperbolic random geometric graphs* and *random hyperbolic graphs*. The model was intended to model complex networks and, in particular, it is motivated by the assumption that the properties of complex networks are the expression of a hidden geometry which expresses the hierarchies among classes of nodes of the network. Krioukov et al. postulate that this geometry is hyperbolic space.

The KPKVB model

Given $\nu \in (0, \infty)$ a fixed constant and a natural number $n > \nu$, we let $R = 2 \log(n/\nu)$, or equivalently $n = \nu \exp(R/2)$. Also, let $\alpha \in (0, \infty)$.

The hyperbolic plane $\mathbb{H}$ is a surface with constant Gaussian curvature -1 . It has several convenient representations (i.e., coordinate maps), including the Poincaré halfplane model, the Poincaré disk model and the Klein disk model. A gentle introduction to hyperbolic geometry and these representations of the hyperbolic plane can for instance be found in [28]. Throughout this paper we will be working with a representation of the hyperbolic plane using *hyperbolic polar coordinates*. That is, a point $p \in \mathbb{H}$ is represented as (r, θ) , where r is the hyperbolic distance between p and the origin O and θ as the angle between the line segment Op and the positive x -axis (Here, when mentioning “the origin” and the angle between the line segment and the positive x -axis, we think of $\mathbb{H}$ embedded as the Poincaré disk in the ordinary Euclidean plane.) We shall denote by $\mathcal{D}_R$ the hyperbolic disk of radius R around the origin O , and by $d_{\mathbb{H}}(u, v)$ we denote the hyperbolic distance between two points $u, v \in \mathbb{H}$.

The vertex set of the KPKVB random graph $G(n; \alpha, \nu)$ consists of n i.i.d. points in $\mathcal{D}_R$ with probability density function

$$f_{n,\alpha,\nu}(r, \theta) = \frac{\alpha \sinh \alpha r}{2\pi(\cosh \alpha R - 1)}, \quad (1)$$

for $0 \leq r < R$ and $0 < \theta \leq 2\pi$. Note that this is a joint probability density function. For the probability density function of the radial coordinate only, we use the notation

$$g_{n,\alpha,\nu}(r) = \frac{\alpha \sinh \alpha r}{\cosh \alpha R - 1}, \quad (2)$$

for $0 \leq r < R$.

When $\alpha = 1$ the distribution of (r, θ) given by (1) is the uniform distribution on $\mathcal{D}_R$. For general $\alpha \in (0, \infty)$ Krioukov et al. [21] call this the *quasi-uniform* distribution on $\mathcal{D}_R$. In fact, for general α it can be viewed as the uniform distribution on a disk of radius R on the hyperbolic plane with curvature $-\alpha^2$.

The KPKVB random graph $G(n; \alpha, \nu)$ is the random graph whose vertex set is a set V_n of n points of chosen i.i.d. according to the (α, R) -quasi uniform distribution, where any two of them are joined by an edge if they are within hyperbolic distance at most R .

Krioukov et al. [21] observed that the distribution of the degrees in $G(n; \alpha, \nu)$ follows a power law with exponent $2\alpha + 1$, for $\alpha \in (1/2, \infty)$. This was verified rigorously by Gugelmann et al. in [15]. Note that for $\alpha \in (1/2, 1)$, this quantity is between 2 and 3, which is in line with numerous observations in networks which arise in applications (see for example [3]). In addition, Krioukov et al. observed, and Gugelmann et al. proved rigorously, that the (local) clustering coefficient of the graph stays bounded away from zero a.s. Here and in the rest of the paper we use the following notation: for a sequence of events $(E_n)_{n \in \mathbb{N}}$, we say that E_n occurs *asymptotically almost surely* (a.s.), if $\mathbb{P}(E_n) \rightarrow 1$ as $n \rightarrow \infty$. Also, at various places we use the shorthand *w.p.* to write *with probability*.

Krioukov et al. [21] observed also that the average degree of $G(n; \alpha, \nu)$ is determined via the parameter ν for $\alpha \in (1/2, \infty)$. This was rigorously verified in [15] too. In particular, they proved that the average degree tends to $2\alpha^2\nu/\pi(\alpha - \frac{1}{2})^2$ in probability.

In [5], it was established that $\alpha = 1$ is the critical point for the emergence of a giant component in $G(n; \alpha, \nu)$. In particular, when $\alpha \in (0, 1)$, the fraction of the vertices contained in the largest component is bounded away from 0 a.s., whereas if $\alpha \in (1, \infty)$, the largest component is sublinear in n a.s. For $\alpha = 1$, the component structure depends on ν . If ν is large enough, then a giant component exists a.s., but if ν is small enough, then a.s. all components are sublinear [5].

In [11] this picture is sharpened. There, the first and the third author showed that the fraction of vertices belonging to the largest component converges in probability to a constant which depends on α and ν . For $\alpha = 1$, the existence of a critical value $\nu_0 \in (0, \infty)$ is established such that when ν crosses ν_0 a giant component emerges a.s. In [16] and [18], the second author together with Kiwi considered the size of the second largest component and showed that when $\alpha \in (1/2, 1)$, a.s., the second largest component has polylogarithmic order with exponent $1/(\alpha - 1/2)$.

Apart from the degree sequence, clustering and component sizes, the graph distances in this model have also been considered. In [16] and [12] a.s. polylogarithmic upper and lower bounds on the diameter of the largest component are shown, and in [24], these were sharpened to show that $\log n$ is the correct order of the diameter. Furthermore, in [1] it is shown that for $\alpha \in (1/2, 1)$ the largest component is what is called an *ultra-small world*: it exhibits doubly logarithmic typical distances.

Results on the global clustering coefficient were obtained in [8], and on the evolution of graphs on more general spaces with negative curvature in [10]. The spectral gap of the Laplacian of this model was studied in [17].

The first and third author together with Bode [6], showed that $\alpha = 1/2$ is the critical value for connectivity: that is, when $\alpha \in (0, 1/2)$, then $G(n; \alpha, \nu)$ is a.a.s. connected, whereas $G(n; \alpha, \nu)$ is a.a.s. disconnected when $\alpha \in (1/2, \infty)$. The second half of this statement is in fact already immediate from the results of Gugelmann et al. [14]: there it is shown that for $\alpha > 1/2$, a.a.s., there are linearly many isolated vertices. For $\alpha = 1/2$, the probability of connectivity tends to a limiting value that is function of ν that is continuous and non-decreasing and that equals one if and only if $\nu \geq \pi$.

Our results

In the present paper, we explore the existence of Hamilton cycles and perfect matchings in the $G(n; \alpha, \nu)$ model. In the light of the result on isolated vertices mentioned above, the question is non-trivial only for $\alpha \leq 1/2$. A perfect matching trivially cannot exist when n is odd. For this reason we find it convenient to switch to considering *near perfect matchings* from now on. That is, matchings that cover all but at most one vertex. (So if n is even a near perfect matching is the same as a perfect matching; and the existence of a Hamilton cycle implies the existence of a near perfect matching.)

Our main result shows that in the regime $\alpha < 1/2$, a.a.s., the existence of a Hamilton cycle as well as of a (near) perfect matching has a non-trivial phase transition in ν .

Theorem 1.1. *For all positive real $\alpha < \frac{1}{2}$, there are constants $\nu_0 = \nu_0(\alpha)$ and $\nu_1 = \nu_1(\alpha)$ such that the following hold. For all $0 < \nu < \nu_0$, the random graph $G(n; \alpha, \nu)$ a.a.s. does not have a near perfect matching (and consequently no Hamilton cycle either). For all $\nu > \nu_1$, $G(n; \alpha, \nu)$ a.a.s. has a Hamilton cycle.*

To our knowledge, this is the first time this problem is considered for the $G(n; \alpha, \nu)$ model. We conjecture that the dependence on ν is sharp.

Conjecture 1.2. *For every $0 < \alpha < 1/2$ there exists a critical $\nu_c = \nu_c(\alpha) > 0$ such that when $\nu < \nu_c$ a.a.s. $G(n; \alpha, \nu)$ has no near perfect matching, whereas if $\nu > \nu_c$ then a.a.s. $G(n; \alpha, \nu)$ has a Hamilton cycle.*

A natural question to ask is what happens in the case $\alpha = 1/2$. Does there exist ν large enough so that the graph a.a.s. becomes Hamiltonian in this case as well?

It would also be interesting to explore the relation of Hamiltonicity with the property of 2-connectivity. If the above conjecture is true, is there a similar behavior for the property of 2-connectivity? If yes, are the corresponding critical constants ν_c equal?

Outline of proof. The proof of Theorem 1.1 has two parts: in a nutshell, in the first part we show that for ν small enough, the number of vertices close to the boundary of the disk of radius R having no neighbor close to the boundary of the disk will be bigger than the total number of vertices relatively close to the center of the disk. Hence, the former vertices would have to be all matched to distinct vertices close to the center of the disk, but that cannot happen. For the second part, we show that for ν large enough, we can tessellate the disk in such a way, so that iteratively, from the boundary towards the center of the disk, we can maintain a set of vertex-disjoint cycles and isolated points, which will eventually be merged close to the center. The fact that ν is large enough makes the density of vertices in each cell of the tessellation high enough so that this procedure terminates successfully.

2. Preliminaries

2.1. Probabilistic preliminaries

To prove [Theorem 1.1](#), we will perform our analysis in the *Poissonization* of the $G(n; \alpha, \nu)$ model. There, the vertex set is the point set of a Poisson point process on $\mathcal{D}_R$ with n points on average. Although the independence that accompanies the Poisson point process facilitates our proofs, when doing standard de-Poissonization, we need to show a slightly stronger version of [Theorem 1.1](#). We give details here.

We denote the Poissonized version of the KPKVB model by $G_{\text{Po}}(n; \alpha, \nu)$. The vertex set of this random graph is the set of points of the Poisson point process $\mathcal{P}_{n,\alpha,\nu}$ on $\mathcal{D}_R$ with intensity $n \cdot \frac{1}{2\pi} g_{n,\alpha,\nu}$. The set of edges of $G_{\text{Po}}(n; \alpha, \nu)$ consists of those pairs of points of $\mathcal{P}_{n,\alpha,\nu}$ which are at hyperbolic distance at most R . Alternatively, the Poissonized KPKVB model $G_{\text{Po}}(n; \alpha, \nu)$ can be constructed as follows. Consider an infinite supply of i.i.d. points $p_1, p_2, \dots$, chosen according to the (α, R) -quasi uniform distribution, and a Poisson random variable $Z \stackrel{\Delta}{=} \text{Po}(n)$. The vertex set of G_{Po} is now the set of points $p_1, \dots, p_Z$ and again we add edges between pairs at hyperbolic distance at most R .

The function $n \cdot \frac{1}{2\pi} g_{n,\alpha,\nu}$ is the *intensity measure* associated with $\mathcal{P}_{n,\alpha,\nu}$. This means in particular that for any Borel subset $A \subseteq \mathcal{D}_R$ the expected number of points that fall in A equals

$$\mu_{n,\alpha,\nu}(A) = n \cdot \frac{1}{2\pi} \int_A g_{n,\alpha,\nu}(r) dr d\theta.$$

We set $\mathcal{P}_{n,\alpha,\nu}(A) := \mathcal{P}_{n,\alpha,\nu} \cap A$; hence $|\mathcal{P}_{n,\alpha,\nu}(A)| \stackrel{\Delta}{=} \text{Po}(\mu_{n,\alpha,\nu}(A))$. An elementary, but key, observation is that conditional on $|\mathcal{P}_{n,\alpha,\nu}| = n$, the process $\mathcal{P}_{n,\alpha,\nu}$ is distributed as V_n . In other words, the probability space of the process V_n can be realized as the space of $\mathcal{P}_{n,\alpha,\nu}$ conditional on $|\mathcal{P}_{n,\alpha,\nu}| = n$.

The following observation is well known. We include its proof here for completeness.

Lemma 2.1. *Let $\mathcal{F}$ be a graph property (formally a family of graphs closed under isomorphism). We then have that*

$$\mathbb{P}(G_{\text{Po}}(n; \alpha, \nu) \in \mathcal{F} \mid |\mathcal{P}_{n,\alpha,\nu}| = n) = 1 - o(1), \text{ if } \mathbb{P}(G_{\text{Po}}(n; \alpha, \nu) \in \mathcal{F}) = 1 - o(n^{-1/2}).$$

Thus, if $\mathbb{P}(G_{\text{Po}}(n; \alpha, \nu) \notin \mathcal{F}) = o(n^{-1/2})$, then $\mathbb{P}(G(n; \alpha, \nu) \notin \mathcal{F}) = o(1)$.

Proof. By Stirling's formula

$$\mathbb{P}(|\mathcal{P}_{n,\alpha,\nu}| = n) = \frac{n^n}{n!} e^{-n} = (1 + o(1)) \frac{n^n}{\sqrt{2\pi n^{n+\frac{1}{2}}} e^{-n}} e^{-n} = (1 + o(1)) \frac{1}{\sqrt{2\pi n}}$$

So as $n \rightarrow \infty$, writing $E_n := \{G_{\text{Po}}(n; \alpha, \nu) \in \mathcal{F}\}$:

$$\mathbb{P}(E_n^c) \geq \mathbb{P}(E_n^c \mid |\mathcal{P}_{n,\alpha,\nu}| = n) \cdot \Theta(n^{-1/2}).$$

Therefore, if $\mathbb{P}(E_n^c) = o(n^{-1/2})$, we deduce that $\mathbb{P}(E_n^c \mid |\mathcal{P}_{n,\alpha,\nu}| = n) = o(1)$. $\square$

We then apply a standard Depoissonization technique: using [Lemma 2.1](#), we will develop our arguments in the Poissonization of the KPKVB model. More precisely, we will show that $G_{\text{Po}}(n; \alpha, \nu)$ satisfies certain events with sufficiently high probability (that is, with probability

at least $1 - o(n^{-1/2})$, and we then use [Lemma 2.1](#) to deduce that $G(n; \alpha, \nu)$ also satisfies them a.s., that is, with probability $1 - o(1)$.

Another useful tool that will allow us to compute expectations of sums over the points of $\mathcal{P}_{n,\alpha,\nu}$ is the *Campbell–Mecke* formula. Let $\mathcal{P}$ be a point process on a metric space $\mathcal{S}$ with density ρ . Let $\mathcal{N} = \mathcal{N}(\mathcal{S})$ be the set of all countable point configurations in $\mathcal{S}$ equipped with the σ -algebra of the point process (generated by the following sets: for any open subset $A \subseteq \mathcal{S}$ and any non-negative integer m consider the subset of $\mathcal{N}$ which consists of all configurations which have exactly m points in A). Now, let $h : \mathbb{R}^k \times \mathcal{N} \rightarrow \mathbb{R}$ be a measurable function. The Palm theory of Poisson point processes on metric spaces [\[22\]](#) yields:

$$\mathbb{E} \left(\sum_{\substack{p_1, \dots, p_k \in \mathcal{P}, \\ \text{distinct}}} h(p_1, \dots, p_k, \mathcal{P}) \right) = \int_{\mathcal{S}} \cdots \int_{\mathcal{S}} \mathbb{E}(h(x_1, \dots, x_k, \mathcal{P})) d\rho(x_1) \cdots d\rho(x_k), \quad (3)$$

where the sum ranges over all those k -tuples of points which contain no repetitions. We also use the following form of Chernoff's bound:

Lemma 2.2. *For a Poisson random variable X with expectation μ , and a positive integer $k \leq \mu$,*

$$\mathbb{P}(X \leq k) \leq e^{-\mu H(\frac{k}{\mu})},$$

where $H(a) = 1 - a + a \ln a$ for $a > 0$; in particular if $\frac{k}{\mu} \leq \frac{1}{2}$ it holds that $H(\frac{k}{\mu}) \geq \frac{1}{2}(1 - \ln 2) > 0$.

A proof can for instance be found in [\[25\]](#).

2.2. Geometric preliminaries

For two points $p_1 = (r_1, \theta_1)$ and $p_2 = (r_2, \theta_2)$ in $\mathcal{D}_R$, we let $\theta(p_1, p_2)$ be their *angular distance* which we define as:

$$\theta(p_1, p_2) = \min\{|\theta_1 - \theta_2|, 2\pi - |\theta_1 - \theta_2|\}.$$

Note that $\theta(p_1, p_2) \in [0, \pi]$.

The hyperbolic law of cosines relates the angular distance $\theta(p_1, p_2)$ between p_1, p_2 with their hyperbolic distance:

$$\cosh(d_{\mathbb{H}}(p_1, p_2)) = \cosh(r_1) \cosh(r_2) - \sinh(r_1) \sinh(r_2) \cos(\theta(p_1, p_2)). \quad (4)$$

Now, for $r_1, r_2 \in [0, R)$, we let $\theta_R(r_1, r_2) \in [0, \pi]$ be such that

$$\cosh(R) = \cosh(r_1) \cosh(r_2) - \sinh(r_1) \sinh(r_2) \cos(\theta_R(r_1, r_2)).$$

For two points p_1, p_2 of radii r_1 and r_2 , respectively, such that $\theta(p_1, p_2) = \theta_R(r_1, r_2)$, we have $d_{\mathbb{H}}(p_1, p_2) = R$. Eq. (4) implies that $d_{\mathbb{H}}(p_1, p_2) \leq R$ if and only if $\theta(p_1, p_2) \leq \theta_R(r(p_1), r(p_2))$.

The ball $B(p)$ of radius R around p inside $\mathcal{D}_R$ is thus defined as

$$B(p) := \{p' \in \mathcal{D}_R : \theta(p, p') \leq \theta_R(r(p), r(p'))\}.$$

If p is a vertex either in the vertex set of $G(n; \alpha, \nu)$ or in the vertex set of $G_{\text{Po}}(n; \alpha, \nu)$, then it is adjacent precisely to any other vertex that belongs to $B(p)$.

It will be convenient for our analysis to express $\theta_R(r_1, r_2)$ explicitly as a function of r_1, r_2 . We will make use of the following lemma from [\[11\]](#), which does this.

Lemma 2.3 ([11], Lemma 28). *There exists a constant $K > 0$ such that for every $\epsilon > 0$ and R sufficiently large, the following holds: for every $r_1, r_2 \in [\epsilon R, R]$ with $r_1 + r_2 > R$, we have*

$$2e^{\frac{1}{2}(R-r_1-r_2)} - Ke^{\frac{3}{2}(R-r_1-r_2)} \leq \theta_R(r_1, r_2) \leq 2e^{\frac{1}{2}(R-r_1-r_2)} + Ke^{\frac{3}{2}(R-r_1-r_2)}.$$

Moreover, if $r_1, r_2 < R - K$, then $\theta_R(r_1, r_2) \geq 2e^{\frac{R-r_1-r_2}{2}}$.

3. Non-existence of perfect matching for sufficiently small ν

The following theorem yields the first part of Theorem 1.1.

Theorem 3.1. *For all positive real $\alpha < \frac{1}{2}$, there is a $\nu_0 = \nu_0(\alpha) > 0$ such that for all $0 < \nu < \nu_0$, the random graph $G_{Po}(n; \alpha, \nu)$ does not have a near perfect matching w.p. $1 - o(n^{-1/2})$.*

Proof. The strategy is as follows. Let $s = \frac{1}{\alpha} > 2$. Let N_s be number of vertices with radial coordinate at least $R - s$ and with no neighbor with radial coordinate at least $R - s$. Let M_s be the number of vertices with radial coordinate at most $R - s$. Hence, M_s is the number of points of $\mathcal{P}_{n,\alpha,\nu}$ inside the disk of radius $R - s$ and N_s is a subset of the annulus $\mathcal{A}_s = \mathcal{D}_R \setminus \mathcal{D}_{R-s}$ of width s . If there is a perfect matching, then $M_s \geq N_s$ because a vertex with no neighbor with radius at least $R - s$ must be matched to a vertex with radius less than $R - s$, so distinct vertices counted by N_s must be matched to distinct vertices counted by M_s . If we can show that M_s and N_s are concentrated around their expectation w.p. $1 - o(n^{-1/2})$ and that $\mathbb{E}M_s = (1 + o(1))c_{M_s}n$ and $\mathbb{E}N_s = (1 + o(1))c_{N_s}n$ as $n \rightarrow \infty$ with $c_{M_s} < c_{N_s}$, then there will be no near perfect matching and hence no Hamilton cycle w.p. $1 - o(n^{-1/2})$.

We observe that $M_s \stackrel{\Delta}{=} \text{Po}(\mu_{n,\alpha,\nu}(\mathcal{D}_{R-s}))$, where

$$\mu_{n,\alpha,\nu}(\mathcal{D}_{R-s}) = n \cdot \frac{\cosh \alpha(R-s) - 1}{\cosh \alpha R - 1} \cdot \frac{1}{2\pi} \int_0^{2\pi} d\theta \sim n \cdot \frac{\frac{1}{2}e^{\alpha(R-s)}}{\frac{1}{2}e^{\alpha R}} = ne^{-\alpha s} = ne^{-1}.$$

Since M_s is Poisson distributed, we deduce that $\mathbb{E}M_s = \text{Var } M_s \sim ne^{-1}$. (Here and elsewhere we write $a_n \sim b_n$ to denote that $a_n/b_n = 1 + o(1)$.) By Chebyshev's inequality, it follows that for all $\epsilon > 0$,

$$\mathbb{P}(|M_s - \mathbb{E}M_s| \geq \epsilon \mathbb{E}M_s) \leq \frac{\text{Var}(M_s)}{\epsilon^2 (\mathbb{E}M_s)^2} = O(n^{-1}) = o(n^{-1/2}).$$

Our goal is now to give a lower bound on $\mathbb{E}N_s$ and an upper bound on the variance of $\text{Var}(N_s)$ (more precisely, $\text{Var}(N_s)/\mathbb{E}^2 N_s = o(n^{-1/2})$), which together will imply concentration of N_s around $\mathbb{E}N_s$. More precisely, we will show that $\text{Var}(N_s)/\mathbb{E}^2 N_s = o(n^{-1/2})$, which implies concentration of N_s around $\mathbb{E}N_s$ (and we will lower bound $\mathbb{E}N_s$).

For a point $p \in \mathcal{D}_R$ we let $\lambda(p, \mathcal{P}_{n,\alpha,\nu})$ denote the indicator random variable which is equal to 1 if and only if $\mathcal{P}_{n,\alpha,\nu}(\check{B}_s(p)) := \mathcal{P}_{n,\alpha,\nu}((B(p) \setminus \{p\}) \cap \mathcal{A}_s) = \emptyset$.

In other words, $\lambda(p, \mathcal{P}_{n,\alpha,\nu})$ is equal to 1 if and only if no point of $\mathcal{P}_{n,\alpha,\nu} \setminus \{p\}$ is contained in $B(p) \cap \mathcal{A}_s$.

We can write

$$N_s = \sum_{p \in \mathcal{P}_{n,\alpha,\nu} \cap \mathcal{A}_s} \lambda(p, \mathcal{P}_{n,\alpha,\nu}).$$

By the Campbell–Mecke formula (3), we have

$$\begin{aligned}\mathbb{E}N_s &= n \cdot \frac{1}{2\pi} \int_{\mathcal{A}_s} \mathbb{E}\lambda((r, \theta), \mathcal{P}_{n,\alpha,v}) g_{n,\alpha,v}(r) dr d\theta \\ &= n \cdot \frac{1}{2\pi} \int_{R-s}^R \int_0^{2\pi} \mathbb{E}\lambda((r, \theta), \mathcal{P}_{n,\alpha,v}) g_{n,\alpha,v}(r) dr d\theta \\ &= n \cdot \int_{R-s}^R \mathbb{E}\lambda((r, 0), \mathcal{P}_{n,\alpha,v}) g_{n,\alpha,v}(r) dr,\end{aligned}\tag{5}$$

where the first equality holds since $\lambda((r, \theta), \mathcal{P}_{n,\alpha,v})$ is identically zero if $r < R - s$, and the last one since $\mathbb{E}\lambda((r, \theta), \mathcal{P}_{n,\alpha,v})$ and $g_{n,\alpha,v}(r)$ are invariant with respect to θ .

We have

$$\check{B}_s((r, 0)) = \{(r', \theta') \in \mathbb{R}^2 : R - s \leq r' < R, |\theta'| \leq \theta_R(r, r')\},$$

and, therefore,

$$\mu_{n,\alpha,v}(\check{B}_s((r, 0))) = n \cdot \int_{R-s}^R \frac{2\theta_R(r, r')}{2\pi} \frac{\alpha \sinh \alpha r'}{\cosh \alpha R - 1} dr'.$$

We can give an asymptotic approximation to this integrand. From Lemma 2.3, we infer that for n large enough, uniformly over all $r, r' \geq R - s$:

$$\theta_R(r, r') = 2e^{\frac{R-r-r'}{2}} (1 + O(e^{R-r-r'})) = 2e^{\frac{R-r-r'}{2}} (1 + O(e^{-R})),$$

and

$$\frac{\sinh \alpha r'}{\cosh \alpha R - 1} \sim e^{-\alpha(R-r')}.$$

Therefore,

$$\begin{aligned}\mu_{n,\alpha,v}(\check{B}_s((r, 0))) &\sim n \cdot e^{-r/2} \frac{\alpha}{\pi} \int_{R-s}^R e^{(1/2-\alpha)(R-r')} dr' = n \cdot e^{-r/2} \frac{\alpha}{\pi} \int_0^s e^{(1/2-\alpha)y} dy \\ &\stackrel{n=ve^{R/2}}{=} v \cdot e^{(R-r)/2} \frac{\alpha}{\pi(1/2-\alpha)} (e^{(1/2-\alpha)s} - 1).\end{aligned}$$

We can bound from above this expression by using $R - r \leq s$ and then plugging in $s = \frac{1}{\alpha}$, yielding the upper bound

$$ve^{s/2} \frac{\alpha}{\pi(1/2-\alpha)} (e^{1/(2\alpha)-1} - 1) =: vc_\alpha.$$

Note that $R - s \leq r$ also implies that for any point $p = (r, \theta)$ with $r \geq R - s$, for n large enough, $\mu_{n,\alpha,v}(\check{B}_s(p)) \leq vc_\alpha$. So for any such n

$$\mathbb{E}\lambda((r, 0), \mathcal{P}_{n,\alpha,v}) = \mathbb{P}(\text{Po}(\mu_{n,\alpha,v}(\check{B}_s(p))) = 0) \geq e^{-vc_\alpha}.$$

If we substitute this into (5), we get the following lower bound:

$$\begin{aligned}\mathbb{E}N_s &\geq ne^{-vc_\alpha} \cdot \int_{R-s}^R g_{n,\alpha,v}(r) dr \\ &= ne^{-vc_\alpha} \cdot \frac{\cosh(\alpha(R)) - \cosh(\alpha(R-s))}{\cosh(\alpha R) - 1} \sim ne^{-vc_\alpha} (1 - e^{-1}).\end{aligned}$$

Since $e > 2$, we have that $e^{-1} < 1 - e^{-1}$. Choose $\nu > 0$ sufficiently small so that

$$e^{-1} < e^{-\nu c_\alpha}(1 - e^{-1}).$$

Thereafter, we choose $\epsilon > 0$ sufficiently small so that

$$e^{-1}(1 + \epsilon) < (1 - \epsilon)e^{-\nu c_\alpha}(1 - e^{-1}).$$

We will use Chebyshev's inequality to show that w.p. $1 - o(n^{-1/2})$

$$N_s \geq (1 - \epsilon)\mathbb{E}N_s.$$

Since $M_s < ne^{-1}(1 + \epsilon)$ w.p. $1 - o(n^{-1/2})$, the union bound implies $M_s < N_s$ w.p. $1 - o(n^{-1/2})$.

We will show that $\text{Var}(N_s) = O(n)$. To bound the variance, let us set

$$\text{Cov}(p_1, p_2) := \mathbb{E}(\lambda(p_1, \mathcal{P}_{n,\alpha,\nu})\lambda(p_2, \mathcal{P}_{n,\alpha,\nu})) - \mathbb{E}(\lambda(p_1, \mathcal{P}_{n,\alpha,\nu}))\mathbb{E}(\lambda(p_2, \mathcal{P}_{n,\alpha,\nu})).$$

We write

$$\text{Var}(N_s) = \mathbb{E} \left(\sum_{\substack{p_1, p_2 \in \mathcal{P}_{n,\alpha,\nu}, \\ \text{distinct}}} \text{Cov}(p_1, p_2) \right) + \mathbb{E} \left(\sum_{p \in \mathcal{P}_{n,\alpha,\nu}} \text{Cov}(p, p) \right).$$

We will use the Campbell–Mecke formula (3) to calculate these sums. For the former one, we have

$$\begin{aligned} & \mathbb{E} \left(\sum_{\substack{p_1, p_2 \in \mathcal{P}_{n,\alpha,\nu}, \\ \text{distinct}}} \text{Cov}(p_1, p_2) \right) \\ &= n^2 \cdot \left(\frac{1}{2\pi} \right)^2 \int_{R-s}^R \int_0^{2\pi} \int_{R-s}^R \int_0^{2\pi} \mathbb{E}(\text{Cov}((r_1, \theta_1), (r_2, \theta_2))) g_{n,\alpha,\nu}(r_1) \\ & \quad \times g_{n,\alpha,\nu}(r_2) dr_1 d\theta_1 dr_2 d\theta_2. \end{aligned}$$

Let $p_1, p_2 \in \mathcal{A}_s$. We claim that for n sufficiently large, $\check{B}_s(p_1) \cap \check{B}_s(p_2) = \emptyset$, if $\theta(p_1, p_2) > 5ve^s/n$. Indeed, Lemma 2.3 implies that for any $r, r' > R-s$, we have $\theta_R(r, r') < 2e^{\frac{1}{2}(R-r-r')}(1 + O(e^{\frac{3}{2}(R-r-r')})) < 2e^{s-R/2}(1 + O(e^{-\frac{3}{2}R})) < 2.5e^{s-R/2}$, for n sufficiently large. Thus, for any such n , if $p \in \check{B}_s(p_1) \cap \check{B}_s(p_2)$, then $\theta(p, p_1), \theta(p, p_2) < 2.5e^{s-R/2}$. This would imply that $\theta(p_1, p_2) < 5e^{s-R/2}$. So if $\theta(p_1, p_2) > 5e^{s-R/2}$, then $\check{B}_s(p_1) \cap \check{B}_s(p_2) = \emptyset$.

This implies that for n sufficiently large, if $\theta(p_1, p_2) > 5ve^s/n$, we have $\mathbb{E}(\text{Cov}(p_1, p_2)) = 0$. Moreover, recalling that the expected number of vertices in an angle at most $5ve^s/n$ around a fixed point is $O(1)$, we get

$$\mathbb{E} \left(\sum_{\substack{p_1, p_2 \in \mathcal{P}_{n,\alpha,\nu}, \\ \text{distinct}}} \text{Cov}(p_1, p_2) \right) = O(n).$$

Regarding the second term, we use that $\mathbb{E} \text{Cov}(p, p) \leq \mathbb{E} \lambda(p, \mathcal{P}_{n,\alpha,\nu})$ and bound

$$\mathbb{E} \left(\sum_{p \in \mathcal{P}_{n,\alpha,\nu}} \text{Cov}(p, p) \right) \leq \mathbb{E} \left(\sum_{p \in \mathcal{P}_{n,\alpha,\nu}} \mathbb{E} \lambda(p, \mathcal{P}_{n,\alpha,\nu}) \right) = \mathbb{E}(N_s) = O(n).$$

Putting both sums of the covariance together we obtain

$$\text{Var}(N_s) = O(n).$$

Chebyshev's inequality yields

$$\mathbb{P}(N_s \geq (1 - \epsilon)\mathbb{E}N_s) \leq \frac{\text{Var}(N_s)}{\epsilon^2 \mathbb{E}^2(N_s)} = O\left(\frac{1}{n}\right) = o(n^{-1/2}),$$

and Theorem 3.1 follows. $\square$

4. Existence of Hamilton cycles for sufficiently large ν

The aim of this section is to prove the existence of a Hamilton cycle in $G_{P_0}(n; \alpha, \nu)$ when ν is large enough with sufficiently high probability.

Theorem 4.1. *For all positive real $\alpha < \frac{1}{2}$, there is a $\nu_1 = \nu_1(\alpha)$ such that for all $\nu > \nu_1$, the random graph $G_{P_0}(n; \alpha, \nu)$ has a Hamilton cycle and hence also a near perfect matching with probability $1 - o(n^{-1/2})$.*

4.1. A useful tiling

We consider the following tiling:

$$T_{i,j} = \{(r, \theta) \in \mathcal{D}_R : R - (i+1)2\ln 2 \leq r < R - i2\ln 2, j\frac{2\pi}{n_i} < \theta \leq (j+1)\frac{2\pi}{n_i}\},$$

where $n_i = n_{i,R} = 2^{4-i+\lfloor \frac{R}{2\ln 2} \rfloor} \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}$, for $i \in \mathbb{N}_0$, $i \leq i_{\max} = \lceil \frac{0.9R}{2\ln 2} \rceil$ and $j \in \mathbb{N}_0$, $j < n_i$. We call i, j *admissible* if they satisfy these constraints. Note that for all admissible i , the parameter n_i is an integer and, in fact, a power of 2, as the exponent $4 - i + \lfloor \frac{R}{2\ln 2} \rfloor$ is an integer. Moreover, for $i \leq i_{\max}$ the exponent is also at least $4 - \frac{0.9R}{2\ln 2} + \frac{R}{2\ln 2} - 1 = 3 + 0.1\frac{R}{2\ln 2} > 0$.

We call the collection of tiles with a fixed given i the i th layer. These are the tiles in the i th annulus where we start counting from zero at the boundary of the disk. Note that there are n_i tiles in the i th layer and the tiling covers the annulus with exterior radius R and interior radius $R - i_{\max}2\ln 2 = (1 + o(1))0.1R$ (in particular, the most interior layer $i_{\max}$ is contained in the smaller disk with radius $\frac{R}{2}$ around the origin). A schematic picture is shown in Fig. 1.

We say that a tile $T_{i',j'}$ is *below* the tile $T_{i,j}$, if $i' \leq i$ and the sector defined by $T_{i,j}$ contains $T_{i',j'}$.

Lemma 4.2 (Adjacency among the Tiles). *For admissible indices i, j , any point $p \in T_{i,j}$ is within distance R from any point p' in any tile below the tile $T_{i,j}$.*

Proof. Let $p = (r, \theta) \in T_{i,j}$ and $p' = (r', \theta') \in T_{i',j'}$ be a vertex in any tile below tile $T_{i,j}$ (in the sense of the statement above). Note that $r' \geq r$ must hold. Then, the angular distance $|\theta - \theta'|_{2\pi}$ between p and p' is at most the angular width of the tile $T_{i,j}$, which is

$$\frac{2\pi}{n_i} = 2^{-3}\pi 2^{i-\lfloor \frac{R}{2\ln 2} \rfloor} \leq 2^{i-1}e^{-\frac{R}{2}}.$$

On the other hand, we know that the radial coordinates satisfy $r \leq R - i2\ln 2$ and $r' < R$. If $r + r' \leq R$, we have adjacency by the triangle inequality. If $r + r' > R$ and using that $r, r' \geq (1 + o(1))0.1R$ (as remarked earlier), we distinguish two cases:

1. If $r, r' < R - K$ (with K as in Lemma 2.3), then by the last part of Lemma 2.3, it holds that

$$\theta_R(r, r') \geq 2e^{\frac{R-r-r'}{2}} \geq 2e^{\frac{i2\ln 2 - R}{2}} = 2^{i+1}e^{-\frac{R}{2}}.$$

Fig. 1. (Partial) tiling in the hyperbolic disk; example of a tile $T_{i,j}$ (colored black) and the tiles which are guaranteed to lie within its neighborhood ball by Lemma 4.2 (colored black and gray).

2. Otherwise we may assume that $r' \geq R - K$ holds, while still $r, r' \geq (1 + o(1))0.1R$. Therefore, $R - r - r' \leq R - (R - K) - (1 + o(1))0.1R = -(1 + o(1))0.1R + K$, hence the error term in Lemma 2.3 is $Ke^{\frac{3}{2}(R-r-r')} = o(e^{\frac{1}{2}(R-r-r')})$ and it follows that

$$\theta_R(r, r') \geq 2e^{\frac{R-r-r'}{2}} - o(e^{\frac{1}{2}(R-r-r')}) > e^{\frac{R-r-r'}{2}} \geq e^{\frac{i2\ln 2 - R}{2}} = 2^i e^{-\frac{R}{2}}.$$

We conclude that $|\theta - \theta'|_{2\pi} \leq \theta_R(r, r')$ from which the claim follows. $\square$

We will denote by $N(T_{i,j})$ the number of points falling into $T_{i,j}$.

Lemma 4.3 (Expected Number of Points in a Tile). *Let $\alpha, v > 0$. For admissible indices i, j , the expected number of points falling into $T_{i,j}$ satisfies:*

$$\mathbb{E}N(T_{i,j}) = \Theta(v2^{i(1-2\alpha)}).$$

Proof. The expected number of points falling into $T_{i,j}$ is given by

$$\begin{aligned} \mu_{n,\alpha,v}(T_{i,j}) &= n \cdot \int_{R-(i+1)2\ln 2}^{R-i2\ln 2} \int_{j\frac{2\pi}{n_i}}^{(j+1)\frac{2\pi}{n_i}} \frac{\alpha \sinh \alpha r}{2\pi(\cosh \alpha R - 1)} d\theta dr \\ &= n \cdot \int_{R-(i+1)2\ln 2}^{R-i2\ln 2} \frac{\alpha \sinh \alpha r}{n_i(\cosh \alpha R - 1)} dr \\ &= n \cdot \frac{\cosh(\alpha(R - i2\ln 2)) - \cosh(\alpha(R - (i+1)2\ln 2))}{n_i(\cosh \alpha R - 1)}. \end{aligned}$$

Since $i \leq i_{\max}$, we have that $R - i2 \ln 2 \geq 0.1R \rightarrow \infty$, and hence

$$\cosh(\alpha(R - i2 \ln 2)) = (1 + o(1)) \frac{1}{2} e^{\alpha(R - i2 \ln 2)},$$

$$\cosh(\alpha(R - (i + 1)2 \ln 2)) = (1 + o(1)) \frac{1}{2} e^{\alpha(R - (i + 1)2 \ln 2)}$$

and

$$\cosh \alpha R = (1 + o(1)) \frac{1}{2} e^{\alpha R}.$$

Furthermore, $n_i = 2^{4-i+\lfloor \frac{R}{2 \ln 2} \rfloor} = \Theta(2^{-i+\frac{R}{2 \ln 2}}) = \Theta(2^{-i} e^{\frac{R}{2}})$. Therefore,

$$\begin{aligned} \mathbb{E}N(T_{i,j}) &= \Theta \left(n \frac{e^{\alpha(R - i2 \ln 2)} - e^{\alpha(R - (i+1)2 \ln 2)}}{2^{-i} e^{\frac{R}{2}} e^{\alpha R}} \right) = \Theta(n 2^i e^{-\frac{R}{2}} e^{-i2\alpha \ln 2} (1 - e^{-2\alpha \ln 2})) \\ &= \Theta(n 2^{i(1-2\alpha)} e^{-\frac{R}{2}}). \end{aligned}$$

Finally, using that $R = 2 \ln \frac{n}{v}$, that is, $n = v e^{\frac{R}{2}}$, yields the claim. $\square$

4.2. A procedure for finding a Hamilton cycle

In this subsection we describe the strategy of our procedure for finding a Hamilton cycle in a graph which is embedded in the hyperbolic disk $\mathcal{D}_R$ and which makes use of the tiling $(T_{i,j})_{0 \leq i \leq i_{\max}, 0 \leq j < n_i}$ defined above. Roughly speaking, the procedure iterates through the layers of the tiling, working upwards from the 0th layer to layer $i_{\max}$, gathering a suitable collection of vertex-disjoint cycles and isolated vertices. When processing the tile $T_{i,j}$, it merges as many vertex-disjoint cycles and isolated points from previous iterations that are below the tile as possible. Once the procedure has reached the maximum layer which is completely contained in the smaller disk with radius $\frac{R}{2}$, the procedure attempts to merge all the remaining cycles and points.

We now describe the procedure in more detail. For each tile $T_{i,j}$ we will define a random variable $D_{i,j}$ called the *demand* of $T_{i,j}$, which will be used later in the probabilistic analysis to show that the procedure terminates successfully. Recall that $N(T_{i,j})$ denotes the number of points in tile $T_{i,j}$ and note that the collection of $N(T_{i,j})$ for admissible i, j are independent Poisson random variables for the Poissonized KPKVB model.

Lemma 4.4 (Cycle Merging, See Fig. 2). *If the vertices strictly below tile $T_{i,j}$ can be covered by x vertex-disjoint cycles and isolated vertices and the number of vertices in tile $T_{i,j}$ is $N(T_{i,j}) = y \geq 3$, then the set of all vertices below $T_{i,j}$ (including those in $T_{i,j}$) can be covered by $\max\{1, x - y + 1\}$ cycles and points.*

In particular, if $y > x$, then the vertices below $T_{i,j}$ can be covered by a single cycle which has $y - x$ edges within $T_{i,j}$.

Proof. If $y = N(T_{i,j}) \geq 3$, then the vertices in $T_{i,j}$ form a cycle by Lemma 4.2. Each of its y edges can be used to merge this cycle in $T_{i,j}$ with a cycle or point strictly below $T_{i,j}$: to pick up a cycle, use an edge $e_i = v_i v_{i+1}$ of the cycle $v_1, \dots, v_y$ in $T_{i,j}$ and choose any edge $e_* = a_* b_*$ from the cycle to be picked up. By Lemma 4.2, the four endpoints form a clique and therefore, we can go along the edges $v_i a_*$, then the cycle to be picked up (without the

Fig. 2. In three steps, three cycles (colored blue) are merged (resulting in the green cycle) by replacing an edge of one cycle by a detour around the other cycle. Note that we zoomed into the part of the disk which matters for the cycle merging. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

edge e_*), and finally along b_*v_{i+1} to bring us back to the cycle in $T_{i,j}$. To pick up a vertex a_* below, we can just use the edges v_ia_* and a_*v_{i+1} instead of v_iv_{i+1} .

If $y \leq x$, then all edges of the original cycle in $T_{i,j}$ will be used and we end up with $x - y + 1 \geq 1$ cycles and points below (and including) $T_{i,j}$. If $y > x$, then all cycles and points strictly below $T_{i,j}$ become part of the original cycle in $T_{i,j}$ and $y - x > 0$ edges of the cycle in $T_{i,j}$ remain unused and part of the final cycle. $\square$

The demand random variables $D_{i,j}$ for admissible i, j are defined in terms of the point counts $N(T_{i,j})$ as follows. For $i = 0$ and $j = 0, \dots, n_0 - 1$ we set:

$$D_{0,j} = \begin{cases} N(T_{0,j}) & \text{if } N(T_{0,j}) \in \{1, 2\}, \\ 0 & \text{otherwise,} \end{cases}$$

and, for $0 < i \leq i_{\max}$ and $j = 0, \dots, n_i - 1$ we set:

$$D_{i,j} = \max\{D_{i-1,2j} + D_{i-1,2j+1} + 3 - N(T_{i,j}), 0\}.$$

In particular $D_{i,j}$ and $D_{i,j'}$ are independent for $j \neq j'$, since they depend on disjoint regions. Also, the $D_{0,j}$ are i.i.d. random variables with values in $\{0, 1, 2\}$ satisfying

$$\mathbb{P}(D_{0,j} = 1) = \mu_0 e^{-\mu_0}, \quad \mathbb{P}(D_{0,j} = 2) = \frac{\mu_0^2}{2} e^{-\mu_0},$$

where we used the notation $\mu_0 := \mu_{n,\alpha,v}(T_{0,0}) \stackrel{\text{Lemma 4.3}}{=} \Theta(v)$.

Lemma 4.5. For admissible indices i, j , if $D_{i,j} = x$, then the vertices below (and in) $T_{i,j}$ can be covered by at most $x + 1$ vertex-disjoint cycles and isolated points (in total).

Moreover, if $i > 0$ and $D_{i,j} = 0$, then the vertices below (and in) $T_{i,j}$ can be covered by exactly one cycle which has at least one edge which is completely contained in $T_{i,j}$.

Proof. The proof is by induction on i . For $i = 0$, the claim is clear because $D_{0,j} = 0$ implies that there is either one cycle or no vertex in $T_{0,j}$.

For $i > 0$, assuming the claim for $i - 1$ we show it for i . By the induction hypothesis, the vertices below (and inside) $T_{i-1,2j}$ ($T_{i-1,2j+1}$, respectively) can be covered by $D_{i-1,2j} + 1$ ($D_{i-1,2j+1} + 1$, respectively) many vertex-disjoint cycles and isolated points. Thus, the vertices

strictly below $T_{i,j}$ can be covered by $D_{i-1,2j} + D_{i-1,2j+1} + 2$ many vertex-disjoint cycles and isolated points in total. If $N(T_{i,j}) \geq 3$, then by [Lemma 4.4](#), the vertices below $T_{i,j}$ can be covered by

$$\begin{aligned} \max\{1, D_{i-1,2j} + D_{i-1,2j+1} + 2 - N(T_{i,j}) + 1\} \leq \\ \max\{D_{i-1,2j} + D_{i-1,2j+1} + 3 - N(T_{i,j}), 0\} + 1 = D_{i,j} + 1. \end{aligned}$$

If $N(T_{i,j}) \leq 2 < 3$, then the points in $T_{i,j}$ just remain as single points and the area in and below $T_{i,j}$ can be covered by at most $D_{i-1,2j} + D_{i-1,2j+1} + 2 + N(T_{i,j}) \leq D_{i,j} + 1$ vertex-disjoint cycles and isolated points: indeed, if $N(T_{i,j}) = 0$ or $N(T_{i,j}) = 1$, the statement is clear. In the case $N(T_{i,j}) = 2$, we have $D_{i,j} + 1 = D_{i-1,2j} + D_{i-1,2j+1} + 3 - 2 + 1 = D_{i-1,2j} + D_{i-1,2j+1} + 2$. By the induction hypothesis, all vertices in the area below tile $T_{i,j}$ can be covered by $D_{i-1,2j} + D_{i-1,2j+1} + 2$ cycles and isolated vertices. If there is indeed any cycle or isolated vertex in the area below tile $T_{i,j}$, then the two vertices inside tile $T_{i,j}$ can be merged with it. So, all vertices in the area in and below tile $T_{i,j}$ can (still) be covered by $D_{i-1,2j} + D_{i-1,2j+1} + 2$ cycles and isolated vertices, which is the value of $D_{i,j} + 1$. Otherwise, all vertices in the area in and below tile $T_{i,j}$ can be covered by 2 (cycles and) isolated vertices as there are only 2 vertices and the claim follows because $D_{i,j} + 1 \geq 2$.

For the second part of the lemma, if $D_{i,j} = 0$, the points in the area in and below $T_{i,j}$ can be covered by one cycle or point. If $i > 0$, then the assumption $D_{i,j} = 0$ and the definition of $D_{i,j}$ imply that there are at least 3 points in $T_{i,j}$. Hence, the vertices in and below $T_{i,j}$ can be covered by exactly one cycle, which will have at least one edge with both endpoints inside $T_{i,j}$ (using that $N(T_{i,j}) > D_{i-1,2j} + D_{i-1,2j+1} + 2$). $\square$

Lemma 4.6. *If $D_{i,j} = 0$ for $i = i_{\max}$ and for all $j = 0, \dots, n_i - 1$, then there is a Hamilton cycle.*

Proof. Firstly, we observe that for $i = i_{\max}$, if $D_{i,j} = 0$, then all vertices in and below $T_{i,j}$ can be covered by one cycle that contains an edge whose endpoints are both in $T_{i,j}$ by [Lemma 4.5](#). Taking such an edge for $T_{i,0}$ and $T_{i,1}$, the four endpoints form a clique by the triangle inequality because all radial coordinates are at most $\frac{R}{2}$ and hence the cycle of $T_{i,1}$ can be taken as a detour to the cycle of $T_{i,0}$ as in the proof of [Lemma 4.4](#). As a result, we have a cycle covering all vertices below $T_{i,0}$ and $T_{i,1}$ and with an edge inside the i th layer. We can repeat this procedure to merge this resulting cycle also with those in $T_{i,2}, \dots, T_{i,n_i-1}$. We will end up with one cycle covering all vertices below all tiles $T_{i,0}, \dots, T_{i,n_i-1}$, and this cycle contains an edge whose endpoints are both in the inner disk with radius $\frac{R}{2}$. The remaining vertices in the inner disk, that are not in any tile, form a clique and in particular can be covered by a cycle. We can again merge this cycle with the one we created earlier via the same trick. $\square$

4.3. Probabilistic lemmas which ensure the a.a.s. successful termination of the procedure

In this subsection we show that the algorithm explained previously works successfully for the Poissonized KPKVB model $G_{\text{Po}}(n; \alpha, \nu)$ with $M \stackrel{\Delta}{=} \text{Po}(n)$ many vertices (the standard De-poissonization of [Lemma 2.1](#) gives then the result in the standard KPKVB model). [Lemma 4.8](#) of this section shows the exponential decay of the demand random variables, which we then use in [Lemma 4.9](#) to show that the demand random variables are simultaneously zero in the maximum layer. Appealing to [Lemma 4.6](#), we can then conclude that this makes the algorithm work.

4.3.1. Sub-exponential tail decay of demand

We first show the following technical lemma:

Lemma 4.7. *For all real $\epsilon \in (0, 1)$, there exists $\kappa = \kappa(\epsilon) > 0$ such that for all $i \in \mathbb{N}$, for all $x \geq \kappa i^2 \ln(1+i)$ we have: $(x+1)e^{-\frac{x}{i^2}} \leq \epsilon$.*

Proof. Pick $\kappa > \max\{\frac{1}{\ln 2}, 3\}$ such that $(\kappa+1)2^{3-\kappa} \leq \epsilon$; this is possible as $\lim_{a \rightarrow \infty} (a+1)2^{3-a} = 0$. We prove the lemma in the following way: in the first step we verify that for $x = \kappa i^2 \ln(1+i)$ we have $(x+1)e^{-\frac{x}{i^2}} \leq \epsilon$, and then we show that the left-hand side of the inequality is monotone decreasing in x (by showing that its derivative with respect to x is negative). Since the right-hand side is independent of x , this clearly implies the lemma.

For the first step, we need to show that $(\kappa i^2 \ln(1+i) + 1)e^{-\kappa \ln(1+i)} \leq \epsilon$. Using that $i^2 \leq (1+i)^2$, $\ln(1+i) \leq 1+i$ and $1 \leq (1+i)^3$, we note that the left-hand side of the inequality can be bounded from above by

$$(\kappa i^2 \ln(1+i) + 1)e^{-\kappa \ln(1+i)} \leq (\kappa+1)(1+i)^3(1+i)^{-\kappa} = (\kappa+1)(1+i)^{3-\kappa}$$

Now, if we plug in $i = 1$, this upper bound is at most ϵ by the choice of κ . The derivative in i of the latter expression is

$$(\kappa+1)(3-\kappa)(1+i)^{2-\kappa},$$

which is negative for $\kappa > 3$ for all $i \geq 1$. Therefore, the upper bound is monotone decreasing in i and hence, for all $i \geq 1$ and $x = \kappa i^2 \ln(1+i)$, we have $(x+1)e^{-\frac{x}{i^2}} \leq \epsilon$, concluding the first step.

For the second step, we need to verify that the derivative of $(x+1)e^{-\frac{x}{i^2}}$ is negative in $x \geq \kappa i^2 \ln(1+i) > 0$: using the assumptions of $\kappa > \frac{1}{\ln 2}$ and $i \geq 1$, we have

$$e^{-\frac{x}{i^2}} \left(1 + (x+1) \left(-\frac{1}{i^2} \right) \right) \leq 1 - \kappa \ln(1+i) - \frac{1}{i^2} \leq 1 - \kappa \ln 2 < 0.$$

The lemma follows. $\square$

We are now ready to state and prove the main lemma of this section.

Lemma 4.8. *There is a constant $c > 0$ such that for $0 < \alpha < \frac{1}{2}$ and v sufficiently large, for all admissible i, j , and all $t \geq 0$:*

$$\mathbb{P}(D_{i,j} \geq t) \leq e^{-ct}.$$

Proof. Set $c = 10$, $c_0 = c + \sum_{i=1}^{\infty} \frac{1}{i^2} < \infty$ and $c_i = c_{i-1} - \frac{1}{i^2}$, for $i > 0$. So, in particular, we have $\infty > c_0 > c_1 > \dots > c = 10 > 0$. The lemma immediately follows from the following claim: for all admissible i, j , for all $t \geq 0$, $\mathbb{P}(D_{i,j} \geq t) \leq e^{-c_i t}$.

We prove this claim by induction on i . For the base case $i = 0$, the claim is clear for $t > 2$ because $D_{0,j} \in \{0, 1, 2\}$, so $\mathbb{P}(D_{0,j} \geq t) = 0 < e^{-c_0 t}$. For $t = 1, 2$, we have

$$\mathbb{P}(D_{0,j} \geq t) \leq \mu_0 e^{-\mu_0} + \frac{\mu_0^2}{2} e^{-\mu_0} = O(v^2) e^{-\Theta(v)},$$

where the equality follows by Lemma 4.3. In particular, by choosing v large enough, it holds that $\mathbb{P}(D_{0,j} \geq t) \leq e^{-c_0 t}$ for $t = 1, 2$.

For the inductive step, assume the statement is true for $i - 1$ with $1 \leq i \leq i_{\max}$. Note that as $D_{i-1,2j}$ and $D_{i-1,2j+1}$ are independent, we can apply the induction hypothesis to $D_{i-1,2j}$ and $D_{i-1,2j+1}$ to get

$$\begin{aligned} \mathbb{P}(D_{i-1,2j} + D_{i-1,2j+1} \geq t) &\leq \sum_{s=0}^t \mathbb{P}(D_{i-1,2j} \geq s) \mathbb{P}(D_{i-1,2j+1} \geq t-s) \\ &\leq \sum_{s=0}^t e^{-c_{i-1}s} e^{-c_{i-1}(t-s)} = (t+1)e^{-c_{i-1}t}. \end{aligned} \quad (6)$$

Define

$$\epsilon = \min \left\{ e^{-3c_0}, \frac{1}{2}(1 - e^{-c}) \right\},$$

let $\kappa = \kappa(\epsilon)$ as in Lemma 4.7 and set

$$t_i := \kappa i^2 \ln(1+i) + 3.$$

We make a case distinction in t .

Case 1: $t \geq t_i$.

Using the definition of $D_{i,j}$ and by (6), we have

$$\begin{aligned} \mathbb{P}(D_{i,j} \geq t) &= \sum_{s=0}^{\infty} \mathbb{P}(N(T_{i,j}) = s) \mathbb{P}(D_{i-1,2j} + D_{i-1,2j+1} \geq t+s-3) \\ &\stackrel{(6)}{\leq} \sum_{s=0}^{\infty} \mathbb{P}(N(T_{i,j}) = s) (t+s-3+1) e^{-c_{i-1}(t+s-3)}. \end{aligned}$$

Now, we can apply Lemma 4.7 to $x = t+s-3 \geq \kappa i^2 \ln(1+i)$ to deduce that

$$(x+1)e^{(c_i - c_{i-1})x} = (x+1)e^{-\frac{x}{i^2}} \leq \epsilon, \text{ which implies } (x+1)e^{-c_{i-1}x} \leq \epsilon e^{-c_i x}.$$

We infer that

$$\begin{aligned} \mathbb{P}(D_{i,j} \geq t) &\leq \sum_{s=0}^{\infty} \mathbb{P}(N(T_{i,j}) = s) \epsilon e^{-c_i(t+s-3)} \\ &= \epsilon e^{3c_i} e^{-c_i t} \sum_{s=0}^{\infty} \mathbb{P}(N(T_{i,j}) = s) e^{-c_i s} \\ &\leq e^{-c_i t} \sum_{s=0}^{\infty} \mathbb{P}(N(T_{i,j}) = s) = e^{-c_i t}, \end{aligned}$$

where the third line follows by choice of ϵ and the definition of the sequence $c_0, c_1, \dots$, which implies that $c_i < c_0$.

Case 2: $t < t_i$.

Let $\mu_i := \mu_{n,\alpha,v}(T_{i,0})$. We first observe that for all $i \in \mathbb{N}_0$:

$$\mu_i \geq (c_i t_i + \ln 2) \frac{2}{1 - \ln 2} \quad (7)$$

and

$$\frac{1}{2} \mu_i \geq t_i + 3. \quad (8)$$

To see that this holds, note that since $\mu_i = \Omega(\nu 2^{i(1-2\alpha)})$ (see Lemma 4.3), we can take any $\nu_* > 0$ and then pick $i_0 = i_0(\nu_*) \in \mathbb{N}$ such that for all $\nu \geq \nu_*$ and all $i \geq i_0$, the claims hold (the right-hand side is independent of ν and grows at most polynomially in i whereas μ_i grows exponentially in i). Then, as the right-hand sides of (7) and (8) are independent of ν , we can pick $\nu_{**} > \nu_*$ large enough such that (7) and (8) also hold for $i = 0, \dots, i_0(\nu_*)$.

Thus, (7) and (8) hold for all $\nu > \nu_{**}$ and all $i \in \mathbb{N}_0$.

We have

$$\begin{aligned} & \mathbb{P}(D_{i,j} \geq t) \\ &= \sum_{j=0}^{\infty} \mathbb{P}(D_{i-1,2j} + D_{i-1,2j+1} = j) \mathbb{P}(D_{i-1,2j} + D_{i-1,2j+1} + 3 - N(T_{i,j}) \\ & \quad \geq t | D_{i-1,2j} + D_{i-1,2j+1} = j) \\ &= \sum_{j=0}^{\infty} \mathbb{P}(D_{i-1,2j} + D_{i-1,2j+1} = j) \mathbb{P}(N(T_{i,j}) \leq j + 3 - t) \\ &= \sum_{j=\max\{t-3,0\}}^{\infty} \mathbb{P}(D_{i-1,2j} + D_{i-1,2j+1} = j) \mathbb{P}(N(T_{i,j}) \leq j + 3 - t) \\ &\leq \sum_{j=\max\{t-3,0\}}^{\infty} \mathbb{P}(D_{i-1,2j} + D_{i-1,2j+1} = j) \mathbb{P}(N(T_{i,j}) \leq j + 3). \end{aligned}$$

We split the sum into two parts: for $j + 3 \leq \frac{1}{2}\mu_i$, we apply Lemma 2.2 with $k = j + 3$ and $\mu = \mu_i$ and hence, $\frac{k}{\mu} = \frac{j+3}{\mu_i} \leq \frac{1}{2}$. Therefore, $H(\frac{k}{\mu}) \geq \frac{1}{2}(1 - \ln 2) > 0$ and we get

$$\mathbb{P}(N(T_{i,j}) \leq j + 3) \leq e^{-\mu_i \frac{1}{2}(1-\ln 2)}.$$

By Eq. (7), it follows that

$$e^{-\mu_i \frac{1}{2}(1-\ln 2)} \leq \frac{1}{2} e^{-c_i t_i} \leq \frac{1}{2} e^{-c_i t}.$$

So, we have for the first part of the sum

$$\sum_{\substack{j=\max\{t-3,0\} \\ j+3 \leq \frac{1}{2}\mu_i}}^{\infty} \mathbb{P}(D_{i-1,2j} + D_{i-1,2j+1} = j) \mathbb{P}(N(T_i) \leq j + 3) \leq e^{-\mu_i \frac{1}{2}(1-\ln 2)} \leq \frac{1}{2} e^{-c_i t}.$$

For the second part, we have $j + 3 > \frac{1}{2}\mu_i$. By Eq. (8), $\frac{1}{2}\mu_i \geq t_i + 3 > t_i$. By (6) and Lemma 4.7 with $x = j \geq \kappa i^2 \ln(1 + i)$ it holds that

$$\mathbb{P}(D_{i-1,2j} + D_{i-1,2j+1} = j) \leq \mathbb{P}(D_{i-1,2j} + D_{i-1,2j+1} \geq j) \leq (j + 1) e^{-c_{i-1} j} \leq \epsilon e^{-c_i j}.$$

Using this inequality, we can also bound from above the second sum:

$$\begin{aligned} & \sum_{\substack{j=\max\{t-3,0\} \\ j+3 > \frac{1}{2}\mu_i}}^{\infty} \mathbb{P}(D_{i-1,2j} + D_{i-1,2j+1} = j) \mathbb{P}(N(T_{i,j}) \leq j + 3) \\ & \leq \sum_{j=t}^{\infty} \epsilon e^{-c_i j} = e^{-c_i t} \epsilon \sum_{j=0}^{\infty} (e^{-c_i})^j \end{aligned}$$

$$\begin{aligned} &= e^{-c_i t} \epsilon \frac{1}{1 - e^{-c_i}} \\ &\leq e^{-c_i t} \epsilon \frac{1}{1 - e^{-c}} \leq \frac{1}{2} e^{-c_i t}. \end{aligned}$$

where the last inequality follows from the choice of ϵ and the fact that $c_i > c$.

By combining both sums, we conclude that also for t as in Case 2, $\mathbb{P}(D_{i,j} \geq t) \leq \frac{1}{2} e^{-c_i t} + \frac{1}{2} e^{-c_i t} = e^{-c_i t}$, and the lemma follows. $\square$

4.3.2. Deriving Theorem 4.1

Finally, Theorem 4.1 is a result of the following lemma together with Lemma 4.6.

Lemma 4.9. *Let $0 < \alpha < \frac{1}{2}$ and v be sufficiently large. Then*

$$\mathbb{P}(D_{i_{\max},j} = 0, \text{ for all } j = 0, \dots, n_{i_{\max}} - 1) = 1 - o(n^{-1/2}).$$

Proof. First, let us recall that the number of tiles in the i th layer is $n_i = 2^{4-i+\lfloor \frac{R}{2 \ln 2} \rfloor}$. For $i = i_{\max} = \lceil \frac{0.9R}{2 \ln 2} \rceil$, it follows that $n_i = \Theta(2^{\frac{0.1R}{2 \ln 2}}) = \Theta(n^{0.1})$. Furthermore, $\mu_i = \Omega(2^{i_{\max}(1-2\alpha)}) = \Omega(n^{0.9(1-2\alpha)})$.

We have

$$\mathbb{P}(\text{for all } j = 0, \dots, n_i - 1 : D_{i,j} = 0) = 1 - \mathbb{P}(D_{i,j} > 0 \text{ for some } j).$$

By the union bound over all tiles in layer $i = i_{\max}$

$$\mathbb{P}(D_{i,j} > 0 \text{ for some } j) \leq \sum_{j=0}^{n_i-1} \mathbb{P}(D_{i,j} > 0).$$

Now we observe that if $D_{i-1,2j} \leq \frac{\mu_i}{100}$ and $D_{i-1,2j+1} \leq \frac{\mu_i}{100}$ and $N(T_{i,j}) \geq \frac{3}{100} \mu_i$ all hold, then $3 + D_{i-1,2j} + D_{i-1,2j+1} - N(T_{i,j}) \leq 3 + \frac{2}{100} \mu_i - \frac{3}{100} \mu_i \leq 0$ since $\mu_i = \Omega(v 2^{i(1-2\alpha)})$. Hence if all three of these conditions hold, then $D_{i,j} \leq 0$. In other words, if $D_{i,j} > 0$, then $D_{i-1,2j} > \frac{\mu_i}{100}$ or $D_{i-1,2j+1} > \frac{\mu_i}{100}$ or $N(T_{i,j}) < \frac{3}{100} \mu_i$.

Therefore,

$$\begin{aligned} \mathbb{P}(D_{i,j} > 0) &\leq \mathbb{P}\left(D_{i-1,2j} > \frac{\mu_i}{100} \text{ or } D_{i-1,2j+1} > \frac{\mu_i}{100} \text{ or } N(T_{i,j}) < \frac{3\mu_i}{100}\right) \\ &\leq \mathbb{P}\left(D_{i-1,2j} > \frac{\mu_i}{100}\right) + \mathbb{P}\left(D_{i-1,2j+1} > \frac{\mu_i}{100}\right) + \mathbb{P}\left(N(T_{i,j}) < \frac{3\mu_i}{100}\right). \end{aligned}$$

For the first two terms, we use Lemma 4.8, taking v sufficiently large, and for the third term, we apply Lemma 2.2. We get

$$\mathbb{P}(D_{i,j} > 0) \leq 2e^{-c \frac{\mu_i}{100}} + e^{-\Omega(\mu_i)} = e^{-\Omega(\mu_i)}.$$

Using that $\mu_i = \Omega(n^{0.9(1-2\alpha)})$, it follows that $\mathbb{P}(D_{i,j} > 0) = O(e^{-\Omega(n^{0.9(1-2\alpha)})})$. Since $n_{i_{\max}} = \Theta(n^{0.1})$, we obtain

$$\mathbb{P}(D_{i,j} > 0 \text{ for some } j) \leq \sum_j \mathbb{P}(D_{i,j} > 0) = O\left(n^{0.1} e^{-\Omega(n^{0.9(1-2\alpha)})}\right) = o(n^{-1/2}),$$

and the lemma follows. $\square$

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- [1] M.A. Abdullah, M. Bode, N. Fountoulakis, Typical distances in a geometric model for complex networks, *Internet Math.* 1 (2017).
- [2] M. Ajtai, J. Komlós, E. Szemerédi, First occurrence of Hamilton cycles in random graphs, *North-Holland Math. Stud.* 115 (1985) 173–175.
- [3] R. Albert, A.-L. Barabási, Statistical mechanics of complex networks, *Rev. Modern Phys.* 74 (1) (2002) 47–97.
- [4] J. Balogh, B. Bollobás, M. Krivelevich, T. Müller, M. Walters, Hamilton cycles in random geometric graphs, *Ann. Appl. Probab.* 21 (3) (2011) 1053–1072.
- [5] M. Bode, N. Fountoulakis, T. Müller, On the largest component of a hyperbolic model of complex networks, *Electron. J. Combin.* 22 (3) (2015) Paper P3.24, 43 pages.
- [6] M. Bode, N. Fountoulakis, T. Müller, The probability that the hyperbolic random graph is connected, *Random Structures Algorithms* 49 (1) (2016) 65–94.
- [7] B. Bollobás, The evolution of sparse graphs, in: *Graph Theory and Combinatorics*, Academic Press, London, 1984, pp. 35–57.
- [8] E. Candellero, N. Fountoulakis, Clustering and the hyperbolic geometry of complex networks, *Internet Math.* 12 (2016) 2–53.
- [9] J. Díaz, D. Mitsche, X. Pérez, Sharp threshold for Hamiltonicity of random geometric graphs, *SIAM J. Discrete Math.* 21 (1) (2007) 57–65.
- [10] N. Fountoulakis, On a geometrization of the chung-lu model for complex networks, 2015, pp. 361–387.
- [11] N. Fountoulakis, T. Müller, Law of large numbers in a hyperbolic model of complex networks, *Ann. Appl. Probab.* 28 (2018) 607–650.
- [12] T. Friedrich, A. Krohmer, On the diameter of hyperbolic random graphs, *SIAM J. Discrete Math.* 32 (2018) 1314–1334.
- [13] A. Frieze, X. Pérez-Giménez, P. Prałat, B. Reiniger, Perfect matchings and Hamiltonian cycles in the preferential attachment model, *Random Structures Algorithms* (2019) 258–288.
- [14] L. Gugelmann, K. Panagiotou, U. Peter, Random hyperbolic graphs: degree sequence and clustering, in: *Conference version in ICALP 2012*. Preprint. Available from <http://arxiv.org/abs/1205.1470>.
- [15] L. Gugelmann, K. Panagiotou, U. Peter, Random hyperbolic graphs: Degree sequence and clustering, in: *Proceedings of the 39th International Colloquium Conference on Automata, Languages, and Programming - Volume Part II, ICALP'12*, Springer-Verlag, Berlin, Heidelberg, 2012, pp. 573–585.
- [16] M.A. Kiwi, D. Mitsche, A bound for the diameter of random hyperbolic graphs, in: R. Sedgewick, M.D. Ward (Eds.), *Proceedings of the Twelfth Workshop on Analytic Algorithmics and Combinatorics, ANALCO 2015*, San Diego, CA, USA, January 4, 2015, SIAM, 2015, pp. 26–39.
- [17] M. Kiwi, D. Mitsche, Spectral gap of random hyperbolic graphs and related parameters, *Ann. Appl. Probab.* 28 (2018) 941–989.
- [18] M.A. Kiwi, D. Mitsche, On the second largest component of random hyperbolic graphs, *SIAM J. Discrete Math.* 33 (4) (2019) 2200–2217.
- [19] J. Komlós, E. Szemerédi, Limit distribution for the existence of Hamiltonian cycles in a random graph, *Discrete Math.* 43 (1983) 55–63.
- [20] A. Korshunov, A solution of a problem of P. Erdős and A. Rényi about Hamilton cycles in non-oriented graphs, *Metody Diskr. Anal. Teoriy Upr. Syst., Sb. Trudov Novosibirsk* 31 (1977) 17–56 (in Russian).
- [21] D. Krioukov, F. Papadopoulos, M. Kitsak, A. Vahdat, M. Boguñá, Hyperbolic geometry of complex networks, *Phys. Rev. E* (3) 82 (3) (2010) 036106, 18.
- [22] G. Last, M. Penrose, *Lectures on the Poisson Process*, IMS Textbooks, Cambridge University Press, 2018.
- [23] T. Müller, X. Pérez-Giménez, N.C. Wormald, Disjoint Hamilton cycles in the random geometric graph, *J. Graph Theory* 68 (4) (2011) 299–322.
- [24] T. Müller, M. Staps, The diameter of KPKVB random graphs, 2017, Available at [arXiv:1707.09555](https://arxiv.org/abs/1707.09555).
- [25] M.D. Penrose, *Random Geometric Graphs*, Oxford University Press, Oxford, 2003.
- [26] L. Pósa, Hamiltonian circuits in random graphs, *Discrete Math.* 14 (1976) 359–364.
- [27] R.W. Robinson, N.C. Wormald, Almost all regular graphs are Hamiltonian, *Random Struct. Algorithms* 5 (1994) 363–374.
- [28] J. Stillwell, *Geometry of Surfaces*, Universitext, Springer-Verlag, New York, Berlin, Heidelberg, 1992.