University of Groningen

Cytotoxic edema associated with hemorrhage predicts poor outcome after traumatic brain injury

Turtzo, L Christine; Luby, Marie; Jikaria, Neekita; Griffin, Allison Diane; Greenman, Danielle; Bokkers, Reinoud P H; Parikh, Gunjan; Peterkin, Nicole; Whiting, Mark; Latour, Lawrence

Published in:
Journal of Neurotrauma

DOI:
[10.1089/neu.2021.0037](https://doi.org/10.1089/neu.2021.0037)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Turtzo, L. C., Luby, M., Jikaria, N., Griffin, A. D., Greenman, D., Bokkers, R. P. H., Parikh, G., Peterkin, N., Whiting, M., & Latour, L. (2021). Cytotoxic edema associated with hemorrhage predicts poor outcome after traumatic brain injury. *Journal of Neurotrauma*, 38(22), 3107-3118. <https://doi.org/10.1089/neu.2021.0037>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

ORIGINAL ARTICLE

IMAGING

Cytotoxic Edema Associated with Hemorrhage Predicts Poor Outcome after Traumatic Brain Injury

L. Christine Turtzo,¹ Marie Luby,¹ Neekita Jikaria,² Allison Diane Griffin,² Danielle Greenman,² Reinoud P.H. Bokkers,³ Gunjan Parikh,^{4,5} Nicole Peterkin,² Mark Whiting,² and Lawrence L. Latour^{1,2,*}

Abstract

Magnetic resonance imaging (MRI) is used rarely in the acute evaluation of traumatic brain injury (TBI) but may identify findings of clinical importance not detected by computed tomography (CT). We aimed to characterize the association of cytotoxic edema and hemorrhage, including traumatic microbleeds, on MRI obtained within hours of acute head trauma and investigated the relationship to clinical outcomes. Patients prospectively enrolled in the Traumatic Head Injury Neuroimaging Classification study (NCT01132937) with evidence of diffusion-related findings or hemorrhage on neuroimaging were included. Blinded interpretation of MRI for diffusion-weighted lesions and hemorrhage was conducted, with subsequent quantification of apparent diffusion coefficient (ADC) values. Of 161 who met criteria, 82 patients had conspicuous hyperintense lesions on diffusion-weighted imaging (DWI) with corresponding regions of hypointense ADC in proximity to hemorrhage. Median time from injury to MRI was 21 (10–30) h. Median ADC values per patient grouped by time from injury to MRI were lowest within 24 h after injury. The ADC values associated with hemorrhagic lesions are lowest early after injury, with an increase in diffusion during the subacute period, suggesting transformation from cytotoxic to vasogenic edema during the subacute post-injury period. Of 118 patients with outcome data, 60 had Glasgow Outcome Scale Extended scores ≤ 6 at 30/90 days post-injury. Cytotoxic edema on MRI (odds ratio [OR] 2.91 [1.32–6.37], $p = 0.008$) and TBI severity (OR 2.51 [1.32–4.74], $p = 0.005$) were independent predictors of outcome. These findings suggest that in patients with TBI who had findings of hemorrhage on CT, patients with DWI/ADC lesions on MRI are more likely to do worse.

Keywords: apparent diffusion coefficient; cytotoxic edema; diffusion-weighted imaging; hemorrhage; magnetic resonance imaging; traumatic brain injury

Introduction

Cerebral edema contributes to the pathogenesis of secondary injury after traumatic brain injury (TBI) and stroke.^{1–3} The failure of energy-dependent processes can lead to cytotoxic edema, which appears on acute magnetic resonance imaging (MRI) as a hyperintense signal on diffusion-weighted imaging (DWI), caused by a correlated decrease in the apparent diffusion coefficient (ADC) of water. With hemorrhagic transformation, a mix of cytotoxic and vasogenic edema occurs, with an evolution toward the latter over 5 to 10 days, accompanied by normalization of the ADC as water mobility increases.^{4–12}

The cause of the vasogenic edema remains an area of active investigation, potentially attributable to the acute immune response and amenable to therapy. Because MRI and especially serial MRI are used rarely in the acute setting for TBI, it is unknown whether traumatic lesions undergo a similar transformation from cytotoxic to vasogenic.

Identification of acute cytotoxic edema after TBI has important clinical implications. While vasogenic edema is, in principle, reversible, lesions identified as cytotoxic on acute MRI may represent non-viable tissue and be an independent predictor of poor outcome. Evidence of large hemorrhagic lesions is a known predictor of poor

¹National Institute of Neurological Disorders and Stroke, National Institutes of Health, Bethesda, Maryland, USA.

²Center for Neuroscience and Regenerative Medicine, Bethesda, Maryland, USA.

³Department of Radiology, Medical Imaging Center, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands.

⁴R Adams Shock Trauma Center and ⁵Division of Neurocritical Care and Emergency Neurology, University of Maryland School of Medicine, Baltimore, Maryland, USA.

*Address correspondence to: Lawrence L. Latour, PhD, National Institute of Neurological Disorders and Stroke, National Institutes of Health, 10 Center Drive, Building 10 Room B1D733, Bethesda, MD 20892, USA E-mail: LatourL@ninds.nih.gov

outcome in moderate and severe TBI, presumably in part because of concomitant parenchymal injury leading to cell death.² In patients with mild TBI, however, the relationship between acute hemorrhage and outcome has not been investigated thoroughly, leading to conflicting findings. Although molecular processes involved in cytotoxic and vasogenic edema are interrelated,² distinct therapeutic approaches would be beneficial for patients with MRI-identified acute cytotoxic edema.

After TBI of all severities, traumatic microbleeds (TMBs) predict poor outcome and may be indicative of primary traumatic vascular injury.^{13,14} We hypothesized that traumatic vascular injury may be associated with early cytotoxic edema, with subsequent evolution toward vasogenic edema. Tissue ADC values in proximity to TMBs or larger traumatic lesions such as contusion or intraparenchymal hemorrhage (IPH) would then be low early after injury. With increasing time from injury, ADC values would be isointense—i.e., normalize, as described in stroke studies. We further hypothesized that cytotoxic edema associated with traumatic vascular injury could predict poor clinical outcome.

The goals of this study are to: (1) describe the concordance of DWI hyperintense lesions with the presence and type of hemorrhage and associated ADC lesions, (2) quantify acute values and temporal evolution of hemorrhage-associated ADC, in particular TMBs, over the first week after TBI, and (3) determine whether cytotoxic edema associated with vascular injury is a predictor of clinical outcome.

Methods

Additional details are available in Supplementary Methods.

Patient population and study design

This is an observational cohort analysis of adult patients with TBI prospectively enrolled in the Institutional Review Board approved Traumatic Head Injury Neuroimaging Classification (THINC) study (NCT01132937) from October 2010 to May 2017. Written informed consent was obtained before any research procedures. Enrolled patients were included for retrospective analysis if they had (i) completed baseline visit with MRI within 48 h of injury, (ii) an imaging target defined as an acute or subacute lesion on DWI, or hemorrhage on MRI or computed tomography (CT), including TMB, contusion, IPH, or subarachnoid hemorrhage, (iii) Glasgow Coma Scale (GCS) score assessed at baseline, and (iv) one-week follow-up MRI (Fig. 1). Optional follow-up visits with Glasgow Outcome Scale Extended (GOSE) occurred at 30 and 90 days.

Image acquisition

Patients were imaged for research purposes on a 1.5T (GE Healthcare, USA) or a 3T (Siemens, USA or Phil-

lips, USA) MRI scanner using a standardized imaging protocol, appropriate for imaging in an acute care setting¹³ and approximately 30 min in duration. The imaging protocol included (i) diffusion tensor imaging and DWI/ADC, (ii) T2*, (iii) T2-weighted fluid attenuated inversion recovery (FLAIR), (iv) three dimensional (3D)-T1 SPGR, (v) two-dimensional (2D)-T1 weighted, (vi) either dynamic susceptibility contrast perfusion or dynamic FLAIR during injection of a single weight-adjusted dose of gadolinium-based contrast agent at 5 mL/sec, (vii) post-contrast 2D-T1, and (viii) post-contrast FLAIR.

Image analysis

Image interpretation and analysis were completed in two stages with blinding to minimize investigator bias: (i) independent raters reviewed imaging for qualitative identification of focal lesions on DWI/ADC to determine their concordance with evidence of blood on T2* imaging and to classify the type of all hemorrhagic lesions observed for each patient, irrespective of association with diffusion changes; and (ii) the quantitative ADC value in areas of focal DWI hyperintensity was assessed by a separate set of trained raters independent of the findings from the qualitative consensus review. Only after quantitative measurements were complete were labels matched for statistical analysis, with each focal lesion assigned to a single type of lesion.

Statistical analysis

Interrater agreement was assessed using the Cohen kappa. Mean or median are reported, and non-parametric testing (Wilcoxon rank-sum and Mann-Whitney *U*) was used unless normality could be established (Kolmogorov-Smirnov test) for univariate analysis; categorical variables are presented as proportions (chi-square or Fisher exact tests). Baseline variables were compared between patients with and without evidence of cytotoxic edema, for all patients and excluding moderate and severe TBI.

Univariate analysis was performed on predictors of residual disability (defined as GOSE score ≤ 6 from latest available of 30- or 90-day follow-up), for all patients and excluding moderate and severe TBI. Stepwise logistic regression was used to test for independence using $p = 0.05$ for entry and $p = 0.1$ for removal. Statistical analysis was performed in IBM SPSS™ (Version 26).

Results

Patient population

One-hundred sixty-one patients met criteria and were included in this study as follows: consent was obtained from 641 patients from October 2010 to May 2017, 575 completed baseline imaging, 225 had an imaging target (defined as acute or subacute lesion on DWI, or

FIG. 1. Flowchart of study patient inclusion and exclusion. THINC, Traumatic Head Injury Neuroimaging Classification study (NCT01132937); MRI, magnetic resonance imaging; GCS, Glasgow Coma Scale; DWI, diffusion-weighted imaging; TMB, traumatic microbleed; GOSE, Glasgow Outcome Scale Extended; TBI, traumatic brain injury.

hemorrhage on MRI or CT), 165 completed follow-up MRI, and four were excluded because of unavailable GCS score (see flowchart in Fig. 1).

Demographics and baseline data are presented in Table 1. Patients were predominantly male (73%). Twenty-nine (18%) were classified as having moderate or severe TBI. Median times from injury to clinical CT and to research MRI were 1.7 (1.1–6.4) and 21 (10–30) h. A

majority of the included patients (59%) had evidence of acute blood on CT, largely attributable to extra-axial hemorrhage. Of the 20% of patients who were CT negative but MRI positive, 94% had TMBs.

Temporal evolution of cytotoxic edema

Figure 2 displays examples of DWI hyperintense lesions on research MRI, stratified by proximity of hemorrhagic

Table 1. Patients Stratified by Absence or Presence of Cytotoxic Edema

	Total (n = 161)	Absence (n = 79)	Presence (n = 82)	p
Demographics				
Age (median, IQR)	54 (33–66)	57 (38–66)	49 (29–65)	0.24
18–30	35 (22%)	13 (16%)	22 (27%)	0.26
31–64	80 (50%)	41 (52%)	39 (48%)	
65 +	46 (29%)	25 (32%)	21 (26%)	
Male	117 (73%)	53 (67%)	64 (78%)	0.15
Race				
White	112 (70%)	50 (63%)	62 (75%)	0.13
African American	38 (23%)	24 (30%)	14 (17%)	
Other	11 (7%)	5 (6%)	6 (7%)	
Clinical Presentation				
Trauma Level I	85 (52%)	38 (48%)	47 (57%)	0.27
Clinical Dx intracranial hemorrhage	92 (57%)	32 (40%)	60 (73%)	< 0.001
Clinical Dx of TBI	114 (70%)	51 (64%)	63 (76%)	0.11
ED discharge disposition				
Floor	54 (38%)	35 (55%)	19 (24%)	0.001
Stepdown unit	5 (3%)	2 (3%)	3 (3%)	
ICU	85 (53%)	27 (34%)	58 (71%)	
Injury cause				
Road traffic accident	62 (39%)	33 (41%)	29 (35%)	0.81
Incidental fall	72 (44%)	34 (43%)	38 (46%)	
Violence/assault	17 (10%)	7 (8%)	10 (12%)	
Other	10 (6%)	5 (6%)	5 (6%)	
Injury mechanism				
Acceleration/deceleration	32 (20%)	18 (23%)	14 (17%)	0.59
Direct impact–blow to head	40 (25%)	19 (24%)	21 (25%)	
Direct impact–head against object	25 (15%)	12 (15%)	13 (15%)	
Fall–ground floor	40 (25%)	21 (26%)	19 (23%)	
Fall from height >1 m	23 (14%)	8 (10%)	15 (18%)	
GCS				
15	112 (70%)	63 (80%)	49 (60%)	0.02
14	31 (19%)	11 (14%)	20 (24%)	
≤13	18 (11%)	5 (6%)	13 (16%)	
LOC reported	100 (62%)	42 (53%)	58 (70%)	0.02
PTA reported	109 (68%)	51 (65%)	58 (71%)	0.50
TBI severity				
Subconcussive	33 (21%)	21 (27%)	12 (15%)	0.12
Mild	99 (61%)	48 (61%)	51 (62%)	
Moderate	21 (13%)	8 (10%)	13 (16%)	
Severe	8 (5%)	2 (2%)	6 (7%)	
Imaging Findings				
CT–Clinical				
Hours–injury to CT (median, IQR)	1.68 (1.1–6.4)	1.66 (1.2–5.4)	1.68 (1.1–7.5)	0.98
CT positive for TBI	95 (59%)	31 (39%)	64 (78%)	<0.001
Intraparenchymal hematoma	38 (24%)	25 (30%)	13 (16%)	0.04
Extra-axial hemorrhage	88 (55%)	28 (35%)	60 (73%)	<0.001
Skull fracture	36 (22%)	13 (17%)	23 (28%)	0.09
MRI				
Hours–injury to MRI (median, IQR)	20.5 (10–30)	17.7 (8–26)	21.9 (15–33)	0.01
Strength–1.5T	49 (30%)	27 (34%)	22 (27%)	0.31
Strength–3T	112 (70%)	52 (66%)	60 (73%)	
Subdural hemorrhage	75 (47%)	23 (29%)	52 (63%)	<0.001
Subarachnoid hemorrhage	68 (42%)	17 (21%)	51 (62%)	<0.001
TMB	101 (63%)	47 (59%)	54 (66%)	0.42
Contusion or IPH	49 (30%)	10 (13%)	39 (48%)	<0.001

IQR, interquartile range; Dx, diagnosis; ED, emergency department; TBI, traumatic brain injury; ICU, intensive care unit; GCS, Glasgow Coma Scale; LOC, loss of consciousness; PTA, post-traumatic amnesia; CT, computed tomography; MRI, magnetic resonance imaging; TMB, traumatic microbleed; IPH, intraparenchymal hematoma. **Bold** indicates $p < 0.05$.

findings on T2* imaging. Most lesions, except those associated with IPH, were small and punctate or linear appearing (Fig. 2A, in reconstructed multi-planar view). One hundred seventeen (73%) patients had lesions conspicuous on DWI, and 82 (51%) had corresponding visible regions of hypointense ADC, suggestive of cytotoxic edema.

Interrater agreement for presence of DWI hyperintensity and ADC hypointensity was $k=0.78$ and $k=0.73$, respectively. Agreement on association with hemorrhagic lesion type varied from $k=0.38$ to 0.8. Approximately half of patients with TMBs, in comparison with ~95% of patients with IPH, also had DWI

hyperintense lesions. Of 33 isolated DWI lesions, 28 (85%) were correspondingly hypointense on ADC.

Figure 3 shows a patient with DWI hyperintensity in proximity to a TMB at initial presentation and follow-up. The TMB conspicuity appears similar on gradient recalled echo (i) and DWI (ii). The lesion on ADC (iii) transitions from hypointense to hyperintense and increases in conspicuity on follow-up T2 FLAIR (iv).

Of 54 patients identified as having DWI lesions associated with TMBs, two were excluded from quantitative analysis because of motion artifact. Values of ADC were not normally distributed. The median ADC value at baseline for all volumes of interest (VOIs) was 0.77 (0.68–0.83), significantly lower than the ADC value at follow-up 0.82 (0.76–0.93), $p < 0.001$. Values of ADC from reference VOIs in deep gray matter were similar between time points ($p = 0.98$). ADC values for all VOIs stratified by hemorrhagic lesion type did not differ from each other at baseline, but significantly increased over time (Fig. 4A).

Lesions with lower ADC values at baseline, suggesting cytotoxic edema, had increased ADC values at follow-up (Fig. 4B). The ADC values for all TMB-associated VOIs were averaged per subject and plotted along with reference VOIs against time from injury to imaging for both baseline and follow-up (Fig. 5A). The per patient average ADC value of DWI hyperintense TMBs imaged within 24 h was 0.70 (0.65–0.80), which was significantly lower than those obtained between 24–72 h, 0.83 (0.79–0.90), $p < 0.001$, and those after 72 h, 0.82 (0.79–0.84), $p < 0.001$ (Fig. 5B). Lesion ADC values did not differ between 24–72 hour and >72 h groupings ($p = 0.801$).

Cytotoxic edema: predictors and clinical outcome

Table 1 is stratified by the presence or absence of cytotoxic edema, regardless of association with hemorrhagic lesions. Univariate predictors of cytotoxic edema in-

cluded GCS score, documented loss of consciousness, and hemorrhagic findings on CT and MRI, but not baseline TBI severity, even for the mild TBI population (Supplementary Table 1).

Clinical outcome assessed by GOSE was available for 118 (73%) patients (19 with 30-day only, 16 with 90-day only, 83 with both), with approximately half having some degree of disability (GOSE score ≤ 6). The univariate predictors of outcome trauma center level, TBI severity, presence of TMB or intraparenchymal hematoma/contusion on MRI, DWI hyperintensity, and ADC hypointensity were associated with disability (Table 2).

Acute findings on CT and MRI field strength were not associated with outcome. The MRI findings, specifically TMBs and DWI hyperintensity and ADC hypointensity associated with TMBs, were predictors for the mild TBI population (Supplementary Table 2). In a stepwise logistic regression that considered trauma center level, TBI severity, evidence of intraparenchymal hemorrhage/contusion or TMBs, cytotoxic edema seen on MRI, and ADC hypointensity, only TBI severity (OR 2.51 [1.32–4.74], $p = 0.005$) and cytotoxic edema (OR 2.91 [1.32–6.37], $p = 0.008$) were independently associated with outcome.

At presentation to the emergency department, 41% (66/161) patients were documented as having polytrauma, with no significant differences in the outcome of cytotoxic edema between those with and without polytrauma ($p = 0.084$). There were no significant differences in GOSE outcomes between those with and without polytrauma, whether the effects were attributed to head injury alone ($p = 0.819$), illness or injury to other parts of the body ($p = 0.522$), or to a mixture of head injury and other ($p = 0.219$). Two patients in the study had cranial surgery (one with hemispherectomy before acute MRI, and another with cranial burr holes after acute MRI), with one reporting a GOSE score ≤ 6 and the other reporting GOSE score > 6 at this study's end point.

FIG. 2. Types of acute traumatic brain injury lesions associated with magnetic resonance imaging findings. White arrows indicate the areas of interest. (A) Parenchymal hyperintensities on diffusion-weighted imaging (DWI), as seen in (i) coronal view, can be associated with acute traumatic microbleeds (TMBs) seen on (ii) gradient recalled echo (GRE) imaging, with (iii) axial DWI hyperintensities colocalizing with (iv) apparent diffusion coefficient (ADC) hypointensities. (B) Parenchymal (i) DWI hyperintensity and colocalizing (ii) ADC hypointensity can also occur adjacent to areas of subarachnoid hemorrhage, as seen on (iii) gradient recalled echo (GRE) imaging. (C) Similar changes can occur in parenchymal adjacent to areas of hemorrhage associated with contusion or intraparenchymal hematoma. (D) Isolated areas hyperintense on DWI and hypointense on ADC may also be observed in the absence of hemorrhage on GRE. (E) Frequency of visually conspicuous hyperintense lesions on DWI and hypointense lesions on ADC stratified by adjacent hemorrhagic lesion or seen in isolation. SAH, subarachnoid hemorrhage; TMB, traumatic microbleed; IPH, intraparenchymal hematoma.

FIG. 3. Representative example of the evolution of magnetic resonance imaging findings associated with a traumatic microbleed from acute presentation to follow-up. **(A)** At 16 h after mild traumatic brain injury, near the traumatic microbleed on (i) gradient recalled echo (GRE), there is a hyperintense area on (ii) diffusion-weighted imaging (DWI) that is hypointense on initial (iii) apparent diffusion coefficient (ADC), indicative of cytotoxic edema. On follow-up **(B)**, one week after injury, this same area remains hyperintense on DWI but is now hyperintense on ADC, suggesting that vasogenic edema has developed. The hemorrhage related to the traumatic microbleed is visible on both initial and follow-up GRE, with associated (iv) fluid-attenuated inversion recovery (FLAIR) hyperintensity.

Discussion

We present evidence of cytotoxic edema in clinically mild patients imaged early after head trauma. Lesions appearing as focal hyperintensities on DWI were found in isolation, but more commonly near areas of hemorrhage seen on T2* MRI, including TMBs. Approximately two-thirds of patients with hyperintense lesions on DWI had corresponding regions of ADC hypointensity, similar to findings seen in acute stroke.

Quantitative analysis of ADC values demonstrated a temporal evolution from low values (visually identifiable as hypointensity) early after injury to elevated values at follow-up, suggesting evolution from cytotoxic to vasogenic edema. Patients with evidence of cytotoxic edema, regardless of the strata of hemorrhagic lesion, were more likely to have residual disability in subsequent months. Collectively, the data suggest that primary traumatic injury to cerebral vasculature can be associated with cytotoxic edema and poor outcome.

During initial enrollment in this protocol, small lesions with DWI hyperintensity and corresponding ADC hypointensity were observed in clinically mild trauma patients with negative CT. These radiological findings were initially interpreted as suggestive either of ischemic stroke or diffuse axonal injury (DAI),^{15–19} resulting in further clinical subspecialty consultation. As cases accumulated,

many of these DWI lesions were seen in proximity to hemorrhagic lesions, but the clinical relevance of this observation in clinically mild patients with TBI remained unclear. The current analysis suggests that when DWI hyperintense/ADC hypointense lesions are seen on MRI early after injury, the patient is at greater risk of incomplete recovery.

The mechanism behind these DWI/ADC changes remains speculative at this point and may be multifactorial. As part of our quantitative analysis, we investigated TMBs in particular, which have become known to signify traumatic vascular injury, serving as distinct lesions where mechanical forces to vessels can result in vascular injury and extravasation of blood into the perivascular space and parenchyma.^{13,14}

Cytotoxic edema detected in proximity to TMBs could reflect any of the following: focal ischemia from interruption in blood flow, anoxic depolarization as a cascade initiated by the presence of blood and free radicals in direct contact with the neuropil, direct traumatic disruption to cellular/axonal elements, or a combination of these mechanisms. A recent positron emission tomography study of cerebral physiology in moderate/severe TBI observed cerebral ischemia in structurally normal tissue in two-thirds of patients within the first 24 h.²⁰ Perihematomal ischemia and edema are often observed in primary IPH²¹ or contusion.²²

FIG. 4. Quantification of apparent diffusion coefficient (ADC) of water for all lesion types. **(A)** Scatter plot of ADC values of baseline presentation versus follow-up visit for all lesion types (one data point is off-scale). **(B)** Change in ADC for all volumes of interest stratified by quartiles based on baseline ADC value (minimum = 0.3954; 25th percentile = 0.6769; median = 0.7610; 75% = 0.8699; maximum = 1.419; $n = 481$). IPH, intraparenchymal hematoma; SAH, subarachnoid hemorrhage; TMB, traumatic microbleed. Color image is available online.

Hemorrhagic lesions after TBI, however, may reside on a continuum of severity. Even relatively discrete lesions such as TMBs can be associated with parenchymal edema.

Direct mechanical injury to parenchyma may result in lesions on DWI MRI. Traditionally, TMBs have been linked with DAI,^{15–19} a hallmark of TBI across injury severities and a significant predictor of outcome.^{23–25} The

FIG. 5. Quantification of apparent diffusion coefficient (ADC) of water for lesions in proximity to traumatic microbleeds. **(A)** The ADC values at baseline and follow-up visits for traumatic microbleeds are plotted based on the time from injury to magnetic resonance imaging (MRI), with reference values shown for comparison. **(B)** Box-whisker plot of ADC values grouped by baseline time from injury to MRI for traumatic microbleeds (* $p < 0.001$ for <24 h versus 24–72 h and versus >72 h). Color image is available online.

DAI may result from shearing forces at the moment of injury, causing extensive damage to neurons, glia, and blood vessels and ultimately resulting in neurologic deficits.^{26,27}

Recent studies call into question whether TMBs are always associated with axonal injury. Across injury severity, TMBs that appear punctate in the axial plane are often revealed as linear structures that may be specific

Table 2. Predictors of Clinical Outcome at 30 or 90 Days

	Total (n = 118)	GOSE ≤6 (n = 60)	GOSE 7 or 8 (n = 58)	p
Demographics				
Age (median, IQR)	47 (37–62)	52 (30–64)	60 (33–66)	
18–30	25 (21%)	14 (23%)	11 (19%)	0.16
31–64	56 (48%)	32 (53%)	24 (41%)	
65 +	37 (31%)	14 (23%)	23 (40%)	
Male	84 (71%)	43 (72%)	41 (70%)	> 0.99
Race				
White	84 (71%)	39 (65%)	45 (78%)	0.13
African American	24 (20%)	15 (25%)	9 (16%)	
Other	10 (9%)	6 (10%)	4 (7%)	
Level I Trauma	59 (50%)	36 (60%)	23 (40%)	0.04
Clinical Presentation				
GCS				
15	83 (70%)	38 (63%)	45 (78%)	0.07
14	21 (18%)	11 (18%)	10 (17%)	
≤13	14 (12%)	11 (18%)	3 (5%)	
TBI severity				
Subconcussive	20 (16%)	6 (10%)	14 (24%)	0.01
Mild	75 (63%)	37 (61%)	38 (65%)	
Moderate	17 (14%)	11 (18%)	6 (10%)	
Severe	6 (5%)	6 (10%)	0 (0%)	
LOC reported	73 (62%)	42 (70%)	31 (53%)	0.08
PTA reported	85 (72%)	44 (73%)	41 (71%)	0.83
Clinical diagnosis intracranial hemorrhage	65 (55%)	33 (55%)	32 (55%)	> 0.99
Imaging Findings				
CT-clinical				
CT positive for TBI	68 (58%)	35 (58%)	33 (57%)	> 0.99
Intraparenchymal hemorrhage	29 (25%)	16 (27%)	13 (22%)	0.67
Extra-axial hemorrhage	65 (55%)	34 (57%)	31 (53%)	0.85
MRI				
Strength–1.5T	36 (30%)	15 (25%)	21 (36%)	0.19
Strength–3T	82 (70%)	45 (75%)	37 (64%)	
Subdural or epidural hemorrhage	55 (46%)	32 (53%)	23 (40%)	0.14
Subarachnoid hemorrhage	53 (45%)	26 (43%)	27 (47%)	0.85
Contusion or IPH	36 (31%)	24 (40%)	12 (21%)	0.02
TMB(s)	78 (66%)	45 (75%)	33 (57%)	0.05
MRI - DWI hyperintense proximate to hemorrhage				
DWI lesion–not stratified by hemorrhage	86 (72%)	51 (85%)	35 (60%)	0.004
DWI Lesion–isolated	25 (21%)	17 (28%)	8 (14%)	0.07
Subdural or epidural hemorrhage	43 (36%)	28 (47%)	15 (26%)	0.02
Subarachnoid hemorrhage	40 (34%)	20 (33%)	20 (35%)	> 0.99
Contusion or IPH	36 (31%)	24 (40%)	12 (21%)	0.02
TMB raumatic(s)	40 (33%)	28 (46%)	12 (20%)	0.004
MRI–DWI hyperintense, ADC isointense	21 (18%)	10 (17%)	11 (19%)	0.81
MRI–ADC hypointense proximate to hemorrhage				
ADC lesion–not stratified by hemorrhage	65 (55%)	41 (68%)	24 (41%)	0.005
ADC lesion–isolated	22 (19%)	16 (27%)	6 (10%)	0.03
Subdural or epidural hemorrhage	12 (10%)	6 (10.0%)	6 (10%)	> 0.99
Subarachnoid hemorrhage	24 (20%)	14 (23%)	10 (17%)	0.49
Contusion or IPH	31 (26%)	21 (35%)	10 (17%)	0.03
TMB(s)	25 (2%)	19 (32%)	6 (10%)	0.006

GOSE, Glasgow Outcome Scale Extended; IQR, interquartile range; GCS, Glasgow Coma Scale; TBI, traumatic brain injury; LOC, loss of consciousness; PTA, post-traumatic amnesia; CT, computed tomography; MRI, magnetic resonance imaging; IPH, intraparenchymal hematoma; DWI, diffusion-weighted imaging; ADC, apparent diffusion coefficient of water; TMB, traumatic microbleed. **Bold** indicates $p < 0.05$

to TBI.¹³ We demonstrated that acute TMBs correspond with post-mortem pathological features consistent with secondary injury surrounding injured vessels, but not with traditional markers of DAI.¹⁴ Other MRI studies have reported limited colocalization of TMBs with diffusion evidence of axonal injury in TBI.^{28,29}

In the region adjacent to TMBs, we observed hyperintense DWI signal colocalizing with ADC hypointensity during the acute period, which can be characteristic of

failures of energy-dependent processes and ischemic tissue. At follow-up, nearly 70% of these TMBs had increased normalized ADC values. Lesions with the lowest ADC values at baseline also had larger changes in ADC values between baseline and follow-up. The increase in ADC surrounding TMBs was similar to the increases seen with IPH and subarachnoid hemorrhage (SAH). In IPH and SAH, extravasated blood triggers secondary injury through a cascade of endothelial swelling, vasospasm,

vasoconstriction, and evolution toward vasogenic edema in the subacute phase of injury.^{22,30,31}

Our results indicate that primary traumatic vascular injury can be associated with cytotoxic edema evolving over time into vasogenic edema. While further investigation is needed to determine the underlying mechanisms of these changes, the patterns of ADC evolution appear strikingly similar to the changes seen on acute and subacute MRI in ischemic stroke.

The ability to detect ADC hypointense (i.e., cytotoxic) lesions after TBI also depends on the timing of acute MRI. The earliest scans detected VOIs with the lowest ADC values, consistent with the hypothesis that cytotoxic edema predominates in these VOIs acutely after injury, while vasogenic edema becomes more prevalent with time. This reinforces the importance of obtaining acute and follow-up scans after injury and has implications for therapeutic interventions targeting the vasculature. Cytotoxic edema early after injury typically results in irreversible cell death unless energy substrates can be restored. Conversely, vasogenic edema is, in principle, reversible when detected early enough to administer treatments to restore blood–brain barrier function and integrity. Left untreated, however, cytotoxic edema may lead to secondary injury and cell death.

Previous studies of diffusion-related changes in TBI were conducted in both animal models and humans, with conflicting results. Animal models showed both overall increases³² and decreases³³ in ADC acutely after injury, while other studies reported a progressive decrease from the acute to subacute period.³⁴ Early studies of DWI MRI in human TBI attributed decreased ADC values between one and 18 days post-injury¹⁶ and reduced white matter diffusion anisotropy within 24 h of injury³⁵ to DAI.

Another investigation of early (<14 day) DWI lesions in the context of DAI found a mixture of lesions that could be differentiated as either increased or decreased diffusion, reflecting vasogenic and cytotoxic edema, respectively.³⁶ A group classifying lesion load as either vasogenic or cytotoxic found that both edema types were associated with poor outcome, despite more severe injury in those with predominantly cytotoxic edema, and saw both cytotoxic and vasogenic lesions in the corpus callosum, an area susceptible to DAI.³⁷ An early serial MRI study and a case report suggest an association between temporal resolution of edema-associated DAI lesions of the corpus callosum and better outcomes.^{38,39}

Several studies examined whole-brain and regional ADC values as predictors of outcome after TBI. Whole-brain ADC values on MRI at 1–23 days post-injury significantly correlated with admission GCS scores,⁴⁰ while others reported significant correlations between injury severity and ADC values of normal-appearing white matter.⁴¹ Another group reported lower ADC values in deep white and gray matter in patients with unfavorable outcome compared with those with favorable outcome,

with all regions except peripheral gray matter showing an increase in ADC relative to controls.⁴²

In contrast, pediatric TBI studies reported reduced ADC values in peripheral white matter predicted poor outcome in severe TBI, while deep white matter ADC values were reduced relative to controls.⁴³ Paradoxically, a subset of this pediatric population excluding the most severely injured patients demonstrated that increased ADC values in peripheral white and gray matter predicted worse long-term neurocognitive outcomes.⁴⁴ Although maturational changes might complicate interpretation of diffusion-related MRI outcomes between adults and children, these divergent findings highlight the difficulty of using early ADC values to predict outcome in TBI.

The TRACK-TBI study reported that contusion and hemorrhagic axonal injury⁴⁵ and decreases in fractional anisotropy⁴⁶ on MRI at two weeks after injury predict poorer GOS-E scores at three months post-injury. Our finding that hemorrhage-associated cytotoxic edema on acute MRI predicts outcome is consistent with those results and suggests there may be therapeutic targets for intervention. Once the biological mechanisms are better understood, future targeted therapies could be administered to those patients presenting with acute hemorrhage and diffusion changes.

Approximately 50% of those presenting to two civilian emergency departments with criteria for mild TBI were not initially recognized as having TBI.⁴⁷ While in sports and the military implementation of standard screening protocols has improved identification of patients, in the general population even many of those recognized as having mild TBI with ongoing symptoms do not receive appropriate education and follow-up care.⁴⁸ This lack of early identification and education, except in the most severely impaired, may contribute to worse outcomes.⁴⁹

Patients found to have small amounts of hemorrhage on CT are more likely to have worse outcome (defined by GOS-E) if evidence of diffusion changes are found on MRI. This prognostic value, in itself, may provide evidence needed to support the clinical utility of MRI in select patients. In the present study, TMBs were the most common finding seen on acute MRI that was not detected on acute CT. This, combined with our previous work demonstrating poorer outcomes in patients with TMBs,¹⁴ suggests that acute MRI may be warranted to screen for microbleeds in CT negative patients, although additional research is needed. Further, there may be targeted therapies that could, one day, be applied to reduce the risk of edema, or, at a minimum, to flag these patients as being potentially in need of early education, connection with outpatient care, and possibly ongoing follow-up to assess their need for rehabilitation services.

In this study, we examined ADC changes in conspicuous DWI lesions during initial hospitalization as opposed to during a subsequent outpatient visit. This allowed us to confirm that: (i) acute ADC values can be used to predict

outcome, and (ii) acute cytotoxic edema associated with lesions such as TMB and IPH may contribute to poor outcome. We also characterized the temporal evolution of ADC values associated with hemorrhage by comparing baseline and follow-up MRI, demonstrating a pattern of ADC evolution that indicates lesion evolution between the acute and subacute periods.

Limitations

Our findings must be interpreted in comparison with the literature by considering the time course of MRI used in the present study, with a median time to baseline MRI of 21 h far earlier than previously described studies, possibly accounting for the ADC evolution differences. There is also the potential for selection bias in our patient cohort, with patients being more likely to return for follow-up visits if they had clinical findings or symptoms.

Cranial surgery was not an exclusion criterion for this study and may have affected outcomes, although GOS-E outcomes were split for the two patients who underwent cranial surgery during their acute visits. Patients with polytrauma were included in this dataset and outcomes analysis, and an association between polytrauma and poorer GOS-E outcomes has been reported previously.⁵⁰ In the present study, patients were asked to attribute their GOS-E answers in the context of their head injury, but difficulties in their doing so may have confounded results.

Whether MRI is quantifiably superior to CT for the detection of TBI-related lesions such as contusions and TMBs is not directly addressed in the present study and is the focus of ongoing research. Additional studies of patients with acute MRI with larger sample sizes are needed to confirm and extend these findings. Further investigation is also required to determine which of the potential underlying biological mechanisms are causing these acute ADC changes.

Conclusion

An emerging body of literature indicates that traumatic vascular injury is associated with chronic pathophysiological processes and poor neurological outcome. The current study, by identifying acute injury processes associated with intracranial hemorrhage on clinically relevant *in vivo* acute and subacute MRI, supports the hypothesis that traumatic vascular injury is a key contributor to the pathogenesis of TBI.

Acknowledgments

We would like to thank our patients and their families for their willingness to participate in clinical research. We are also grateful for the clinical and administrative teams who have supported the THINC study.

Authors' Contributions

L.C.T., M.L., R.P.H.B., G.P., and L.L.L. contributed to the study conceptualization and design. N.J., A.G., and D.G.

participated in the data collection. L.C.T., R.P.H.B., G.P., and L.L.L. provided neurological and/or neuroimaging expertise. L.C.T., M.L., N.J., A.G., D.G., N.P., and L.L.L. analyzed and interpreted the data. L.C.T., M.L., N.J., D.G., N.P., M.W., and L.L.L. drafted the manuscript and/or figures. All authors critically revised the manuscript.

Funding Information

Funding for this work was provided by Intramural Research Program of the National Institute of Neurological Disorders and Stroke, National Institutes of Health, and by the Department of Defense through the Center for Neuroscience and Regenerative Medicine.

Author Disclosure Statement

No competing financial interests exist.

Supplementary Material

Supplementary Methods
Supplementary Table S1
Supplementary Table S2

References

- Winkler, E.A., Minter, D., Yue, J.K., and Manley, G.T. (2016). Cerebral edema in traumatic brain injury: pathophysiology and prospective therapeutic targets. *Neurosurg. Clin. N. Am.* 27, 473–488.
- Jha, R.M., Kochanek, P.M., and Simard, J.M. (2019). Pathophysiology and treatment of cerebral edema in traumatic brain injury. *Neuropharmacology* 145, 230–246.
- Bramlett, H.M., and Dietrich, W.D. (2004). Pathophysiology of cerebral ischemia and brain trauma: similarities and differences. *J. Cereb. Blood Flow Metab.* 24, 133–150.
- Ito, J., Marmarou, A., Barzo, P., Fatouros, P., and Corwin, F. (1996). Characterization of edema by diffusion-weighted imaging in experimental traumatic brain injury. *J. Neurosurg.* 84, 97–103.
- Schlaug, G., Siewert, B., Benfield, A., Edelman, R.R., and Warach, S. (1997). Time course of the apparent diffusion coefficient (ADC) abnormality in human stroke. *Neurology* 49, 113–119.
- Lansberg, M.G., Thijs, V.N., O'Brien, M.W., Ali, J.O., de Crespigny, A.J., Tong, D.C., Moseley, M.E., and Albers, G.W. (2001). Evolution of apparent diffusion coefficient, diffusion-weighted, and T2-weighted signal intensity of acute stroke. *AJNR Am J Neuroradiol* 22, 637–644.
- van Everdingen, K.J., van der Grond, J., Kappelle, L.J., Ramos, L.M., and Mali, W.P. (1998). Diffusion-weighted magnetic resonance imaging in acute stroke. *Stroke* 29, 1783–1790.
- Nagesh, V., Welch, K.M., Windham, J.P., Patel, S., Levine, S.R., Hearshen, D., Peck, D., Robbins, K., D'Oliveira, L., Soltanian-Zadeh, H., and Boska, M.D. (1998). Time course of ADCw changes in ischemic stroke: beyond the human eye! *Stroke* 29, 1778–1782.
- Hsia, A.W., Luby, M., Cullison, K., Burton, S., Armonda, R., Liu, A.H., Leigh, R., Nadareishvili, Z., Benson, R.T., Lynch, J.K., and Latour, L.L. (2019). Rapid apparent diffusion coefficient evolution after early revascularization. *Stroke* 50, 2086–2092.
- Provenzale, J.M., and Sorensen, A.G. (1999). Diffusion-weighted MR imaging in acute stroke: theoretic considerations and clinical applications. *AJR Am. J. Roentgenol.* 173, 1459–1467.
- Muir, K.W., Buchan, A., von Kummer, R., Rother, J., and Baron, J.C. (2006). Imaging of acute stroke. *Lancet Neurol.* 5, 755–768.
- Gonzalez, R.G. (2012). Clinical MRI of acute ischemic stroke. *J. Magn. Reson. Imaging* 36, 259–271.
- Chiara Ricciardi, M., Bokkers, R.P., Butman, J.A., Hammoud, D.A., Pham, D.L., Warach, S., and Latour, L.L. (2017). Trauma-specific brain abnormalities in suspected mild traumatic brain injury patients identified in the first 48 hours after injury: a blinded magnetic resonance imaging comparative study including suspected acute minor stroke patients. *J. Neurotrauma* 34, 23–30.
- Griffin, A.D., Turtzo, L.C., Parikh, G.Y., Tolpygo, A., Lodato, Z., Moses, A.D., Nair, G., Perl, D.P., Edwards, N.A., Dardzinski, B.J., Armstrong, R.C., Ray-Chaudhury, A., Mitra, P.P., and Latour, L.L. (2019). Traumatic

- microbleeds suggest vascular injury and predict disability in traumatic brain injury. *Brain* 142, 3550–3564.
15. Scheid, R., Preul, C., Gruber, O., Wiggins, C., and von Cramon, D.Y. (2003). Diffuse axonal injury associated with chronic traumatic brain injury: evidence from T2*-weighted gradient-echo imaging at 3 T. *AJNR Am. J. Neuroradiol.* 24, 1049–1056.
  16. Liu, A.Y., Maldjian, J.A., Bagley, L.J., Sinson, G.P., and Grossman, R.I. (1999). Traumatic brain injury: diffusion-weighted MR imaging findings. *AJNR Am. J. Neuroradiol.* 20, 1636–1641.
  17. Liu, J., Kou, Z., and Tian, Y. (2014). Diffuse axonal injury after traumatic cerebral microbleeds: an evaluation of imaging techniques. *Neural Regen. Res.* 9, 1222–1230.
  18. Kinoshita, T., Moritani, T., Hiwatahi, A., Wang, H.Z., Shrier, D.A., Numa-guchi, Y., and Westesson, P.L. (2005). Conspicuity of diffuse axonal injury lesions on diffusion-weighted MR imaging. *Eur. J. Radiol.* 56, 5–11.
  19. Hahnel, S., Stippich, C., Weber, I., Darm, H., Schill, T., Jost, J., Friedmann, B., Heiland, S., Blatow, M., and Meyding-Lamade, U. (2008). Prevalence of cerebral microhemorrhages in amateur boxers as detected by 3T MR imaging. *AJNR Am. J. Neuroradiol.* 29, 388–391.
  20. Launey, Y., Fryer, T.D., Hong, Y.T., Steiner, L.A., Nortje, J., Veenith, T.V., Hutchinson, P.J., Ercole, A., Gupta, A.K., Aigbirio, F.I., Pickard, J.D., Coles, J.P., and Menon, D.K. (2020). Spatial and temporal pattern of ischemia and abnormal vascular function following traumatic brain injury. *JAMA Neurol.* 77, 339–349.
  21. Ironside, N., Chen, C.J., Ding, D., Mayer, S.A., and Connolly, E.S., Jr. (2019). Perihematomal edema after spontaneous intracerebral hemorrhage. *Stroke* 50, 1626–1633.
  22. Kurland, D., Hong, C., Aarabi, B., Gerzanich, V., and Simard, J.M. (2012). Hemorrhagic progression of a contusion after traumatic brain injury: a review. *J. Neurotrauma* 29, 19–31.
  23. Husiman, T.A.G.M., Schwamm, H., Schaefer, P.W., Koroshetz, W.J., Shetty-Alva, N., Ozsunar, Y., Wu, O., and Sorensen, A.G. (2004). Diffusion tensor imaging as potential biomarker of white matter injury in diffuse axonal injury. *AJNR Am. J. Neuroradiol.* 25, 370–376.
  24. Spitz, G., Alway, Y., Gould, K.R., and Ponsford, J.L. (2017). Disrupted white matter microstructure and mood disorders after traumatic brain injury. *J. Neurotrauma* 34, 807–815.
  25. van Eijck, M.M., Schoonman, G.G., van der Naalt, J., de Vries, J., and Roks, G. (2018). Diffuse axonal injury after traumatic brain injury is a prognostic factor for functional outcome: a systematic review and meta-analysis. *Brain Inj.* 32, 395–402.
  26. Johnson, V.E., Stewart, W., and Smith, D.H. (2013). Axonal pathology in traumatic brain injury. *Exp. Neurol.* 246, 35–43.
  27. Adams, J.H., Graham, D., Scott, G., Parker, L.S., and Doyle, D. (1980). Brain damage in fatal non-missile head injury. *J. Clin. Pathol.* 33, 1132–1145.
  28. Andreasen, S.H., Andersen, K.W., Conde, V., Dyrby, T.B., Puonti, O., Kammergaard, L.P., Madsen, C.G., Madsen, K.H., Poulsen, I., and Siebner, H.R. (2020). Limited colocalization of microbleeds and microstructural changes after severe traumatic brain injury. *J. Neurotrauma* 37, 581–592.
  29. Jolly, A.E., Balaet, M., Azor, A., Friedland, D., Sandrone, S., Graham, N.S.N., Zimmerman, K., and Sharp, D.J. (2021). Detecting axonal injury in individual patients after traumatic brain injury. *Brain* 144, 92–113.
  30. Francoeur, C.L., and Mayer, S.A. (2016). Management of delayed cerebral ischemia after subarachnoid hemorrhage. *Crit. Care* 20, 277.
  31. Budohoski, K.P., Guilfoyle, M., Helmy, A., Huuskonen, T., Czosnyka, M., Kirollos, R., Menon, D.K., Pickard, J.D., and Kirkpatrick, P.J. (2014). The pathophysiology and treatment of delayed cerebral ischaemia following subarachnoid haemorrhage. *J. Neurol. Neurosurg. Psychiatry* 85, 1343–1353.
  32. Hanstock, C.C., Faden, A.I., Bendall, M.R., and Vink, R. (1994). Diffusion-weighted imaging differentiates ischemic tissue from traumatized tissue. *Stroke* 25, 843–848.
  33. Alsop, D.C., Murai, H., Detre, J.A., McIntosh, T.K., and Smith, D.H. (1996). Detection of acute pathologic changes following experimental traumatic brain injury using diffusion-weighted magnetic resonance imaging. *J. Neurotrauma* 13, 515–521.
  34. Barzo, P., Marmarou, A., Fatouros, P., Hayasaka, K., and Corwin, F. (1997). Contribution of vasogenic and cellular edema to traumatic brain swelling measured by diffusion-weighted imaging. *J. Neurosurg.* 87, 900–907.
  35. Arfanakis, K., Haughton, V.M., Carew, J.D., Rogers, B.P., Dempsey, R.J., and Meyerand, M.E. (2002). Diffusion tensor MR imaging in diffuse axonal injury. *AJNR Am. J. Neuroradiol.* 23, 794–802.
  36. Hergan, K., Schaefer, P.W., Sorensen, A.G., Gonzalez, R.G., and Huisman, T.A. (2002). Diffusion-weighted MRI in diffuse axonal injury of the brain. *Eur. Radiol.* 12, 2536–2541.
  37. Hudak, A.M., Peng, L., Marquez de la Plata, C., Thottakara, J., Moore, C., Harper, C., McColl, R., Babcock, E., and Diaz-Arrastia, R. (2014). Cytotoxic and vasogenic cerebral oedema in traumatic brain injury: assessment with FLAIR and DWI imaging. *Brain Inj.* 28, 1602–1609.
  38. Tokutomi, T., Hirohata, M., Miyagi, T., Abe, T., and Shigemori, M. (1997). Posttraumatic edema in the corpus callosum shown by MRI. *Brain Edema* X 70, 80–83.
  39. Takayama, H., Kobayashi, M., Sugishita, M., and Mihara, B. (2000). Diffusion-weighted imaging demonstrates transient cytotoxic edema involving the corpus callosum in a patient with diffuse brain injury. *Clin. Neurol. Neurosurg.* 102, 135–139.
  40. Shanmuganathan, K., Gullapalli, R.P., Mirvis, S.E., Roys, S., and Murthy, P. (2004). Whole-brain apparent diffusion coefficient in traumatic brain injury: correlation with Glasgow Coma Scale score. *AJNR Am. J. Neuroradiol.* 25, 539–544.
  41. Goetz, P., Blamire, A., Rajagopalan, B., Cadoux-Hudson, T., Young, D., and Styles, P. (2004). Increase in apparent diffusion coefficient in normal appearing white matter following human traumatic brain injury correlates with injury severity. *J. Neurotrauma* 21, 645–654.
  42. Hou, D.J., Tong, K.A., Ashwal, S., Oyoyo, U., Joo, E., Shutter, L., and Obenaus, A. (2007). Diffusion-weighted magnetic resonance imaging improves outcome prediction in adult traumatic brain injury. *J. Neurotrauma* 24, 1558–1569.
  43. Galloway, N.R., Tong, K.A., Ashwal, S., Oyoyo, U., and Obenaus, A. (2008). Diffusion-weighted imaging improves outcome prediction in pediatric traumatic brain injury. *J. Neurotrauma* 25, 1153–1162.
  44. Babikian, T., Tong, K.A., Galloway, N.R., Freier-Randall, M.C., Obenaus, A., and Ashwal, S. (2009). Diffusion-weighted imaging predicts cognition in pediatric brain injury. *Pediatr. Neurol.* 41, 406–412.
  45. Yuh, E.L., Mukherjee, P., Lingsma, H.F., Yue, J.K., Ferguson, A.R., Gordon, W.A., Valadka, A.B., Schnyer, D.M., Okonkwo, D.O., Maas, A.I., Manley, G.T., and TRACK-TBI Investigators. (2013). Magnetic resonance imaging improves 3-month outcome prediction in mild traumatic brain injury. *Ann. Neurol.* 73, 224–235.
  46. Yuh, E.L., Cooper, S.R., Mukherjee, P., Yue, J.K., Lingsma, H.F., Gordon, W.A., Valadka, A.B., Okonkwo, D.O., Schnyer, D.M., Vassar, M.J., Maas, A.I., Manley, G.T., and TRACK-TBI Investigators. (2014). Diffusion tensor imaging for outcome prediction in mild traumatic brain injury: a TRACK-TBI study. *J. Neurotrauma* 31, 1457–1477.
  47. Cota, M.R., Moses, A.D., Jikaria, N.R., Bittner, K.C., Diaz-Arrastia, R.R., Latour, L.L., and Turtzo, L.C. (2019). Discordance between documented criteria and documented diagnosis of traumatic brain injury in the emergency department. *J. Neurotrauma* 36, 1335–1342.
  48. Seabury, S.A., Gaudette, E., Goldman, D.P., Markowitz, A.J., Brooks, J., McCrea, M.A., Okonkwo, D.O., Manley, G.T., Adeoye, O., Badjatia, N., Boase, K., Bodien, Y., Bullock, M.R., Chesnut, R., Corrigan, J.D., Crawford, K., Diaz-Arrastia, R., Dikmen, S., Duhaime, A.C., Ellenbogen, R., Feeser, V.R., Ferguson, A., Foreman, B., Gardner, R., Giacino, J., Gonzalez, L., Gopinath, S., Gullapalli, R., Hemphill, J.C., Hotz, G., Jain, S., Korley, F., Kramer, J., Kreitzer, N., Levin, H., Lindsell, C., Machamer, J., Madden, C., Martin, A., McAllister, T., Merchant, R., Mukherjee, P., Nelson, L., Noel, F., Palacios, E., Perl, D., Puccio, A., Rabinowitz, M., Robertson, C., Rosand, J., Sander, A., Satris, G., Schnyer, D., Stein, M., Taylor, S., Temkin, N., Toga, A., Valadka, A., Vassar, M., Vespa, P., Wang, K., Yue, J., Yuh, E., and Zafonte, R. (2018). Assessment of follow-up care after emergency department presentation for mild traumatic brain injury and concussion. *JAMA Network Open* 1.
  49. Nelson, L.D., Temkin, N.R., Dikmen, S., Barber, J., Giacino, J.T., Yuh, E., Levin, H.S., McCrea, M.A., Stein, M.B., Mukherjee, P., Okonkwo, D.O., Robertson, C.S., Diaz-Arrastia, R., Manley, G.T., and the T-TBI.I., Adeoye, O., Badjatia, N., Boase, K., Bodien, Y., Bullock, M.R., Chesnut, R., Corrigan, J.D., Crawford, K., Duhaime, A.C., Ellenbogen, R., Feeser, V.R., Ferguson, A., Foreman, B., Gardner, R., Gaudette, E., Gonzalez, L., Gopinath, S., Gullapalli, R., Hemphill, J.C., Hotz, G., Jain, S., Korley, F., Kramer, J., Kreitzer, N., Lindsell, C., Machamer, J., Madden, C., Martin, A., McAllister, T., Merchant, R., Noel, F., Palacios, E., Perl, D., Puccio, A., Rabinowitz, M., Rosand, J., Sander, A., Satris, G., Schnyer, D., Seabury, S., Sherer, M., Taylor, S., Toga, A., Valadka, A., Vassar, M.J., Vespa, P., Wang, K., Yue, J.K., and Zafonte, R. (2019). Recovery after mild traumatic brain injury in patients presenting to US Level I trauma Centers: a Transforming Research and Clinical Knowledge in Traumatic Brain Injury (TRACK-TBI) study. *JAMA Neurol.* 76, 1049–1059.
  50. Yue, J.K., Satris, G.G., Dalle Ore, C.L., Huie, J.R., Deng, H., Winkler, E.A., Lee, Y.M., Vassar, M.J., Taylor, S.R., Schnyer, D.M., Lingsma, H.F., Puccio, A.M., Yuh, E.L., Mukherjee, P., Valadka, A.B., Ferguson, A.R., Markowitz, A.J., Okonkwo, D.O., Manley, G.T., and the TRACK-TBI Investigators. (2020). Polytrauma is associated with increased three- and six-month disability after traumatic brain injury: a TRACK-TBI pilot study. *Neurotrauma Rep.* 1, 32–41.