University of Groningen

Does the impaired postural control in Parkinson's disease affect the habituation to non-sequential external perturbation trials?

Spiandor Beretta, Victor; Carpenter, Mark Gregory; Barbieri, Fabio Augusto; Rocha Santos, Paulo Cezar; Orcioli-Silva, Diego; Pereira, Marcelo Pinto; Bucken Gobbi, Lilian Teresa

Published in:
Clinical Biomechanics

DOI:
[10.1016/j.clinbiomech.2021.105363](https://doi.org/10.1016/j.clinbiomech.2021.105363)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Spiandor Beretta, V., Carpenter, M. G., Barbieri, F. A., Rocha Santos, P. C., Orcioli-Silva, D., Pereira, M. P., & Bucken Gobbi, L. T. (2021). Does the impaired postural control in Parkinson's disease affect the habituation to non-sequential external perturbation trials? *Clinical Biomechanics*, 85, [105363]. <https://doi.org/10.1016/j.clinbiomech.2021.105363>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Does the impaired postural control in Parkinson's disease affect the habituation to non-sequential external perturbation trials?

Victor Spiandor Beretta^a, Mark Gregory Carpenter^b, Fabio Augusto Barbieri^c,
Paulo Cezar Rocha Santos^{a,d}, Diego Orcioli-Silva^a, Marcelo Pinto Pereira^a,
Lilian Teresa Bucken Gobbi^{a,*}

^a São Paulo State University (Unesp), Institute of Biosciences, Graduate Program in Movement Sciences, Posture and Gait Studies Laboratory (LEPLO), Rio Claro, Brazil

^b School of Kinesiology, The University of British Columbia, Vancouver, British Columbia, Canada

^c São Paulo State University (Unesp), School of Sciences, Graduate Program in Movement Sciences, Human Movement Research Laboratory (MOVI-LAB), Bauru, Brazil

^d Center for Human Movement Sciences, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands

ARTICLE INFO

Keywords:

Movement disorders
Balance control
Adaptation
Support-base translation
Center of pressure

ABSTRACT

Background: How people with Parkinson's disease habituate their postural response to unpredictable translation perturbation is not totally understood. We compared the capacity to change the postural responses after unexpected external perturbation and investigated the habituation plateaus of postural responses to non-sequential perturbation trials in people with Parkinson's disease and healthy older adults.

Methods: In people with Parkinson's disease ($n = 37$) and older adults ($n = 20$), sudden posterior support-surface translational were applied in 7 out of 17 randomized trials to ensure perturbation unpredictability. Electromyography and center of pressure parameters of postural response were analyzed by ANOVAs (Group vs. Trials). Two simple planned contrasts were performed to determine at which trial the responses first significantly habituate, and by which trials the habituation plateaus.

Findings: Older adults demonstrated a first response change in trial 5 and habituation plateaus after trial 4, while for people with Parkinson's disease, the first change occurred in trial 2 and habituation plateau after trial 5 observed by center of pressure range. People with Parkinson's disease demonstrated a greater center of pressure range in trial 1 compared to older adults. Independent of trial, people with Parkinson's disease vs. older adults demonstrated a greater ankle muscle co-activation and recovery time.

Interpretation: Despite the greater center of pressure range in the first trial, people with Parkinson's disease can habituate to unpredictable perturbations. This is reflected by little, to no difference in the time-course of adaptation for all but 2 parameters that showed only marginal differences between people with Parkinson's disease and older adults.

1. Introduction

Adequate postural response to a balance perturbation is important to avoid falls and involves both predictive and reactive neuromuscular responses (Horak et al., 2005, 1997; Mochizuki et al., 2008). Reactive balance responses are triggered after the perturbation onset, and are modulated by perturbation characteristics as well as the individual's central set (Horak et al., 1997; Mochizuki et al., 2008). Due to the neurodegenerative process, Parkinson's disease (PD) affects the excitatory/inhibitory cortical control impairing reactive postural responses

(Park et al., 2015). The most remarkable postural changes in people with PD vs. healthy controls are greater displacement of center of mass (CoM) and pressure (CoP) (Horak et al., 2005), which are likely related to an abnormal modulation of muscle response (Dimitrova et al., 2004a; Horak et al., 2005), reflecting excessive activity of antagonistic muscles and muscle co-activation (Bloem, 1992; Carpenter et al., 2004; Dimitrova et al., 2004b; Horak et al., 1992; Lang et al., 2019).

Impaired neuromuscular control in PD affects the ability to adapt/habituate predictive and reactive postural responses to repetitive postural perturbations (Nanhoe-Mahabier et al., 2012; Smith et al.,

* Corresponding author at: São Paulo State University (Unesp), Institute of Biosciences, Posture and Gait Studies Laboratory (LEPLO), avenida 24-A, 1515, Bela Vista, 13506-900 Rio Claro/SP, Brazil.

E-mail address: lilian.gobbi@unesp.br (L.T.B. Gobbi).

<https://doi.org/10.1016/j.clinbiomech.2021.105363>

Received 30 June 2020; Accepted 19 April 2021

Available online 22 April 2021

0268-0033/© 2021 Elsevier Ltd. All rights reserved.

2012). The habituation to repetitive perturbations indicates a versatility marker of motor control (Oude Nijhuis et al., 2009; Van Ooteghem et al., 2017), which reflects the capacity of an individual to identify and cope with changing circumstances in an optimal and safe strategy (Chong et al., 2000, 1999; Horak et al., 1997). In postural control studies, habituation is observed as decreased muscle activity and CoM/CoP displacements by repetitive exposure to stimuli (Bloem et al., 1998; Nanhoe-Mahabier et al., 2012).

The current knowledge of habituation to postural perturbation in PD is inconsistent (Bloem et al., 1998; Nanhoe-Mahabier et al., 2012; Visser et al., 2010). While data suggest that people with PD habituate the postural responses (decrease medial gastrocnemius - MG - muscle amplitude) after ~3 trials vs. no habituation in older adults (Bloem et al., 1998), other evidence suggests a similar level of postural habituation (decreases in soleus amplitude and trunk flexion after 8 repeated perturbations) in both PD and older adults (Visser et al., 2010). On the contrary, a delay of the postural response habituation in PD compared with older adults (trials 5 vs. 2, respectively) was observed by a decrease in the CoM displacement (Nanhoe-Mahabier et al., 2012). This delay of the postural response habituation to perturbation in PD is reasonable considering the neurodegenerative process, which affects the capacity to identify relevant stimuli and generate adequate responses based on prior experience (Horak et al., 1997; Nanhoe-Mahabier et al., 2012). In addition to those inconsistent results, prior perturbations were induced by rotation of support-base and were repeated in sequence (Bloem et al., 1998; Nanhoe-Mahabier et al., 2012) or under different conditions of the support-base in the same experiment (translation and rotation) (Visser et al., 2010). It is conceivable that continuous repetition of perturbations would affect the habituation of the postural responses. To the best of our knowledge, it has not yet been examined whether PD can affect the habituation responses to unpredictable and non-sequential perturbation trials, situations more commonly experienced during daily life tasks. Therefore, we aimed (1) to investigate the capacity to change the reactive postural responses after unexpected external perturbation between people with PD and healthy older adults (CG), and (2) to investigate the habituation plateaus of postural responses to non-sequential trials of translations of the support-surface in both people with PD and CG. We hypothesized that both groups habituate the electromyographic (EMG) and CoP parameters, but CG vs. PD would present a more rapid change after unexpected external perturbation and habituation plateaus in fewer trials. We also expected that CG would demonstrate a lower EMG and CoP activity in reactive postural response (demonstrating better balance recovery) (Nanhoe-Mahabier et al., 2012; Visser et al., 2010). These postural changes would be expected mainly by a decrease in the range of CoP (to avoid the CoP positioning close to the limits of stability).

2. Methods

2.1. Participants

Thirty-seven people with PD (PDG) (diagnoses based on criteria determined by the UK Brain Bank) and twenty CG participated in this study. All subjects had not participated previously in other studies that induced displacement of the support base.

We included people with PD in Hoehn & Yahr scale (H&Y) scores <3, and all individuals over 60 years of age. The exclusion criteria were: (i) cognitive decline (Mini-Mental State Examination – MMSE <24; (Brucki et al., 2003)); (ii) use of any medication that causes side effects in the balance; (iii) orthopedic, musculoskeletal and/or visual impairments that would affect the performance of the protocol; (iv) presence of any uncontrolled disease that could affect peripheral sensory function (e.g., diabetes); and (v) presence of labyrinthitis. The clinical and postural control evaluations were performed in the “ON” state of the specific PD medication (approximately 1 h after medication intake). The participants signed an informed consent form approved by the research ethics

committee of the São Paulo State University at Rio Claro–Brazil (CAAE:52534316.1.0000.5465).

2.2. Clinical evaluation

PDG was assessed by an experienced researcher using the Unified Parkinson's Disease Rating Scale (UPDRS; Fahn et al., 1987) and the H&Y scale adapted version (Schenkman et al., 2001). Levodopa Equivalent Dose (LED) was calculated (Tomlinson et al., 2010).

2.3. Postural control evaluation

Participants wore a safety harness and stood with feet side-by-side on a force plate fixed to a translating platform. Participants stood for 17 trials each of 20 s duration, with a resting period of 60 s between each trial to allow the participants to move and avoid prolonged periods of static standing. To guarantee the perturbations were unpredictable, the subjects experienced perturbations (15 cm/s of velocity and 5 cm of displacement in the posterior direction) in only 7 out of the 17 randomized trials; the remaining 10 trials were control trials where no perturbation was experienced. The number of trials with perturbation was chosen because previous studies indicated that the habituation plateaus in people with PD occurs within the first 7 trials (Nanhoe-Mahabier et al., 2012). In these trials, the perturbation was triggered at a random time within 5 to 15 s period of the 20 s stance trial.

2.4. Experimental apparatus and procedures

A Trigno™ Wireless System (Delsys, Inc. Boston, MA, USA – 2000 Hz) and a force plate (AccuGait, Advanced Mechanical Technologies, Boston, MA, USA – 200 Hz) were used to acquire the EMG and CoP parameters, respectively. Trigno sensors were positioned on the biceps femoris (BF), vastus medialis (VM), MG and tibialis anterior (TA) of the dominant lower limb for the posture (CG) and of the most affected limb (PDG – determined by UPDRS items) (Barbieri et al., 2016; Beretta et al., 2015) following the SENIAM recommendations (Hermens et al., 2000).

2.5. Data analysis

The perturbation onset and the EMG were synchronized through an accelerometer (Trigno™ Wireless System – Delsys, Inc. Boston, MA, USA – 148.15 Hz) positioned in the force plate by a time vector. The perturbation start time was determined by the moment at which the acceleration was greater than the mean plus two standard deviations of the baseline.

2.5.1. Analysis of EMG parameters

EMG signals were band-pass filtered (20–450 Hz), rectified, and low-pass filtered (50 Hz) using a dual-pass Butterworth filter. In addition, the system had a rejection rate of 80 db, and the resolution of the A/D board was 16 bits. The onset of muscle activity corresponded to the first point where the value was greater than the mean + 2 standard deviations of the baseline (750 and 500 ms before the perturbation) by at least 30 ms (Cleworth et al., 2016). All points of interest were determined by a semi-automatic algorithm (Cleworth et al., 2016). Temporal aspects of the EMG were measured as the onset latency and the time to peak (TTP) of BF and MG (Cleworth et al., 2016; de Freitas et al., 2010). To analyze the EMG amplitude, the magnitude of muscle activation was calculated by the area below the curve (integral - iEMG) for the reactive period, measured over a 250 ms time-window after the muscle activity onset (de Freitas et al., 2010; Santos et al., 2010) (Fig. 1-a). The iEMG was normalized by the iEMG baseline ($100 + (100 * ((iEMG_{\text{reactive period}} - iEMG_{\text{baseline}}) / iEMG_{\text{baseline}}))$) (de Freitas et al., 2010). After the normalization, the co-activation of BF/VM and MG/TA pairs were determined through the ratio $((iEMG_{\text{antagonist}} / iEMG_{\text{agonist}}) * 100)$ (de Freitas et al., 2010) in the reactive period.

Fig. 1. Illustration of interest points in the analysis. The figure represents the behavior of a PDG participant in the same trial: a) EMG analysis of MG activity; b) CoP analysis.

2.5.2. CoP analysis

The CoP onset was determined when the CoP displacement was greater than the mean + 2 standard deviations relative to the baseline. Reactive postural responses were analyzed for a period between the CoP onset and 700 ms post-perturbation (Fig. 1-b). In the reactive period, we analyzed the range and the peak of CoP in the anterior-posterior direction. Temporal aspects of the CoP were measured as the TTP and the recovery time to a stable position. The recovery time was defined as the time interval between CoP onset and the instant when the CoP variability (standard deviation during one second) after perturbation was less than or equal to the CoP variability at baseline (determined one second before perturbation) (Beretta et al., 2019).

2.6. Statistical analysis

The statistical analysis was performed using SPSS 21.0. Demographic characteristics were compared through Student's t-test. Analysis of reactive postural response was determined by a two-way ANOVA with Group (PDG x CG) and Trial (1x2x3x4x5x6x7) as factor. Trial main effects or Group by Trial interactions were followed up post-hoc using simple planned contrasts. Contrast 1 was performed to determine the earliest trial where significant changes relative to the first trial were observed (Trial 1 vs. 2–7). Contrast 2 was performed to analyze the moment when individuals reached the habituation plateaus, after which no further significant changes compared to the last trial were observed (Trial 7 vs. Trials 2–6). Post-hoc comparisons using Student's t-test were also applied to significant Group by Trial interactions to compare group differences for Trial 1. Significance was set at $P < 0.05$.

3. Results

3.1. Participants' characteristics

Statistical analysis indicated that both groups presented similar demographic and global cognition characteristics (Table 1).

3.2. Reactive postural response

3.2.1. Temporal parameters

ANOVA revealed a Trial main effect for the recovery time ($F_{(6,294)} = 9.619$, $P < 0.001$, $\eta_p^2 = 0.164$) (Fig. 2-a), onset latency of BF ($F_{(6,324)} = 7.729$, $P < 0.001$, $\eta_p^2 = 0.125$) (Fig. 2-b) and MG muscles ($F_{(6,318)} = 5.071$, $P < 0.001$, $\eta_p^2 = 0.087$) (Fig. 2-c), TTP of CoP ($F_{(6,306)} = 25.314$, P

Table 1

Mean and standard deviations of participants' characteristics.

	PDG (n = 37)	CG (n = 20)	P-value
Sex (Male/Female)	17/20	8/12	0.666
Age (years)	70.89 ± 8.36	70.50 ± 5.35	0.831
Body Weight (kg)	70.76 ± 11.69	70.56 ± 10.48	0.949
Body Height (cm)	161.64 ± 9.37	162.19 ± 6.92	0.816
MMSE (0–30)	27.75 ± 2.05	28.65 ± 1.63	0.098
UPDRS III (0–108)	25.41 ± 10.28		
H&Y (stage)	2.0(1–3)		
LED (mg/day)	456.29 ± 318.99		
PD duration (years)	5.03 ± 3.60		

UPDRS = Unified Parkinson's Disease Rating Scale; H&Y = Hoehn & Yahr; MMSE = Mini-Mental State Examination; LED = Levodopa Equivalent Dose.

< 0.001 , $\eta_p^2 = 0.332$) (Fig. 2-d), and for the TTP of BF ($F_{(6,306)} = 3.342$, $P = 0.003$, $\eta_p^2 = 0.062$) (Fig. 2-e) and MG ($F_{(6,192)} = 4.275$, $P < 0.001$, $\eta_p^2 = 0.118$) (Fig. 2-f). Compared to trial 1, all individuals had significantly lower values of the TTP of CoP by trial 2 ($P < 0.001$), recovery time by trial 3 ($P = 0.028$), onset latency of BF and MG in trial 2 ($P = 0.023$) and 4 ($P = 0.037$) respectively, and TTP of BF and MG in trial 3 ($P = 0.015$) and 2 ($P = 0.001$), respectively. Compared to trial 7, all individuals had significantly longer TTP of CoP by trial 2 ($P = 0.025$), longer recovery time by trial 5 ($P = 0.014$), greater onset latency of BF and MG by trials 2 ($P = 0.006$) and 5 ($P = 0.021$), respectively, and longer TTP of BF by trial 4 ($P = 0.018$). There was a main effect of Group for recovery time ($F_{(1,49)} = 18.890$, $P < 0.001$, $\eta_p^2 = 0.278$), indicating a longer recovery time for PDG compared to CG (Table 2). Means of significant parameters are also demonstrated in Table 2 and means of non-significant parameters are presented as supplementary material (Supplementary Table S1).

3.2.2. Amplitude of EMG and CoP parameters

ANOVA indicated a Group by Trial interaction for the range of CoP ($F_{(6,324)} = 4.466$, $P < 0.001$, $\eta_p^2 = 0.076$) (Fig. 3-a) and co-activation of BF/VM ($F_{(6,324)} = 2.366$, $P = 0.030$, $\eta_p^2 = 0.042$) (Fig. 3-b). For the first trial analyses, there was a greater range of CoP in PDG vs. CG ($t_{(54)} = 3.549$, $P = 0.001$), but no difference between groups was revealed for the co-activation of BF/VM ($t_{(55)} = -0.142$, $P = 0.888$). PDG demonstrated a more rapid change for range of CoP, with the first significant change observed in trial 2 vs. 1 ($P = 0.005$) compared to trial 5 vs. 1 ($P = 0.035$) in CG. CG also presented lower co-activation of BF/VM muscles in trial 6 vs. 1 ($P = 0.034$). In addition, when trial 7 was contrasted with other trials, PDG showed a greater range of CoP in trial 4 ($P = 0.024$) and CG showed a greater range of CoP in trial 3 ($P = 0.021$) and greater co-activation of BF/VM in trial 2 ($P = 0.026$). ANOVA revealed Trial main effect for the iEMG of BF ($F_{(6,306)} = 2.777$, $P = 0.012$, $\eta_p^2 = 0.052$) (Fig. 3-c) and iEMG of TA ($F_{(6,288)} = 2.927$, $P = 0.009$, $\eta_p^2 = 0.057$) (Fig. 3-d). Individuals presented higher values of iEMG of BF and TA in trials 6 and 2 in contrast to trial 1 ($P = 0.020$ and $P = 0.026$, respectively). In addition, participants demonstrated higher values of iEMG of BF and TA in trials 6 and 2 compared to trial 7 ($P = 0.006$ and $P = 0.001$, respectively). ANOVA demonstrated Group main effect for iEMG of BF ($F_{(1,51)} = 6.493$, $P = 0.014$, $\eta_p^2 = 0.113$) and for co-activation of MG/TA ($F_{(1,54)} = 4.308$, $P = 0.043$, $\eta_p^2 = 0.074$). CG vs. PDG presented greater iEMG of BF, and lower co-activation of MG/TA (Table 2). Means of the significant

parameters are also demonstrated in Table 2 and means of non-significant parameters are presented as supplementary material (Supplementary Table S1).

4. Discussion

This study investigated the capacity to change the reactive postural responses after unexpected external perturbation between PDG and CG and investigated the habituation plateaus of postural responses to non-sequential trials of external translations of the support-surface in both groups. People with PD present a modest impairment on postural control in situations with external perturbation compared to CG. The greater range of CoP in the first trial and the greater MG/TA co-activation, greater recovery time, and lower iEMG of BF in the unpredictable translation of the support base in people with PD may reflect the central nervous system inability in controlling the muscle activity and in coordinating the muscles in people with PD vs. CG (Bloem, 1992; Dimitrova et al., 2004b; Horak et al., 1992). Indeed, this inadequate muscle control might be associated with the typical rigidity and bradykinesia symptoms in PD, which could impair the balance maintenance, mainly in more challenging situations, such as after external perturbation (Horak et al., 2005; Peterson and Horak, 2016).

The unexpected faster capacity in people with PD to change the postural response after the first trial than older adults (2nd and 5th trials, respectively) is in line with previous data (Bloem et al., 1998). The results may be explained by the 36% higher range of CoP in PDG vs. CG in the 1st trial (indicative of worse postural control to perturbation), which reflects in a greater room for postural adjustments in a subsequent trial (Bloem et al., 1998). However, the worse postural control during the first trial in people with PD needs to be considered with caution because the differences between groups were observed only in the range of CoP.

As expected, both PDG and CG changed postural control when they were exposed to a repeated perturbation, demonstrating an ability to habituate to perturbations. Habituation was observed by a gradual decrease in time to activate muscle (the onset latency of BF and MG, and TTP of BF and MG), in magnitude of muscle activation (iEMG of TA and BF muscles), and CoP (recovery time to the stable position, TTP, and the range of CoP). The observation that people with PD could modulate and habituate to unexpected and unpredictable perturbation is in line with other studies that demonstrated the capacity of people with PD to

Fig. 2. Trial main effect for recovery time (a), onset latency of BF (b) and MG (c) muscles; TTP of CoP (d), BF (e) and MG (f) muscles in reactive postural responses. Note: # Significant difference for trial 1; & Significant difference for trial 7; Solid lines indicate differences in the group by trial interaction and; dashed lines indicate differences in the Trial main effect.

Fig. 3. Group*Trials interaction for range of CoP (a) and Co-activation BF/VM (b) and; Trial main effect for iEMG of BF (c) and TA (d) in reactive adjustments. Note: *Significant difference between groups (PDG and CG); # Significant difference for trial 1; & Significant difference for trial 7; Solid lines indicate differences in the group by trial interaction and; dashed lines indicate differences in the Trial main effect.

change and to habituate the postural response after perturbation (Bloem et al., 1998; Nanhoe-Mahabier et al., 2012). However, previous studies presented the perturbation in blocks (sequence of perturbation trials) and by the rotation of support-base (Bloem et al., 1998; Nanhoe-Mahabier et al., 2012), and a new aspect of our study is that perturbations were presented in non-sequential trials and by the support base translation. One possibility is that the people with PD could somehow retain, at least for a short while, the information from the previous perturbations (Duncan et al., 2014; Van Ooteghem et al., 2017). The postural response adaptations of young adults have been evidenced independently of the perturbation exposure in sequence, suggesting that for the adaptations, the number of perturbations is important (Duncan et al., 2014). The similarities between our data and literature may be related to the characteristics of support-base perturbation, since both types of perturbations required similar involvement of muscle groups and strategies (Duncan et al., 2014; Visser et al., 2010).

Only slight and maybe not meaningful differences were observed in time to habituation plateau in people with PD. Although the delay in habituation in people with PD was expected (Nanhoe-Mahabier et al., 2012), the slight delay in reaching the habituation plateaus needs to be considered with caution since: (a) it was observed in only two parameters (range of CoP and co-activation of the BF/VM); (b) the close difference in these parameters between PDG vs. CG (5th and 4th trials, respectively) for CoP range. While PD impairs the cortico-basal networks and brainstem structures that are responsible for coordinating muscles and for adapting the response to sequential perturbation series (Nanhoe-Mahabier et al., 2012; Van Ooteghem et al., 2017), people with PD may be compensating for such impairments by using different brain areas (de Kam et al., 2014; Peterson and Horak, 2016). The habituation to repetitive perturbations reflects the individual's capacity to identify and cope with changing circumstances in an optimal and safe strategy (Chong et al., 2000, 1999; Horak et al., 1997). The habituation in few trials could indicate less difficulty, and adaptation of the postural responses (Oude Nijhuis et al., 2009). Thus, it is plausible that procedural

learning may have influenced the habituation capacity and may affect the postural response (Duncan et al., 2014; Muslimovic et al., 2007; Nanhoe-Mahabier et al., 2012). Procedural learning supports the idea that the subjects acquire and improve the postural control as the results of repeated perturbation exposures, decreasing the exaggerated responses evidenced during the first trial (Bloem et al., 1998; Nanhoe-Mahabier et al., 2012; Oude Nijhuis et al., 2009). The modest delay in habituation in people with PD may support the idea that, although people with PD demonstrated the ability to retain the improvement of the postural response after practice (Van Ooteghem et al., 2017), procedural learning is less efficient in people with PD than older adults (Krebs et al., 2001; Muslimovic et al., 2007).

Such slight delay in habituation in people with PD is likely linked to the deficits in the basal ganglia function (Jacobs and Horak, 2007; Peterson and Horak, 2016), since this area is associated with learning, control of movements and modulation of postural adjustments (Jacobs and Horak, 2007; Lester et al., 2017; Wilkinson et al., 2009). Such impairments in basal ganglia involve a PD-typical imbalanced excitatory/inhibitory cortical control (Peterson and Horak, 2016; Takakusaki, 2017), impairing the neural drive to synergize and antagonize muscles, which may result in accentuated co-activation during postural control (Papegaaij et al., 2014; Takakusaki et al., 2004). This concept supports the apparent 19% greater MG/TA co-activation observed in PDG vs. CG (Table 2).

Study limitations include lack of analysis of the perturbation start with the CoP parameters, albeit our protocol allowed us to investigate the temporal and amplitude responses to perturbation by using behavioral (CoP) and control (EMG) strategies. Although the slight delay in habituation plateaus was observed through spatial parameters (range of CoP), additional parameters, such as virtual time to contact (spatio-temporal proximity of the CoP to the limits of stability), would add a greater overview of PD-impairments related to postural response to perturbation (Slobounov et al., 2009, 1997). Also, by selecting a short 250 ms window (same period of the reactive adjustments) to determine

Table 2

Mean and standard deviations of temporal and amplitude parameters of EMG and CoP in reactive postural response for the significant Group by Trial interaction and main effect of Group and/or Trial.

		Trials							Group*Trials Interaction	Group Effect	Trial Effect
		1	2	3	4	5	6	7			
Range of CoP (cm)	PDG	13.9 ± 3.5	11.8 ± 3.4	10.4 ± 3.3	10.5 ± 3.1	9.9 ± 3.0	9.3 ± 3.4	8.8 ± 2.7	PDG: 1 > 2 ^a ; 4 > 7 ^b CG: 1 > 5 ^a ; 3 > 7 ^b	–	–
	CG	10.7 ± 2.6	11.0 ± 2.4	10.2 ± 2.7	9.4 ± 2.3	9.0 ± 2.4	8.4 ± 2.1	8.4 ± 2.0			
TTP of CoP (ms)	PDG	519.4 ± 86.2	441.2 ± 86.0	425.2 ± 71.4	411.5 ± 81.0	426.1 ± 76.2	406.4 ± 65.1	396.7 ± 73.1	ns	ns	1 > 2 ^a 2 > 7 ^b
	CG	518.5 ± 97.0	391.0 ± 81.0	384.0 ± 95.6	399.5 ± 82.6	388.5 ± 72.3	403.0 ± 70.9	384.0 ± 53.5			
Recovery time (s)	PDG	3.8 ± 1.0	3.3 ± 0.8	3.3 ± 0.9	3.1 ± 0.8	3.2 ± 1.1	2.8 ± 1.0	2.7 ± 1.0	ns	PDG > CG	1 > 3 ^a 5 > 7 ^b
	CG	3.1 ± 0.7	3.1 ± 1.0	2.8 ± 1.3	2.5 ± 0.6	2.3 ± 0.7	2.2 ± 0.9	2.0 ± 0.7			
Onset latency BF (ms)	PDG	243.3 ± 172.3	191.1 ± 97.0	186.1 ± 76.1	186.4 ± 86.8	169.2 ± 56.4	168.3 ± 46.1	155.3 ± 57.4	ns	ns	1 > 2 ^a 2 > 7 ^b
	CG	249.0 ± 134.2	193.5 ± 71.3	153.5 ± 39.1	173.0 ± 33.9	171.0 ± 49.1	173.0 ± 30.6	166.5 ± 31.3			
Onset latency MG (ms)	PDG	179.4 ± 57.6	150.3 ± 45.3	146.1 ± 47.5	143.1 ± 55.8	142.2 ± 41.7	126.1 ± 22.8	124.7 ± 34.7	ns	ns	1 > 4 ^a 5 > 7 ^b
	CG	134.2 ± 59.7	143.7 ± 44.0	140.0 ± 50.2	126.8 ± 31.6	131.1 ± 35.3	125.3 ± 32.7	116.3 ± 33.2			
TTP of BF (ms)	PDG	233.8 ± 175.5	208.1 ± 198.0	185.9 ± 186.3	183.8 ± 175.0	204.3 ± 139.1	178.4 ± 171.3	152.2 ± 136.0	ns	ns	1 > 3 ^a 4 > 7 ^b
	CG	254.4 ± 172.1	212.5 ± 192.3	136.3 ± 87.6	201.3 ± 138.5	102.5 ± 88.1	144.4 ± 91.8	93.8 ± 59.0			
TTP of MG (ms)	PDG	296.4 ± 289.5	255.9 ± 314.8	206.4 ± 220.2	147.0 ± 151.7	145.0 ± 135.4	142.7 ± 134.1	183.2 ± 228.7	ns	ns	1 > 2 ^a
	CG	358.3 ± 254.1	110.0 ± 80.9	110.0 ± 80.1	167.5 ± 227.0	154.2 ± 125.9	85.0 ± 50.9	207.5 ± 248.1			
iEMG BF (% baseline)	PDG	393.0 ± 304.4	401.1 ± 297.4	365.9 ± 278.7	346.0 ± 216.0	380.3 ± 407.2	424.8 ± 361.7	360.8 ± 297.6	ns	PDG < CG	1 < 6 ^a 6 > 7 ^b
	CG	591.6 ± 525.8	656.8 ± 656.0	635.1 ± 582.4	514.6 ± 298.2	766.3 ± 511.2	751.0 ± 646.8	631.5 ± 656.6			
iEMG TA (% baseline)	PDG	491.5 ± 376.6	582.4 ± 476.8	566.7 ± 561.3	547.1 ± 513.9	474.6 ± 414.5	552.3 ± 558.0	492.6 ± 385.9	ns	ns	1 < 2 ^a 2 > 7 ^b
	CG	548.1 ± 593.8	854.8 ± 717.7	420.7 ± 275.7	545.4 ± 361.3	466.2 ± 372.9	399.1 ± 303.7	429.3 ± 355.1			
Co-activation BF/VM (%)	PDG	54.1 ± 21.3	55.3 ± 25.2	55.5 ± 23.8	51.0 ± 23.2	54.5 ± 23.6	57.2 ± 26.0	57.3 ± 25.0	PDG: ns CG: 1 > 6 ^a ; 2 < 7 ^b	–	–
	CG	56.2 ± 20.3	58.4 ± 24.1	48.5 ± 26.0	45.0 ± 19.0	46.1 ± 28.7	39.3 ± 29.3	43.7 ± 24.9			
Co-activation MG/TA (%)	PDG	61.6 ± 25.5	59.7 ± 25.7	59.1 ± 21.9	52.1 ± 26.0	60.1 ± 24.5	57.7 ± 23.2	57.5 ± 21.2	ns	PDG > CG	ns
	CG	57.8 ± 22.9	53.0 ± 29.1	53.0 ± 32.7	47.3 ± 26.7	50.1 ± 31.5	42.1 ± 24.0	41.5 ± 24.9			

TTP = time to peak; CoP = Center of Pressure; iEMG = integral of electromyography; BF = biceps femoris; VM = vastus medialis; MG = medial head of gastrocnemius; TA = tibialis anterior; PDG = Parkinson's disease group; CG = Control group; ns = not significant; a = indicated the difference in the Contrast 1 analysis; b = indicated the difference in the Contrast 2 analysis. Note: Means of parameters that demonstrated non-significant difference are presented as supplementary material.

the baseline (usual approach adopted by the literature (de Freitas et al., 2010; Santos et al., 2010)), EMG could have had the influence of small bursts of muscle activity. Therefore, future studies should consider longer time-windows (e.g., 500 ms - Cleworth et al., 2016) and include additional analysis to provide more robust information regarding the between-group differences of shapes of postural responses to perturbation over time (e.g., exponential decay functions). Such analysis would detail the group-differences in behavioral aspects of the habituation over time. The difference in postural response between groups in just a few parameters may be explained because our participants with PD are involved in a physical exercise program and because of the lower number of perturbation exposures (Smania et al., 2010). Also, the low number of trials allowed only to identify initial habituation plateaus; thus, future studies with more trials are needed to identify the full habituation of postural responses. Besides, while participants did not know whether a perturbation would or would not occur, they were aware of the direction of the perturbation, which could limit the unpredictability of the task in terms of direction. Future studies should examine the involvement of brain areas during the habituation of postural adjustments to the perturbation in PD to the neuromechanism and brain networks involved in the task (Jacobs and Horak, 2007, 2006; Mochizuki et al., 2008). In addition, understanding the role of anti-parkinsonian medication on postural habituation of people with PD

would provide information about the responsiveness of postural response to medication. Despite such limitations, our result could support rehabilitation strategies to improve postural control in PD. Future studies would implement exercise intervention to facilitate the habituation. Furthermore, cue and additional sensorial information could be implemented in those strategies.

5. Conclusions

Despite the greater range of CoP in the first trial compared to healthy older adults, people with PD can habituate to unpredictable postural perturbations. This capacity to habituate is reflected by little, to no difference in the time-course of adaptation for all but 2 parameters (range of CoP and co-activation of the BF/VM) that showed only marginal differences between people with PD and older adults. Therefore, there was only a slight and maybe not meaningful delay to habituate to perturbation in people with PD.

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.clinbiomech.2021.105363>.

Declaration of Competing Interest

None.

Acknowledgments

This work was supported by the São Paulo Research Foundation (FAPESP) [grant number #2016/00503-0], National Council for Scientific and Technological Development (CNPq) [#142057/2017-7; #309045/2017-7] and financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) [Finance Code 001].

References

- Barbieri, F.A., Polastri, P.F., Baptista, A.M., Lirani-Silva, E., Simieli, L., Orcioli-Silva, D., Beretta, V.S., Gobbi, L.T.B., 2016. Effects of disease severity and medication state on postural control asymmetry during challenging postural tasks in individuals with Parkinson's disease. *Hum. Mov. Sci.* 46, 96–103. <https://doi.org/10.1016/j.humov.2015.12.009>.
- Beretta, V.S., Gobbi, L.T.B., Lirani-Silva, E., Simieli, L., Orcioli-Silva, D., Barbieri, F.A., 2015. Challenging postural tasks increase asymmetry in patients with Parkinson's disease. *PLoS One* 10, e0137722. <https://doi.org/10.1371/journal.pone.0137722>.
- Beretta, V.S., Vitorio, R., dos Santos, P.C.R., Orcioli-Silva, D., Gobbi, L.T.B., 2019. Postural control after unexpected external perturbation: effects of Parkinson's disease subtype. *Hum. Mov. Sci.* 64, 12–18. <https://doi.org/10.1016/j.humov.2019.01.001>.
- Bloem, B.R., 1992. Postural instability in Parkinson's disease. *Clin. Neurol. Neurosurg.* 94 (Suppl), S41–S45. [https://doi.org/10.1016/0303-8467\(92\)90018-x](https://doi.org/10.1016/0303-8467(92)90018-x).
- Bloem, B.R., van Vugt, J.P., Beckley, D.J., Remler, M.P., Roos, R.A., 1998. Habituation of lower leg stretch responses in Parkinson's disease. *Electroencephalogr. Clin. Neurophysiol.* 109, 73–77. [https://doi.org/10.1016/s0924-980x\(97\)00073-8](https://doi.org/10.1016/s0924-980x(97)00073-8).
- Brucki, S.M., Nitrini, R., Caramelli, P., Bertolucci, P.H., Okamoto, I.H., 2003. Suggestions for utilization of the mini-mental state examination in Brazil. *Arq. Neuropsiquiatr.* 61, 777–781. <https://doi.org/10.1590/S0004-282X2003000500014>.
- Carpenter, M.G., Allum, J.H., Honegger, F., Adkin, A.L., Bloem, B.R., 2004. Postural abnormalities to multidirectional stance perturbations in Parkinson's disease. *J. Neurol. Neurosurg. Psychiatry* 75, 1245–1254. <https://doi.org/10.1136/jnnp.2003.021147>.
- Chong, R.K., Jones, C.L., Horak, F.B., 1999. Postural set for balance control is normal in Alzheimer's but not in Parkinson's disease. *J. Gerontol. Ser. A Biol. Sci. Med. Sci.* 54, M129–M135. <https://doi.org/10.1093/gerona/54.3.m129>.
- Chong, R.K., Horak, F.B., Woollacott, M.H., 2000. Parkinson's disease impairs the ability to change set quickly. *J. Neurol. Sci.* 175, 57–70. [https://doi.org/10.1016/s0022-510x\(00\)00277-x](https://doi.org/10.1016/s0022-510x(00)00277-x).
- Cleworth, T.W., Chua, R., Inglis, J.T., Carpenter, M.G., 2016. Influence of virtual height exposure on postural reactions to support surface translations. *Gait Posture* 47, 96–102. <https://doi.org/10.1016/j.gaitpost.2016.04.006>.
- Dimitrova, D., Horak, F.B., Nutt, J.G., 2004a. Postural muscle responses to multidirectional translations in patients with Parkinson's disease. *J. Neurophysiol.* 91, 489–501. <https://doi.org/10.1152/jn.00094.2003>.
- Dimitrova, D., Nutt, J., Horak, F.B., 2004b. Abnormal force patterns for multidirectional postural responses in patients with Parkinson's disease. *Exp. Brain Res.* 156, 183–195. <https://doi.org/10.1007/s00221-003-1770-4>.
- Duncan, C.A., Langlois, R.G., Albert, W.J., MacKinnon, S.N., 2014. The habituation of human postural responses to platform perturbations. *Int. J. Ind. Ergon.* 44, 874–881. <https://doi.org/10.1016/j.ergon.2014.06.005>.
- Fahn, S., Elton, R., UPDRS, M. of the, 1987. The unified Parkinson's disease rating scale. In: Fahn, S., Marsden, C.D., Calne, D.B., Goldstein, M. (Eds.), *Recent Developments in Parkinson's Disease*. Mcmillan Health Care Information, Florham Park, NJ, pp. 153–163.
- de Freitas, P.B., Knight, C.A., Barela, J.A., 2010. Postural reactions following forward platform perturbation in young, middle-age, and old adults. *J. Electromyogr. Kinesiol.* 20, 693–700. <https://doi.org/10.1016/j.jelekin.2009.11.009>.
- Hermens, H.J., Frenkers, B., Disselhorst-Klug, C., Rau, G., 2000. Development of recommendations for SEMG sensors and sensor placement procedures. *J. Electromyogr. Kinesiol.* 10, 361–374. [https://doi.org/10.1016/S1050-6411\(00\)00027-4](https://doi.org/10.1016/S1050-6411(00)00027-4).
- Horak, F.B., Nutt, J.G., Nashner, L.M., 1992. Postural inflexibility in parkinsonian subjects. *J. Neurol. Sci.* 111, 46–58. [https://doi.org/10.1016/0022-510X\(92\)90111-W](https://doi.org/10.1016/0022-510X(92)90111-W).
- Horak, F.B., Henry, S.M., Shumway-Cook, A., 1997. Postural perturbations: new insights for treatment of balance disorders. *Phys. Ther.* 77, 517–533. <https://doi.org/10.1093/ptj/77.5.517>.
- Horak, F.B., Dimitrova, D., Nutt, J.G., 2005. Direction-specific postural instability in subjects with Parkinson's disease. *Exp. Neurol.* 193, 504–521. <https://doi.org/10.1016/j.expneurol.2004.12.008>.
- Jacobs, J.V., Horak, F.B., 2006. Abnormal proprioceptive-motor integration contributes to hypometric postural responses of subjects with Parkinson's disease. *Neuroscience* 141, 999–1009. <https://doi.org/10.1016/j.neuroscience.2006.04.014>.
- Jacobs, J.V., Horak, F.B., 2007. Cortical control of postural responses. *J. Neural Transm.* 114, 1339–1348. <https://doi.org/10.1007/s00702-007-0657-0>.
- de Kam, D., Nonnekes, J., Oude Nijhuis, L.B., Geurts, A.C.H., Bloem, B.R., Weerdesteyn, V., 2014. Dopaminergic medication does not improve stepping responses following backward and forward balance perturbations in patients with Parkinson's disease. *J. Neurol.* 261, 2330–2337. <https://doi.org/10.1007/s00415-014-7496-3>.
- Krebs, H.I., Hogan, N., Hening, W., Adamovich, S.V., Poizner, H., 2001. Procedural motor learning in Parkinson's disease. *Exp. Brain Res.* 141, 425–437. <https://doi.org/10.1007/s002210100871>.
- Lang, K.C., Hackney, M.E., Ting, L.H., McKay, J.L., 2019. Antagonist muscle activity during reactive balance responses is elevated in Parkinson's disease and in balance impairment. *PLoS One* 14, e0211137. <https://doi.org/10.1371/journal.pone.0211137>.
- Lester, M.E., Cavanaugh, J.T., Foreman, K.B., Shaffer, S.W., Marcus, R., Dibble, L.E., 2017. Adaptation of postural recovery responses to a vestibular sensory illusion in individuals with Parkinson disease and healthy controls. *Clin. Biomech.* 48, 73–79. <https://doi.org/10.1016/j.clinbiomech.2017.07.008>.
- Mochizuki, G., Sibley, K.M., Esposito, J.G., Camilleri, J.M., McIlroy, W.E., 2008. Cortical responses associated with the preparation and reaction to full-body perturbations to upright stability. *Clin. Neurophysiol.* 119, 1626–1637. <https://doi.org/10.1016/j.clinph.2008.03.020>.
- Muslimovic, D., Post, B., Speelman, J.D., Schmand, B., 2007. Motor procedural learning in Parkinson's disease. *Brain* 130, 2887–2897. <https://doi.org/10.1093/brain/awm211>.
- Nanhoe-Mahabier, W., Allum, J.H., Overeem, S., Borm, G.F., Oude Nijhuis, L.B., Bloem, B.R., 2012. First trial reactions and habituation rates over successive balance perturbations in Parkinson's disease. *Neuroscience* 217, 123–129. <https://doi.org/10.1016/j.neuroscience.2012.03.064>.
- Oude Nijhuis, L.B., Allum, J.H., Borm, G.F., Honegger, F., Overeem, S., Bloem, B.R., 2009. Directional sensitivity of “first trial” reactions in human balance control. *J. Neurophysiol.* 101, 2802–2814. <https://doi.org/10.1152/jn.90945.2008>.
- Papegaai, S., Taube, W., Baudry, S., Otten, E., Hortobagyi, T., 2014. Aging causes a reorganization of cortical and spinal control of posture. *Front. Aging Neurosci.* 6, 28. <https://doi.org/10.3389/fnagi.2014.00028>.
- Park, J.H., Kang, Y.J., Horak, F.B., 2015. What is wrong with balance in Parkinson's disease? *J. Mov. Disord.* 8, 109–114. <https://doi.org/10.14802/jmd.15018>.
- Peterson, D.S., Horak, F.B., 2016. Neural control of walking in people with parkinsonism. *Physiol. Sci.* 31, 95–107. <https://doi.org/10.1152/physiol.00034.2015>.
- Santos, M.J., Kanekar, N., Aruin, A.S., 2010. The role of anticipatory postural adjustments in compensatory control of posture: 1. Electromyographic analysis. *J. Electromyogr. Kinesiol.* 20, 388–397. <https://doi.org/10.1016/j.jelekin.2009.06.006>.
- Schenkman, M., Wei Zhu, C., Cutson, T.M., Whetten-Goldstein, K., 2001. Longitudinal evaluation of economic and physical impact of Parkinson's disease. *Parkinsonism Relat. Disord.* 8, 41–50. [https://doi.org/10.1016/s1353-8020\(00\)00079-1](https://doi.org/10.1016/s1353-8020(00)00079-1).
- Slobounov, S., Cao, C., Jaiswal, N., Newell, K.M., 2009. Neural basis of postural instability identified by VTC and EEG. *Exp. Brain Res.* 199, 1–16. <https://doi.org/10.1007/s00221-009-1956-5>.
- Slobounov, S.M., Slobounova, E.S., Newell, K.M., 1997. Virtual time-to-collision and human postural control. *J. Mot. Behav.* 29, 263–281. <https://doi.org/10.1080/00222899709600841>.
- Smania, N., Corato, E., Tinazzi, M., Stanzani, C., Fiaschi, A., Girardi, P., Gandolfi, M., 2010. Effect of balance training on postural instability in patients with idiopathic Parkinson's disease. *Neurorehabil. Neural Repair* 24, 826–834. <https://doi.org/10.1177/1545968310376057>.
- Smith, B.A., Jacobs, J.V., Horak, F.B., 2012. Effects of magnitude and magnitude predictability of postural perturbations on preparatory cortical activity in older adults with and without Parkinson's disease. *Exp. Brain Res.* 222, 455–470. <https://doi.org/10.1007/s00221-012-3232-3>.
- Takakusaki, K., 2017. Functional Neuroanatomy for posture and gait control. *J. Mov. Disord.* 10, 1–17. <https://doi.org/10.14802/jmd.16062>.
- Takakusaki, K., Saitoh, K., Harada, H., Kashiwayanagi, M., 2004. Role of basal ganglia-brainstem pathways in the control of motor behaviors. *Neurosci. Res.* 50, 137–151. <https://doi.org/10.1016/j.neures.2004.06.015>.
- Tomlinson, C.L., Stowe, R., Patel, S., Rick, C., Gray, R.E.C.C., 2010. Systematic review of levodopa dose equivalency reporting in Parkinson's disease. *Mov. Disord.* 25, 2649–2653. <https://doi.org/10.1002/mds.23429>.
- Van Ooteghem, K., Frank, J.S., Horak, F.B., 2017. Postural motor learning in Parkinson's disease: the effect of practice on continuous compensatory postural regulation. *Gait Posture* 57, 299–304. <https://doi.org/10.1016/j.gaitpost.2017.06.021>.
- Visser, J.E., Oude Nijhuis, L.B., Janssen, L., Bastiaanse, C.M., Borm, G.F., Duysens, J., Bloem, B.R., 2010. Dynamic posturography in Parkinson's disease: diagnostic utility of the “first trial effect”. *Neuroscience* 168, 387–394. <https://doi.org/10.1016/j.neuroscience.2010.03.068>.
- Wilkinson, L., Khan, Z., Jahanshahi, M., 2009. The role of the basal ganglia and its cortical connections in sequence learning: evidence from implicit and explicit sequence learning in Parkinson's disease. *Neuropsychologia* 47, 2564–2573. <https://doi.org/10.1016/j.neuropsychologia.2009.05.003>.