University of Groningen

Prognostic value of the FUT family in acute myeloid leukemia

Dai, Yifeng; Cheng, Zhiheng; Pang, Yifan; Jiao, Yang; Qian, Tingting; Quan, Liang; Cui, Longzhen; Liu, Yan; Si, Chaozeng; Chen, Jinghong

Published in:
Cancer Gene Therapy

DOI:
[10.1038/s41417-019-0115-9](https://doi.org/10.1038/s41417-019-0115-9)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Dai, Y., Cheng, Z., Pang, Y., Jiao, Y., Qian, T., Quan, L., Cui, L., Liu, Y., Si, C., Chen, J., Ye, X., Chen, J., Shi, J., Wu, D., Zhang, X., & Fu, L. (2020). Prognostic value of the FUT family in acute myeloid leukemia. *Cancer Gene Therapy*, 27(1-2), 70-80. <https://doi.org/10.1038/s41417-019-0115-9>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Prognostic value of the *FUT* family in acute myeloid leukemia

Yifeng Dai^{1,2} · Zhiheng Cheng^{2,3,4,5} · Yifan Pang⁶ · Yang Jiao⁷ · Tingting Qian^{1,4} · Liang Quan^{1,4} · Longzhen Cui³ · Yan Liu³ · Chaozeng Si⁸ · Jinghong Chen⁴ · Xu Ye¹ · Jingqi Chen⁴ · Jinlong Shi⁹ · Depei Wu¹⁰ · Xinyou Zhang¹¹ · Lin Fu^{1,4,5}

Received: 30 April 2019 / Revised: 30 May 2019 / Accepted: 1 June 2019 / Published online: 17 June 2019
© The Author(s), under exclusive licence to Springer Nature America, Inc. 2019

Abstract

Genetic abnormalities are more frequently viewed as prognostic markers in acute myeloid leukemia (AML) in recent years. Fucosylation, catalyzed by fucosyltransferases (FUTs), is a post-translational modification that widely exists in cancer cells. However, the expression and clinical implication of the *FUT* family (*FUT1-11*) in AML has not been investigated. From the Cancer Genome Atlas database, a total of 155 AML patients with complete clinical characteristics and *FUT1-11* expression data were included in our study. In patients who received chemotherapy alone showed that high expression levels of *FUT3*, *FUT6*, and *FUT7* had adverse effects on event-free survival (EFS) and overall survival (OS) (all $P < 0.05$), whereas high *FUT4* expression had favorable effects on EFS and OS (all $P < 0.01$). However, in the allogeneic hematopoietic stem cell transplantation (allo-HSCT) group, we only found a significant difference in EFS between the high and low *FUT3* expression subgroups ($P = 0.047$), while other *FUT* members had no effect on survival. Multivariate analysis confirmed that high *FUT4* expression was an independent favorable prognostic factor for both EFS (HR = 0.423, $P = 0.001$) and OS (HR = 0.398, $P < 0.001$), whereas high *FUT6* expression was an independent risk factor for both EFS (HR = 1.871, $P = 0.017$) and OS (HR = 1.729, $P = 0.028$) in patients who received chemotherapy alone. Moreover, we found that patients with low *FUT4* and high *FUT6* expressions had the shortest EFS and OS ($P < 0.05$). Our study suggests that high expressions of *FUT3/6/7* predict poor prognosis, high *FUT4* expression indicates good prognosis in AML; *FUT6* and *FUT4* have the best prognosticating profile among them, but their effects could be neutralized by allo-HSCT.

Introduction

Acute myeloid leukemia (AML) is a heterogeneous group of hematopoietic malignancies originating from myeloid progenitors or myeloid-primed multipotential progenitors. It is characterized by the leukemia cells' unchecked

These authors contributed equally: Yifeng Dai, Zhiheng Cheng

These authors jointly supervised this work: Depei Wu, Xinyou Zhang, Lin Fu

✉ Lin Fu
fulin022@126.com

¹ Department of Hematology, The Second Affiliated Hospital of Guangzhou Medical University, 510260 Guangzhou, China

² Department of Pathology and Medical Biology, University Medical Center Groningen, University of Groningen, Groningen, Netherlands

³ Translational Medicine Center, Huaihe Hospital of Henan University, 475000 Kaifeng, China

⁴ Translational Medicine Center, The Second Affiliated Hospital of Guangzhou Medical University, 510260 Guangzhou, China

⁵ Department of Hematology, Huaihe Hospital of Henan University, 475000 Kaifeng, China

⁶ Department of Medicine, William Beaumont Hospital, Royal Oak, MI 48073, USA

⁷ Life Sciences Institute and Innovation Center for Cell Signaling Network, Zhejiang University, 310058 Hangzhou, China

⁸ Department of Operations and Information Management, China-Japan Friendship Hospital, 100029 Beijing, China

⁹ Department of Biomedical Engineering, Chinese PLA General Hospital, 100853 Beijing, China

¹⁰ Department of Hematology, The First Affiliated Hospital of Soochow University, 215006 Suzhou, China

¹¹ Department of Hematology, The Second Clinical Medical College (Shenzhen People's Hospital), Jinan University, 518020 Shenzhen, China

proliferation, abnormal differentiation, and aggressive infiltration of peripheral blood, bone marrow, or other tissues [1, 2]. The leukemogenesis and the clinical behavior of AML is dominated by the intricate molecular abnormalities and cytogenetics—many of them have been identified as independent prognostic markers, inducing abnormal expression of downstream genes and disrupting the transcriptional regulation systems [3, 4]. For instance, *TET2* and *FLT3* mutations may work synergistically in stimulating the leukemia cell proliferation, which explains their roles as poor prognostic indicators [5, 6]. On the other hand, another two common genetic abnormalities, *CEBPA* and *NPM1* mutations, indicate good prognosis in AML [7, 8]. Gene expression abnormalities exert profound impacts on therapy response and treatment outcomes. Previous studies observed that in AML, high expressions of *FHL2*, *iASPP*, *DOK4*, and *DOK5* act as adverse prognostic factors, and high *DOK7* expression is a good prognostic factor, however, their prognostic effects might overcome by allogeneic hematopoietic stem cell transplantation (allo-HSCT) [9, 10]. Therefore, researchers have been looking for the missing pieces in AML genetic expression profiling to improve the risk classification and individualized treatment for patients.

Fucosylation is a post-translational modification widely existing in proliferating cancer cells, catalyzed by fucosyltransferases (FUTs) [11, 12]. FUT family, including FUT1 to FUT11, are fucosylation synthases which are responsible for adding fucose to oligosaccharide chains of glycolipids, oligosaccharides, and glycoproteins [11, 13]. Fucosylated oligosaccharides participate in multiple cell–cell interactions during cell development, differentiation, and malignant transformation [13, 14]. Knocking down the *FUT* genes can potentially inhibit the biosynthesis of certain oligosaccharide chains on tumor cell surface, making them desirable therapeutic targets [15, 16]. According to previous studies, high expressions of *FUT1* and *FUT2* in human ovarian carcinoma-derived RMG-1 cells can promote cell proliferation and resistance to anticancer drugs [17, 18]. Increasing *FUT4* or *FUT7* expression can accelerate hepatocellular carcinoma cell proliferation [19, 20]. Decreasing *FUT3* or *FUT6* expression suppress colon carcinoma cell proliferation [21]. In addition, *FUT8* is highly expressed in the very aggressive human hepatocellular carcinoma cell line HCCLM3 [22].

Nevertheless, the clinical and prognostic value of the *FUT* family in AML remains unknown. The purpose of this study is to explore the impact of the *FUT* family's expression levels on the survival of AML patients who either received chemotherapy alone or followed by allo-HSCT.

Patients and Methods

Patients

The Cancer Genome Atlas (TCGA) database (<https://cancergenome.nih.gov/>) was screened for patients with complete *FUT* family (*FUT1-11*) expression data [23]. A total of 155 AML patients were included in this study, 84 patients received chemotherapy only and 71 patients received allo-HSCT. The baseline clinical and molecular characteristics, follow-up survival data were public accessible from the database, including age, sex, race, white blood cell (WBC) counts, peripheral blood (PB) and bone marrow (BM) blast percentages, French-American-British (FAB) subtypes, karyotype, cytogenetic risk categorization, and the frequencies of known recurrent genetic mutations/fusions. All patients provided informed consent. This study was approved by the Washington University Human Studies Committee.

Statistical analysis

Descriptive statistics were used to display the clinical and molecular characteristics of patients. The Chi-square test and the Mann–Whitney *U* test were used to compare categorical data and numerical data between groups, respectively. Event-free survival (EFS) and overall survival (OS) were the endpoints of this study. EFS was defined as the time from diagnosis to the first event, including death, and relapse, or was censored at the last follow up. OS was defined as the time from diagnosis to death from any cause or the last follow-up. Kaplan–Meier method and the log-rank test were used to estimate and compare survival. Multivariate Cox proportional hazard models were constructed for EFS and OS using a limited backward elimination procedure [24]. A two-tailed $P < 0.05$ was considered statistically significant. All above statistical analyses were performed by the SPSS 24.0 statistical software, and the GraphPad Prism 7.0 software was used for graphics.

Results

The prognostic value of *FUT* family's expression levels in AML

In order to estimate the prognostic value of *FUT* family in AML patients who underwent chemotherapy alone or received allo-HSCT, both treatment groups were divided into high and low expression subgroups by the median expression levels of each *FUT* members (*FUT1-11*). The

Table 1 Comparison of EFS and OS between the high and low expression levels of the *FUT* family

Variables	EFS		OS	
	χ^2	P-value	χ^2	P-value
Chemotherapy-only group				
<i>FUT1</i> (high vs. low)	1.015	0.314	1.429	0.232
<i>FUT2</i> (high vs. low)	0.194	0.660	0.222	0.638
<i>FUT3</i> (high vs. low)	7.595	0.006	6.497	0.011
<i>FUT4</i> (high vs. low)	11.910	0.001	13.273	<0.001
<i>FUT5</i> (high vs. low)	0.733	0.392	0.694	0.405
<i>FUT6</i> (high vs. low)	5.365	0.021	4.141	0.042
<i>FUT7</i> (high vs. low)	6.863	0.009	8.967	0.003
<i>FUT8</i> (high vs. low)	0.010	0.919	0.311	0.577
<i>FUT10</i> (high vs. low)	0.000	0.993	0.083	0.774
<i>FUT11</i> (high vs. low)	0.106	0.745	0.015	0.901
Allo-HSCT group				
<i>FUT1</i> (high vs. low)	0.950	0.330	1.419	0.234
<i>FUT2</i> (high vs. low)	0.141	0.707	1.268	0.260
<i>FUT3</i> (high vs. low)	3.931	0.047	0.739	0.390
<i>FUT4</i> (high vs. low)	0.314	0.575	0.835	0.361
<i>FUT5</i> (high vs. low)	0.432	0.511	0.144	0.704
<i>FUT6</i> (high vs. low)	0.726	0.394	0.067	0.795
<i>FUT7</i> (high vs. low)	0.027	0.870	0.541	0.462
<i>FUT8</i> (high vs. low)	0.002	0.968	0.015	0.902
<i>FUT10</i> (high vs. low)	0.414	0.520	0.045	0.833
<i>FUT11</i> (high vs. low)	0.134	0.714	0.326	0.568

EFS event-free survival, OS overall survival, Allo-HSCT allogeneic hematopoietic stem cell transplantation

differences of EFS and OS between high and low expression subgroups were presented in Table 1. In the chemotherapy-only group, high expression levels of *FUT3*, *FUT6*, and *FUT7* adversely affected both EFS and OS (all $P < 0.05$, Fig. 1a, b, e, f), while high *FUT4* expression had favorable effects on EFS and OS (all $P < 0.01$, Fig. 1c, d). However, in the allo-HSCT group, only *FUT3* expression affected EFS ($P = 0.047$), while the expression levels of other *FUT* members were independent of survival.

Relationship between *FUT3/4/6/7* expressions and other patient characteristics in the chemotherapy-only group

As shown in Table 2 and Table 3, the clinical and molecular characteristics of high and low *FUT3/4/6/7* expression subgroups were compared. The *FUT3*^{high} group had more FAB-M1 ($P = 0.010$) and fewer FAB-M4 patients ($P = 0.040$), more patients with complex karyotype ($P = 0.024$), and more frequent *TP53* mutation ($P = 0.004$). No significant differences were observed in age, gender and race distributions, WBC count, BM blasts, PB blasts, other FAB

subtypes, risk stratification, frequencies of other genetic mutations (*FLT3-ITD*, *NPM1*, *DNMT3A*, *IDH1/IDH2*, *RUNX1*, *NRAS/KRAS*, *TET2*, and *MLL*), or relapse rates between the *FUT3*^{high} and *FUT3*^{low} groups.

Compared with the *FUT4*^{low} subgroup, the *FUT4*^{high} subgroup were younger ($P = 0.033$), had more FAB-M4 patients ($P = 0.002$), more patients with normal karyotype ($P = 0.029$) or *MLL* fusion ($P = 0.024$), but higher relapse ratio ($P = 0.042$). No significant differences were noticed in gender or race distributions, WBC count, BM blasts, PB blasts, other FAB subtypes, risk stratification, or the frequencies of other genetic mutations (*FLT3-ITD*, *NPM1*, *DNMT3A*, *IDH1/IDH2*, *RUNX1*, *NRAS/KRAS*, *TET2*, and *TP53*) between the two subgroups.

Compared with *FUT6*^{low} subgroup, the *FUT6*^{high} subgroup had higher percentage of PB blasts ($P = 0.004$), more FAB-M1 and fewer FAB-M4 patients (all $P = 0.040$), more patients with complex karyotype ($P = 0.024$), fewer good-risk patients ($P = 0.013$), and more frequent *TP53* mutations ($P = 0.024$). There were no significant differences in age, gender and race distributions, WBC count, BM blasts, other FAB subtypes, frequency of other genetic mutations (*FLT3-ITD*, *NPM1*, *DNMT3A*, *IDH1/IDH2*, *RUNX1*, *NRAS/KRAS*, *TET2*, and *MLL*), or relapse rates between the two subgroups.

In regard to *FUT7* expression, the *FUT7*^{high} subgroup had more WBC count ($P = 0.038$), more patients with normal karyotype ($P = 0.029$), fewer patients with *CBFβ-MYH11* or *RUNX1-RUNX1T1* (all $P = 0.026$), and fewer good-risk patients ($P < 0.001$) than the low expression counterpart. Similar to the other three *FUTs*, no significant differences were found in age, gender and race distributions, BM and PB blasts, FAB subtypes, frequency of common genetic mutations (*FLT3-ITD*, *NPM1*, *DNMT3A*, *IDH1/IDH2*, *RUNX1*, *NRAS/KRAS*, *TET2*, *TP53*, and *MLL*), and relapse rates between the two *FUT7* expression subgroups.

Multivariate analysis of possible prognostic factors in the chemotherapy-only group

To further evaluate the prognostic significance of *FUT3/4/6/7* in patients who received chemotherapy alone, we selected multiple variables, including the expression levels of *FUT3/4/6/7* (high vs. low), age (≥ 60 vs. < 60 years), WBC count (≥ 15 vs. $< 15 \times 10^9/L$), BM blasts (≥ 70 vs. $< 70\%$), PB blasts (≥ 20 vs. $< 20\%$), *FLT3-ITD* (positive vs. negative), *NPM1* (mutated vs. wild), *DNMT3A* (mutated vs. wild), *RUNX1* (mutated vs. wild), and *TP53* (mutated vs. wild), to construct multivariate Cox proportional hazard models (Table 4).

Results indicated that high *FUT4* expression was an independent favorable factor for EFS (HR = 0.423, $P = 0.001$) and OS (HR = 0.398, $P < 0.001$), whereas high *FUT6* expression, age ≥ 60 years, BM blasts $\geq 70\%$, and

Fig. 1 Kaplan-Meier curves of EFS and OS in patients who received chemotherapy alone by univariate analysis. **a, b** High *FUT3* expressers had shorter EFS and OS than the low expressers. **c, d** High *FUT4* expressers had longer EFS and OS than the low expressers. **e, f** High

FUT6 expressers had shorter EFS and OS than the low expressers. **g, h** High *FUT7* expressers had shorter EFS and OS than the low expressers

Table 2 Comparison of the clinical and molecular characteristics between high and low *FUT3* and *FUT4* expression subgroups among patients who received chemotherapy alone

Characteristics	<i>FUT3</i>			<i>FUT4</i>		
	High (<i>n</i> = 42)	Low (<i>n</i> = 42)	<i>P</i> -value	High (<i>n</i> = 42)	Low (<i>n</i> = 42)	<i>P</i> -value
Age/years, median (range)	68.0 (34.0, 88.0)	64.0 (22.0, 82.0)	0.354 [*]	63.0 (25.0, 81.0)	70.5 (22.0, 88.0)	0.033 [*]
Age group/ <i>n</i> (%)			0.157 [§]			0.345 [§]
<60 years	10 (23.8)	16 (38.1)		15 (35.7)	11 (26.2)	
≥60 years	32 (76.2)	26 (61.9)		27 (64.3)	31 (73.8)	
Gender/ <i>n</i> (%)			0.512 [§]			0.126 [§]
Male	21 (50.0)	24 (57.1)		26 (61.9)	19 (45.2)	
Female	21 (50.0)	18 (42.9)		16 (38.1)	23 (54.8)	
Race/ <i>n</i> (%)						0.450 [§]
Caucasian	31 (73.8)	32 (76.2)	0.801 [§]	33 (78.6)	30 (71.4)	
Others	11 (26.2)	10 (23.8)		9 (21.4)	12 (28.6)	
WBC/× 10 ⁹ /L, median (range)	11.0 (0.7, 171.9)	18.7 (1.0, 297.4)	0.231 [*]	16.1 (2.1, 131.5)	10.5 (0.7, 297.4)	0.458 [*]
BM blasts/%, median (range)	72.5 (32.0, 98.0)	71.5 (30.0, 99.0)	0.865 [*]	73.5 (30.0, 98.0)	69.5 (32.0, 99.0)	0.366 [*]
PB blasts/%, median (range)	49.0 (0.0, 97.0)	16.0 (0.0, 98.0)	0.075 [*]	23.0 (0.0, 97.0)	24.0 (0.0, 98.0)	0.921 [*]
FAB subtypes/ <i>n</i> (%)						
M0	5 (11.9)	2 (4.8)	0.433 [§]	1 (2.4)	6 (14.3)	0.109 [§]
M1	15 (35.7)	5 (11.9)	0.010 [§]	9 (21.4)	11 (26.2)	0.608 [§]
M2	10 (23.8)	11 (26.2)	0.801 [§]	9 (21.4)	12 (28.6)	0.450 [§]
M4	6 (14.3)	14 (33.3)	0.040 [§]	16 (38.1)	4 (9.5)	0.002 [§]
M5	3 (7.1)	9 (21.4)	0.061 [§]	7 (16.7)	5 (11.9)	0.533 [§]
M6	0 (0.0)	1 (2.4)	1.000 [§]	0 (0.0)	1 (2.4)	1.000 [§]
M7	3 (7.1)	0 (0.0)	0.241 [§]	0 (0.0)	3 (7.1)	0.241 [§]
Cytogenetics/ <i>n</i> (%)						
Normal	18 (42.9)	22 (52.4)	0.382 [§]	25 (59.5)	15 (35.7)	0.029 [§]
t(9;22)/ <i>BCR-ABL1</i>	1 (2.4)	0 (0.0)	1.000 [§]	0 (0.0)	1 (2.4)	1.000 [§]
inv(16)/ <i>CBFβ-MYH11</i>	1 (2.4)	5 (11.9)	0.202 [§]	2 (4.8)	4 (9.5)	0.676 [§]
Complex	9 (21.4)	2 (4.8)	0.024 [§]	3 (7.1)	8 (19.0)	0.106 [§]
11q23/ <i>MLL</i>	1 (2.4)	2 (4.8)	1.000 [§]	3 (7.1)	0 (0.0)	0.241 [§]
-7/7q-	3 (7.1)	0 (0.0)	0.241 [§]	0 (0.0)	3 (7.1)	0.241 [§]
t(8;21)/ <i>RUNX1-RUNX1T1</i>	3 (7.1)	3 (7.1)	1.000 [§]	4 (9.5)	2 (4.8)	0.676 [§]
Others	6 (14.3)	8 (19.0)	0.558 [§]	5 (11.9)	9 (21.4)	0.242 [§]
Risk/ <i>n</i> (%)						
Good	4 (9.5)	8 (19.0)	0.212 [§]	6 (14.3)	6 (14.3)	1.000 [§]
Intermediate	21 (50.0)	25 (59.5)	0.381 [§]	25 (59.5)	21 (50.0)	0.381 [§]
Poor	16 (38.1)	8 (19.0)	0.053 [§]	11 (26.2)	13 (31.0)	0.629 [§]
<i>FLT3</i> / <i>n</i> (%)			0.606 [§]			0.291 [§]
<i>FLT3</i> -ITD	9 (21.4)	6 (14.3)		9 (21.4)	6 (14.3)	
<i>FLT3</i> -TKD	4 (9.5)	3 (7.1)		5 (11.9)	2 (4.8)	
Wild type	29 (69.0)	33 (78.6)		28 (66.7)	34 (81.0)	
<i>NPM1</i> / <i>n</i> (%)			0.102 [§]			0.243 [§]
Mutation	10 (23.8)	17 (40.5)		16 (38.1)	11 (26.2)	
Wild type	32 (76.2)	25 (59.5)		26 (61.9)	31 (73.8)	
<i>DNMT3A</i> / <i>n</i> (%)			0.221 [§]			0.807 [§]
Mutation	14 (33.3)	9 (21.4)		11 (26.2)	12 (28.6)	
Wild type	28 (66.7)	33 (78.6)		31 (73.8)	30 (71.4)	

Table 2 (continued)

Characteristics	<i>FUT3</i>			<i>FUT4</i>		
	High (<i>n</i> = 42)	Low (<i>n</i> = 42)	<i>P</i> -value	High (<i>n</i> = 42)	Low (<i>n</i> = 42)	<i>P</i> -value
<i>IDH1/IDH2/n</i> (%)						
Mutation	7 (16.7)	8 (19.0)	0.776 [§]	8 (19.0)	7 (16.7)	0.776 [§]
Wild type	35 (83.3)	34 (81.0)		34 (81.0)	35 (83.3)	
<i>RUNX1/n</i> (%)						0.079 [§]
Mutation	9 (21.4)	5 (11.9)	0.242 [§]	4 (9.5)	10 (23.8)	0.242 [§]
Wild type	33 (78.6)	37 (88.1)		38 (90.5)	32 (76.2)	
<i>NRAS/KRAS/n</i> (%)			0.533 [§]			0.533 [§]
Mutation	5 (11.9)	7 (16.7)	0.106 [§]	7 (16.7)	5 (11.9)	0.106 [§]
Wild type	37 (88.1)	35 (83.3)		35 (83.3)	37 (88.1)	
<i>TET2/n</i> (%)			0.106 [§]			0.332 [§]
Mutation	8 (19.0)	3 (7.1)	0.004 [§]	7 (16.7)	4 (9.5)	0.004 [§]
Wild type	34 (81.0)	39 (92.9)		35 (83.3)	38 (90.5)	
<i>TP53/n</i> (%)			0.004 [§]			0.106 [§]
Mutation	10 (23.8)	1 (2.4)	0.746 [§]	3 (7.1)	8 (19.0)	0.746 [§]
Wild type	32 (76.2)	41 (97.6)		39 (92.9)	34 (81.0)	
<i>MLL</i>			0.746 [§]			0.024 [§]
Positive	5 (11.9)	6 (14.3)	0.498 [§]	9 (21.4)	2 (4.8)	0.498 [§]
Negative	37 (88.1)	36 (85.7)		33 (78.6)	40 (95.2)	
Relapse/ <i>n</i> (%)			0.498 [§]			0.042 [§]
Yes	17 (40.5)	14 (33.3)	0.042 [§]	20 (47.6)	11 (26.2)	0.042 [§]
No	25 (59.5)	28 (66.7)		22 (52.4)	31 (73.8)	

WBC white blood cell, BM bone marrow, PB peripheral blood, FAB French American British

*denotes Mann–Whitney *U* test

§denotes chi-square test

TP53 mutation were independent risk factors for both EFS and OS (all $P < 0.05$). In addition, *DNMT3A* mutation was an independent risk prognostic factor for EFS ($P = 0.026$).

The combined prognostic effect *FUT4* and *FUT6* expression on AML

To further analyze the combined prognostic effects of the two *FUT* members with independent prognostic value, the chemotherapy-only group was further stratified according to the median expression levels of both *FUT4* and *FUT6*. We observed that the *FUT4*^{high}*FUT6*^{low} and *FUT4*^{high}*FUT6*^{high} subgroups had significantly longer EFS and OS than the *FUT4*^{low}*FUT6*^{high} group, especially the former (all $P < 0.01$, Fig. 2a, b).

Discussion

In our study, we found that high expressions of *FUT3/6/7* were poor prognostic indicators, and high *FUT4* expression was a good prognostic factor in AML patients who received chemotherapy alone, but their prognostic effects were not

observed in patients who underwent allo-HSCT, suggesting that allo-HSCT may neutralize their prognostic impact.

The relationship between the *FUTs* and tumorigenesis has been described in multiple solid tumors. High expressions of *FUT3* and *FUT6* can promote the expression of Sialy Lewis X (SLe^x), which in turn has a positive impact on the growth and invasion of oral squamous cell carcinoma. Inhibiting fucosylation may be useful to keep cancer stem cells from proliferation and metastatic spread [25]. In human pancreatic cancer cell lines, the N-fucosylation-related genes, *FUT3*, *FUT5*, and *FUT6*, are activated and may explain the elevated fucosylation on the cell surface and promote cancer cell metastasis [26]. The migratory ability of the Capan-1 cells is decreased with downregulated *FUT3*, making it a suitable target for reducing the metastatic virulence of pancreatic cancer [27]. In prostate adenocarcinoma, however, the expression of *FUT3* is lower than benign prostate alteration [28]. Moreover, in colorectal carcinoma patients, low expression of *FUT3* is associated with tumor infiltration and distant metastasis [29]. The dual role of *FUT3* in tumor cells suggests that different kinds of tumor modulate *FUT3* expression differently to acquire the most beneficial and adaptable characteristics during their

Table 3 Comparison of clinical and molecular characteristics between the high and low *FUT6* and *FUT7* expression subgroups among patients who only received chemotherapy alone

Characteristics	<i>FUT6</i>			<i>FUT7</i>		
	High (<i>n</i> = 42)	Low (<i>n</i> = 42)	<i>P</i> -value	High (<i>n</i> = 42)	Low (<i>n</i> = 42)	<i>P</i> -value
Age/years, median (range)	66.0 (22.0, 82.0)	67.0 (25.0, 88.0)	0.329*	66.0 (33.0, 88.0)	67.0 (22.0, 82.0)	0.671*
Age group/ <i>n</i> (%)			1.000§			0.637§
<60 years	13 (31.0)	13 (31.0)		12 (28.6)	14 (33.3)	
≥60 years	29 (69.0)	29 (69.0)		30 (71.4)	28 (66.7)	
Gender/ <i>n</i> (%)			0.512§			0.274§
Male	21 (50.0)	24 (57.1)		25 (59.5)	20 (47.6)	
Female	21 (50.0)	18 (42.9)		17 (40.5)	22 (52.4)	
Race/ <i>n</i> (%)			0.208§			0.450§
Caucasian	29 (69.0)	34 (81.0)		33 (78.6)	30 (71.4)	
Others	13 (31.0)	8 (19.0)		9 (21.4)	12 (28.6)	
WBC/× 10 ⁹ /L, median (range)	21.6 (0.7, 297.4)	12.2 (1.0, 131.5)	0.620*	28.5 (0.7, 297.4)	9.7 (1.0, 171.9)	0.038*
BM blasts/%, median (range)	74.0 (30.0, 99.0)	70.5 (32.0, 98.0)	0.561*	77.0 (30.0, 99.0)	69.5 (32.0, 95.0)	0.312*
PB blasts/%, median (range)	48.5 (0.0, 98.0)	12.0 (0.0, 97.0)	0.004*	33.0 (0.0, 98.0)	20.0 (0.0, 97.0)	0.635*
FAB subtypes/ <i>n</i> (%)						
M0	4 (9.5)	3 (7.1)	1.000§	3 (7.1)	4 (9.5)	1.000§
M1	14 (33.3)	6 (14.3)	0.040§	11 (26.2)	9 (21.4)	0.608§
M2	13 (31.0)	8 (19.0)	0.208§	9 (21.4)	12 (28.6)	0.450§
M4	6 (14.3)	14 (33.3)	0.040§	11 (26.2)	9 (21.4)	0.608§
M5	3 (7.1)	9 (21.4)	0.061§	8 (19.0)	4 (9.5)	0.212§
M6	0 (0.0)	1 (2.4)	1.000§	0 (0.0)	1 (2.4)	1.000§
M7	2 (4.8)	1 (2.4)	1.000§	0 (0.0)	3 (7.1)	0.241§
Cytogenetics/ <i>n</i> (%)						
Normal	23 (54.8)	17 (40.5)	0.190§	25 (59.5)	15 (35.7)	0.029§
t(9;22)/ <i>BCR-ABL1</i>	1 (2.4)	0 (0.0)	1.000§	0 (0.0)	1 (2.4)	1.000§
inv(16)/ <i>CBFβ-MYH11</i>	1 (2.4)	5 (11.9)	0.202§	0 (0.0)	6 (14.3)	0.026§
Complex	9 (21.4)	2 (4.8)	0.024§	5 (11.9)	6 (14.3)	0.746§
11q23/ <i>MLL</i>	1 (2.4)	2 (4.8)	1.000§	2 (4.8)	1 (2.4)	1.000§
-7/7q-	3 (7.1)	0 (0.0)	0.241§	1 (2.4)	2 (4.8)	1.000§
t(8;21)/ <i>RUNX1-RUNX1T1</i>	1 (2.4)	5 (11.9)	0.202§	0 (0.0)	6 (14.3)	0.026§
Others	3 (7.1)	11 (26.2)	0.019§	9 (21.4)	5 (11.9)	0.242§
Risk/ <i>n</i> (%)						
Good	2 (4.8)	10 (23.8)	0.013§	0 (0.0)	12 (28.6)	<0.001§
Intermediate	22 (52.4)	24 (57.1)	0.661§	27 (64.3)	19 (45.2)	0.079§
Poor	16 (38.1)	8 (19.0)	0.053§	13 (31.0)	11 (26.2)	0.629§
<i>FLT3</i> / <i>n</i> (%)			0.109§			0.291§
<i>FLT3</i> -ITD	11 (26.2)	4 (9.5)		9 (21.4)	6 (14.3)	
<i>FLT3</i> -TKD	4 (9.5)	3 (7.1)		5 (11.9)	2 (4.8)	
Wild type	27 (64.3)	35 (83.3)		28 (66.7)	34 (81.0)	
<i>NPM1</i> / <i>n</i> (%)			0.102§			0.102§
Mutation	17 (40.5)	10 (23.8)		17 (40.5)	10 (23.8)	
Wild type	25 (59.5)	32 (76.2)		25 (59.5)	32 (76.2)	
<i>DNMT3A</i> / <i>n</i> (%)			0.221§			0.087§
Mutation	14 (33.3)	9 (21.4)		15 (35.7)	8 (19.0)	
Wild type	28 (66.7)	33 (78.6)		27 (64.3)	34 (81.0)	

Table 3 (continued)

Characteristics	<i>FUT6</i>			<i>FUT7</i>		
	High (<i>n</i> = 42)	Low (<i>n</i> = 42)	<i>P</i> -value	High (<i>n</i> = 42)	Low (<i>n</i> = 42)	<i>P</i> -value
<i>IDH1/IDH2/n</i> (%)			0.776 [§]			0.776 [§]
Mutation	7 (16.7)	8 (19.0)		8 (19.0)	7 (16.7)	
Wild type	35 (83.3)	34 (81.0)		34 (81.0)	35 (83.3)	
<i>RUNX1/n</i> (%)			0.242 [§]			0.558 [§]
Mutation	5 (11.9)	9 (21.4)		8 (19.0)	6 (14.3)	
Wild type	37 (88.1)	33 (78.6)		34 (81.0)	36 (85.7)	
<i>NRAS/KRAS/n</i> (%)			0.212 [§]			0.533 [§]
Mutation	4 (9.5)	8 (19.0)		7 (16.7)	5 (11.9)	
Wild type	38 (90.5)	34 (81.0)		35 (83.3)	37 (88.1)	
<i>TET2/n</i> (%)			0.106 [§]			0.332 [§]
Mutation	8 (19.0)	3 (7.1)		4 (9.5)	7 (16.7)	
Wild type	34 (81.0)	39 (92.9)		38 (90.5)	35 (83.3)	
<i>TP53/n</i> (%)			0.024 [§]			0.746 [§]
Mutation	9 (21.4)	2 (4.8)		5 (11.9)	6 (14.3)	
Wild type	33 (78.6)	40 (95.2)		37 (88.1)	36 (85.7)	
<i>MLL</i>			0.746 [§]			0.106 [§]
Positive	5 (11.9)	6 (14.3)		8 (19.0)	3 (7.1)	
Negative	37 (88.1)	36 (85.7)		34 (81.0)	39 (92.9)	
Relapse/ <i>n</i> (%)			0.821 [§]			0.821 [§]
Yes	15 (35.7)	16 (38.1)		16 (38.1)	15 (35.7)	
No	27 (64.3)	26 (61.9)		26 (61.9)	27 (64.3)	

WBC white blood cell, BM bone marrow, PB peripheral blood, FAB French American British

*denotes Mann–Whitney *U* test

§denotes chi-square test

Table 4 Multivariate analysis of the potential prognostic factors of EFS and OS in AML patients who received chemotherapy alone

Variables	EFS		OS	
	HR (95%CI)	<i>P</i> -value	HR (95%CI)	<i>P</i> -value
<i>FUT3</i> (high vs. Low)		NS		NS
<i>FUT4</i> (high vs. Low)	0.423 (0.250, 0.715)	0.001	0.398 (0.239, 0.664)	<0.001
<i>FUT6</i> (high vs. Low)	1.871 (1.116, 3.137)	0.017	1.729 (1.060, 2.821)	0.028
<i>FUT7</i> (high vs. Low)		NS		NS
Age (≥60 vs. <60 years)	3.041 (1.629, 5.678)	<0.001	2.550 (1.386, 4.693)	0.003
WBC (≥15 vs. <15 × 10 ⁹ /L)	1.694 (0.975, 2.943)	0.062		NS
BM blasts (≥70 vs. <70%)	2.518 (1.487, 4.265)	0.001	2.126 (1.281, 3.530)	0.004
PB blasts (≥20 vs. <20%)		NS		NS
<i>FLT3-ITD</i> (positive vs. negative)		NS		NS
<i>NPM1</i> (mutated vs. wild)		NS		NS
<i>DNMT3A</i> (mutated vs. wild)	1.927 (1.082, 3.429)	0.026		NS
<i>RUNX1</i> (mutated vs. wild)	1.993 (0.954, 4.164)	0.066		NS
<i>TP53</i> (mutated vs. wild)	2.645 (1.207, 5.795)	0.015	2.230 (1.094, 4.545)	0.027

EFS Event-free survival, OS Overall survival, HR hazard ratio, CI confidence interval, WBC white blood cell, BM bone marrow, PB peripheral blood

establishment in the primary site and later during metastasis [30]. We found that high *FUT3* expression was more likely to occur in AML patients with complex karyotype and

TP53 mutation, yet its negative prognostic impact on AML was independent, suggesting that *FUT3* may also play a positive role in leukemogenesis.

Fig. 2 Comparison of EFS and OS between patients with low or high *FUT4* and *FUT6* expression in univariate analysis. **a, b** Patients in the *FUT4*^{low}*FUT6*^{high} groups had shorter EFS and OS than those in the *FUT4*^{high}*FUT6*^{low} and *FUT4*/*FUT6*^{high} groups

FUT7 controls the progression of the cell cycle via the PLCγ/extracellular signal-regulated kinase signaling pathway in hepatocellular carcinoma [31]. Its overexpression also promotes the occurrence of non-small cell lung cancer by activating the *EGFR/AKT/mTOR* signaling pathway [32]. In AML, our study pointed out that *FUT7* overexpression might also be a tumor-promoting gene, despite its coexistence with other well-known good prognostic factors, such as *CBFβ-MYH11* and *RUNX1-RUNX1T1*.

It has been reported that *FUT6* exhibits a crucial role in mediating the α1,3-fucosylation of glycoprotein during colorectal carcinoma cell and hepatocellular carcinoma cell proliferation [31, 33]. Overexpression of *FUT6* supports the bony metastases of prostate cancer, a biological function alien to both *FUT3* and *FUT7* [34]. On the contrary, Li et al. found that the overexpression of *FUT6*, activated by miR-106b downregulation, can significantly reduce the invasion, migration, and proliferation of human breast cancer cells [35], suggesting that *FUT6* may take part in different biological processes in different cancers. We observed that high *FUT6* expression was an independent adverse prognostic factor for AML patients, and it was associated with complex karyotype, poor cytogenetic risk, and *TP53* mutation, indicating that *FUT6* may be a potential therapeutic target for AML.

FUT4 is highly expressed in both colorectal adenocarcinoma cells and myeloid cells [36]. Studies have shown that high expression of *FUT4* is associated with poor survival in patients with colorectal and lung cancer [37, 38]. In breast cancer, *FUT4* activates the *PI3K/Akt* signaling pathway, leading to multidrug resistance [13]. MiR-26a/26b, miR-125a-5p, and miR-224-3p may also be involved in various *FUT4*-mediated tumorigenic processes [15, 25, 37]. Different from the above results in solid tumor, we found that high expression of *FUT4* was an independent good prognostic factor in AML, and it was associated with younger age and normal karyotype. This indicated that the role of *FUT4* in various diseases may also be different.

Furthermore, we found that AML patients with low *FUT4* expression and high *FUT6* expression had the shortest EFS and OS compared to other patients, indicating that the combination of *FUT4* and *FUT6* may be a better tool in AML prognostication. The detailed biological interactions between *FUT4* and *FUT6* are still unknown, but our results suggested that they may play different roles in leukemogenesis although they belong to the same gene family.

In multivariate analysis, age ≥60 years and BM blasts ≥70% were also independent risk factors for AML patients, consistent with previous findings that older AML patients generally have unfavorable prognosis due to poorer baseline performance status, higher mutation burden, and poorer tolerance to chemotherapy, and that abnormal proliferation of BM blasts has apparent adverse effects on survival [39, 40]. We found that mutations in *DNMT3A* and *TP53* were also independent risk factors for AML patients, consistent with other studies [41, 42]. Our results, while confirming previous study results, will help to refine risk stratification and provide evidence for precision medicine in AML.

In summary, our results have shown that high expressions of *FUT3/6/7* indicate poor prognosis, and high *FUT4* expression predicts good prognosis in AML, but their prognostic effects on survival may be neutralized by allo-HSCT. Moreover, *FUT4* and *FUT6* are stronger prognosticators for AML compared with the other *FUTs*, and their combined predictive effect is more prominent. Our study is limited by its small sample size and our results require larger prospective cohorts to verify. The mechanism of the combined action of *FUT4* and *FUT6* in AML needs to be further delineated.

Acknowledgements This work was supported by grants from the National Natural Science Foundation of China (81500118 and 61501519), the China Postdoctoral Science Foundation funded project (2016M600443), and Jiangsu Province Postdoctoral Science Foundation funded project (1701184B).

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

- Goardon N, Marchi E, Atzberger A, Quek L, Schuh A, Soneji S, et al. Coexistence of LMPP-like and GMP-like leukemia stem cells in acute myeloid leukemia. *Cancer Cell*. 2011;19:138–52.
- Zhang J, Gu Y, Chen B. Mechanisms of drug resistance in acute myeloid leukemia. *Onco Targets Ther*. 2019;12:1937–45.
- Mrozek K, Marcucci G, Nicolet D, Maharry KS, Becker H, Whitman SP, et al. Prognostic significance of the European LeukemiaNet standardized system for reporting cytogenetic and molecular alterations in adults with acute myeloid leukemia. *J Clin Oncol*. 2012;30:4515–23.
- Zhu YM, Wang PP, Huang JY, Chen YS, Chen B, Dai YJ, et al. Gene mutational pattern and expression level in 560 acute myeloid leukemia patients and their clinical relevance. *J Transl Med*. 2017;15:178.
- Wakita S, Yamaguchi H, Omori I, Terada K, Ueda T, Manabe E, et al. Mutations of the epigenetics-modifying gene (DNMT3a, TET2, IDH1/2) at diagnosis may induce FLT3-ITD at relapse in de novo acute myeloid leukemia. *Leukemia*. 2013;27:1044–52.
- Su L, Gao SJ, Li W, Tan YH, Cui JW, Hu RP. NPM1, FLT3-ITD, CEBPA, and c-kit mutations in 312 Chinese patients with de novo acute myeloid leukemia. *Hematology*. 2014;19:324–8.
- Fasan A, Haferlach C, Alpermann T, Jeromin S, Grossmann V, Eder C, et al. The role of different genetic subtypes of CEBPA mutated AML. *Leukemia*. 2014;28:794–803.
- Shamaa S, Laimon N, Aladle DA, Azmy E, Elghannam DM, Salem DA, et al. Prognostic implications of NPM1 mutations and FLT3 internal tandem duplications in Egyptian patients with cytogenetically normal acute myeloid leukemia. *Hematology*. 2014;19:22–30.
- Cheng Z, Dai Y, Pang Y, Jiao Y, Zhao H, Zhang Z, et al. Enhanced expressions of FHL2 and iASPP predict poor prognosis in acute myeloid leukemia. *Cancer Gene Ther*. 2019;26:17–25.
- Zhang L, Li R, Hu K, Dai Y, Pang Y, Jiao Y, et al. Prognostic role of DOK family adapters in acute myeloid leukemia. *Cancer Gene Ther*. <https://doi.org/10.1038/s41417-018-0052-z>.
- Ma B, Simala-Grant JL, Taylor DE. Fucosylation in prokaryotes and eukaryotes. *Glycobiology*. 2006;16:158r–184r.
- Varki A, Cummings RD, Esko JD, Freeze HH, Stanley P, Marth JD, et al. Symbol nomenclature for glycan representation. *Proteomics*. 2009;9:5398–9.
- Cheng L, Luo S, Jin C, Ma H, Zhou H, Jia L. FUT family mediates the multidrug resistance of human hepatocellular carcinoma via the PI3K/Akt signaling pathway. *Cell Death Dis*. 2013;4:e923.
- Liu YC, Yen HY, Chen CY, Chen CH, Cheng PF, Juan YH, et al. Sialylation and fucosylation of epidermal growth factor receptor suppress its dimerization and activation in lung cancer cells. *Proc Natl Acad Sci USA*. 2011;108:11332–7.
- Liang L, Gao C, Li Y, Sun M, Xu J, Li H, et al. miR-125a-3p/FUT5-FUT6 axis mediates colorectal cancer cell proliferation, migration, invasion and pathological angiogenesis via PI3K-Akt pathway. *Cell Death Dis*. 2017;8:e2968.
- Weston BW, Hiller KM, Mayben JP, Manousos GA, Bendt KM, Liu R, et al. Expression of human alpha(1,3)fucosyltransferase antisense sequences inhibits selectin-mediated adhesion and liver metastasis of colon carcinoma cells. *Cancer Res*. 1999;59:2127–35.
- Iwamori M, Tanaka K, Kubushiro K, Lin B, Kiguchi K, Ishiwata I, et al. Alterations in the glycolipid composition and cellular properties of ovarian carcinoma-derived RMG-1 cells on transfection of the alpha1,2-fucosyltransferase gene. *Cancer Sci*. 2005;96:26–30.
- Zhao Y, Lin B, Hao YY. The effects of Lewis(y) antigen content on drug resistance to carboplatin in ovarian cancer line RMG-I. *Prog Biochem Biophys*. 2008;35:1175–82.
- Yang XS, Zhang ZB, Jia SA, Liu YJ, Wang XQ, Yan Q. Overexpression of fucosyltransferase IV in A431 cell line increases cell proliferation. *Int J Biochem. Cell B*. 2007;39:1722–30.
- Wang QY, Guo P, Duan LL, Shen ZH, Chen HL. alpha-1,3-Fucosyltransferase-VII stimulates the growth of hepatocarcinoma cells via the cyclin-dependent kinase inhibitor p27Kip1. *Cell Mol Life Sci*. 2005;62:171–8.
- Hiller KM, Mayben JP, Bendt KM, Manousos GA, Senger K, Cameron HS, et al. Transfection of alpha(1,3)fucosyltransferase antisense sequences impairs the proliferative and tumorigenic ability of human colon carcinoma cells. *Mol Carcinog*. 2000;27:280–8.
- Kang X, Wang N, Pei C, Sun L, Sun R, Chen J, et al. Glycan-related gene expression signatures in human metastatic hepatocellular carcinoma cells. *Exp Ther Med*. 2012;3:415–22.
- Cancer Genome Atlas Research Network, Ley TJ, Miller C, Ding L, Raphael BJ, Mungall AJ, et al. Genomic and epigenomic landscapes of adult de novo acute myeloid leukemia. *N Engl J Med*. 2013;368:2059–74.
- Cheson BD, Bennett JM, Kopecky KJ, Buchner T, Willman CL, Estey EH, et al. Revised recommendations of the international working group for diagnosis, standardization of response criteria, treatment outcomes, and reporting standards for therapeutic trials in acute myeloid leukemia. *J Clin Oncol*. 2003;21:4642–9.
- Desiderio V, Papagerakis P, Tirino V, Zheng L, Matossian M, Prince ME, et al. Increased fucosylation has a pivotal role in invasive and metastatic properties of head and neck cancer stem cells. *Oncotarget*. 2015;6:71–84.
- Gao HF, Wang QY, Zhang K, Chen LY, Cheng CS, Chen H, et al. Overexpressed N-fucosylation on the cell surface driven by FUT3, 5, and 6 promotes cell motilities in metastatic pancreatic cancer cell lines. *Biochem Biophys Res Commun*. 2019;511:482–9.
- Zhan L, Chen L, Chen Z. Knockdown of FUT3 disrupts the proliferation, migration, tumorigenesis and TGF- β induced EMT in pancreatic cancer cells. *Oncol Lett*. 2018;16:924–30.
- Vasconcelos JLD, Ferreira SD, de Lima ALR, Rego MJBD, Bandeira ARG, Cavalcanti CDB, et al. Comparing the Immunoreexpression of FUT3 and FUT6 between Prostatic Adenocarcinoma and Benign Prostatic Hyperplasia. *Acta Histochem Cytoc*. 2013;46:105–9.
- Petretti T, Kemmner W, Schulze B, Schlag PM. Altered mRNA expression of glycosyltransferases in human colorectal carcinomas and liver metastases. *Gut*. 2000;46:359–66.
- do Nascimento JC, Ferreira Sde A, Vasconcelos JL, da Silva-Filho JL, Barbosa BT, Bezerra MF, et al. Fut3 role in breast invasive ductal carcinoma: Investigating its gene promoter and protein expression. *Exp Mol Pathol*. 2015;99:409–15.
- Li D, Sun H, Bai G, Wang W, Liu M, Bao Z, et al. alpha-1,3-Fucosyltransferase-VII siRNA inhibits the expression of SLex and hepatocarcinoma cell proliferation. *Int J Mol Med*. 2018;42:2700–8.

32. Liang JX, Gao W, Cai L. Fucosyltransferase VII promotes proliferation via the EGFR/AKT/mTOR pathway in A549 cells. *Oncotargets Ther.* 2017;10:3971–8.
33. Pan S, Liu Y, Liu Q, Xiao Y, Liu B, Ren X, et al. HOTAIR/miR-326/FUT6 axis facilitates colorectal cancer progression through regulating fucosylation of CD44 via PI3K/AKT/mTOR pathway. *Biochim Biophys Acta, Mol Cell Res.* 2019;1866:750–60.
34. Li J, Guillebon AD, Hsu JW, Barthel SR, Dimitroff CJ, Lee YF, et al. Human fucosyltransferase 6 enables prostate cancer metastasis to bone. *Br J Cancer.* 2013;109:3014–22.
35. Li N, Liu Y, Miao Y, Zhao L, Zhou H, Jia L. MicroRNA-106b targets FUT6 to promote cell migration, invasion, and proliferation in human breast cancer. *IUBMB life.* 2016;68:764–75.
36. Taniguchi A, Suga R, Matsumoto K. Expression and transcriptional regulation of the human α 1, 3-fucosyltransferase 4 (FUT4) gene in myeloid and colon adenocarcinoma cell lines. *Biochem Biophys Res Commun.* 2000;273:370–6.
37. Li Y, Sun Z, Liu B, Shan Y, Zhao L, Jia L. Tumor-suppressive miR-26a and miR-26b inhibit cell aggressiveness by regulating FUT4 in colorectal cancer. *Cell Death Dis.* 2017;8:e2892.
38. Ogawa J, Inoue H, Koide S. Expression of α -1,3-fucosyltransferase type IV and VII genes is related to poor prognosis in lung cancer. *Cancer Res.* 1996;56:325–9.
39. Schoen MW, Woelich SK, Braun JT, Reddy DV, Fesler MJ, Petruska PJ, et al. Acute myeloid leukemia induction with cladribine: Outcomes by age and leukemia risk. *Leuk Res.* 2018;68:72–78.
40. Bacher U, Haferlach C, Alpermann T, Kern W, Schnittger S, Haferlach T. Comparison of genetic and clinical aspects in patients with acute myeloid leukemia and myelodysplastic syndromes all with more than 50% of bone marrow erythropoietic cells. *Haematologica.* 2011;96:1284–92.
41. Kumar D, Mehta A, Panigrahi MK, Nath S, Saikia KK. DNMT3A (R882) mutation features and prognostic effect in acute myeloid leukemia in coexistent with NPM1 and FLT3 mutations. *Hematol Oncol Stem Cell Ther.* 2018;11:82–89.
42. Welch JS. Patterns of mutations in TP53 mutated AML. *Best Pr Res Clin Haematol.* 2018;31:379–83.