University of Groningen

The Rgg1518 transcriptional regulator is a necessary facet of sugar metabolism and virulence in *Streptococcus pneumoniae*

Shlla, Bushra; Gazioglu, Ozcan; Shafeeq, Sulman; Manzoor, Irfan; Kuipers, Oscar P; Ulijasz, Andrew; Hiller, N Luisa; Andrew, Peter W; Yesilkaya, Hasan

Published in:
Molecular Microbiology

DOI:
[10.1111/mmi.14788](https://doi.org/10.1111/mmi.14788)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Shlla, B., Gazioglu, O., Shafeeq, S., Manzoor, I., Kuipers, O. P., Ulijasz, A., Hiller, N. L., Andrew, P. W., & Yesilkaya, H. (2021). The Rgg1518 transcriptional regulator is a necessary facet of sugar metabolism and virulence in *Streptococcus pneumoniae*. *Molecular Microbiology*, 116(3), 996-1008.
<https://doi.org/10.1111/mmi.14788>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

RESEARCH ARTICLE

WILEY

The Rgg1518 transcriptional regulator is a necessary facet of sugar metabolism and virulence in *Streptococcus pneumoniae*

Bushra Shlla^{1,2} | Ozcan Gazioglu¹ | Sulman Shafeeq³ | Irfan Manzoor³ |
Oscar P. Kuipers ³ | Andrew Ulijasz⁴ | N. Luisa Hiller⁵ | Peter W. Andrew¹ |
Hasan Yesilkaya ¹

¹Department of Respiratory Sciences,
University of Leicester, Leicester, UK

²Department of Biology, College of Science,
University of Mosul, Mosul, Iraq

³Molecular Genetics, University of
Groningen, Groningen, The Netherlands

⁴Department of Microbiology and
Immunology, Loyola University Chicago,
Maywood, Illinois, USA

⁵Department of Biological Sciences,
Carnegie Mellon University, Pittsburgh,
Pennsylvania, USA

Correspondence

Hasan Yesilkaya, Department of Respiratory
Sciences, University of Leicester, Leicester,
LE1 7RH, UK.
Email: hy3@le.ac.uk

Funding information

National Institute of Allergy and Infectious
Diseases, Grant/Award Number: R01
AI135060-01A1 and R01 AI139077-01A1

Abstract

Rggs are a group of transcriptional regulators with diverse roles in metabolism and virulence. Here, we present work on the Rgg1518/SHP1518 quorum sensing system of *Streptococcus pneumoniae*. The activity of Rgg1518 is induced by its cognate peptide, SHP1518. In vitro analysis showed that the Rgg1518 system is active in conditions rich in galactose and mannose, key nutrients during nasopharyngeal colonization. Rgg1518 expression is highly induced in the presence of these sugars and its isogenic mutant is attenuated in growth on galactose and mannose. When compared with other Rgg systems, Rgg1518 has the largest regulon on galactose. On galactose it controls up- or downregulation of a functionally diverse set of genes involved in galactose metabolism, capsule biosynthesis, iron metabolism, protein translation, as well as other metabolic functions, acting mainly as a repressor of gene expression. Rgg1518 is a repressor of capsule biosynthesis, and binds directly to the capsule regulatory region. Comparison with other Rggs revealed inter-regulatory interactions among Rggs. Finally, the *rgg1518* mutant is attenuated in colonization and virulence in a mouse model of colonization and pneumonia. We conclude that Rgg1518 is a virulence determinant that contributes to a regulatory network composed of multiple Rgg systems.

KEYWORDS

capsule synthesis, Rgg, *Streptococcus pneumoniae*, sugar metabolism, transcriptional regulation, virulence

1 | INTRODUCTION

Streptococcus pneumoniae (pneumococcus) causes a diverse array of infections. It can exist as an asymptomatic colonizer of the upper respiratory tract or can spread to middle ear, lower respiratory tract, blood, and central nervous system, causing diseases such as otitis media, pneumonia, septicemia, and meningitis, respectively (Weiser et al., 2018). *S. pneumoniae* is the leading agent of lower respiratory infection morbidity and mortality globally, causing more deaths than those caused by other major respiratory infectious agents combined

in 2016 (Troeger et al., 2017). Unfortunately, there is a rising trend in the number of antibiotic-resistant pneumococcal strains. The danger posed by the rising antibiotic resistance was recognized by the World Health Organization, hence *S. pneumoniae* has been declared as a priority pathogen for which effective antibiotics are urgently needed.

The study of how the pneumococcus alters its phenotype in health and disease, and in different tissue sites, can reveal novel drug targets. *S. pneumoniae* encounters diverse environmental conditions in its human host, including fluctuations in oxygen and

metal concentration, differences in concentration and type of nutrients, and changes in temperature (Hajaj et al., 2017; Sahin-Yilmaz & Naclerio, 2011; Shafeeq et al., 2011). This microbe has an innate capacity to sense and respond to changing environmental stimuli, employing various mechanisms including regulatory RNAs, stand-alone “one-component” regulators, and two-component regulatory systems (TCS) (Cuevas et al., 2017; Gómez-Mejía et al., 2018; Kadam et al., 2017; Wilton et al., 2015; Zhi et al., 2018). These mechanisms modulate an array of essential cellular functions, including acquisition and metabolism of nutrients, detoxification of reactive oxygen species, incorporation of genetic material, and expression of virulence factors.

Quorum sensing (QS) systems play a key role in sensing and responding to the environment. These multi-component regulatory mechanisms coordinate bacterial population responses to diverse stimuli (Cuevas et al., 2017; Kadam et al., 2017; Motib et al., 2019). The operation of these cell-cell communication systems relies on secretion and subsequent detection of signaling molecules by the producing and neighboring cells. The response directs community-level synchronization of transcription, enabling microbes to adopt a phenotype consistent with the requirements of the specific host niche (Rutherford & Bassler, 2012). The pneumococcal QS systems can be divided broadly into two major categories: Autoinducer-2 (AI-2) and peptide-mediated QS systems. AI-2 signaling molecule synthesis relies on the metabolic enzyme LuxS, which catalyzes the conversion of S-ribosyl-homocysteine into homocysteine. It was shown by Trapetti et al that AI-2 QS signaling is mediated through a fructose-specific phosphoenolpyruvate-phosphotransferase system (FruA), and that AI-2 triggers upregulation of galactose catabolism through the Leloir pathway, ultimately enhancing capsule biosynthesis (Trapetti et al., 2017). In addition, the LuxS/AI-2 QS system plays an important role in biofilm synthesis, competence, and fratricide (Trapetti et al., 2017).

Peptide-mediated pneumococcal QS systems are divided into three main groups based on peptide sequence features, transporters they utilize, and receptors; these are: (i) double glycine processing site peptides, (ii) peptides associated with the RRNPP (Rap, Rgg, NprR, PlcR, and PrgX) superfamily of quorum-sensing proteins, and (iii) lanthionine-containing peptides (Aggarwal et al., 2020; Cook et al., 2013).

Pneumococcal double glycine peptides, such as the competence stimulating peptide (CSP) and the bacteriocin inducing peptide (BIP), contain a conserved N-terminal leader sequence that terminates commonly in Gly-Gly residues. These peptides are exported after cleavage of the leader sequence via peptidase-containing ABC transporters. After secretion, some double-glycine processed peptides induce a response in the neighboring cell by signaling via two-component systems (TCS) (Håvarstein et al., 1995; de Saizieu et al., 2000; Wang et al., 2020). The peptide interacts with the histidine kinase of the TCS, stimulates its autophosphorylation and transphosphorylation of the cognate response regulator, and initiate changes in the transcriptional composition of the cell.

Cyclic peptides, such as lanthionine-containing peptides, undergo post-translational modifications that result in the introduction of thioether amino acids lanthionine and methyllanthionine (Willey & van der Donk, 2007). Their distinctive structure is formed by LanBC or LanM enzymes, when serine or threonine residues in the pro-peptide are dehydrated and linked to cysteine thiols. The peptide is exported by dedicated LanT transporters (Koponen et al., 2002; Sen et al., 1999; Willey & van der Donk, 2007) and in various cases, autoregulation via a two-component regulatory system also occurs, as exemplified by nisin autoinduction via NisRK in *Lactococcus lactis* (Kuipers et al., 1995).

Two RRNPP type QS systems have been studied in *S. pneumoniae*: Tpr/Phr and Rgg/SHP regulators (Cuevas et al., 2017; Hoover et al., 2015; Motib et al., 2019; Zhi et al., 2018). The Tpr (transcription factor regulated by Phr peptide) and Rgg (Regulator gene of glycosyltransferase) transcription factors are positively regulated by their cognate peptides Phr (phosphatase regulator) and SHP (short hydrophobic peptides), respectively. The Tprs are part of the PlcR family, the best-characterized pair, TprA/PhrA, play a critical role in pneumococcal colonization and virulence (Motib et al., 2019). Multiple Rgg regulators have been characterized in pneumococcus and other streptococcal species. The model that emerges is that SHPs are cleaved into active forms during export from precursor polypeptides that are typically shorter than 35 residues (Aggarwal et al., 2014; Aggarwal et al., 2020; Zhi et al., 2018). Multiple streptococcal species export SHPs through an ABC transporter, PptAB, whose homolog in *Enterococcus faecalis* exports sex pheromones (Chang & Federle, 2016; Varahan et al., 2014). The exact mechanism(s) directing SHPs to their transporter remains unclear. The mature peptides are re-imported into the cell via an oligopeptide permease system, where they interact with the cognate Rgg regulator, altering the cell's transcriptional state (Aggarwal et al., 2020; Chang et al., 2011; Fleuchot et al., 2011; Zhi et al., 2018).

The pneumococcal pangenome encodes eight *rgg* homologs. In type 2 D39 strain there are six Rgg paralogues. Previously we characterized Rgg1952, Rgg144/SHP144, and Rgg939/SHP939 (Bortoni et al., 2009; Junges et al., 2017; Zhi et al., 2018). Rgg1952 and Rgg144 are part of the core genome. Rgg1952 is important in oxidative stress resistance. The Rgg144/SHP144 system represses capsule biosynthesis and induces VP1, a peptide that promotes biofilm development and virulence (Cuevas et al., 2017; Zhi et al., 2018). Its regulon contains genes involved in replication, recombination, translation, and nucleotide transport and metabolism (Zhi et al., 2018). It also controls multiple sugar transporters and its regulon is sugar-specific, with a broad response in mannose and a restricted response in galactose. Finally, Rgg939 is part of the accessory genome (Junges et al., 2017; Zhi et al., 2018). The diversity of genes regulated by Rgg939/SHP939 varies in response to environmental conditions, with an extensive regulon during growth on mannose and a limited regulon in galactose (Junges et al., 2017; Zhi et al., 2018). The regulon is predicted to function in cell division, iron transport, cell membrane biogenesis, and metabolism among others. Rgg939 also regulates capsule synthesis. While Zhi et al. (2018) suggested that Rgg939 can directly inhibit expression from the capsule locus,

Junges et al. (2017), showed that Rgg939 expression positively regulates the expression of genes involved in the production of compounds present in capsular polysaccharides of serotypes 12A and 12F. Rgg939/SHP939 signaling downregulated biofilm development on A549 epithelial cells (Junges et al., 2017). In murine models of disease, Rgg939/SHP939 was shown to be associated with a reduction in bacterial fitness in a pneumonia model (Junges et al., 2017), or alternatively, increased nasopharyngeal colonization and virulence in carriage and pneumonia models (Zhi et al., 2018).

In this study, we investigated the role of Rgg1518 (SPD_1518). This Rgg appears to be at the intersection of multiple RRNPP regulators, including Rgg144, Rgg939, and TprA. It is part of the accessory pneumococcal genome; present in approximately half of strains. We identified its peptide pheromone SHP1518, whose induction requires Rgg1518. Rgg1518 is highly induced on mannose and galactose, with an extensive regulon on galactose controlling the expression of genes involved in sugar transport, galactose metabolism, and capsule synthesis. Finally, Rgg1518 promotes colonization and contributes to lung disease.

2 | RESULTS

2.1 | Rgg1518 promotes growth on galactose and mannose

Previous studies on pneumococcal peptide-mediated RRNPP family QS systems, namely Rgg144/SHP144, Rgg939/SHP939, and TprA/PhrA, indicated their role in host derived sugar metabolism (Motib et al., 2019; Zhi et al., 2018). Due to its homology to other Rggs, we hypothesized that Rgg1518 may also play a role in host-derived sugar metabolism, such as for galactose and mannose, which are plentiful in N- and O-linked glycans (King, 2010). To determine if Rgg1518 has a role in nutrient utilization, wild-type and Δ rgg1518 strains were grown in Brain heart infusion (BHI) and chemically defined medium (CDM) supplemented either with glucose, galactose, or mannose. No difference in growth profiles was detected in BHI (Supporting information Figure S1; $p > .05$). Likewise, the strains grew equally well on CDM supplemented with glucose (Figure 1). In contrast, on galactose and mannose, the growth rate ($0.037 \pm 0.002 \text{ hr}^{-1}$ and $0.093 \pm 0.004 \text{ hr}^{-1}$, $n = 6$) and yield of Δ rgg1518 was significantly lower than that of the wild type D39 ($0.150 \pm 0.001 \text{ hr}^{-1}$, and $0.111 \pm 0.0028 \text{ hr}^{-1}$, $n = 6$, respectively), and the complemented strain rescued the wild-type growth ($0.14 \pm 0.001 \text{ hr}^{-1}$; Figure 1). Notably, on galactose, the growth of the Δ rgg1518 displayed not only a lower max OD but also an extended lag phase. We conclude that *rgg1518* plays an important role in the utilization of galactose and mannose.

2.2 | The promoter for *rgg1518* is induced by galactose and mannose

The growth attenuation on galactose and mannose indicated a potential role for Rgg1518 on the metabolism of these sugars. To

FIGURE 1 Growth profiles of pneumococcal strains in chemically defined medium supplemented with 55 mM glucose, galactose, or mannose. Experiment was repeated using three replicates of six independent biological samples; vertical bars indicate standard error of means (SEM). Significant differences were seen by comparing the growth rates of mutant strains to the wild type D39 using one-way ANOVA and Dunnett's multiple comparisons test (* $p < .05$, **** $p < .0001$)

investigate this further, we determined if the *rgg1518* promoter (P) could be induced by galactose and mannose using LacZ reporter strains. Strains were grown in CDM-Glucose and then transferred to CDM with no sugar, or a single sugar for 1 hr. No significant induction of *Prgg1518::lacZ*-wild-type was observed on glucose ($176 \pm 3.0 \text{ MU}$, $n = 3$) relative to the no sugar control ($172 \pm 2 \text{ MU}$, $n = 3$) ($p > .05$). In contrast, a significant promoter induction was detected on galactose ($969 \pm 6 \text{ MU}$, $n = 3$), and mannose ($503 \pm 13 \text{ MU}$, $n = 3$; $p < .001$ and $p < .01$, for galactose and mannose) compared to no sugar control. These results show that the *rgg1518* promoter is inducible by galactose and mannose (Figure 2).

2.3 | Rgg1518 and SHP1518 form a quorum-sensing system

Out of six Rgg homologous present in type 2 D39 pneumococcal strain, only Rgg144/SHP144 and Rgg939/SHP939 were demonstrated to be peptide-mediated QS systems (Zhi et al., 2018). A recent study in D39 reported a putative *shp1518* gene encoding for a 33 aa long peptide (MGFKYLNLPKNSGFLIWSWIQLIWFETWFWG) (Wang et al., 2020). The gene is located 93 nucleotides away from *rgg1518* and inversely transcribed relative to *rgg1518*. We tested whether Rgg1518 and SHP1518 form a QS system. To this end, we tested the induction of *Pshp1518::lacZ* in wild type and Δ *rgg1518* backgrounds (Figure 3a). Induction of *Pshp1518* required Rgg1518, as the deletion of *rgg1518* led to a 3.75-fold decrease in *Pshp1518* driven β -galactosidase activity relative to wild-type. Moreover, similar to *Prgg1518*, the *Pshp1518* activity was significantly induced in galactose but not glucose. These results demonstrate that SHP1518 is induced by galactose and positively regulated by Rgg1518.

Next, we identified an active form of SHP1518. We generated synthetic 12- (IQLIWFETWFWG) and 16- (IWSWIQLIWFETWFWG) residues C-terminal truncations of SHP1518. These peptides were evaluated for their ability to induce *Pshp1518* in wild type and Δ *rgg1518*. The reporter assay was performed on glucose, rather than galactose, to reduce the induction of *Pshp1518* by the native SHP1518. The 12 aa peptide did not induce *Pshp1518* (data not shown). In contrast, the 16aa peptide showed a gradual

dose-dependent induction of *Pshp1518* relative to control. The difference in induction was significant and even at 5 nM and at 50 nM there was more than twofold increase relative to the no peptide control ($p < .01$). Furthermore, the induction by the synthetic peptide required the presence of Rgg1518 (Figure 3b). We conclude that SHP1518 serves as the cognate peptide for Rgg1518.

SHPs have been defined and grouped based on (i) the presence of lysine or arginine residues in their N-terminus and glutamate or aspartate residues at their C-terminus, and (ii) encoded by a gene upstream of an *rgg* homolog and divergently transcribed. SHP1518 fits these criteria. SHPs are further classified into three groups based on the presence of aspartate (Group 1) or glutamate (Group 2) in the C-terminus (Ibrahim et al., 2007). Additionally, Group 3 peptides meet the criteria for SHP but are encoded by genes overlapping the end of the *rgg* genes in a convergent orientation (Fleuchot et al., 2011). Unlike Group 3, the *shp1518* sequence does not overlap with *rgg1518*. Thus, SHP1518 can be classified as Group 2 based on its position on the locus and the glutamate residue in the C-terminus.

2.4 | Role of Rgg1518 in the regulation of other pneumococcal Rggs

Previously, we reported the interaction between different Rgg/SHP systems, and a summary of regulon overlap among different Rggs on galactose and mannose have been graphically

FIGURE 2 Induction analysis of *rgg1518* promoter by different sugars using *lacZ* reporter assay. Strains were grown in chemically defined medium (CDM)-Glucose and then transferred to CDM with no sugar, or a single sugar for 1 hr. Miller unit is expressed as nmol p-nitrophenol/min/ml. The values are the average of three independent experiments, each with three replicates, and vertical bars indicate standard error of means (SEM; ** $p < .01$ and *** $p < .001$ compared to the expression on galactose)

FIGURE 3 Expression levels (in Miller units) of pneumococcal transcriptional *lacZ*-fusion to the promoter region of *shp1518* (*Pshp1518::lacZ*) in wild type and Δ *rgg1518* in glucose and galactose (a), and in the absence and presence of synthetic SHP1518 peptide in cultures grown in chemically defined medium supplemented with glucose (b)

illustrated in Figure 4a and Supporting information Figure S3. Specifically, we found that the presence of non-cognate Rgg regulators is needed for maximal induction of SHP144 and SHP939 (Zhi et al., 2018). Further, in our previous work, both Rgg144 and Rgg939 were found to influence the transcription of adjacent to Rgg1518 (SPD_1513-1517); which here we show is also controlled by Rgg1518 (Zhi et al., 2018; Table 1).

To further corroborate the interactions within the Rgg network, a *Prgg1518-lacZ* construct was introduced into Δ rgg1518, Δ rgg144, Δ rgg939, and the wild-type D39, generating the reporter strains *Prgg1518::lacZ- Δ rgg1518*, *Prgg1518::lacZ- Δ rgg144*, *Prgg1518::lacZ- Δ rgg939*, and *Prgg1518::lacZ-wt*, respectively. β -Galactosidase activity in the reporter strains was measured in CDM supplemented with galactose. The β -galactosidase activity was lower for *Prgg1518::lacZ- Δ rgg1518* (29.23 ± 1.97 MU, $n = 3$) compared to the activity in the wild type (970 ± 16 MU, $n = 3$), showing that Rgg1518 is required for its own expression ($p < .0001$; Figure 4). In addition, full induction of *Prgg1518* required

FIGURE 4 Analysis of regulatory interactions among Rggs. (a) A Venn diagram showing the overlap between three pneumococcal Rggs. The diagram was produced by merging the differentially regulated genes on galactose and mannose for each Rgg by using VENNY (<http://bioinfogp.cnb.csic.es/tools/venny/index.html>). (b) The activity of β -galactosidase, driven by *rgg1518* putative promoter in different mutant strain backgrounds as measured in chemically defined medium-Gal, is expressed as nmol p-nitrophenol/min/ml. The values are the average of three independent experiments each with three replicates vertical bars indicate standard error of means (SEM; **** $p < .0001$ compared to its expression in the wild type D39)

the presence of Rgg144 and Rgg939, as the induction of *Prgg1518* was significantly lower in the Δ rgg144 and Δ rgg939 compared to the induction in the wild type (*Prgg1518::lacZ-wt*) ($p < .0001$). We conclude that multiple Rggs form a regulatory network when pneumococci are metabolizing galactose.

2.5 | Characterization of the Rgg1518 regulon

In the respiratory tract, the pneumococcus uses both galactose and mannose from host glycans as a source of energy (King, 2010). Thus, we employed microarray analysis to compare gene expression between a wild-type and a strain with a deletion in *rgg1518* (Δ rgg1518). On galactose, we observed differential expression of 378 genes. Under these conditions 80 were downregulated and 298 were upregulated in the mutant (Supporting information Table S1). This suggests that Rgg1518/SHP1518 acts mainly as a repressor, at least in the condition tested (CDM-galactose, mid-log phase). These genes are organized into 61 operons with an annotated role in the synthesis of capsule, fatty acid, hypothetical proteins, sugar and metal transporters, proteins involved in galactose metabolism as well as several transcriptional regulators. Moreover, among the highly regulated genes (4–9 fold difference) is the adjacent locus, encoding SPD_1513-1517. A selection of differentially expressed loci in Δ rgg1518 relative to the wild type are shown in Table 1. On mannose, the size of the regulon ($n = 51$) was smaller than that of galactose: 44 genes were downregulated and 7 were upregulated in the mutant, representing six operons (Supporting information Table S2). Notably, the expression of genes adjacent to *rgg1518* (SPD_1513-SPD_1517) was downregulated in the mutant relative to the wild type on mannose. It is very likely that the expression of some of these genes is affected indirectly by Rgg1518, hence additional studies are needed to determine genes directly controlled by Rgg1518.

2.6 | Rgg1518 is a repressor of capsule synthesis on galactose

Capsular polysaccharide (CPS) is an essential pneumococcal virulence determinant (Hyams et al., 2010). The microarray data revealed that Rgg1518 negatively regulates the capsule, as shown by the up-regulation of the capsule synthesis locus (SPD_0315-SPD_0328) in Δ rgg1518 on galactose (Tables 1 and S1). The metabolism of galactose generates precursors for capsule synthesis (Carvalho et al., 2011). The type 2 capsule contains glucuronic acid (Kenne et al., 1975), therefore we quantified glucuronic acid in pneumococcal strains grown in CDM supplemented either with galactose or glucose (Figure 5). On galactose, the production of capsular polysaccharide was significantly higher in Δ rgg1518 ($105 \pm 4 \mu\text{g}/10^9$ CFU, $n = 6$) than the wild type ($63 \pm 3 \mu\text{g}/10^9$ CFU, $n = 6$), and the complemented mutant rescued wild type levels ($63 \pm 4 \mu\text{g}/10^9$ CFU, $n = 6$; $p < .001$; Figure 5a). On glucose, there was no significant difference

TABLE 1 Some of the notable genomic loci that are found to be differentially expressed in Δ rgg1518 relative to the wild type D39 grown in chemically defined medium supplemented with galactose

Locus tag	Fold difference	Putative function
SPD_0113-SPD_0124	2.09-8.53	Hypothetical protein
SPD_0145-SPD_0150	2.27-3.66	CAAX amino terminal protease family protein
SPD_0179-SPD_0182	2.18-6.06	Lipoprotein synthesis
SPD_0192-SPD_0219	2.66-8.71	Ribosomal protein synthesis
SPD_0315-SPD_0334	2.28-6.36	Capsule locus synthesis
SPD_0378-SPD_0390	2.16-7.19	Fatty acid biosynthesis
SPD_0404-SPD_0408	2.08-3.09	Acetolactate synthase synthesis
SPD_0705-SPD_0710	2.16-2.48	Cell morphology/division
SPD_0750-SPD_0757	2.34-4.33	Membrane protein synthesis
SPD_0760-SPD_0766	2.14-5.72	FeS assembly
SPD_0905-SPD_0913	2.4-266	Genetic information processing
SPD_0942-SPD_0951	-2.06-2.27	UDP-N-acetyl-D-mannosaminuronic acid dehydrogenase
SPD_0989-SPD_0991	3.62-3.83	Protein synthesis
SPD_1050-SPD_1053	3.46-4.73	Tagatose pathway, galactose catabolism
SPD_1344-SPD_1349	2.04-2.88	acetyltransferase, GNAT family protein synthesis
SPD_1473-SPD_1480	2.63-5.48	Cell division
SPD_1512-SPD_1517	2.64-9.48	Membrane protein synthesis
SPD_1667-SPD_1671	2.38-7.63	Oligopeptide ABC transporter
SPD_1683-SPD_1698	2.02-5.53	tRNA biosynthesis
SPD_1726-SPD_1730	2.1-4.81	Pneumolysin/hypothetical protein synthesis
SPD_1985-SPD_1989	-2.1-2.14	Mannose transport and metabolism
SPD_1942-SPD_1949	2.36-2.6	Hypothetical protein
SPD_2066-SPD_2069	2.55-5.84	DNA replication

Note: These were selected based on the number of effected genes in a given genomic locus. “-” shows downregulation.

in the production of glucuronic acid among the wild type, Δ rgg1518 and Δ rgg1518c (60 ± 6 , 59 ± 6 , and 60 ± 7 μ g/ 10^9 CFU, $n = 6$, respectively; $p > .05$; Figure 5b). These results show that Rgg1518 is a repressor of capsule synthesis on galactose.

2.7 | Role of inactivated rgg1518 in capsule locus gene regulation

Having established Rgg1518's involvement in the regulation of capsule synthesis, next we tested if the capsule locus is directly regulated by Rgg1518. To do this, we employed a LacZ reporter of the capsule promoter in wild type and Δ rgg1518 backgrounds. The β -galactosidase activities in the reporter strains were measured in strains grown in CDM supplemented with galactose or glucose. The activity in $P_{cps}::lacZ$ - Δ rgg1518 (69 ± 2 MU/ 10^9 CFU, $n = 3$) was more than four times that of the wild type (16 ± 1 MU/ 10^9 CFU, $n = 3$; ($p < .0001$; Figure 5c), supporting the view that Rgg1518 is a repressor of the *cps* locus.

In the next step, we employed electrophoretic mobility shift assay (EMSA) to investigate whether Rgg1518 directly interacts

with the capsule locus. The results showed that the mobility pattern of DNA-protein complexes was different from that of DNA alone, showing the direct interaction of Rgg1518 with the promoter of the *cps* locus (Figure 6). There was no detectable change in mobility when Rgg1518 was incubated with *gyrB*, indicating that the Rgg1518-*cps* interaction is specific.

To identify the Rgg1518 binding box, we hypothesized that the Rgg-box would be shared across multiple genes regulated in galactose. Thus, we selected 14 operons from our microarray analysis, all differentially expressed on galactose. We aligned the 225 to 260 bp upstream of the first gene in the operon and used MEME to identify conserved motifs (Bailey et al., 2015) (Supporting information Figure S2). We identified two putative binding motifs (motif 1: AAAGGTG and motif 2 CCGTAAAA), located near the core promoter elements as shown in Figure 6a. To investigate if either of these motifs was indeed the binding site for Rgg1518, DNA probes lacking either motif 1 or motif 2 were generated by splicing overlap extension, and Rgg1518 binding to these modified DNA probes was determined by EMSA. The absence of motif 2 did not affect Rgg1518 binding. In contrast, elimination of motif 1 abolished Rgg1518 binding. These data demonstrate a direct interaction between Rgg1518 and the *cps* promoter,

FIGURE 5 The quantification of glucuronic acid levels in pneumococcal strains grown in chemically defined medium supplemented either with galactose (a) or glucose (b). Influence of Rgg1518 on the expression of the capsule locus (c) The activity of β -galactosidase, driven by cps putative promoter in different mutant strains, is expressed as nmol p-nitrophenol/min/ml. Vehicle refers to no sugar control. The activity was not measured in Δ rgg1518 strain on glucose. The values are the average of at least three independent experiments each with three replicates vertical bars indicate standard error of means (SEM). **** $p < .0001$

and establish that motif 1 is required for binding of Rgg1518 to the *Pcps* promoter. In addition to the regulatory region of capsule locus, motif 1 has been identified in the regulatory regions of several genes represented in Rgg1518 regulon. The significance of this motif in the Rgg1518 regulon needs to be experimentally verified.

2.8 | Contribution of Rgg1518 to pneumococcal colonization and virulence

To explore the contribution of Rgg1518 to carriage, we employed the murine CD1 pneumococcal colonization model. One hour after infection, the bacterial load in the nasopharyngeal tissue was similar for all strains ($p > .05$; Figure 7a). However, at 3 and 7 days post-infection, the colony counts for Δ rgg1518 ($\text{Log}_{10} 1.45 \pm 0.09$ CFU/mg and 0.95 ± 0.104 CFU/mg, respectively, $n = 5$) was significantly lower than that of wild type ($\text{Log}_{10} 2.69 \pm 0.27$ CFU/mg and $\text{Log}_{10} 2.98 \pm 0.17$ CFU/mg $n = 4$). Moreover, there was no significant difference in the counts of the complemented strain, Δ rgg1518c, and the wild type ($p < .001$) at 3 ($\text{Log}_{10} 2.83 \pm 0.07$ CFU/mg, $n = 5$) and 7 days ($\text{Log}_{10} 2.85 \pm 0.087$ CFU/mg $n = 5$) post-infection ($p > .05$). These results clearly prove the involvement of Rgg1518 in pneumococcal colonization of the murine nasopharynx.

Colonization can extend beyond the nasopharynx causing pneumonia. To investigate the role of Rgg1518 in lung disease, we employed a murine model where pneumococci are inoculated in a high volume, such that they invade lung tissue. We analyzed mice inoculated with either wild type or the Δ rgg1518 strain for survival and bacterial load. The median survival time of mice infected intranasally with Δ rgg1518 ($123.3 \text{ hr} \pm 18.2$, $n = 10$) was significantly longer than the wild type ($47 \text{ hr} \pm 3.1$ $n = 10$) and Δ rgg1518c ($60.4 \text{ hr} \pm 26.9$, $n = 5$) infected groups ($p < .01$; Figure 7b). Further, there was no significant difference in survival time between the wild type and Δ rgg1518c ($p > .05$). Similarly, the bacterial counts for the Δ rgg1518 infected cohort were significantly lower at 24- and 48-hr post-infection ($\text{Log}_{10} 1.62 \pm 0.74$ CFU/ml, $\text{Log}_{10} 2.8 \pm 1.15$ CFU/ml, respectively, $n = 10$) in comparison with the wild type infected cohort at 24 ($\text{Log}_{10} 4.65 \pm 0.92$ CFU/ml, $n = 10$) and 48-hr post-infection ($\text{Log}_{10} 7.55 \pm 0.53$ CFU/ml, $n = 10$; $p < .05$; $p < .05$, $p < .01$; Figure 7c). The progression of bacteremia in a cohort infected with the complemented strain Δ rgg1518c at 24-hr ($\text{Log}_{10} 3.78 \pm 1.06$ CFU/ml, $n = 5$) and 48-hr post-infection ($\text{Log}_{10} 5.44 \pm 1.38$ CFU/ml, $n = 5$) was similar to that of wild type infected cohort ($p > .05$). Together, these findings demonstrate that Rgg1518 promotes colonization and pathogenesis of the pneumococcus.

3 | DISCUSSION

In fluctuating host environments, community-level synchronization of transcription enables *S. pneumoniae* to display a phenotype consistent with the requirements of the specific host niche. In this manner, peptide-mediated communication is critical to colonization and pathogenesis. In this study, we present the transcriptional regulator Rgg1518 and its cognate peptide SHP1518. The Rgg1518 regulator is induced in galactose and mannose, required for galactose and mannose utilization, and a negative regulator of capsule expression. It is critically positioned in a circuit of multiple Rgg communication systems implicated in the response to host carbohydrates. Finally, Rgg1518 promotes colonization and virulence.

FIGURE 6 Direct interaction of Rgg1518 with the capsule (*cps*) locus. Concentrations stated at the top of the gel refer to the recombinant Rgg1518. (a) The putative promoter region of the *cps* locus is shown. The predicted Rgg binding motifs 1 and 2 are shown in blue and red, respectively. Italic sequence is the -35 , and the underlined is the -10 sequence. (b) Electrophoretic mobility shift assay (EMSA) analysis showing the direct interaction of recombinant His-tagged Rgg1518 protein with the promoter of *cps* locus. (c) EMSA indicating the absence of the putative motif 1 abolishes the interaction of recombinant Rgg1518 with the probe. (d) EMSA indicating the absence of motif 2 did not affect the Rgg1518 binding to the probe. In each EMSA image, lane 1 contains only DNA probe as shown with a dotted line. In b, c and d “C+” represents 10 nM of FAM-labeled *gyrB* amplicons incubated with 0.6 μ M recombinant Rgg1518 while “C-” shows *gyrB* amplicons without Rgg1518. In c and d, “Cap” shows intact *csp* locus incubated with 0.6 mM Rgg1518. DNA-protein interaction is seen as an upward shift of DNA probe, represented by a black arrow

This work suggests that Rgg1518 is part of an interconnected set of peptide-mediated regulators. Rgg144/SHP144 and Rgg939/SHP939 play an important role in sugar metabolism and capsule synthesis and have a major impact on colonization and virulence (Junges et al., 2017; Motib et al., 2019; Zhi et al., 2018). In this study, we demonstrate that Rgg1518/SHP1518 is part of a larger network of Rgg regulators. First, Rgg144 and Rgg939 induce *rgg1518* promoter activity. Second, there is overlap in the regulons for Rgg1518, Rgg144, and Rgg939 (Supporting information Figures S3 and 4a), prominently they all include the locus immediately upstream of Rgg1518 (SPD1513-SPD1517), the VP1 operon, and the capsular locus. While all these regulators play an important role in host-derived sugar metabolism, their roles differ in response to specific sugars. Rgg144 and Rgg939 are induced the highest by mannose and then galactose, while Rgg1518 is stimulated mainly by galactose (Junges et al., 2017; Zhi et al., 2018). These data suggest that these

regulator-peptide systems are integrated, their relative roles are carbohydrate dependent, and their combined effects shape the bacterial transcriptional response to host nutrient availability.

Many questions remain regarding the connections and architecture of this multi-Rgg network. The systems have variable overlap not only in the sugars where the activity is most prominent, but also their distribution in the pangenome, and the nature and function of their regulons. While Rgg144 is part of the core genome, Rgg939 and Rgg1518 are part of the accessory genome, with distributions of approximately 54% and 38% of the strains within sequenced pneumococcal genomes, respectively (Zhi et al., 2018). It is tempting to speculate that the distribution pattern of Rggs may be linked to fitness and pathogenic potential in diverse niches such as lungs, blood, middle ear, and brain. The Rgg1518 regulon is larger than the regulon of Rgg144 or Rgg939 and is consistent with a master regulator. In galactose Rgg1518 controls, directly or indirectly, the expression of

FIGURE 7 In vivo test of pneumococcal strains. (a) $\Delta rgg1518$ is impaired in nasopharyngeal colonization. Approximately 5×10^5 CFU of pneumococci was used to inoculate mice. Five mice were culled approximately 1 hr after infection (day 0), day 3 and day 7, CFU/mg of pneumococci were defined by serial dilutions of homogenized nasopharyngeal. Each bar represents the average data collected from five mice. (b) The percent survival of mice infected intranasally with approximately 2×10^6 pneumococci/mouse. There are significant differences in percent survival of mice infected with the wild-type, complemented mutant and mutant strains using Mantel-Cox test. (c) Development of bacteremia after intranasal infection. Each column represents the average data for wild type D39 and mutant strains obtained from ten mice, while complemented strains were from five mice. Error bars illustrate the standard error of the mean. There are significant differences in bacterial counts between wild type and mutant strains using one-way ANOVA and Tukey's multiple comparisons test, (* $p < .05$ and ** $p < .01$, *** $p < .001$, **** $p < .0001$)

genes with diverse functions, notably those coding for carbohydrate metabolism and transport, capsule, fatty acid synthesis, iron transporters, and protein translation.

Regarding sugar metabolism, Rgg1518 appears to decrease overall galactose metabolism. The pneumococcus utilizes galactose via the Leloir and tagatose catabolic pathways, and, in galactose, Rgg1518 represses genes in both of these pathways. Further,

the utilization of galactose leads to the generation of mixed acid products mainly via the activity of pyruvate formate lyase (Bidossi et al., 2012; Paixão et al., 2015; Yesilkaya et al., 2009). Rgg1518 also downregulates pyruvate formate lyase and transporters linked to galactose.

The connection between Rggs and capsule is clear for multiple Rgg systems. Here we show that Rgg1518 downregulates the capsule. In previous work, we demonstrated that the Rgg144 and Rgg939 systems also contributed to capsule synthesis repression in strain D39. The conditions where each Rgg is at play may vary. Rgg144 and Rgg939 repress capsule in mannose, but not on galactose (Zhi et al., 2018). In contrast, Rgg1518 inhibits the capsule in galactose. In mannose, only one gene from the capsule locus is downregulated by Rgg1518 (very likely due to the instability of the transcript on mannose). Thus, the role of Rgg1518 in capsule may be linked to galactose metabolism, and not mannose. In this study, we demonstrated a direct interaction of recombinant Rgg1518 with the capsule locus promoter at the "AAAGGTG" sequence. While we have previously shown the binding of Rgg144 and Rgg939 to the capsule locus promoter, their binding sites have not been identified. Thus, interesting questions remain regarding the overlap among Rgg-binding motifs on the capsule promoter, their role in diverse sugars, and whether there is a synergistic or antagonistic interaction among these Rggs for capsule locus expression.

The pneumococcal capsule has antiphagocytic and anticomplement activities (Janoff et al., 1999). Furthermore, by significantly preventing entrapment of *S. pneumoniae* in airway mucus, it decreases mucus-mediated clearance (Nelson et al., 2007). It has also been shown that capsule is essential for virulence and promotes colonization (Geno et al., 2015; Nelson et al., 2007). Contrary to the expectation, increased capsule synthesis in the mutant did not elevate the virulence, but rather we observed a decrease in the virulence potential of mutant as well as its colonization in the nasopharynx. This apparent inconsistency of our results and the perceived function of capsule in colonization and virulence can be due to a number of different reasons. First, previous studies reporting the importance of capsule in colonization obtained their results using isogenic mutants devoid of capsule or the wild-type strains. On the other hand, in our study, we compared a capsule overexpressing mutant due to the *rgg1518* mutation with the wild type. Therefore, the excess capsule production may hinder colonization. Second, the in vivo conditions may be different than the in vitro assay conditions that indicated overexpression of capsule in the mutant. Finally, decreases in capsule have been linked to increase adhesion, likely via increase exposure of pneumococcal receptors on the bacterial surface (Hammerschmidt et al., 2005), which may, in turn, contribute to colonization and dissemination. We propose that the attenuation of virulence and colonization in the Rgg1518 mutant is likely caused by the impact of Rgg1518 on multiple metabolic processes, where the capsule is one contributor. Specifically, in natural infections downregulation of the capsule may increase cell adhesion and in doing so contribute to increased colonization and virulence.

4 | EXPERIMENTAL PROCEDURES

4.1 | Bacterial strains and growth conditions

The *S. pneumoniae* and *Escherichia coli* strains and plasmids constructed used in this study are listed in Supporting information Table S3. BHI broth medium (Oxoid) or blood agar base (BAB) medium (Oxoid) supplemented with 5% (v/v) defibrinated horse blood were used to grow *S. pneumoniae* at 37°C. When required, different antibiotics such as spectinomycin (100 µg/ml), kanamycin (200 µg/ml) and tetracycline (15 µg/ml) were used for supplementing the growth medium. Pneumococcal strains were grown anaerobically in a CDM supplemented with 55 mM of selected sugars (Motib et al., 2019).

4.2 | Synthetic peptides

Several synthetic peptides were used to determine the inducibility of Shp1518 (Table S4). The synthetic peptides were reconstituted as 6 mM stocks in dimethyl sulfoxide (DMSO) and stored at -80°C.

4.3 | Construction of genetically modified strains

SOEing protocol (splicing by overlap extension) was used for mutagenesis using the primers listed in Supporting information Table S4 (Al-Bayati et al., 2017). Briefly, a spectinomycin resistance cassette flanked by the DNA fragments of *rgg1518* generated through two-step PCR was transformed into *S. pneumoniae* type 2 D39 strain using CSP and the transformants were selected on BAB containing spectinomycin. The mutation was confirmed by PCR and sequencing. Genetic complementation of mutant strains was performed to eliminate the possibility polar effect of the mutation using a previously described protocol using pCEP (Guiral et al., 2006).

4.4 | RNA extraction from bacterial cells, cDNA synthesis, and quantitative RT-PCR

Pneumococcal RNA extraction was done using the Trizol method (Yesilkaya et al., 2009). SuperScript III reverse transcriptase (Invitrogen) was used to perform the first-strand cDNA synthesis according to the manufacturer's instructions using DNase-treated RNA (~1 µg) and random primers (Invitrogen), in 20 µl. qRT-PCR was done using SensiMix™ SYBR® Hi-ROX kit (Bioline, UK). The transcription level of selected genes was normalized to the expression level of the house-keeping *gyrB* gene. The fold difference in expression was calculated by C_T method (Livak & Schmittgen, 2001).

4.5 | Microarray analysis

Pneumococcal strains were grown to early exponential phase, OD₆₀₀ approximately 0.3, in CDM supplemented with either

55 mM galactose or mannose. Three biological replicates were used for analysis. The MicroPrep software package was used to obtain the microarray data from the slides. CyberT implementation of a variant of *t*-test (<http://bioinformatics.biol.rug.nl/cybert/index.shtml>) was performed and false discovery rates (FDRs) were calculated (van Hijum et al., 2003). For differentially expressed genes, $p < .001$ and $FDR < 0.05$ were considered for significance threshold. For the determination of differentially expressed genes, a Bayesian p -value of $< .001$ and a fold-change cut-off of two was applied. All other procedures for the DNA microarray experiments and data analysis were performed as described before (Hajaj et al., 2017). Microarray data have been submitted to GEO (Gene Expression Omnibus) database under the accession number GSE154887.

4.6 | Construction of *lacZ* fusions and β -galactosidase assay

The putative promoter region of *rgg1518* was amplified and ligated into the pPP2 plasmid using the primers listed in Supporting information Table S4 (Halfmann et al., 2007). The recombinant DNA molecule was transformed into *E. coli* TOP10 chemically competent cells for propagation in LA supplemented with ampicillin (100 µg/ml). After sequencing, the recombinant pPP2 was transformed into *S. pneumoniae* wild type D39 or selected mutants.

To evaluate promoter induction, strains were grown in CDM-glucose and then transferred to CDM with no sugar, or a single sugar for 1 hr. The β -galactosidase activity was determined as previously described and the activity level was expressed for 10⁹ CFU/ml.

4.7 | Cloning and expression of recombinant proteins

To obtain recombinant Rgg1518, *rgg1518* was cloned into pLEICS-01 (Protex, University of Leicester). The construct was transformed into *E. coli* BL21 (DE3) pLysS for protein expression at 37°C. The expression was induced with 1 mM IPTG. The 10 preppacked gravity flow column His GraviTrap™ (GE Healthcare, UK) was used to purify protein. To confirm protein identity, the purified recombinant proteins were analyzed by matrix-assisted laser desorption ionization-time of flight (MALDI-TOF) mass spectrometry (PNAAC, University of Leicester).

4.8 | Electrophoretic mobility shift assay (EMSA)

The presence of regulatory elements in the putative promoter regions of target genes was determined using the bacterial promoter prediction tool (BPROM) (Solovyev & Salamov, 2011) and the motifs were localized using (MEME) (Bailey & Elkan, 1994). For this analysis, we used the upstream regions of differentially regulated selected genes in Δ rgg1518 predicted to be part of an operon including SPD_0315, SPD_1053, SPD_1634, SPD_1342, SPD_1461,

SPD_1127, SPD_0113, SPD_0122, SPD_0193, SPD_0197, SPD_0208, SPD_0126, SPD_0341, SPD_0378, SPD_0404, and SPD_1512.

The genetic fragment containing the putative promoter sites of target genes were amplified using primers containing 5'FAM fluorescent tags (Supporting information Table S4), and the amplicons were purified using a commercial kit. Varying concentrations of recombinant Rgg1518 was incubated in 20 µl containing binding buffer (20 mM Tris-HCl pH 7.5, 30 mM KCl, 1 mM DTT, 1 mM EDTA pH 8.0, and 10% v/v glycerol), the DNA probe (50 nM), and peptide to a final concentration of 2 µM. The reaction mixture was incubated at room temperature for 20 min. An aliquot (8 µl) of reaction mixture was loaded on 5% (v/v) polyacrylamide gel buffered with 20 mM potassium phosphate, pH 7.5, for 60 min at 200 V. All gel shifts were detected by fluorescence imaging using a Typhoon PhosphorImager.

4.9 | Glucuronic acid assay

Capsular polysaccharide (CPS) production was quantified by measuring the amount of glucuronic acid, which is found in type 2 pneumococcal capsule (Najmuldeen et al., 2019; Zhi et al., 2018). Briefly, the pneumococcal cultures (500 µl) grown to late exponential phase in CDM supplemented with different sugars were mixed with 100 µl of 1% (v/v) Zwittergent 3-14 detergent (Sigma-Aldrich) in 100 mM citric acid (pH 2.0) and the mixture was kept at 50°C for 20 min. The polysaccharides were precipitated with 1 ml of absolute ethanol. The pellet was dissolved in 200 µl distilled water, and added with 1,200 µl 12.5 mM borax (Sigma) in H₂SO₄. The mixture was mixed thoroughly, boiled for 5 min. Then 20 µl 0.15% 3-hydroxydiphenol (Sigma) was added. The absorbance of the mixture at 520 nm was recorded, and the glucuronic acid content of the samples was quantified in comparison to a standard curve generated with the known concentrations of glucuronic acid (Sigma).

4.10 | In vivo studies

This study used 8–10 weeks old female CD1 mice (Charles River, UK). Standard inoculum was prepared as described previously (Kahya et al., 2017). To administer the dose, mice were lightly anesthetized with 2.5% (v/v) Isoflurane (Isocare, UK) over oxygen (1.4 to 1.6 L/min), in an anesthetic box. To initiate infection, approximately 2×10^6 CFU of *S. pneumoniae* were administered equally dropwise into the nostrils. Mice were watched periodically for disease signs (hunched, piloerect, or lethargy) for 7 days. As soon they reached the lethargic stage, mice were sacrificed humanely. The time to reach lethargic stage was considered as survival time (Al-Bayati et al., 2017). The median survival times of mice were examined statistically by using the Mann-Whitney *U* test. To follow the progression of bacteremia, 20 µl of venous blood from the tail vein was taken 24- and 48-hr after infection. The colonies were enumerated by plating the serial dilutions of the blood on blood agar plates. To evaluate Rgg1518's contribution to colonization, each mouse was

given approximately 5×10^5 CFU of *S. pneumoniae* in 20 µl PBS. The bacterial counts in the nasopharynx were determined by plating the serial dilutions of tissue homogenates (Kahya et al., 2017).

ACKNOWLEDGMENTS

We are grateful for the support from the NIH (R01 AI139077-01A1 to NLH and R01 AI135060-01A1 to AU), also to the Eberly Family Trust to NLH. We also gratefully acknowledge the valuable support of the Preclinical Facility staff for the *in vivo* experiments in Leicester.

CONFLICT OF INTEREST

The authors declare no competing financial interests.

AUTHOR CONTRIBUTIONS

Conceived and designed the experiments: H. Yesilkaya, P.W. Andrew, N.L. Andrew, A. Ulijasz, O.P. Kuipers, and S. Shafeeq. H. Yesilkaya, B. Shlla, SS, and I. Manzoor analyzed the data. All authors contributed to the writing of the manuscript.

ETHICS APPROVAL

In vivo studies were carried out under the UK Home Office approved project (60/4327) and personal licenses (I4142DBBE). The procedures were approved by the local ethic committee. The animals were placed in individually ventilated cages in a controlled environment. The mice were repeatedly monitored after infection in order to minimize suffering.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

ORCID

Oscar P. Kuipers <https://orcid.org/0000-0001-5596-7735>

Hasan Yesilkaya <https://orcid.org/0000-0003-3617-1437>

REFERENCES

- Aggarwal, C., Jimenez, J.C., Nanavati, D. & Federle, M.J. (2014) Multiple length peptide-pheromone variants produced by *Streptococcus pyogenes* directly bind Rgg proteins to confer transcriptional regulation. *Journal of Biological Chemistry*, 289, 22427–22436. <https://doi.org/10.1074/jbc.M114.583989>
- Aggarwal, S.D., Yesilkaya, H., Dawid, S. & Hiller, N.L. (2020) The pneumococcal social network. *PLOS Pathogens*, 16, e1008931.
- Al-Bayati, F.A.Y., Kahya, H.F.H., Damianou, A., Shafeeq, S., Kuipers, O.P., Andrew, P.W. et al. (2017) Pneumococcal galactose catabolism is controlled by multiple regulators acting on pyruvate formate lyase. *Scientific Reports*, 7, 43587. <https://doi.org/10.1038/srep43587>
- Bailey, T.L. & Elkan, C. (1994) Fitting a mixture model by expectation maximization to discover motifs in biopolymers. *Proceedings of the International Conference on Intelligent Systems for Molecular Biology*, 2, 28–36.
- Bailey, T.L., Johnson, J., Grant, C.E. & Noble, W.S. (2015) The MEME suite. *Nucleic Acids Research*, 43, W39–W49. <https://doi.org/10.1093/nar/gkv416>
- Bidossi, A., Mulas, L., Decorosi, F., Colomba, L., Ricci, S., Pozzi, G. et al. (2012) A functional genomics approach to establish the complement

- of carbohydrate transporters in *Streptococcus pneumoniae*. *PLoS One*, 7, e33320. <https://doi.org/10.1371/journal.pone.0033320>
- Bortoni, M.E., Terra, V.S., Hinds, J., Andrew, P.W. & Yesilkaya, H. (2009) The pneumococcal response to oxidative stress includes a role for Rgg. *Microbiology*, 155, 4123–4134. <https://doi.org/10.1099/mic.0.028282-0>
- Carvalho, S.M., Kloosterman, T.G., Kuipers, O.P. & Neves, A.R. (2011) CcpA ensures optimal metabolic fitness of *Streptococcus pneumoniae*. *PLoS One*, 6, e26707. <https://doi.org/10.1371/journal.pone.0026707>
- Chang, J.C. & Federle, M.J. (2016) PptAB exports Rgg quorum-sensing peptides in *Streptococcus*. *PLoS One*, 11, e0168461. <https://doi.org/10.1371/journal.pone.0168461>
- Chang, J.C., LaSarre, B., Jimenez, J.C., Aggarwal, C. & Federle, M.J. (2011) Two group A streptococcal peptide pheromones act through opposing Rgg regulators to control biofilm development. *PLoS Pathogens*, 7, e1002190.
- Cook, L.C., LaSarre, B. & Federle, M.J. (2013) Interspecies communication among commensal and pathogenic streptococci. *mBio*, 4, e00382-13. <https://doi.org/10.1128/mBio.00382-13>
- Cuevas, R.A., Eutsey, R., Kadam, A., West-Roberts, J.A., Woolford, C.A., Mitchell, A.P. et al. (2017) A novel streptococcal cell-cell communication peptide promotes pneumococcal virulence and biofilm formation. *Molecular Microbiology*, 105, 554–571. <https://doi.org/10.1111/mmi.13721>
- De Saizieu, A., Gardès, C., Flint, N., Wagner, C., Kamber, M., Mitchell, T.J. et al. (2000) Microarray-based identification of a novel *Streptococcus pneumoniae* regulon controlled by an autoinduced peptide. *Journal of Bacteriology*, 182, 4696–4703.
- Fleuchot, B., Gitton, C., Guillot, A., Vidic, J., Nicolas, P., Besset, C. et al. (2011) Rgg proteins associated with internalized small hydrophobic peptides: a new quorum-sensing mechanism in streptococci. *Molecular Microbiology*, 80, 1102–1119. <https://doi.org/10.1111/j.1365-2958.2011.07633.x>
- Geno, K.A., Gilbert, G.L., Song, J.Y., Skovsted, I.C., Klugman, K.P., Jones, C. et al. (2015) Pneumococcal capsules and their types: past, present, and future. *Clinical Microbiology Reviews*, 34, e00320-20. <https://doi.org/10.1128/CMR.00024-15>
- Gómez-Mejía, A., Gámez, G. & Hammerschmidt, S. (2018) *Streptococcus pneumoniae* two-component regulatory systems: the interplay of the pneumococcus with its environment. *International Journal of Medical Microbiology*, 308, 722–737. <https://doi.org/10.1016/j.ijmm.2017.11.012>
- Guiral, S., Hénard, V., Laaberki, M.-H., Granadel, C., Prudhomme, M., Martin, B. et al. (2006) Construction and evaluation of a chromosomal expression platform (CEP) for ectopic, maltose-driven gene expression in *Streptococcus pneumoniae*. *Microbiology*, 152, 343–349. <https://doi.org/10.1099/mic.0.28433-0>
- Hajaj, B., Yesilkaya, H., Shafeeq, S., Zhi, X., Benisty, R., Tchalah, S. et al. (2017) CodY regulates thiol peroxidase expression as part of the pneumococcal defense mechanism against H₂O₂ stress. *Frontiers in Cellular and Infection Microbiology*, 7, 210. <https://doi.org/10.3389/fcimb.2017.00210>
- Halfmann, A., Hakenbeck, R. & Brückner, R. (2007) A new integrative reporter plasmid for *Streptococcus pneumoniae*. *FEMS Microbiology Letters*, 268, 217–224. <https://doi.org/10.1111/j.1574-6968.2006.00584.x>
- Hammerschmidt, S., Wolff, S., Hocke, A., Rosseau, S., Müller, E. & Rohde, M. (2005) Illustration of pneumococcal polysaccharide capsule during adherence and invasion of epithelial cells. *Infection and Immunity*, 73, 4653–4667. <https://doi.org/10.1128/IAI.73.8.4653-4667.2005>
- Håvarstein, L.S., Coomaraswamy, G. & Morrison, D.A. (1995) An unmodified heptadecapeptide pheromone induces competence for genetic transformation in *Streptococcus pneumoniae*. *Proceedings of the National Academy of Sciences USA*, 92, 11140–11144. <https://doi.org/10.1073/pnas.92.24.11140>
- Hoover, S.E., Perez, A.J., Tsui, H.C., Sinha, D., Smiley, D.L., DiMarchi, R.D. et al. (2015) A new quorum-sensing system (TprA/PhrA) for *Streptococcus pneumoniae* D39 that regulates a lantibiotic biosynthesis gene cluster. *Molecular Microbiology*, 97, 229–243. <https://doi.org/10.1111/mmi.13029>
- Hyams, C., Camberlein, E., Cohen, J.M., Bax, K. & Brown, J.S. (2010) The *Streptococcus pneumoniae* capsule inhibits complement activity and neutrophil phagocytosis by multiple mechanisms. *Infection and Immunity*, 78, 704–715. <https://doi.org/10.1128/IAI.00881-09>
- Ibrahim, M., Guillot, A., Wessner, F., Algaron, F., Besset, C., Courtin, P. et al. (2007) Control of the transcription of a short gene encoding a cyclic peptide in *Streptococcus thermophilus*: a new quorum-sensing system? *Journal of Bacteriology*, 189, 8845–8854.
- Janoff, E.N., Fasching, C., Orenstein, J.M., Rubins, J.B., Opstad, N.L. & Dalmasso, A.P. (1999) Killing of *Streptococcus pneumoniae* by capsular polysaccharide-specific polymeric IgA, complement, and phagocytes. *Journal of Clinical Investigation*, 104, 1139–1147. <https://doi.org/10.1172/JCI6310>
- Junges, R., Salvadori, G., Shekhar, S., Åmdal, H.A., Periselmanis, J.N., Chen, T. et al. (2017) A quorum-sensing system that regulates *Streptococcus pneumoniae* biofilm formation and surface polysaccharide production. *mSphere*, 13, e00324-17. <https://doi.org/10.1128/mSphere.00324-17>
- Kadam, A., Eutsey, R.A., Rosch, J., Miao, X., Longwell, M., Xu, W. et al. (2017) Promiscuous signaling by a regulatory system unique to the pandemic PMEN1 pneumococcal lineage. *PLoS Pathogens*, 13, e1006339. <https://doi.org/10.1371/journal.ppat.1006339>
- Kahya, H.F., Andrew, P.W. & Yesilkaya, H. (2017) Deacetylation of sialic acid by esterases potentiates pneumococcal neuraminidase activity for mucin utilization, colonization and virulence. *PLoS Pathogens*, 13, e1006263. <https://doi.org/10.1371/journal.ppat.1006263>
- Kenne, L., Lindberg, B. & Svensson, S. (1975) The structure of capsular polysaccharide of the pneumococcus type II. *Carbohydrate Research*, 40, 69–75. [https://doi.org/10.1016/S0008-6215\(00\)82669-5](https://doi.org/10.1016/S0008-6215(00)82669-5)
- King, S.J. (2010) Pneumococcal modification of host sugars: a major contributor to colonization of the human airway? *Molecular Oral Microbiology*, 25, 15–24. <https://doi.org/10.1111/j.2041-1014.2009.00564.x>
- Koponen, O., Tolonen, M., Qiao, M., Wahlström, G., Helin, J. & Saris, P.E.J. (2002) NisB is required for the dehydration and NisC for the lantionine formation in the post-translational modification of nisin. *Microbiology*, 148, 3561–3568. <https://doi.org/10.1099/00221287-148-11-3561>
- Kuipers, O.P., Beerthuyzen, M.M., Ruyter, P.G.G.A.D., Luesink, E.J. & Vos, W.M.D. (1995) Autoregulation of nisin biosynthesis in *Lactococcus lactis* by signal transduction. *Journal of Biological Chemistry*, 270, 27299–27304. <https://doi.org/10.1074/jbc.270.45.27299>
- Livak, K.J. & Schmittgen, T.D. (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2^{(-Delta Delta C(T))} method. *Methods*, 25, 402–408.
- Motib, A.S., Al-Bayati, F.A.Y., Manzoor, I., Shafeeq, S., Kadam, A., Kuipers, O.P. et al. (2019) TprA/PhrA quorum sensing system has a major effect on pneumococcal survival in respiratory tract and blood, and its activity is controlled by CcpA and GlnR. *Frontiers in Cellular and Infection Microbiology*, 9, 326. <https://doi.org/10.3389/fcimb.2019.00326>
- Najmuldeen, H., Alghamdi, R., Alghofaili, F. & Yesilkaya, H. (2019) Functional assessment of microbial superoxide dismutase isozymes suggests a differential role for each isozyme. *Free Radical Biology and Medicine*, 134, 215–228. <https://doi.org/10.1016/j.freeradbiomed.2019.01.018>
- Nelson, A.L., Roche, A.M., Gould, J.M., Chim, K., Ratner, A.J. & Weiser, J.N. (2007) Capsule enhances pneumococcal colonization by limiting mucus-mediated clearance. *Infection and Immunity*, 75, 83–90. <https://doi.org/10.1128/IAI.01475-06>

- Paixão, L., Oliveira, J., Veríssimo, A., Vinga, S., Lourenço, E.C., Ventura, M.R. et al. (2015) Host glycan sugar-specific pathways in *Streptococcus pneumoniae*: Galactose as a key sugar in colonisation and infection. *PLoS One*, 10, e0127483. <https://doi.org/10.1371/journal.pone.0121042>
- Rutherford, S.T. & Bassler, B.L. (2012) Bacterial quorum sensing: its role in virulence and possibilities for its control. *Cold Spring Harbor Perspectives in Medicine*, 2, a012427. <https://doi.org/10.1101/cshperspect.a012427>
- Sahin-Yilmaz, A. & Naclerio, R.M. (2011) Anatomy and physiology of the upper airway. *Proceedings of the American Thoracic Society*, 8, 31–39. <https://doi.org/10.1513/pats.201007-050RN>
- Sen, A.K., Narbad, A., Horn, N., Dodd, H.M., Parr, A.J., Colquhoun, I. et al. (1999) Post-translational modification of nisin. *European Journal of Biochemistry*, 261, 524–532.
- Shafeeq, S., Yesilkaya, H., Kloosterman, T.G., Narayanan, G., Wandel, M., Andrew, P.W. et al. (2011) The cop operon is required for copper homeostasis and contributes to virulence in *Streptococcus pneumoniae*. *Molecular Microbiology*, 81, 1255–1270. <https://doi.org/10.1111/j.1365-2958.2011.07758.x>
- Solovyev, V. & Salamov, A. (2011) Automatic annotation of microbial genomes and metagenomic sequences. In: Li, R. (Ed.) *Metagenomics and its applications in agriculture, biomedicine and environmental studies*. Nova Science Publishers, pp. 61–78.
- Trappetti, C., McAllister, L.J., Chen, A., Wang, H., Paton, A.W., Oggioni, M.R. et al. (2017) Autoinducer 2 signaling via the phosphotransferase FruA drives galactose utilization by *Streptococcus pneumoniae*, resulting in hypervirulence. *Mbio*, 8, e02269-16.
- Troeger, C., Forouzanfar, M., Rao, P.C., Khalil, I., Brown, A., Swartz, S. et al. (2017) Estimates of the global, regional, and national morbidity, mortality, and aetiologies of lower respiratory tract infections in 195 countries: a systematic analysis for the Global Burden of Disease Study 2015. *The Lancet Infectious Diseases*, 17, 1133–1161. [https://doi.org/10.1016/S1473-3099\(17\)30396-1](https://doi.org/10.1016/S1473-3099(17)30396-1)
- van Hijum, S.A.F.T. van García de la Nava, J., Trelles, O., Kok, J. & Kuipers, O.P. (2003) MicroPreP: a cDNA microarray data pre-processing framework. *Applied Bioinformatics*, 2, 241–244.
- Varahan, S., Harms, N., Gilmore, M.S., Tomich, J.M. & Hancock, L.E. (2014) An ABC transporter is required for secretion of peptide sex pheromones in *Enterococcus faecalis*. *Mbio*, 5, e01726-14. <https://doi.org/10.1128/mBio.01726-14>
- Wang, C.Y., Medlin, J.S., Nguyen, D.R., Disbennett, W.M. & Dawid, S. (2020) Molecular determinants of substrate selectivity of a pneumococcal Rgg-regulated peptidase-containing ABC transporter. *Mbio*, 11, e02502-19. <https://doi.org/10.1128/mBio.02502-19>
- Weiser, J.N., Ferreira, D.M. & Paton, J.C. (2018) *Streptococcus pneumoniae*: transmission, colonization and invasion. *Nature Reviews Microbiology*, 16, 355–367. <https://doi.org/10.1038/s41579-018-0001-8>
- Wiley, J.M. & van der Donk, W.A. (2007) Lantibiotics: peptides of diverse structure and function. *Annual Review of Microbiology*, 61, 477–501. <https://doi.org/10.1146/annurev.micro.61.080706.093501>
- Wilton, J., Acebo, P., Herranz, C., Gómez, A. & Amblar, M. (2015) Small regulatory RNAs in *Streptococcus pneumoniae*: discovery and biological functions. *Frontiers in Genetics*, 6, 126. <https://doi.org/10.3389/fgene.2015.00126>
- Yesilkaya, H., Spissu, F., Carvalho, S.M., Terra, V.S., Homer, K.A., Benisty, R. et al. (2009) Pyruvate formate lyase is required for pneumococcal fermentative metabolism and virulence. *Infection and Immunity*, 77, 5418–5427. <https://doi.org/10.1128/IAI.00178-09>
- Zhi, X., Abdullah, I.T., Gazioglu, O., Manzoor, I., Shafeeq, S., Kuipers, O.P. et al. (2018) Rgg-Shp regulators are important for pneumococcal colonization and invasion through their effect on mannose utilization and capsule synthesis. *Scientific Reports*, 8, 6369. <https://doi.org/10.1038/s41598-018-24910-1>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Shlla, B., Gazioglu, O., Shafeeq, S., Manzoor, I., Kuipers, O.P., Ulijasz, A., et al. (2021) The Rgg1518 transcriptional regulator is a necessary facet of sugar metabolism and virulence in *Streptococcus pneumoniae*. *Molecular Microbiology*, 116, 996–1008. <https://doi.org/10.1111/mmi.14788>