University of Groningen

Dosing of losartan in men versus women with heart failure with reduced ejection fraction

Ferreira, Joao Pedro; Konstam, Marvin A.; McMurray, John J. V.; Butler, Javed; Girerd, Nicolas; Rossignol, Patrick; Sharma, Abhinav; Voors, Adriaan A.; Lam, Carolyn S. P.; Packer, Milton

Published in:
European Journal of Heart Failure

DOI:
[10.1002/ejhf.2255](https://doi.org/10.1002/ejhf.2255)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Ferreira, J. P., Konstam, M. A., McMurray, J. J. V., Butler, J., Girerd, N., Rossignol, P., Sharma, A., Voors, A. A., Lam, C. S. P., Packer, M., & Zannad, F. (2021). Dosing of losartan in men versus women with heart failure with reduced ejection fraction: the HEAAL trial. *European Journal of Heart Failure*, 23(9), 1477-1484. <https://doi.org/10.1002/ejhf.2255>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Dosing of losartan in men versus women with heart failure with reduced ejection fraction: the HEAAL trial

João Pedro Ferreira^{1*}, Marvin A. Konstam², John J.V. McMurray³, Javed Butler⁴, Nicolas Girerd¹, Patrick Rossignol¹, Abhinav Sharma⁵, Adriaan A. Voors⁶, Carolyn S.P. Lam⁷, Milton Packer^{8,9}, and Faiez Zannad¹

¹Université de Lorraine, INSERM, Centre d'Investigations Cliniques Plurithématique 1433, UMR 1116, CHRU de Nancy, F-CRIN INI-CRCT Cardiovascular and Renal Clinical Trialists, Nancy, France; ²The CardioVascular Center of Tufts Medical Center, Boston, MA, USA; ³Institute of Cardiovascular and Medical Sciences, BHF Glasgow Cardiovascular Research Centre, University of Glasgow, Glasgow, UK; ⁴Department of Medicine, University of Mississippi Medical Center, Jackson, MS, USA; ⁵Division of Cardiology, McGill University Health Centre, McGill University, Montreal, Canada; ⁶University of Groningen, University Medical Center Groningen, Groningen, The Netherlands; ⁷National Heart Center Singapore, and Duke National University of Singapore, Singapore; ⁸Baylor University Medical Center, Dallas, TX, USA; and ⁹Imperial College, London, UK

Received 4 March 2021; revised 25 May 2021; accepted 25 May 2021; online publish-ahead-of-print 17 June 2021

Aims

In heart failure with reduced ejection fraction (HFrEF), guidelines recommend up-titration of angiotensin-converting enzyme inhibitors (ACEi) and angiotensin receptors blockers (ARBs) to the maximum tolerated dose. However, some studies suggest that women might need lower doses of ACEi/ARBs than men to achieve similar treatment benefit.

Methods and results

The HEAAL trial compared low vs. high dose of losartan. We reassessed the efficacy and safety of high- vs. low-dose in men vs. women using Cox models and machine learning algorithms. The mean age was 66 years and 30% of patients were women. Men appeared to have benefited more from high-dose than from low-dose losartan, whereas women appeared to have responded similarly to low and high doses [hazard ratio (95% confidence interval) comparing high- vs. low-dose losartan for the composite outcome of all-cause death or all-cause hospitalization: 0.89 (0.81–0.98) in men and 1.10 (0.95–1.28) in women; interaction $P = 0.018$]. Female sex clustered along with older age, ischaemic heart failure, New York Heart Association class III/IV, and estimated glomerular filtration rate <60 mL/min. Patients with these features had a poorer response to high-dose losartan. Subgroup analyses supported no benefit from high-dose losartan in patients with poorer kidney function and severe heart failure symptoms.

Conclusions

Compared with men, women might need lower doses of losartan to achieve similar treatment benefit. However, beyond sex, other factors (e.g. kidney function, age, and symptoms) may influence the response to high-dose losartan, suggesting that sex-based subgroup findings may be biased by other confounders.

*Corresponding author. Centre d'Investigation Clinique 1433 module Plurithématique, CHRU Nancy – Hôpitaux de Brabois, Institut Lorrain du Cœur et des Vaisseaux Louis Mathieu, 4 rue du Morvan, 54500 Vandœuvre les Nancy, France. Tel: +33 3 83157315, Fax: +33 3 83157324, Email: j.ferreira@chru-nancy.fr

Graphical Abstract

Treatment effect of high- vs. low-dose losartan in the HEAAL trial. In HEAAL, men appeared to have benefited more from high-dose (compared with low-dose) losartan, whereas women appeared to have responded similarly to low or high doses. ACH, all-cause hospitalization; ACM, all-cause mortality; CI confidence interval; HR, hazard ratio.

Keywords

Sex differences • Heart failure • Treatment dose • Dose–response • Machine learning

Introduction

Heart failure guidelines recommend up-titration of angiotensin-converting enzyme inhibitors (ACEi) and angiotensin receptors blockers (ARBs), to the maximum tolerated dose, in men and women with heart failure with reduced ejection fraction (HFrEF).^{1,2} The recommendations rely on trial validated effective doses; however, clinicians may face many challenges in achieving such doses.^{3,4} However, most HFrEF trials used escalating doses, up to a maximum trial defined single dose or maximum tolerated dose. Only the HEAAL (Effects of High-Dose vs. Low-Dose Losartan on Clinical Outcomes in Patients with Heart Failure) and ATLAS (Comparative Effects of Low and High Doses of the Angiotensin-Converting Enzyme Inhibitor, Lisinopril, on Morbidity and Mortality in Chronic Heart Failure) trials compared high vs. low treatment doses in well-powered outcome trials, concluding that high-dose losartan and lisinopril are superior to the corresponding low doses.^{5,6}

Compared with men, women generally have a smaller body size with higher percentage of body fat. These factors may affect drug metabolism including the duration of action and peak plasma concentrations, which may influence efficacy and safety of certain drugs.⁷ If drug metabolism can be influenced by sex, it can also be affected by age, renal function, body mass index, concomitant treatments, and disease severity.^{8–12} Compared to men, women with HFrEF are more symptomatic, usually older, more often obese, have poorer kidney function and more comorbid conditions, and are less frequently treated with guideline-directed therapy.¹³ All

these factors may influence the response to treatments (including ACEi and ARBs), independently of biological sex.

A study using observational data from two independent cohorts including men and women with HFrEF, suggested that women might need lower doses of ACEi or ARBs than men to obtain similar treatment benefit.¹⁴ Using data from the HEAAL trial, we aim to test whether the response to high- vs. low-dose losartan differed between men and women, and assess whether other factors such as age, kidney function and clinical presentation (which may differ between men and women) could have influenced the response to high- vs. low-dose losartan. Additionally, as exploratory analysis, we have also analysed published data from ATLAS to interpret the effect of high- vs. low-dose lisinopril in men and women with HFrEF.

Methods

Study design and population

HEAAL was an international, multicentre, double-blind, event-driven trial, comparing the effect of two doses of losartan 150 vs. 50 mg/day among 3846 patients with symptomatic heart failure [New York Heart Association (NYHA) class II–IV], a left ventricular ejection fraction (LVEF) of $\leq 40\%$, stable cardiovascular medical therapy for at least 2 weeks, and known intolerance to ACEi.⁵ The primary endpoint was a composite of all-cause death or admission for heart failure. In the present study, we have also analysed the composite of cardiovascular death or admission for heart failure, the composite of all-cause death or all-cause hospitalization, cardiovascular death, and all-cause death,

although the trial was underpowered for the study of fatal outcomes in isolation. The median follow-up time was 4.7 (3.5–5.5) years. The study conformed to the principles outlined in the Declaration of Helsinki and was approved by the ethics committee or institutional review board of every site. All patients provided their written informed consent before randomization. HEAAL is registered with ClinicalTrials.gov, number NCT00090259.

We did not have access to the ATLAS trial database, and, as exploratory analysis, we have used the published results for subgroup analysis (forest plots) depicting the treatment effect of high- vs. low-dose lisinopril in men and women for the outcomes of all-cause death and the composite outcome of all-cause death or all-cause hospitalization.⁶ ATLAS included 3164 patients with NYHA class II–IV and LVEF $\leq 30\%$. Patients were randomly assigned to receive either low doses (2.5–5.0 mg/day) or high doses (32.5–35 mg/day) of lisinopril for 39–58 months. ATLAS was underpowered to study the effect of high- vs. low-dose lisinopril on all-cause death because the event rate was lower than expected for this outcome. For this reason and before the blind was broken, the Steering Committee recognized that a composite outcome of time-to-first hospitalization for any reason or all-cause death would provide power to adequately test the study hypothesis.⁶ The trial was conducted in accordance with the Declaration of Helsinki. The protocol was approved by the institutional review boards of all participating hospitals. Written informed consent was obtained from all patients.

Statistical analysis

Descriptive statistics and comparison of the characteristics between women and men were performed using categorical and continuous variables with the appropriate tests. Survival analyses were performed by intention-to-treat. We used time-to-event methods (Kaplan–Meier curves and Cox proportional hazards models) to compare the high- vs. low-dose losartan groups for the studied outcomes. The heterogeneity of hazard ratio (HR) estimates for subgroups was tested by including an interaction term between treatment and subgroup in the Cox model (e.g. treatment-by-sex interaction). Latent class analysis (LCA) is an unsupervised ‘machine learning’ method that allows the grouping of individuals into ‘classes’ with similar characteristics within each ‘class’ and different characteristics between ‘classes’. LCA was performed using maximum-likelihood estimation to identify the most common patterns using nine variables for a range of two to five subgroups. The variables were selected based on statistically and clinically significant sex differences found in Table 1, and included sex (women vs. men), age (≤ 70 vs. > 70 years), body mass index (BMI ≤ 30 vs. > 30 kg/m²), ischaemic aetiology for heart failure (yes vs. no), hypertension (yes vs. no), NYHA class (II vs. III or IV), LVEF ($< 35\%$ vs. $\geq 35\%$), atrial fibrillation (yes vs. no), and estimated glomerular filtration rate (eGFR < 60 vs. ≥ 60 mL/min/1.73 m²). The optimal number of classes was determined using the first minima of the Bayesian Information Criterion (BIC) and Akaike Information Criterion (AIC) with a condition to the percentage of patients in each class to be at least 20% of the total. Based on these criteria, the optimal number of classes was three. Probabilities of membership in each subgroup for every LCA variable were used to determine the most likely subgroup for each patient.

As exploratory analysis, a fixed-effect meta-analysis with inverse variance weights was conducted using the estimated HR [95% confidence interval (CI)] from the HEAAL trial and the sex subgroup estimates (HR and 95% CI) published in the ATLAS trial report.⁶

Table 1 Characteristics of women and men enrolled in the HEAAL trial

Characteristics	Women	Men	P-value
<i>n</i>	1143	2691	
Age, years	66.4 $\pm$ 11.6	63.3 $\pm$ 11.6	<0.001
Age > 70 years	455 (39.8%)	814 (30.2%)	<0.001
White race	678 (59.3%)	1643 (61.1%)	0.31
Atrial fibrillation	291 (25.5%)	779 (28.9%)	0.028
Ischaemic heart disease	647 (56.6%)	1809 (67.2%)	<0.001
Hypertension	763 (66.8%)	1529 (56.8%)	<0.001
Diabetes mellitus	367 (32.1%)	832 (30.9%)	0.47
NYHA class III/IV	415 (36.3%)	759 (28.2%)	<0.001
LVEF, %	32.0 $\pm$ 6.6	31.4 $\pm$ 6.5	0.006
LVEF <35%	734 (64.2%)	1867 (69.4%)	0.002
SBP, mmHg	129.1 $\pm$ 18.7	124.8 $\pm$ 17.5	<0.001
SBP <110 mmHg	241 (21.1%)	703 (26.1%)	<0.001
Heart rate, bpm	75.7 $\pm$ 16.2	73.9 $\pm$ 15.8	0.002
Heart rate >75 bpm	503 (44.0%)	1021 (37.9%)	<0.001
BMI, kg/m ²	29.0 $\pm$ 10.7	28.8 $\pm$ 9.5	0.52
<25	407 (35.6%)	827 (30.7%)	<0.001
25–30	384 (33.6%)	1156 (43.0%)	
>30	352 (30.8%)	708 (26.3%)	
eGFR, mL/min/1.73 m ²	65.4 $\pm$ 22.9	70.8 $\pm$ 21.8	<0.001
<60	521 (45.6%)	867 (32.2%)	<0.001
Potassium, mmol/L	4.5 $\pm$ 0.5	4.5 $\pm$ 0.5	0.39
<4	184 (16.1%)	435 (16.2%)	0.78
4–5	846 (74.0%)	1970 (73.2%)	
>5	113 (9.9%)	286 (10.6%)	
Haemoglobin, g/dL	13.1 $\pm$ 1.4	14.2 $\pm$ 1.5	<0.001
Anaemia	220 (19.2%)	480 (17.8%)	0.30
Beta-blocker stratum	755 (66.1%)	2003 (74.4%)	<0.001
High-dose losartan	583 (51.0%)	1338 (49.7%)	0.47

BMI, body mass index; eGFR, estimated glomerular filtration rate; LVEF, left ventricular ejection fraction; NYHA, New York Heart Association; SBP, systolic blood pressure.

In ATLAS, the treatment effect in men and women was obtained from the published forest plots with an approximation of the estimated HR and corresponding 95% CI for the reported outcomes of all-cause death and all-cause death or all-cause hospitalization. Statistical heterogeneity of the treatment effect between the HEAAL and ATLAS studies was assessed based on the *P*-value derived from Cochran's Q test. The latter was also used to test for treatment-by-sex interactions.¹⁵

Statistical analyses were performed using Stata®, version 16 (Stata Corp, College Station, TX, USA).

Results

Characteristics of the patients by sex

In HEAAL, 30% (*n* = 1143) of the patients were women. Compared with men, women had several different clinical characteristics. Notably, women were older, had more frequent history of hypertension and less frequent history of ischaemic heart disease

Table 2 Events, event rates, and treatment effect in women and men enrolled in the HEAAL trial

Outcomes by sex	Events		Event rate		HR (95% CI)	P-value	Interaction P-value
	Low dose	High dose	Low dose ^a	High dose ^a			
CV death or HF hospitalization							
Overall (n = 3834)	771/1913 (40.3%)	698/1921 (36.3%)	10.7 (10–11.5)	9.3 (8.7–10)	0.88 (0.79–0.97)	0.011	
Women (n = 1143)	188/560 (33.6%)	192/583 (32.9%)	8.3 (7.2–9.6)	8.4 (7.3–9.6)	1.01 (0.82–1.23)	–	0.098
Men (n = 2691)	583/1353 (43.1%)	506/1338 (37.8%)	11.8 (10.9–12.8)	9.7 (8.9–10.6)	0.83 (0.73–0.93)	–	
All-cause death or HF hospitalization							
Overall (n = 3834)	889/1913 (46.5%)	828/1921 (43.1%)	12.4 (11.6–13.2)	11.1 (10.3–11.8)	0.90 (0.82–0.99)	0.03	
Women (n = 1143)	219/560 (39.1%)	228/583 (39.1%)	9.7 (8.5–11)	9.9 (8.7–11.3)	1.03 (0.85–1.24)	–	0.089
Men (n = 2691)	670/1353 (49.5%)	600/1338 (44.8%)	13.6 (12.6–14.7)	11.6 (10.7–12.5)	0.85 (0.76–0.95)	–	
All-cause death or all-cause hospitalization							
Overall (n = 3834)	1269/1913 (66.3%)	1237/1921 (64.4%)	22.8 (21.6–24.1)	21.6 (20.4–22.8)	0.95 (0.88–1.03)	0.22	
Women (n = 1143)	333/560 (59.5%)	363/583 (62.3%)	18.5 (16.6–20.6)	20.6 (18.6–22.8)	1.10 (0.95–1.28)	–	0.018
Men (n = 2691)	936/1353 (69.2%)	874/1338 (65.3%)	24.9 (23.4–26.6)	22 (20.6–23.5)	0.89 (0.81–0.98)	–	
CV death							
Overall (n = 3834)	478/1913 (25.0%)	448/1921 (23.3%)	5.9 (5.4–6.4)	5.4 (4.9–5.9)	0.92 (0.81–1.05)	0.22	
Women (n = 1143)	123/560 (22.0%)	118/583 (20.2%)	4.9 (4.1–5.9)	4.7 (3.9–5.6)	0.95 (0.73–1.22)	–	0.78
Men (n = 2691)	335/1353 (26.2%)	330/1338 (24.7%)	6.3 (5.7–7)	5.7 (5.1–6.4)	0.91 (0.78–1.05)	–	
All-cause death							
Overall (n = 3834)	665/1913 (34.8%)	635/1921 (33.1%)	8.2 (7.6–8.8)	7.6 (7.1–8.3)	0.94 (0.84–1.05)	0.25	
Women (n = 1143)	163/560 (29.1%)	171/583 (29.3%)	6.5 (5.6–7.6)	6.8 (5.8–7.9)	1.03 (0.83–1.28)	–	0.28
Men (n = 2691)	502/1353 (37.1%)	464/1338 (34.7%)	8.9 (8.2–9.7)	8 (7.3–8.8)	0.90 (0.79–1.02)	–	

CI, confidence interval; CV, cardiovascular; HF, heart failure; HR, hazard ratio.

^aEvent rates expressed in 100 person-years.

(all $P < 0.05$). Women were more symptomatic, had more frequently a BMI >30 kg/m², and had poorer kidney function including a higher proportion of patients with eGFR <60 mL/min/1.73 m² (all $P < 0.05$) (Table 1). In ATLAS, 21% ($n = 648$) of the patients were women and the mean age was 64 ± 10 years.⁶

Event rates and treatment effect of high- vs. low-dose losartan by sex

Men had higher event rates than women for all the studied outcomes (Table 2). Men appeared to have benefited more from high-dose (compared with low-dose) losartan, whereas women appeared to have responded similarly to low or high doses. The HR (95% CI) for the primary composite outcome of all-cause death or heart failure hospitalization was 0.85 (0.76–0.95) in men and 1.03 (0.85–1.24) in women (interaction $P = 0.089$). The HR (95% CI) for the composite outcome of cardiovascular death or heart failure hospitalization was 0.83 (0.73–0.93) in men and 1.01 (0.82–1.23) in women (interaction $P = 0.098$). The HR (95% CI) for the composite outcome of all-cause death or all-cause hospitalization was 0.89 (0.81–0.98) in men and 1.10 (0.95–1.28) in women (interaction $P = 0.018$; Table 2 and Graphical Abstract).

Safety and discontinuation of high- vs. low-dose losartan by sex

The occurrence of hypotension, hyperkalaemia and renal failure was not different between women and men (Table 3). Women were more likely than men to discontinue treatment due to adverse events with high-dose losartan: 14/560 (2.5%) women in

the low-dose group and 27/583 (4.6%) women in the high-dose group vs. 44/1353 (3.3%) men in the low-dose group and 40/1338 (3.0%) men in the high-dose group (interaction $P = 0.071$). Women were more likely than men to discontinue treatment due to any cause if taking high-dose losartan: 82/397 (20.7%) women in the low-dose group and 99/412 (24.0%) women in the high-dose group vs. 203/851 (23.9%) men in the low-dose group and 182/874 (20.8%) men in the high-dose group (interaction $P = 0.072$; Table 3).

Latent class analysis and additional subgroup analyses

Women clustered together (in class 3) with age >70 years, ischaemic heart failure, NYHA class III/IV, atrial fibrillation, and eGFR <60 mL/min/1.73 m². Hypertension and LVEF $<35\%$ were also important components but had more weight in class 2 and 1, respectively (Figure 1 and online supplementary Table S1). The comparison of the patients' characteristics across the three classes follows the same pattern described above and are presented in the online supplementary Table S2.

The event rates and the effect of high- vs. low-dose losartan across the three classes are presented in the online supplementary Table S3. Patients in class 3 had the highest event rates (worst prognosis) and experienced a similar effect of high- and low-dose losartan. Patients in class 1 had the lowest event rates (best prognosis) and experienced a similar effect of high- and low-dose losartan. Patients in class 2 were the only ones that responded to high-dose vs. low-dose losartan, with patients in class 2 receiving low-dose losartan having a similar prognosis to patients assigned to

Table 3 Safety events and percentage of drug compliance in women and men enrolled in the HEAAL trial

Safety event	Events		OR (95% CI)	Interaction P-value
	Low dose	High dose		
Hypotension				
Women (n = 1143)	39/560 (7.0%)	55/583 (9.4%)	1.39 (0.91–2.13)	0.85
Men (n = 2691)	106/1353 (7.8%)	148/1338 (11.1%)	1.46 (1.13–1.9)	
Hyperkalaemia				
Women (n = 1143)	7/560 (1.3%)	18/583 (3.1%)	2.52 (1.04–6.07)	0.39
Men (n = 2691)	15/1353 (1.1%)	23/1338 (1.7%)	1.56 (0.81–3)	
Renal failure				
Women (n = 1143)	45/560 (8.0%)	49/583 (8.4%)	1.05 (0.69–1.6)	0.29
Men (n = 2691)	101/1353 (7.5%)	134/1338 (10.0%)	1.38 (1.05–1.81)	
Drug discontinuation due to AE				
Women (n = 1143)	14/560 (2.5%)	27/583 (4.6%)	1.89 (0.98–3.65)	0.071
Men (n = 2691)	44/1353 (3.3%)	40/1338 (3.0%)	0.92 (0.59–1.42)	
Discontinued drug during FU and alive at date of last news				
Women (n = 1143)	82/397 (20.7%)	99/412 (24.0%)	1.22 (0.87–1.69)	0.072
Men (n = 2691)	203/851 (23.9%)	182/874 (20.8%)	0.84 (0.66–1.05)	

AE, adverse event; CI, confidence interval; FU, follow-up; OR, odds ratio.

AEs were reported while patients were taking treatment and for 14 days after the end of treatment. AEs were defined by the investigators. Renal failure was defined using the terms from the Medical Dictionary for Regulatory Activities (version 12): renal failure, renal failure acute, renal failure chronic, renal impairment, and pre-renal failure.

Figure 1 Latent class analysis membership in the HEAAL trial. The figure represents the probability of each individual patient to belong to class 1, 2 or 3. The ‘class membership probability’ (with 0 representing 0% chance of class membership assignment and 1 representing 100% chance of class membership assignment) can be seen in the y-axis, and each class is represented in the x-axis. Women clustered together (in class 3) with age >70 years, ischaemic heart failure (HF), New York Heart Association (NYHA) III/IV, atrial fibrillation (Afib), and estimated glomerular filtration rate (eGFR) <60 mL/min/1.73 m². Hypertension and left ventricular ejection fraction (LVEF) <35% were also important components in class 3 but had more weight in class 2 and 1, respectively. BMI, body mass index.

class 3 and patients in class 2 receiving high-dose losartan having a similar prognosis to patients in class 1. For example, the HR (95% CI) for the composite outcome of all-cause death or heart failure hospitalization was 0.98 (0.84–1.13) in class 3, 0.93 (0.78–1.1) in class 2, and 0.74 (0.62–0.88) in class 1 (interaction $P = 0.048$; Figure 2 and online supplementary Table S3).

Additional individual subgroup analyses are presented in online supplementary Table S4. In concordance with the LCA, the individual subgroup analyses suggest that patients with poorer kidney function defined by an eGFR <60 mL/min/1.73 m² and with severe heart failure symptoms defined by a NYHA class III or IV, might have experienced no benefit from high-dose losartan.

Figure 2 Treatment effect by sex (*upper panel*) and latent class analysis 'class' membership (*lower panel*) in the HEAAL trial. (*Upper panel*) Treatment-by-sex interaction: men on high-dose losartan responded better than men on low-dose losartan, whereas women responded similarly to low- or high-dose losartan. (*Lower panel*) Treatment-by-class (determined with latent class analysis) interaction: class 1 had predominance of younger men, with mild symptoms and preserved renal function (see also *Figure 1*); patients grouped in class 1 responded favourably to high-dose losartan compared with patients grouped in class 1 and receiving low-dose losartan. Class 2 had predominance of hypertensive and obese patients with heart failure of ischaemic aetiology (see also *Figure 1*); patients grouped in class 2 had the best prognosis and the effect of high- or low-dose losartan was similar; patients in class 2 (regardless of the randomly assigned losartan dose) had a favourable prognosis, similar to the prognosis of patients in class 1 receiving high-dose losartan. Class 3 had a patient predominance of elderly women, severely symptomatic, and with poor renal function (see also *Figure 1*); patients grouped in class 3 had the poorest prognosis and the effect of high- or low-dose losartan was similar. In resume, the response to high-dose losartan was driven by class 1 patients, while class 2 had good prognosis and class 3 poor prognosis regardless of losartan dose. CV, cardiovascular; HF, heart failure; Low, low-dose; High, high-dose.

Exploratory meta-analysis with individual patient data from HEAAL and published data from ATLAS

In ATLAS, the HR (95% CI) comparing high-dose vs. low-dose lisinopril for all-cause death was 0.89 (0.80–1.00) in men and 1.07 (0.82–1.36) in women. For the composite of all-cause death or all-cause hospitalization, the corresponding HR (95% CI) was 0.91 (0.82–0.99) in men and 0.82 (0.71–0.99) in women. The subgroup interaction *P*-values were not reported in the manuscript.⁶ Meta-analysing the treatment effects found in HEAAL with those derived from the forest plots in ATLAS, we found that men might have obtained more benefit from high-dose losartan and lisinopril than women, who seemed to have obtained similar benefit from

high or low dose for the outcome of all-cause death; the dose treatment-by-sex heterogeneity *P*-value was =0.11 (online supplementary *Figure S1, upper panel*). For the composite of all-cause death or all-cause hospitalizations, women responded similarly to men with high-dose lisinopril in the ATLAS trial (online supplementary *Figure S1, lower panel*).

Discussion

The present study shows that men appeared to have benefited more from high-dose (compared with low-dose) losartan, whereas women appeared to have responded similarly to low or high doses. However, the sex differences in losartan dose response are likely not due to sex alone but to concomitant conditions, particularly

poorer renal function, older age, and more severe heart failure symptoms among women, which may have limited the tolerability to high-dose losartan and caused more frequent drug discontinuation among women. These findings have clinical implications because they suggest that high-dose losartan may benefit men more than women, likely because the tolerability to high-treatment doses may be limited in women, due to comorbid conditions.

In HFrEF populations, high-dose ACEi and ARBs have been shown to provide clinical benefit compared with low-dose. This evidence comes essentially from two well-powered trials: HEAAL and ATLAS. In HEAAL, 3846 HFrEF patients were randomized to either losartan 150 mg/day (high-dose) or losartan 50 mg/day (low-dose). Compared with low-dose, high-dose losartan resulted in a 10% lower rate of the primary composite outcome of all-cause death or heart failure hospitalization. In ATLAS, 3164 HFrEF patients were randomly assigned to either lisinopril 32.5 to 35 mg/day (high-dose) or 2.5 to 5.0 mg/day (low-dose). Compared with low-dose, high-dose lisinopril resulted in a 15% lower rate of all-cause death or heart failure hospitalization (to use the same outcome as described for HEAAL) and a 12% reduction in the outcome of all-cause death or all-cause hospitalization (the outcome for which the trial had adequate power).

Multinational observational studies suggested that the optimal dose of ACEi and ARBs (and beta-blockers) may be different between men and women with HFrEF, with women having the lowest risk of adverse outcomes at lower doses (half the guideline-recommended doses) compared with men, with no further decrease in risk at higher doses.¹⁴ Causality cannot be established from observational studies. It is possible that women tolerated lower doses than men due to reasons other than sex (e.g. lower body weight, poorer renal function, older age). The hypothesis that factors other than sex may play an important role in the dose–response is sound because women with HFrEF are usually older, lighter and with poorer kidney function than men.

Patients with kidney disease have higher risk of developing hyperkalaemia and transient elevations in creatinine that may lead clinicians to withhold or permanently stop the treatment,¹⁶ which has been associated with poor outcomes.¹⁷ Patients with HFrEF and kidney impairment are also less likely to receive guideline-recommended doses of ACEi and ARBs.¹⁸ Requirement for dose adaptation, higher risk of adverse events, and decreased efficacy have also been found with other drug classes in patients with kidney disease.^{12,19,20} In the HEAAL trial, women had poorer renal function which might have contributed for higher rates of drug discontinuation and poorer response to high-dose losartan.

Heart failure symptoms, as evaluated by the NYHA functional class, also could have played an important role in dose–response in the HEAAL trial. Patients with mild symptoms (NYHA class II) might have responded better to high-dose losartan than patients with severe symptoms (NYHA class III or IV).⁵ These findings are consistent with the published forest plots in ATLAS, where patients in NYHA class II seem to have benefited more from high-dose lisinopril than patients in class II or IV.⁶ It is possible that patients with severe symptoms benefit from lower doses of treatment, that may be up titrated upon improvement of symptoms.

Other factors such as age, can also influence ACEi/ARB up-titration and response.¹⁸

The interplay between several characteristics may be better captured in our LCA from the HEAAL trial, where patients included in class 3 (comprising predominantly elderly women with atrial fibrillation, poor renal function, and severe symptoms) had the poorest prognosis and no increased benefit from high-dose losartan, as compared to lower-dose. The LCA method supports the interplay of multiple factors when it comes to dose–response. In HEAAL the response to high dose losartan (vs. low dose) seems to have been driven by hypertensive and obese patients assigned to class 2, whereas patients in class 1, who were younger, with mild symptoms and preserved renal function, had the best prognosis overall and did not experience any additional benefit from high-dose losartan. Other studies have used similar methods to identify patient ‘clusters’ with difference characteristics, prognosis, and treatment response.^{21,22}

Furthermore, in the ATLAS trial, compared with women, men might also have benefited more from high-dose lisinopril for all-cause death (with similar between-sex response for the composite of all-cause hospitalizations or all-cause death). These findings independently support a greater benefit from high-dose losartan and lisinopril in men. It is important to highlight that high-dose losartan (and lisinopril) did not have a deleterious impact among women, but the effect of high and low dose was similar in women, whereas in men high dose might have provided greater benefit. This might have been linked to factors other than sex, as above discussed.

Reports from both observational and trial data show that women are often not treated with guideline-recommended therapies.^{13,23} Our data show that, beyond sex, a multitude of factors influencing drug response and tolerability play an important role in dose–response; and, in daily clinical judgment, frequently performed under uncertainty in a case-by-case manner.

Limitations

Several limitations should be acknowledged in our study. This was a post-hoc analysis of the HEAAL trial which was underpowered to assess the treatment effect within sex subgroups, and these results should be regarded as hypothesis generating. We have performed multiple interaction tests, which might have increased the likelihood of chance findings. Unfortunately, we were unable to access the individual patient data from the ATLAS trial – as checked with the sponsor of the ATLAS study, patient agreements are lacking in some regions and the data are incomplete. Treatment non-adherence and discontinuation are often underreported in trials, and the real discontinuation rates might have been higher than here reported.²⁴ Finally, our findings apply only to the HEAAL and ATLAS trials with losartan and lisinopril dosing, respectively.

Conclusions

Compared with men, women might need lower doses of losartan to achieve similar treatment benefit. However, beyond sex, other

factors (e.g. kidney function, age, and symptoms) may influence the response to high-dose losartan, suggesting that sex-based subgroup findings may be biased by other confounders.

Supplementary Information

Additional supporting information may be found online in the Supporting Information section at the end of the article.

Conflict of interest: M.A.K. reports data monitoring board: Boehringer Ingelheim, Amgen, Luitpold, Pfizer; consultant: LivaNova, SC Pharma, Merck, Takeda, Arena, Cardurian. F.Z. reports personal fees from Applied Therapeutics, Bayer, Boehringer, Boston Scientific, Novartis, Janssen, Cellprothera and CVRx, advisory board personal fees from AstraZeneca, Vifor Fresenius, Cardior, Cereno Pharmaceutical, Corvidia, Merck, Myokardia, NovoNordisk and Owkin, stock options at G3Pharmaceutical, and being the founder of CardioRenal and the Global Cardiovascular Clinical Trialist Forum. All other authors have nothing to disclose.

References

1. Yancy CW, Jessup M, Bozkurt B, Butler J, Casey DE Jr, Colvin MM, Drazner MH, Filippatos G, Fonarow GC, Givertz MM, Hollenberg SM, Lindenfeld J, Masoudi FA, McBride PE, Peterson PN, Stevenson LW, Westlake C. 2016 ACC/AHA/HFSA focused update on new pharmacological therapy for heart failure: an update of the 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Failure Society of America. *J Am Coll Cardiol* 2016;**68**:1476–1488.
2. Ponikowski P, Voors AA, Anker SD, Bueno H, Cleland JG, Coats AJ, Falk V, Gonzalez-Juanatey JR, Harjola VP, Jankowska EA, Jessup M, Linde C, Nihoyannopoulos P, Parissis JT, Pieske B, Riley JP, Rosano GM, Ruilope LM, Ruschitzka F, Rutten FH, van der Meer P. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: the Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC). Developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur J Heart Fail* 2016;**18**:891–975.
3. Marti CN, Fonarow GC, Anker SD, Yancy C, Vaduganathan M, Greene SJ, Ahmed A, Januzzi JL, Gheorghiade M, Filippatos G, Butler J. Medication dosing for heart failure with reduced ejection fraction – opportunities and challenges. *Eur J Heart Fail* 2019;**21**:286–296.
4. Packer M, Metra M. Guideline-directed medical therapy for heart failure does not exist: a non-judgmental framework for describing the level of adherence to evidence-based drug treatments for patients with a reduced ejection fraction. *Eur J Heart Fail* 2020;**22**:1759–1767.
5. Konstam MA, Neaton JD, Dickstein K, Drexler H, Komajda M, Martinez FA, Riegger GA, Malbecq W, Smith RD, Guptha S, Poole-Wilson PA. Effects of high-dose versus low-dose losartan on clinical outcomes in patients with heart failure (HEAAL study): a randomised, double-blind trial. *Lancet* 2009;**374**:1840–1848.
6. Packer M, Poole-Wilson PA, Armstrong PW, Cleland JG, Horowitz JD, Massie BM, Ryden L, Thygesen K, Uretsky BF. Comparative effects of low and high doses of the angiotensin-converting enzyme inhibitor, lisinopril, on morbidity and mortality in chronic heart failure. ATLAS Study Group. *Circulation* 1999;**100**:2312–2318.
7. Soldin OP, Mattison DR. Sex differences in pharmacokinetics and pharmacodynamics. *Clin Pharmacokinet* 2009;**48**:143–157.
8. Hockings N, Ajayi AA, Reid JL. Age and the pharmacokinetics of angiotensin converting enzyme inhibitors enalapril and enalaprilat. *Br J Clin Pharmacol* 1986;**21**:341–348.
9. Neubeck M, Fliser D, Pritsch M, Weisser K, Fliser M, Nussberger J, Ritz E, Mutschler E. Pharmacokinetics and pharmacodynamics of lisinopril in advanced renal failure. Consequence of dose adjustment. *Eur J Clin Pharmacol* 1994;**46**:537–543.
10. Trobec K, Kerec Kos M, von Haehling S, Springer J, Anker SD, Lainscak M. Pharmacokinetics of drugs in cachectic patients: a systematic review. *PLoS One* 2013;**8**:e79603.
11. Mordi IR, Ouwerkerk W, Anker SD, Cleland JG, Dickstein K, Metra M, Ng LL, Samani NJ, van Veldhuisen DJ, Zannad F, Voors AA, Lang CC. Heart failure treatment up-titration and outcome and age: an analysis of BIOSTAT-CHF. *Eur J Heart Fail* 2021;**23**:436–444.
12. Ferreira JP, Abreu P, McMurray JJV, van Veldhuisen DJ, Swedberg K, Pocock SJ, Vincent J, Lins K, Rossignol P, Pitt B, Zannad F. Renal function stratified dose comparisons of eplerenone versus placebo in the EMPHASIS-HF trial. *Eur J Heart Fail* 2019;**21**:345–351.
13. Dewan P, Rørth R, Jhund PS, Shen L, Raparelli V, Petrie MC, Abraham WT, Desai AS, Dickstein K, Køber L, Mogensen UM, Packer M, Rouleau JL, Solomon SD, Swedberg K, Zile MR, McMurray JJV. Differential impact of heart failure with reduced ejection fraction on men and women. *J Am Coll Cardiol* 2019;**73**:29–40.
14. Santema BT, Ouwerkerk W, Tromp J, Sama IE, Ravera A, Regitz-Zagrosek V, Hillege H, Samani NJ, Zannad F, Dickstein K, Lang CC, Cleland JG, Ter Maaten JM, Metra M, Anker SD, van der Harst P, Ng LL, van der Meer P, van Veldhuisen DJ, Meyer S, Lam CSP, Voors AA. Identifying optimal doses of heart failure medications in men compared with women: a prospective, observational, cohort study. *Lancet* 2019;**394**:1254–1263.
15. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;**21**:1539–1558.
16. Agrawal H, Aggarwal K, Littrell R, Velagapudi P, Turagam MK, Mittal M, Alpert MA. Pharmacological and non pharmacological strategies in the management of coronary artery disease and chronic kidney disease. *Curr Cardiol Rev* 2015;**11**:261–269.
17. Rossignol P, Lainscak M, Crespo-Leiro MG, Laroche C, Piepoli MF, Filippatos G, Rosano GMC, Savarese G, Anker SD, Seferovic PM, Ruschitzka F, Coats AJS, Mebazaa A, McDonagh T, Sahuquillo A, Penco M, Maggioni AP, Lund LH. Unravelling the interplay between hyperkalaemia, renin-angiotensin-aldosterone inhibitor use and clinical outcomes. Data from 9222 chronic heart failure patients of the ESC-HFA-EORP Heart Failure Long-Term Registry. *Eur J Heart Fail* 2020;**22**:1378–1389.
18. Ouwerkerk W, Voors AA, Anker SD, Cleland JG, Dickstein K, Filippatos G, van der Harst P, Hillege HL, Lang CC, Ter Maaten JM, Ng LL, Ponikowski P, Samani NJ, van Veldhuisen DJ, Zannad F, Metra M, Zwinderman AH. Determinants and clinical outcome of uptitration of ACE-inhibitors and beta-blockers in patients with heart failure: a prospective European study. *Eur Heart J* 2017;**38**:1883–1890.
19. Ferreira JP, Rossello X, Pocock SJ, Zannad F, Spironolactone dose in heart failure with preserved ejection fraction: findings from TOPCAT. *Eur J Heart Fail* 2020;**22**:1615–1624.
20. Ferreira JP, Mehta C, Sharma A, Nissen SE, Rossignol P, Zannad F. Alogliptin after acute coronary syndrome in patients with type 2 diabetes: a renal function stratified analysis of the EXAMINE trial. *BMC Med* 2020;**18**:165.
21. Uijl A, Savarese G, Vaartjes I, Dahlström U, Brügts JJ, Linsens GC, van Empel V, Rocca HB, Asselbergs FW, Lund LH, Hoes AW, Koudstaal S. Identification of distinct phenotypic clusters in heart failure with preserved ejection fraction. *Eur J Heart Fail* 2021;**23**:973–982.
22. Kao DP, Lewsey JD, Anand IS, Massie BM, Zile MR, Carson PE, McKelvie RS, Komajda M, McMurray JJ, Lindenfeld J. Characterization of subgroups of heart failure patients with preserved ejection fraction with possible implications for prognosis and treatment response. *Eur J Heart Fail* 2015;**17**:925–935.
23. Greene SJ, Butler J, Albert NM, DeVore AD, Sharma PP, Duffy CI, Hill CL, McCague K, Mi X, Patterson JH, Spertus JA, Thomas L, Williams FB, Hernandez AF, Fonarow GC. Medical therapy for heart failure with reduced ejection fraction: the CHAMP-HF Registry. *J Am Coll Cardiol* 2018;**72**:351–366.
24. Breckenridge A, Aronson JK, Blaschke TF, Hartman D, Peck CC, Vrijens B. Poor medication adherence in clinical trials: consequences and solutions. *Nat Rev Drug Discov* 2017;**16**:149–150.